

VU Research Portal

Gender and cultural understandings in medical nonindicated interventions

Saharso, S.; Coene, Gily

published in

Clinical Ethics

2019

DOI (link to publisher)

[10.1177/1477750919836642](https://doi.org/10.1177/1477750919836642)

document version

Peer reviewed version

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Saharso, S., & Coene, G. (2019). Gender and cultural understandings in medical nonindicated interventions: A critical discussion of attitudes towards nontherapeutic male circumcision and hymen (re)construction. *Clinical Ethics*, 14(1), 33-41. <https://doi.org/10.1177/1477750919836642>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Gender and cultural understandings in medical nonindicated interventions: A critical discussion of attitudes toward nontherapeutic male circumcision and hymen (re)construction

Gily Coene¹  and Sawitri Saharso²

Abstract

Hymen (re)construction and nontherapeutic male circumcision are medical nonindicated interventions that give rise to specific ethical concerns. In Europe, hymen (re)construction is generally more contested among medical professionals than male circumcision. Yet, from a standard biomedical framework, guided by the principles of autonomy, beneficence, nonmaleficence, and justice, circumcision of boys is, as this article explains, more problematic than hymen (re-) construction. While there is a growing debate on the acceptability of infant circumcision, in the case of competent minors and adults the surgery is not questioned. In the case of hymenoplasty, usually requested by a competent patient, it is recommended to only perform the operation after extensive counseling and if there are compelling conditions. The article further explores why attitudes of medical professionals toward both surgeries diverge and seeks to explain how this is largely informed by gendered and socio-cultural understandings. The article further raises critical questions on medical paternalism and the role of counseling.

Keywords

Gender, culture, male circumcision, hymen (re)construction, ethical decision-making, medicine

Introduction

Hymen (re)construction—also described as hymenoplasty, hymenorrhaphy, hymen repair, or revirgination—and male circumcision—typically the removal of part or all of the penile foreskin—are medical surgeries that give rise to specific ethical concerns. Both surgeries are usually performed for cultural and religious reasons, although in rare cases circumcision can also be medically indicated.^{1,2} While hymenoplasty is usually requested by the patient herself, male circumcision is mainly performed on infants and boys upon parental request. While the latter is an irreversible intervention, hymenoplasty has no lasting bodily effects. Yet, nontherapeutic circumcision is widely performed and accepted, while hymenoplasty is a much more controversial technique and medical professionals in Europe are mostly against it.^{3–5}

This raises critical questions about why these two surgeries are regarded so differently. Scholars in

medical ethics, medical professionals, and associations largely agree that decisions in medicine should be guided by the principles of autonomy, beneficence, nonmaleficence, and justice, as devised in the classical work of Beauchamp and Childress.⁶ Yet, as we aim to demonstrate, assessing both surgeries within this framework displays some puzzling inconsistencies.

This article first outlines the main ethical issues that arise in scholarly debates on hymen (re)construction and male circumcision. As these are not only discussed in

¹Faculty of Arts & Philosophy, RHEA – Centre of Excellence on Gender, Diversity and Intersectionality, Vrije Universiteit Brussel, Brussel, Belgium

²Universiteit voor Humanistiek, Utrecht, Netherlands

Corresponding author:

Gily Coene, Faculty of Arts & Philosophy, RHEA – Centre of Excellence on Gender, Diversity and Intersectionality, Vrije Universiteit Brussel, Pleinlaan 2, Brussel 1050, Belgium.

Email: gily.coene@vub.be

medical ethics, we reviewed publications that were selected from medical (PubMed and Medline Database), sociologically, culturally, and philosophically oriented databases, including the Social Sciences Citation Index, Academic Search Elite (EBSCOhost), Sociological Abstracts (Illumina), and Philpapers.⁷ Next, we explore how standard ethical principles apply to both surgeries and conclude that the divergent attitudes toward both surgeries also rely on gendered socio-cultural understandings and the different symbolic meanings that are attributed to both surgeries.

This article does not discuss the foundations or usefulness of a principled ethical framework, nor does it engage with debates on whether such frameworks are cross-culturally applicable or theoretical issues about moral universalism and the foundations of normative ethics. It is assumed that such frameworks are largely accepted to guide clinical practice. Yet, by pointing out how socio-cultural understandings impact on ethical decisions, it aims to stimulate a more critical understanding of ethical decision-making in clinical practice.

Hymen(re)construction: Sexual autonomy and double standards

In many countries around the world, and contrary to Western popular belief not only in Muslim cultures, the norm is that women abstain from premarital sex.^{8–11} An intact hymen and loss of blood during sexual intercourse in the wedding night is traditionally considered as proof of virginity. Yet, many women do not bleed when they have intercourse for the first time¹² and the hymen is not, as is often believed, a membrane separating the vagina from the outside world, but “a thin, bloodless, elastic mucosa surrounding the opening of the vaginal introitus.”¹¹ This piece of mucosa can also be ruptured by doing sport, or by inserting a tampon, and, because of its elasticity does not have to show signs of defloration after penetration. Therefore, it is a myth that doctors can establish whether a young woman is still a virgin.^{13,14} Although there is technically not necessarily a perforated membrane to restore, physicians get requests to perform a surgery to restore the hymen. In communities in which premarital virginity is expected, the consequences when a young woman is suspected of having lost her virginity can be severe, both for herself and for her family. In Turkey, the most frequent cause of attempted suicide among young women is hymen examination¹⁵ and hymen (re)constructions are said to have reduced the number of honor killings in Egypt.¹⁶

Throughout Europe hymen (re)constructions are performed in hospitals and private clinics, usually after (more or less extensive) counselling.^{3,5,14,17,18} The most common methods of surgery described in

the literature are temporary hymen suture and hymenoplasty.^{11,18} The first method comes down to applying a few stitches in the remnants of the hymen, resulting in a small opening. As the stitches hold their tensile strength for only about a week, this method is only suited for women who intend to have the wedding within that week. The second method is more lasting and requires real wound healing of the hymen. Both methods are done under local anesthesia and patients are discharged on the day of the surgery.

In some countries, medical associations have developed guidelines on hymen surgery. According to the Dutch Society of Obstetrics and Gynecology, hymen (re)construction should not be performed, unless it appears, after counseling, that no other option offers a solution. Detailed instructions are given about what the counseling should entail. In case the request for help is inconsistent with the physician's personal views, the patient should be referred to someone else.¹⁹

Some surveys further show that the surgery is a very contested practice among medical professionals. In Sweden, almost half (46.9%) of the consulted GPs and gynecologists fully agreed with the statement that under no circumstance physicians should write a virginity certificate and similarly 42% agreed that they never should perform a hymen restoration. The others agreed that under certain conditions, they would be prepared to assist with this. The researchers conclude that the main value conflict experienced by the respondents is “on the one hand helping patients in distress (or saving lives) and on the other hand the importance of standing up against suppressive and patriarchal norms.”²⁰ Another Swedish survey that included not only gynecologists and midwives, but also youth welfare and social officers suggests that this value conflict is experienced more generally among Swedish health practitioners.³ In the Netherlands 73% of the surveyed gynecologists said they would never perform a hymen (re)construction.²¹ In Switzerland, a questionnaire was sent to 100 clinics for gynecology in public hospitals. While a minority (16.7%) refused to perform a hymen (re)construction, more than half of the clinics (64.3%) reported that they always (28.6%) or mostly (35.7%) granted the request.⁵ In a study that surveyed the views on hymen (re)construction from midwives, predominantly from Europe, the majority of the respondents (86%) agreed with the statement that virginity examinations and hymen operations are part of violence against women and only 8% believed hymen (re)construction is justifiable.²² A survey among gynecologists in Flanders (Belgium) showed that they were very divided about whether hymen (re)construction violated the right to physical integrity of the woman; roughly a third agreed, a third was neutral, and a third disagreed.¹⁸ The study also asked whether physicians had actually

received a request for hymen (re)construction. Of those who had, half of them (52.5%) had performed the procedure. The most frequently mentioned arguments to comply with the woman's request were respecting her autonomy (65.3%) and the risky situation in which she was involved (44.9%). For those who decided not to perform the surgery their main reasons were that it was not a medically indicated procedure (41.5%), because it would keep up the virginity myth (45.3%) and also the double sexual standards (49.1%). Almost half of them (49.1%) considered the procedure a violation of the patient's bodily integrity.

The referred to guidelines and surveys indicate that, generally, medical professionals condemn the virginity rule as infringing on the sexual autonomy of women and as representing double sexual standards. Therefore, and also because the procedure is not medically indicated, they believe hymen (re)construction should not be performed. Some see the surgery as also infringing on the patient's bodily integrity. If nevertheless some physicians are in favor of operating, this is because they wish to respect the woman's autonomy and take into account the reprisals they expect the woman has to face. Even the Dutch guidelines that go furthest in meeting the woman's needs, with their many provisos and the extensive counseling that is required, make clear that the Dutch professionals only grudgingly agree to perform the surgery.

Common ethical arguments contra hymen (re)construction are that there is no medical indication, that performing the operation is becoming an accomplice in deceit, that the operation contributes to keeping up the myth that all women have a hymen that bleeds with the first coitus, that it perpetuates gender inequality, that it amounts to discrimination of women and violates women's right of autonomy and their right to bodily integrity. Monika Christianson and Carola Eriksson²² for instance argue that "virginity control and hymen 'reconstructions' are elements of patriarchy, whereby violence and control are employed to subordinate women." They recognize that if the woman's life is at stake hymen (re)constructions may be justifiable, but generally they consider hymen (re)construction as an instrument in the control of women's sexuality and therefore an "inhumane and degrading treatment and (...) a form of gender discrimination that flies in the face of human rights principles for women."²² The proper way of action, therefore, is to empower women so that they are better able to resist virginity control and hymen (re)constructions. Others do not dispute that requests for hymen (re)constructions follow from patriarchal norms that discriminate against females but give priority to the well-being of the individual patient and her autonomy^{16,23-25} or have other principled arguments pro operating that

outweigh arguments contra. Regarding hymen (re)construction as deceit, Pablo de Lora²⁶ discusses a common objection against the deception argument: if hymen (re)construction is wrong, because it is deceit, then all cosmetic surgery is unethical. De Lora, however, thinks hymen (re)construction is not like any other form of cosmetic surgery. Normally, the plastic surgeon actually does change the appearance of a nose, breast, or belly. In that sense, it is no deceit, but a transformation; the deceit is in not telling that this new appearance is not given by nature. Hymen (re)construction, however, does not restore chastity. The surgery helps the woman to convince someone else that she is still a virgin. Therefore, hymen (re)construction is according to De Lora a form of deceitful cosmetic surgery, which makes the physician an accomplice in an act that is wrong. Yet, there may be other things to consider that justify performing the surgery.²⁶ Alain Vande Putte²⁷ refers to these "other things" when arguing that showing respect for the autonomy and bodily integrity of the patient "may override considerations of truthfulness towards third parties interested in controlling these aspects of the patient's life." Another overriding concern is formulated by Lee Seng Khoo and Vasco Senna-Fernandes,²⁸ who believe that physicians should refrain from moral judgment:

We have to understand the human need to belong, to be accepted and loved by their subcultural group with their own values and beliefs. We cannot attack others' religious beliefs even if they conflict with our own and cannot violate basic human rights by withholding medical or surgical treatment.

Comparing hymen (re)construction and bloodless treatment for Jehovah's Witnesses, Niklas Juth and Niels Lynøe²⁹ also discuss how the medical profession should deal with requests for medical treatments that follow from minority cultural or religious beliefs. For Jehovah's Witnesses blood transfusions are prohibited by their religion and to avoid sanctions, patients from this religion therefore ask for bloodless treatment, i.e. medical interventions that are not routinely offered to patients in healthcare. When discussing whether hymen (re)constructions are a medicalization of social problems, while bloodless treatments are not, they conclude that there is no relevant difference and, in any case, it is not obvious that social problems may not be solved through medical interventions.²⁹ Many authors also notice that performing a hymen (re)construction does not mean that the physician agrees with the virginity rule and that a consequence of performing the surgery on a large scale would be to undermine the rule.²⁷⁻²⁹ The previous argument implies that hymen (re)construction should not be banned (presuming that society rejects the virginity rule).

Yet, one could also argue that offering hymen (re)constructions expresses support for the virginity rule and therefore should be banned. Another argument, made by Juth and Lynøe, against a ban is that society is signaling repudiation of the virginity rule by helping young women to deceive its representatives.²⁹

Nontherapeutic male circumcision: Genital autonomy and gender injustice

Historically, circumcision—the surgical removal of some, or all, of the foreskin—is found among different religious, cultural, and geographical communities.^{30,31} Although the practice dates back to Ancient history, it became rapidly medicalized during the 19th and 20th century, especially in Anglo-Saxon countries, where it was initially recommended to prevent masturbation and sexually transmitted diseases such as syphilis.³² In countries like the US, where neonatal circumcision became routinely performed, antircircumcision activists challenged the surgery as medically unnecessary and potentially harmful.^{33,34} While the American Academy of Pediatrics (AAP) no longer recommends routine neonatal circumcision, it considers the surgery to be a low risk procedure that helps to prevent urinary tract infections, acquisition of HIV, and transmissions of some STIs and penile cancer—a rare, but deadly disease. Because circumcision in infancy is associated with fewer complications, the AAP recommends that the surgery should be available upon parental request.³⁵ In response to the AAP statement, physicians from different European countries, however, declared that the assessment of evidence and the policy statement of the AAP had a cultural bias: if infant circumcision would not have been such a widely accepted practice in the US, benefits and harms would have been assessed in a very different way.³⁶ These critiques do not dispute that circumcision confers certain health benefits, but question whether these can possibly justify infant circumcision or instead would, from a medical standpoint, require the boy's consent. As Frisch and Earp³⁷ explain:

(...) the question is not whether certain health benefits may in fact ensue from the sheer surgical removal of the foreskin, but whether, in light of alternative, less invasive, means of achieving the same desired health outcomes, the benefits are sufficient to outweigh the costs, harms, and other disadvantages (i.e. 'risks'), some of which may be subjective in nature and therefore difficult to quantify.

Yet, in the past decade, based on three studies that indicated that a man's risk of contracting HIV through

peno-vaginal penetrative sexual intercourse could be reduced by approximately 60%, circumcision is increasingly promoted by international health agencies as a global health strategy to prevent heterosexually transmitted HIV/AIDS. Although the WHO and UNAIDS recommend "voluntary" circumcision and to obtain informed consent, in the case of minors it is only suggested to follow "local regulations."³⁸ While the conclusiveness of the evidence on which the recommendation is based is disputed,³⁹ mass campaigns have been set up to promote and offer male circumcision for free, predominantly in the sub-Saharan region. Meanwhile, some studies point to the adverse impacts of circumcision, indicating how it enhances sexual risk behavior,⁴⁰ entails higher complication rates than previously accepted,^{41,42} and long-term adverse effects on sexual functioning, such as decreased sexual pleasure, lower orgasm intensity, discomfort, and pain.⁴³ Even when the surgery is performed in sterile settings by experienced professionals, complications, although relatively low, can still be very serious, including the loss of (a part of) the penis and even death.^{41,42}

In contrast to an earlier wave of bioethical debates that mainly focused on neonatal circumcision in countries like the US, male circumcision is now also increasingly debated in Europe, in the context of politicized debates on multiculturalism and religious accommodation.⁴⁴ In 2001, following a case where six asylum-seeking Bosnian boys were circumcised in a reception center and became badly infected, Sweden was the first to adopt a legislation that restricted the legality of circumcisions of minors to those who were performed by a licensed doctor.⁴⁵ In 2010, the Dutch Federation of Physicians declared that the circumcision of boys was in conflict with the rights of the child, and recommended that surgeries should only be allowed when performed by a medical doctor.⁴⁶ In 2012, in considering the case of a four-year-old Muslim boy whose circumcision had resulted in complications, a German district court in Cologne declared that nontherapeutic circumcision of minors was unlawful offence. The ruling initiated fierce protests, especially from Jewish and Muslim communities. Some spokesmen denounced the court's ruling as an act of anti-Semitism⁴⁷ and Germany responded to the protests by adopting a law that explicitly allows for religious circumcision on parental request, including its performance by nonmedical practitioners.⁴⁸ Yet, also in 2013, the Parliamentary Assembly of the Council of Europe adopted a resolution concerning "Children's Right to Physical Integrity" in which circumcision of young boys is considered as a violation of the physical integrity of children, comparable to female genital mutilation and early childhood medical interventions on intersex children.⁴⁹ The statement evoked fierce

criticisms from minority groups and, in a later resolution, the Assembly stresses that there is no consensus among Member States on circumcision and recommends not to allow ritual circumcision of children unless it is practiced by trained and skilled persons, in appropriate medical and health conditions.⁵⁰ In 2017, the Belgian Advisory Committee on Bioethics issued an advice to no longer refund the surgery by public social security.⁵¹ In 2018, a bill is proposed in Iceland to ban male circumcision.⁵²

In political discourses and scholarly debates, male circumcision is frequently compared with female genital cutting/mutilation (FGC/M).^{53,54} The international community considers FGM/C as a violation of human rights, gender-based violence, and a form of child abuse. According to the WHO, all surgeries that involve the partial or complete removal of external female genitalia as well as other injuries (like pricking, piercing, incising, scraping, and cauterization) are harmful to the health of women and a form of gender-based violence.⁵⁵ The WHO considers male circumcision and FGM as substantially different: while FGM is seen as a manifestation of deep-rooted gender inequality and linked to a reduction in women's sexual desire and an irreversible loss of their capability for sexual functioning, male circumcision, on the other hand, is mainly associated with health benefits and considered as a gender neutral and nondiscriminatory practice.³⁸ Nevertheless, as a number of studies point out, there is often a symbolic overlap in meanings and rationalizations of female and male genital cutting.⁵⁶ Although it is difficult to generalize about the meanings of male circumcision, it cannot be ignored that it is not a gender neutral practice, but one that often serves to establish manhood and male power and thus maintains patriarchal relations, oppressive gender norms, and hierarchies.^{57,58} With regard to harmfulness, some types of FGM/C—particularly those that are considered under type 4—are also less physically harmful than male circumcision. Yet the former are criminalized, while the latter are unregulated.⁵⁹ In identifying harmful cultural practices as practices that only negatively affect girls and women (predominantly in the South), human rights discourses have largely neglected that boys and other genders could also be harmed by cultural practices. Some authors therefore claim that the unequal human rights and legal recognition of physical and sexual integrity of girls and boys constitute in itself a form of gender injustice.^{53,60} Yet, some draw attention to similar harms and meanings of female and male genital cutting in in order to advocate a more repressive approach of male circumcision⁵³ while others use the comparison to endorse a more tolerating approach of—less harmful—forms of FGM/C.⁶¹

Concerns about the protection of the genital integrity of children have also been fed by a growing opposition to so-called normalization surgeries performed on intersex children. While such surgeries are medically nonindicated, they are medically performed to make intersex children fit into the strict gender binary.^{56,62} Whereas female, male, and intersexed genital alterations are increasingly problematized from a children's rights perspective,⁶³ circumcision of male adults or minors that are able to consent is largely perceived as unproblematic and is even recommended as a valuable alternative.⁶⁴

Discussion: Autonomy, beneficence, nonmaleficence, and justice

In the debates on hymen (re)construction that were previously revised, two, or three, positions can be distinguished. While all agree that the request for a hymen (re)construction is inspired by a patriarchal norm, some believe that hymen (re)construction should therefore not be performed, whereas others believe that it may be performed, because the woman's well-being and autonomy are paramount. This, however, should always be combined with counseling to make women aware of this patriarchal norm.^{14,24} Furthermore, some believe that at the same time public measures should be taken aimed at cultural reform of the minority groups in question.^{4,23} Yet, a common argument that is raised is that requests for hymen (re)construction are inspired by a patriarchal norm that infringes on women's autonomy. Request by women to restore their hymen is perceived as not truly autonomous and counseling is therefore recommended. The counseling practices that were reported in the literature, however, far extend the usual conditions of informed consent. In the Netherlands, the guideline of the Dutch Society of Obstetrics and Gynecology gives detailed instructions about what the counseling should entail: informing the patient about her body; if necessary with the use of a mirror, find out about the meaning of virginity for her, her partner, and their families; if possible draw the partner into the counseling, and check on risky sexual behavior, incest, and trauma that may be covered up or aggravated by the procedure. Although counseling can be perceived as enhancing the autonomy of the patient, it can also be used to influence patient's decisions imposing upon them other values and norms.⁶⁵ In debates on abortion, counseling requirements were often opposed by feminist as they were perceived to aim at changing women's minds, create feelings of guilt, and undermining women's right to decide.⁶⁶ We notice, further, that in the debates on hymen (re)construction the surgery is not compared

to arguably harmful cultural practices in the West, like cosmetic surgery, which pose comparable questions around autonomy and choice.^{67–69}

Furthermore, in debates on hymen (re)construction, harm comes to mean social and physical reprisals, like being expelled from home or being murdered, and well-being the absence thereof. Hymen (re)construction to enhance self-worth is only discussed in the case of rape victims. The idea that other motives but fear, like a desire to be part of a community or to express a cultural identity might inspire the requests, is virtually absent. This is also reflected in the common argument that there is no medical indication for hymen (re)construction. Medical need is then reduced to illness, injury, or physical impairment, hence, resting on a biological understanding of health.

In contrast, the reasons to request for a nontherapeutic male circumcision, whether by parents or the persons themselves, are generally not questioned. Ethical issues mainly arise on the conditions of proxy consent: as young children cannot decide for themselves, parents or legal guardians are supposed to act in the best interests of the child and to give their consent to medical treatment, yet their decisions may not put the physical health, well-being, or life of the child in danger. Since health benefits only become relevant later in life while the surgery entails pain, discomfort, risks for complications for the child and is irreversible, nontherapeutic infant circumcision is not justifiable for mere health preventive reasons. Yet, children's well-being does not only involve physical concerns. To be part of a community, to be able to express a religious or cultural identity, and to experience social acceptance and positive self-worth are also important and are largely accepted to justify other medically nonindicated surgeries on children. Cosmetic and corrective surgeries are seen as justifiable if they are in the child's best, immediate or future, interest. However, as is for instance argued in the case of "normalizing" surgeries of intersexed children, what is considered to be in the child's best interests can also rely on oppressive gendered norms.⁵⁶

In the case of hymen (re)construction, it is often argued that the operation should only be performed if the women fear violence and physical sanctions, such as honor killings. However, hymen "repair" can spare the young woman in question a lot of psychological and social suffering, like shame or not being able to marry. While male circumcision may not have the same effect in terms of becoming exposed to physical violence or honor killing, there is also a huge social pressure to perform the surgery. Furthermore, there is hardly any evidence on medical complications to temporary hymen suture, and while hymenoplasty is a more invasive technique, medical complications are

rare. Yet, harms and risks of male circumcision have remained unrecognized for a long time, because of the cultural acceptance of the practice and its gendered meanings, including the endurance of pain as a sign of physical strength and manhood.^{57,58}

Conclusion

Autonomy and bodily integrity are values that are protected by law, human rights instruments, and medical ethical standards. Therapeutic interventions require informed consent, where possible, by the person herself. In the case of adult persons, autonomy is seen as a standard condition, which implies that persons should be regarded as capable of making autonomous decisions, unless a person is severely mentally handicapped or suffers from a psychiatric disease that deteriorates this capacity. Where minors are concerned, there is also a growing tendency to respect their capacities for autonomous decision-making. In the case of very young children, parents or legal guardians are supposed to decide in the best interests of the child. Opinions on what is in the best interests of the child diverge, as these are based on socio-cultural values and meanings. With regard to nontherapeutic surgeries, difficult questions therefore not only arise on how to balance harms and benefits, but also on what counts as relevant to assess harm and well-being. What counts as bodily harm and well-being cannot be disconnected from personal values, social meanings, and cultural identities.

Yet, this recognition is also behind the demand for informed consent and patients' rights to decide for themselves. However, when young and adult women ask for a hymen (re)construction, this is often refused or only performed after extensive counselling. By contrast, requests for male circumcision by adults or minors who have the maturity to understand the implications of a procedure are not problematized. Thus, medical paternalism apparently only comes up with regard to women's requests, and more particularly those of migrant women.

As both concern genital surgeries, they bring gendered and cultural assumptions about the sexual body to the foreground. Male circumcision is more culturally acceptable in western countries whereas hymen repair is almost exclusively conceived as an issue of immigrant minorities, and Muslims in particular. Hymen (re)construction is problematized from a women's rights perspective and condemned as stemming from patriarchal norms. Male circumcision is seldom discussed from a gender perspective, and although increasingly problematized from a children's rights perspective, circumcision of adults and older boys is also increasingly suggested as an acceptable alternative. It thus seems that

autonomy and bodily integrity are regarded as sufficiently protected by the informed consent of the patient in the case of circumcision, but not in the case of hymen (re)construction.


Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The research was supported by a sabbatical grant for the first author of the Flemish Research Foundation, FWO.

ORCID iD

Gily Coene  <http://orcid.org/0000-0002-8158-0718>

References and notes

- Malone P and Steinbrecher H. Medical aspects of male circumcision. *BMJ* 2007; 335: 1206–1209.
- Morris BJ, Wamai RG, Hankins CA, et al. Estimation of country-specific and global prevalence of male circumcision. *Popul Health Metr* 2016; 14: 4.
- Essén B, Blomkvist A, Helström L, et al. The experience and responses of Swedish health professionals to patients requesting virginity restoration (hymen repair). *Reprod Health Matters* 2010; 18: 38–46.
- Juth N and Lynöe N. Zero tolerance against patriarchal norms? A cross-sectional study of Swedish physicians' attitudes towards young females requesting virginity certificates or hymen restoration. *J Med Ethics* 2015; 41: 215–219.
- Tschudin S, Schuster S, Dumont dos Santos D, et al. Restoration of virginity: women's demand and health care providers' response in Switzerland. *J Sex Med* 2013; 10: 2334–2342.
- Beauchamp TL and Childress JF. *Principles of Biomedical Ethics*. 5th ed. Oxford: Oxford University Press, 2001.
- We limited ourselves to the time period 2006–2016 and included articles written in Dutch, German, or English that focused on male circumcision, or hymen repair, respectively, and ethical, legal, gender related and cultural aspects. We used multiple search terms and excluded strictly anthropological descriptions of the tradition and technical medical explanations. This resulted in over 200 articles on male circumcision and almost 100 articles on hymen repair. Some of the most relevant are included in the references.
- Ogland CP, Xu X, Bartkowski JP, et al. The association of religion and virginity status among Brazilian adolescents. *J Adolesc Health* 2011; 48: 651–653.
- Roberts H. Reconstructing virginity in Guatemala. *Lancet* 2006; 367: 1227–1228.
- Thornberry E. Virginity testing, history, and the Nostalgia for custom in contemporary South Africa. *Afr Stud Rev* 2015; 58: 129–148.
- Wei SY, Li Q, Li SK, et al. A new surgical technique of hymenoplasty. *Int J Gynaecol Obstet* 2015; 130: 14–18.
- Loeber O. Over het zwaard en de schede; bloedverlies en pijn bij de eerste coïtus Een onderzoek bij vrouwen uit diverse culturen. *Tijdschr Seksuologie* 2008; 32: 129–137.
- Loeber O. Wrestling with the hymen: consultations and practical solutions. *Eur J Contracept Reprod Health Care* 2015; 20: 128–135.
- Van Moorst BR, van Lunsen RH, van Dijken DK, et al. Backgrounds of women applying for hymen reconstruction, the effects of counselling on myths and misunderstandings about virginity, and the results of hymen reconstruction. *Eur J Contracept Reprod Health Care* 2012; 17: 93–105.
- Gürsoy E and Vural G. Nurses' and midwives' views on approaches to hymen examination. *Nurs Ethics* 2003; 10: 485–496.
- Cook RJ and Dickens BM. Hymen reconstruction: ethical and legal issues. *Int J Gynaecol Obstet* 2009; 107: 266–269.
- Amy JJ. Certificates of virginity and reconstruction of the hymen. *Eur J Contracept Reprod Health Care* 2008; 13: 111–113.
- Heyerick M and Van de Wiele B. *Kennis, attitude en praktijken van Vlaamse gynaecologen ten aanzien van maagdenvliesherstellingen*. Master Thesis, Universiteit Gent, Belgium, 2012.
- NVOG. Hymen reconstructie. Voorgestelde gedragslijn. Datum Goedkeuring: 2004-05-01, Versie: 1.0, <http://nvog-documenten.nl/> (accessed 26 February 2018).
- Juth N and Lynöe N. Zero tolerance against patriarchal norms? A cross-sectional study of Swedish physicians' attitudes towards young females requesting virginity certificates or hymen restoration. *J Med Ethics* 2015; 41: 218.
- Van Lunsen RHW and Van Moorst BR. De gynaecoloog, het hymen, maagdelijkheid en verzoeken tot hymenherstel. *Nederlands Tijdschr Obstet Gynaecol* 2012; 125: 368–374.
- Christianson M and Eriksson C. Acts of violence: virginity control and hymen (re)construction. *Br J Midwifery* 2014; 22: 344–352.
- Earp BD. Hymen 'restoration' in cultures of oppression: how can physicians promote individual patient welfare without becoming complicit in the perpetuation of unjust social norms? *J Med Ethics* 2014; 40: 431.
- Saharso S. Feminisme versus multiculturalisme? De casus van maagdenvlieshersteloperaties. In: G Coene and C Longman (eds) *Eigen Emancipatie Eerst? Over de Rechten en Representatie van Vrouwen in een Multiculturele Samenleving*. Gent: Academia Press, 2005, pp.39–54.
- Wild V, Poulin H, McDougall CW, et al. Hymen reconstruction as pragmatic empowerment? Results of a qualitative study from Tunisia. *Soc Sci Med* 2015; 147: 54–61.
- De Lora P. Is multiculturalism bad for health care? The case for re-virgination. *Theor Med Bioeth* 2015; 36: 141–166.

27. Vande Putte A. *Bio-Ethische Vraagstukken Rond Maagdenvliesherstel*. Masterproef voorgedragen tot het behalen van Master in de Wijsbegeerte, Universiteit Gent, Belgium, 2015.
28. Khoo LS and Senna-Fernandes V. Hymenoplasty and virginity – an issue of socio-cultural morality and medical ethics. *PMFA News* 2015; 3. Available at: <https://www.entandaudiologynews.com/media/2745/pmfadj16-hymen-khoo-senna-fernandes.pdf>.
29. Juth N and Lynøe N. Are there morally relevant differences between hymen restoration and bloodless treatment for Jehovah's witnesses? *BMC Med Ethics* 2014; 15: 89.
30. Gollagher D. *Circumcision: A History of the World's Most Controversial Surgery*. New York: Basic Books, 2001.
31. Aggleton P. "Just a snip"? a social history of male circumcision. *Reprod Health Matters* 2007; 15: 15–21.
32. Carpenter LM. On remedicalisation: male circumcision in the United States and Great Britain. *Sociol Health Illn* 2010; 32: 613–630.
33. Sardi LM. The male neonatal circumcision debate: social movements, sexual citizenship, and human rights. *Societies without Borders* 2011; 6: 304–329.
34. Kennedy A and Sardi LM. The male anti-circumcision movement: ideology, privilege, and equity in social media. *Societies without Borders* 2016; 11: 1–30.
35. American Association of Paediatrics, Task Force on Circumcision. Circumcision policy statement. *Pediatrics* 2012; 130: 585–586.
36. Frisch M, et al. Cultural bias in the AAP's 2012 technical report and policy statement on male circumcision. *Pediatrics* 2013; 131: 796–800.
37. Frisch M and Earp B. Circumcision of male infants and children as a public health measure in developed countries: a critical assessment of recent evidence. *Glob Public Health* 2016; 10: 626–641.
38. World Health Organization and Joint United Nations Programme on HIV/AIDS. *Male Circumcision: Global Trends and Determinants of Prevalence, Safety and Acceptability*. Geneva: World Health Organization, 2007.
39. Bell K. HIV prevention: making male circumcision the 'right' tool for the job. *Glob Public Health* 2015; 10: 552–572.
40. Mukama T, et al. Perceptions about medical male circumcision and sexual behaviours of adults in Rural Uganda: a cross sectional study. *Pan Afr Med J* 2015; 22: 354.
41. Krill AJ, Lane SP and Palmer JS. Complications of circumcision. *Sci World J* 2011; 11: 2458–2468.
42. Lawal TA and Olapade-Olaopa EO. Circumcision and its effects in Africa. *Transl Androl Urol* 2017; 6: 149–157.
43. Bronselaer GA, et al. Male circumcision decreases penile sensitivity as measured in a large cohort. *BJU Int* 2013; 11: 820–827.
44. Coene G. Male circumcision: the emergence of a harmful cultural practice in the West? In: M Fusaschi and G Cavatorta (eds) *FGM/C, From Medicine to Critical Anthropology*. Torino: Meti Edizioni, 2018, pp.133–150.
45. Schiratzki J. Banning god's law in the name of the holy body – the Nordic position on ritual male circumcision. *Fam Law* 2011; 5: 35–53.
46. KNMG. Standpunt niet-therapeutische circumcisie bij minderjarige jongens, <https://www.knmg.nl/advies-richtlijnen/dossiers/jongensbesnijdenis.htm> (2010, accessed 26 April 2018).
47. Levey GB. Thinking about infant male circumcision after the cologne court decision. *Glob Discourse* 2013; 3: 326–331.
48. Aurenque D and Wiesing U. German law on circumcision and its debate: how an ethical and legal issue turned political. *Bioethics* 2015; 29: 203–210.
49. Council of Europe. Children's Right to Physical Integrity. Resolution 1952 (2013) Final Version, <http://assembly.coe.int/nw/xml/XRef/Xref-XML2HTML-en.asp?fileid=20174&lang=en> (2013, accessed 26 April 2018).
50. Council of Europe. Freedom of religion and living together in a democratic society. Resolution 2076, <http://assembly.coe.int/nw/xml/XRef/Xref-XML2HTML-en.asp?fileid=22199&lang=en> (2015, accessed 26 April 2018).
51. Belgian Advisory Committee on Bioethics. Opinion no. 70 of 8 May 2017 on the ethical aspects or nonmedical circumcision, Belgian Advisory Committee on Bioethics, <https://www.health.belgium.be/en/opinion-no-70-ethical-aspects-nonmedical-circumcision> (2017, accessed 26 April 2018).
52. Ciric J. Bill proposes banning circumcision. *Iceland Review*, 19 February, <http://icelandreview.com/news/2018/02/19/bill-proposes-banning-circumcision> (2018, accessed 26 April 2018).
53. Darby R and Svoboda J. A rose by any other name? Rethinking the similarities and differences between male and female genital cutting. *Med Anthropol Q* 2007; 21: 301–323.
54. Earp BD. Female genital mutilation and male circumcision: toward an autonomy-based ethical framework. *Medicoleg Bioeth* 2015; 5: 89–104.
55. World Health Organization. Eliminating female genital mutilation. An interagency statement. OHCHR, UNDP, UNECA, UNESCO, UNFPA, UNHCR, UNIFEM, WHO, http://apps.who.int/iris/bitstream/10665/43839/1/9789241596442_eng.pdf (2008, accessed 26 April 2018).
56. Earp BD and Steinfeld R. Gender and genital cutting: a new paradigm. In: T Giménez Barbat (ed) *Gifted Women, Fragile Men*. Brussels: Euromind Monographs, 2017. Available at <http://euromind.global/brian-d-earp-and->
57. Zoske J. Male circumcision: a gender perspective. *J Men's Stud* 1998; 6: 189–208.
58. Fox M and Thomson M. Foreskin is a feminist issue. *Aust Feminist Stud* 2009; 24: 195–210.
59. Geisheker JV. The completely unregulated practice of male circumcision: human rights' abuse enshrined in law. *New Male Stud* 2013; 2: 18–45.
60. Johnsdotter S. Girls and boys as victims: asymmetries and dynamics in European public discourses on genital modifications in children. In: M Fusaschi and G Cavatorta (eds) *FGM/C, From Medicine to Critical Anthropology*. Torino: Meti Edizioni, pp.31–47.
61. Arora KS and Jacobs AJ. Female genital alteration: a compromise solution. *J Med Ethics* 2016; 42: 148–154.

62. Ammaturo FR. Intersexuality and the right to bodily integrity: critical reflections on female genital cutting, circumcision, and intersex normalizing surgeries in Europe. *Soc Legal Stud* 2016; 25: 591–610.
63. Svoboda SJ. Promoting genital autonomy by exploring commonalities between male, female, intersex, and cosmetic female genital cutting. *Glob Discourse* 2013; 3: 237–255.
64. In the case in *Re L and B* in the UK, the judge ruled that the circumcision of two Muslim boys, as requested by the father and opposed by the mother, had to be postponed until the boys could make their own individual choice, once they had the maturity and insight to appreciate the consequences and longer term effects of the decisions which they reach. See *Re L and B (Children)*; England and Wales High Court (Family Division) decisions, HTLM version of the Judgment, <http://www.bailii.org/ew/cases/EWHC/Fam/2016/849.html> (2016, accessed 26 April 2018).
65. Ayuandini S and Duyvendak JW. Becoming (more) Dutch as medical recommendations: how understandings of national identity enter the medical practice of hymenoplasty consultations. *Nations and Nationalism* 2018; 24 (2): 390–411.
66. Woodcock S. Abortion counselling and the informed consent dilemma. *Bioethics* 2011; 5: 495–504.
67. Chambers C. Are breast implants better than female genital mutilation? Autonomy, gender equality and Nussbaum's political liberalism. *Crit Rev Int Soc Polit Phil* 2004; 7: 1–33.
68. Davis K. *Reshaping the Female Body: The Dilemma of Cosmetic Surgery*. New York: Routledge, 1995.
69. Jeffreys S. *Beauty and Misogyny: Harmful Cultural Practices in the West*. New York: Routledge, 2014.