VU Research Portal

Handgrip Strength Cannot Be Assumed a Proxy for Overall Muscle Strength

Yeung, Suey S.Y.; Reijnierse, Esmee M.; Trappenburg, Marijke C.; Hogrel, Jean Yves; McPhee, Jamie S.; Piasecki, Mathew; Sipila, Sarianna; Salpakoski, Anu; Butler-Browne, Gillian; Pääsuke, Mati; Gapeyeva, Helena; Narici, Marco V.; Meskers, Carel G.M.; Maier, Andrea B.

published in

Journal of the American Medical Directors Association
2018

DOI (link to publisher)

[10.1016/j.jamda.2018.04.019](https://doi.org/10.1016/j.jamda.2018.04.019)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Yeung, S. S. Y., Reijnierse, E. M., Trappenburg, M. C., Hogrel, J. Y., McPhee, J. S., Piasecki, M., Sipila, S., Salpakoski, A., Butler-Browne, G., Pääsuke, M., Gapeyeva, H., Narici, M. V., Meskers, C. G. M., & Maier, A. B. (2018). Handgrip Strength Cannot Be Assumed a Proxy for Overall Muscle Strength. *Journal of the American Medical Directors Association*, 19(8), 703-709. <https://doi.org/10.1016/j.jamda.2018.04.019>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

JAMDA

journal homepage: www.jamda.com

Original Study

Handgrip Strength Cannot Be Assumed a Proxy for Overall Muscle Strength

Suey S.Y. Yeung MSc^{a,b}, Esmee M. Reijnierse PhD^b, Marijke C. Trappenburg MD, PhD^{c,d}, Jean-Yves Hogrel PhD^e, Jamie S. McPhee PhD^f, Mathew Piasecki PhD^g, Sarianna Sipilä PhD^h, Anu Salpakoski PhDⁱ, Gillian Butler-Browne PhD^e, Mati Pääsuke PhD^j, Helena Gapeyeva MD, PhD^j, Marco V. Narici PhD^k, Carel G.M. Meskers MD, PhD^{a,l}, Andrea B. Maier MD, PhD^{a,b,*}

^a Department of Human Movement Sciences, AgeAmsterdam, Faculty of Behavioural and Movement Sciences, Amsterdam Movement Sciences, Vrije Universiteit, Amsterdam, The Netherlands

^b Department of Medicine and Aged Care, AgeMelbourne, The Royal Melbourne Hospital, The University of Melbourne, Melbourne, Australia

^c Department of Internal Medicine, Section of Gerontology and Geriatrics, VU University Medical Center, Amsterdam, The Netherlands

^d Department of Internal Medicine, Amstelland Hospital, Amstelveen, The Netherlands

^e Institute of Myology, Paris, France

^f School of Healthcare Science, Manchester Metropolitan University, Manchester, United Kingdom

^g MRC-ARUK Centre of Excellence for Musculoskeletal Ageing Research, Clinical Metabolic and Molecular Physiology, University of Nottingham, Royal Derby Hospital Centre, Derby, United Kingdom

^h Gerontology Research Center, Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

ⁱ Peurunka Oy, Laukaa, Finland

^j Institute of Sport Sciences and Physiotherapy, University of Tartu, Tartu, Estonia

^k Institute of Physiology, Department of Biomedical Sciences, University of Padova, Padova, Italy

^l Department of Rehabilitation Medicine, VU University Medical Center, Amsterdam, The Netherlands

A B S T R A C T

Keywords:
Muscle strength
knee extension strength
aged
geriatric assessment

Objectives: Dynapenia, low muscle strength, is predictive for negative health outcomes and is usually expressed as handgrip strength (HGS). Whether HGS can be a proxy for overall muscle strength and whether this depends on age and health status is controversial. This study assessed the agreement between HGS and knee extension strength (KES) in populations differing in age and health status.

Design: Data were retrieved from 5 cohorts.

Setting and Participants: Community, geriatric outpatient clinics, and a hospital. Five cohorts (960 individuals, 49.8% male) encompassing healthy young and older individuals, geriatric outpatients, and older individuals post hip fracture were included.

Measures: HGS and KES were measured according to the protocol of each cohort. Pearson correlation was performed to analyze the association between HGS and KES, stratified by sex. HGS and KES were standardized into sex-specific z scores. The agreement between standardized HGS and standardized KES at population level and individual level were assessed by intraclass correlation coefficients (ICC) and Bland-Altman analysis.

Results: Pearson correlation coefficients were low in healthy young (male: 0.36 to 0.45, female: 0.45) and healthy older individuals (male: 0.35 to 0.37, female: 0.44), and moderate in geriatric outpatients (male and female: 0.54) and older individuals post hip fracture (male: 0.44, female: 0.57) ($P < .05$, except for male older individuals post hip fracture [$P = .07$]). Intraclass correlation coefficient values were poor to moderate in all populations (ie, healthy young individuals [0.41, 0.45], healthy older individuals [0.37, 0.41, 0.44], geriatric outpatients [0.54], and older individuals post hip fracture [0.54]). Bland-Altman analysis showed that within the same population of age and health status, agreement between HGS and KES varied on individual level.

This study was supported by the seventh framework program MYOAGE (HEALTH-2007-2.4.5-10), the United Kingdom Medical Research Council (MR/K025252/1) as part of the Lifelong Health and Wellbeing initiative, the Dutch Technology Foundation STW, and The Ministry of Education and Culture, Kela-The Social Insurance Institution of Finland, Juho Vainio Foundation. This study has received funding from the European Union's Horizon 2020 research and innovation

programme under the Marie Skłodowska-Curie grant agreement No 675003. <http://www.birmingham.ac.uk/panini>

The authors declare no conflicts of interest.

* Address correspondence to Andrea B. Maier MD, PhD, Department of Medicine and Aged Care, University of Melbourne, Melbourne Health, The Royal Melbourne Hospital, City Campus, Level 6 North, 300 Grattan St, Parkville, Victoria 3050.

E-mail address: andrea.maier@mh.org.au (A.B. Maier).

Conclusions: At both population and individual level, HGS and KES showed a low to moderate agreement independently of age and health status. HGS alone should not be assumed a proxy for overall muscle strength.

© 2018 AMDA – The Society for Post-Acute and Long-Term Care Medicine.

Measurement of muscle strength is an important part of the comprehensive geriatric assessment¹ because of its predictive validity for decline in cognition, mobility, and functional status in community-dwelling older individuals.^{2–4} Low muscle strength, known as dynapenia, is also associated with an increased risk of postoperative complications, prolonged length of stay, and mortality in hospitalized or postsurgical patients.^{5,6} Muscle strength is part of the diagnostic criteria for sarcopenia, which is defined as low muscle mass and low muscle function (muscle strength and/or physical performance), depending on the applied definition.⁷

In clinical practice, quantification of muscle strength in older individuals is predominantly assessed by measuring handgrip strength (HGS) as the measurement is simple and the device is portable and inexpensive.⁷ In addition to HGS, muscle strength can be assessed by measurement of knee extension strength (KES). This method is, however, more technically challenging and not widely accessible.⁸ It has been shown that the decline of muscle strength with chronological age is greater for the lower limb muscles than that of the upper limb.^{9–11} Previous studies showed a high association between HGS and KES among healthy individuals aged 18–90 years^{12–14} and a low association among geriatric outpatients.¹⁵ Furthermore, previous studies used correlation coefficients quantifying the degree to which 2 variables are related on a population level, but not at individual level.

The aim of this study was to assess the agreement between HGS and KES in various populations of individuals differing in age and health status at population and individual level.

Methods

Study Design

Data were derived from 5 cohorts including 960 individuals encompassing healthy young and older individuals, geriatric outpatients, and older individuals post hip fracture.

MyoAge cohort

Healthy young and older individuals were derived from the European MyoAge cohort. The study rationale and design is reported in detail elsewhere.¹⁶ The MyoAge cohort included healthy young (aged 18–30 years) and older individuals (aged 69–81 years) recruited from 5 centers located in the United Kingdom (Manchester), France (Paris), The Netherlands (Leiden), Estonia (Tartu), and Finland (Jyväskylä). Exclusion criteria included inability to walk for a distance of 250 m, being institutionalized, morbidities (neurologic disorders, metabolic diseases, rheumatoid arthritis, recent malignancy, heart failure, coagulation diseases, chronic obstructive pulmonary disease), using immunosuppressive drugs, insulin, and anticoagulants, fracture over the previous year, immobilization for 1 week over the previous 3 months, and orthopedic surgery during the past 2 years or still causing pain or physical limitation. All study centers adopted the same standardized operation procedure to perform the measurements of muscle strength. In the present analysis, data on HGS and KES were available in 181 healthy young individuals and 320 healthy older individuals.

Manchester Metropolitan University cohort

This cohort encompasses healthy young and older male individuals aged 18–40 years or 60–90 years who were recruited as part of a study investigating the nature and extent of motor unit changes in the vastus lateralis of individuals.¹⁷ Young individuals were recruited from the university and local communities around Manchester, United Kingdom. Older individuals were recruited from the local community. Exclusion criteria were recent history of leg bone fracture, diagnosis with any form of cancer or a stroke within the past 2 years, immobilization for more than 5 days within the past 4 weeks, diagnosis of any neuromuscular disease or dementia at any time, not living independently, and body mass index <18 or >35 kg/m². In the present analysis, data on HGS and KES were available in 42 young and 108 older individuals.

Dehydroepiandrosterone in older individuals cohort

This cohort examining oral dehydroepiandrosterone in older individuals (DHEAge) included healthy female and male individuals aged 60–80 years.¹⁸ Individuals attended geriatric consultations in a geriatric outpatient clinic for various symptoms related to aging such as fatigue, memory complaints, pain, and anxiety. Data was collected before the administration of dehydroepiandrosterone. Exclusion criteria included diseases such as dementia, major depressive state, cardiovascular disorder, respiratory deficiency, Parkinson's disease, endocrine disorder, and antecedent of hormone-dependent cancer. In the present analysis, data on HGS and KES were available in 68 female individuals.

Geriatric outpatients

This inception cohort included community-dwelling older individuals referred due to mobility problems to a geriatric outpatient clinic in a middle-sized teaching hospital (Bronovo Hospital, The Hague, The Netherlands).¹⁹ The comprehensive geriatric assessment included questionnaires and measurements of physical and cognitive function and was performed by trained nurses or medical staff. In the present analysis, data on HGS and KES were available in 163 outpatients.

Promoting Mobility after Hip Fracture cohort

This cohort includes community-dwelling older individuals aged 60 years and older with a hip fracture operated at the Central Finland Central Hospital, Finland.²⁰ Individuals were asked to participate in a randomized controlled trial investigating the effects of a rehabilitation program aiming to restore mobility and functional capacity. Baseline measurements were performed after individuals were discharged home from hospital, on average 65 ± 21 days after hip fracture operation. Exclusion criteria included being institutionalized or confined to bed at the time of the fracture, Mini-Mental State Examination score of <18 points, alcoholism, severe cardiovascular, pulmonary or progressive disease, para- or tetraplegic, or severe depression. In the present analysis, data on HGS and KES were available in 78 individuals.

Characteristics of the Different Cohorts

Demographics of individuals were assessed by questionnaires in the MyoAge, Promoting Mobility after Hip Fracture (ProMo), and

Manchester Metropolitan University (MMU) cohort and by medical charts in the DHEAge cohort and geriatric outpatients. In all cohorts, body weight was measured to the nearest 0.1 kg and height to the nearest 1 mm (to the nearest centimeter for DHEAge cohort). Body composition was assessed by dual-energy X-ray absorptiometry (MyoAge, DHEAge, and MMU cohorts), or by direct segmental multi-frequency bioelectrical impedance analysis (geriatric outpatients and ProMo cohort). Fat mass percentage and lean mass percentage were calculated as total fat mass and total lean mass as percentage of total body mass, respectively. Appendicular lean mass percentage was calculated as the sum of lean mass in all 4 limbs as percentage of total body mass. Gait speed was assessed by the 6-minute (MyoAge cohort), 4-m (MMU cohort and geriatric outpatients), and 10-m walking test (ProMo cohort). Gait speed was expressed in meters per second. Gait speed was not performed in the DHEAge cohort.

Measurement of HGS

HGS was measured using an isometric hand dynamometer (MyoAge cohort and geriatric outpatients: JAMAR, Sammons Preston, Inc, Bolingbrook, IL; MMU cohort: JAMAR, Patterson Medical, Warrenville, IL; DHEAge cohort: Baseline dynamometer; ProMo cohort: Good Strength dynamometer, Metitur Ltd, Palokka, Finland). For the MyoAge cohort, MMU cohort and geriatric outpatients, individuals were instructed to maintain an upright standing position with their arms along the side while holding the dynamometer. For the DHEAge cohort, HGS was assessed according to the American Society of Hand Therapists instructions with individuals being seated and elbow flexed at 90 degrees without support.²¹ For the ProMo cohort, individuals were seated with elbow being supported and flexed at 90 degrees. Three trials were performed²² for left and right hands for all the cohorts except in the ProMo cohort in which HGS were measured from the dominant hand. There was a rest period between trials. For all cohorts, the best performance of all trials was used for analysis and expressed in kilograms.

Measurement of KES

KES was measured using knee extension dynamometer chairs [MyoAge cohort: custom-built devices in the United Kingdom, Estonia, and Finland; Forcelink B.V. (Culemborg, The Netherlands) in The Netherlands, and an isokinetic dynamometer (Biodex system 3 Pro, Biodex Medical System Inc, Shirley, NY) in France; MMU cohort: custom-built dynamometer; DHEAge cohort: an isokinetic dynamometer (Biodex Medical Systems Inc, Shirley, NY); geriatric outpatients: Forcelink B.V. (Culemborg, The Netherlands); ProMo cohort: a Good Strength dynamometer chair (Metitur Ltd, Palokka, Finland)].

For the MyoAge cohort, 3 trials of isometric maximal voluntary contraction strength measurements of knee extension were performed on the dominant leg with a rest of 90 seconds between efforts. For the MMU cohort, 3 trials were performed on the right leg with short rest intervals. In the DHEAge cohort, a 3-second maximum isometric strength measurement was performed for each leg. In geriatric outpatients, individuals were asked to push with maximal effort against a cuff positioned just above the talocrural joint. Three trials were performed for each leg. For the abovementioned cohorts, individuals were seated with knees in 90 degrees and the best performance of all trials was used for analysis and expressed in Newton meters. For the ProMo cohort, KES was measured in the fractured and nonfractured side in a sitting position with a knee angle of 120 degrees. Three maximal efforts were conducted, separated by 30 seconds rest. The best result of the nonfractured side was used for further analysis and expressed in Newton.

Ethical Approval

Each study has been approved by the local ethical committees and has been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. All individuals gave written informed consent, except for geriatric outpatients for whom the need for individuals, informed consent was waived by the ethical committee since the study was based on regular care.

Statistical Analysis

Continuous variables with a normal distribution were presented as mean (standard deviation [SD]) or if not normally distributed as median (interquartile range). Categorical variables were presented as number (*n*) and percentage (%).

Analyses were performed stratified by cohort and age, next to a pooled analysis of the 5 cohorts. At population level, Pearson correlation was performed to analyze the overall association between HGS and KES using the absolute values of maximal HGS and maximal KES, stratified by sex. Pearson correlation coefficient (*r*) from 0.3 to 0.5 was considered as low, 0.5 to 0.7 as moderate, and 0.7 to 0.9 as high.²³ For the pooled analysis, data of the ProMo cohort was excluded because KES was presented in a different unit (Newton) than the other cohorts (Newton meters).

To allow comparison between HGS and KES because of different units, HGS and KES were standardized into sex- and country-specific *z* scores for the MyoAge cohort and sex-specific *z* scores for the other cohorts. Standardization of HGS and KES in each cohort allows comparison between cohorts, even with the use of different assessment methods. For the pooled analysis, cohort-sex-specific *z* scores of HGS and cohort-sex-specific *z* scores of KES from the five cohorts were used.

Intraclass correlation analysis was carried out to examine the relative agreement between the *z* scores of HGS and *z* scores of KES. Intraclass correlation coefficient (ICC) values were calculated using a 2-way mixed model of consistency²⁴ and interpreted as excellent (0.90 or higher), good (0.75 to 0.90), moderate (0.50 to 0.75), or poor (below 0.50).²⁵ At individual level, Bland and Altman analysis were used to assess the agreement between *z* scores of HGS and *z* scores of KES and to visually display the individual dispersion patterns.²⁶ Differences in *z* scores of HGS and *z* scores of KES and the 95% limits of agreement (LOA) (mean difference ± 1.96 SD) were calculated.

Data were analyzed using SPSS v 24.0 (SPSS Inc, Chicago, IL). Visualization of results was performed using GraphPad Prism 5.01.

Results

Characteristics of Different Cohorts

Table 1 shows the characteristics of different cohorts, stratified by age. Most of the individuals were living independently (86.3%–100%), and a low percentage of individuals had excessive alcohol use (0%–14.0%) or were a current smoker (0%–15.4%). The prevalence of multimorbidity and polypharmacy was higher in geriatric outpatients and individuals post hip fracture compared with healthy individuals. HGS and KES were lower in geriatric outpatients and older individuals post hip fracture compared with healthy individuals.

Agreement of HGS and Knee Extension Strength at Population Level

A low to moderate positive correlation was found between HGS and KES, stratified by cohort and age and in the pooled analysis ($P < .05$; $P = .067$ in male older adults post hip fracture) (Table 2 and Supplementary Figure 1). ICC values between *z* scores of HGS and *z*

Table 1
Characteristics of Different Cohorts Stratified by Age

	MyoAge Cohort		MMU Cohort		DHEAge Cohort	Geriatric Outpatients	ProMo Cohort
	Young N = 181	Old N = 320	Young N = 42	Old N = 108	N = 68	N = 163	N = 78
Sociodemographics							
Age, years	23.4 (2.9)	74.4 (3.2)	26.2 (4.4)	72.8 (6.7)	65.7 (2.7)	81.7 (7.2)	79.8 (7.0)
Male, n (%)	85 (47.0)	161 (50.3)	42 (100)	108 (100)	0 (0)	64 (39.3)	18 (23.1)
Independent living,* n (%)	181 (100)	320 (100)	42 (100)	108 (100)	68 (100)	138 (86.3)	78 (100)
Lifestyle factors							
Excessive alcohol use,† n (%)	22 (12.2)	28 (8.8)	1 (2.4)	15 (14.0)	0 (0)	7 (4.3)	0 (0)
Current smoking, n (%)	23 (12.7)	14 (4.4)	0 (0)	4 (3.7)	0 (0)	21 (15.4)	7 (9.0)
Health characteristics							
Multimorbidity,‡ n (%)	0 (0)	56 (17.5)	0 (0)	13 (12.3)	0 (0)	60 (38.2)	68 (87.2)
Polypharmacy,§ n (%)	0 (0)	23 (7.2)	0 (0)	29 (27.3)	0 (0)	98 (61.6)	61 (78.2)
Body composition							
Height, m	1.73 (0.09)	1.67 (0.09)	1.79 (0.06)	1.73 (0.06)	1.61 (0.07)	1.67 (0.10)	1.61 (0.09)
BMI, kg/m ²	22.8 (3.0)	25.6 (3.3)	25.2 (4.4)	25.9 (4.1)	25.3 (3.5)	25.8 (4.6)	25.1 (3.5)
Fat mass, %	23.7 (9.1)	30.5 (8.1)	17.6 (9.1)	26.2 (9.9)	33.6 (6.7)	31.8 (10.1)	31.1 (6.5)
Lean mass, %	72.8 (9.1)	66.6 (8.3)	79.3 (8.8)	70.8 (9.7)	63.1 (6.6)	63.5 (8.8)	68.3 (8.0)
ALM, %	33.1 (4.7)	28.6 (4.1)	38.7 (4.3)	32.8 (5.5)	23.8 (2.8)	28.0 (4.6)	28.0 (2.3)
Physical performance							
Gait speed, m/s	1.85 (0.30)	1.49 (0.23)	1.28 (0.19)	1.09 (0.32)	Not available	0.75 (0.28)	0.88 (0.26)
HGS, kg (male)	52.7 (9.3)	40.3 (7.7)	53.2 (9.2)	38.7 (7.9)	Not applicable	32.9 (5.5)	28.5 (7.3)
HGS, kg (female)	33.0 (5.1)	25.9 (4.9)	Not applicable	Not applicable	26.7 (4.5)	21.5 (4.9)	17.1 (6.7)
KES, Nm (male)	249.0 (59.8)	156.6 (42.2)	249.3 (74.6)	141.1 (44.6)	Not applicable	111.1 (42.5)	285.3 (91.7)*
KES, Nm (female)	149.4 (35.9)	96.1 (25.0)	Not applicable	Not applicable	118.0 (31.5)	61.6 (21.7)	218.9 (81.9)*

ALM, appendicular lean mass; BMI, body mass index.

All values are presented as mean (SD) unless indicated otherwise.

*Defined as living at home or serviced apartment.

†Defined as >21 units/week of alcohol for males and >14 units/week of alcohol for females.

‡Defined as ≥2 diseases including MyoAge cohort: hypertension, cardiovascular events, noninsulin-dependent diabetes mellitus, mild chronic obstructive pulmonary disease, osteoarthritis, arterial surgery, and thyroid disease; Geriatric outpatients: hypertension, myocardial infarction, stroke, diabetes, diabetes mellitus, chronic obstructive pulmonary disease, cancer, Parkinson disease, and rheumatoid arthritis/osteoarthritis.

§Defined as ≥5 medications.

||Assessed by the 6-minute (MyoAge cohort), 4-m (MMU cohort and geriatric outpatients), and 10-m walking test (ProMo cohort).

*Presented as Newton.

scores of KES were poor to moderate, indicating low relative agreement (below 0.8 for all cohorts) (Table 2).

Agreement of Handgrip Strength and Knee Extension Strength at Individual Level

The 95% limits of agreement (LOA) of the differences between z score of HGS and z score of KES were larger in MyoAge cohort, MMU cohort, and DHEAge cohort compared with geriatric outpatients and ProMo cohort, indicating that the agreement between HGS and KES is

lower among healthy individuals compared with geriatric outpatients and older individuals post hip fracture (Table 2 and Figure 1). For each cohort, there were individuals with low agreement between HGS and KES (ie, z score of HGS and z score of KES outside the 95% LOA: healthy young: 0% to 6.1%, healthy old: 2.9% to 5.6%, geriatric outpatients: 6.1% and older individuals post hip fracture 3.8%). Pooled analysis showed that there were 5.1% of individuals with z score of HGS and z score of KES outside the 95% LOA (Figure 1). Since HGS and KES have been standardized into z scores, mean differences between z scores of HGS and z scores of KES were zero for all cohorts.

Table 2
Agreement of HGS and KES Stratified by Cohort and Age

	MyoAge Cohort		MMU Cohort		DHEAge Cohort	Geriatric Outpatients	ProMo Cohort	Pooled
	Young N = 181	Old N = 320	Young N = 42	Old N = 108	N = 68	N = 163	N = 78	N = 960
Pearson correlation*								
R (male)	0.36†	0.35†	0.45†	0.37†	NA	0.54†	0.44	0.67†
R (female)	0.45†	0.44†	NA	NA	0.44†	0.54†	0.57†	0.69†
ICC								
ICC	0.41	0.41	0.45	0.37	0.44	0.54	0.54	0.44
95% CI	0.27–0.52	0.32–0.50	0.17–0.66	0.19–0.52	0.22–0.61	0.42–0.64	0.36–0.68	0.39–0.49
Bland-Altman, 95% LOA								
Lower	–2.09	–2.09	–2.06	–2.21	–2.08	–1.88	–1.87	–2.04
Upper	2.09	2.09	2.06	2.21	2.08	1.88	1.87	2.04

CI, confidence interval; NA, not applicable; R, Pearson correlation coefficient.

Pearson correlation was performed to analyze the overall association between HGS and KES using the absolute values of maximal HGS and maximal KES, stratified by sex. ICC was performed for standardized HGS and standardized KES (sex- and country specific z scores for MyoAge and sex-specific z scores for other cohorts). Bland-Altman analysis was performed for standardized HGS minus standardized KES. LOA was calculated by the mean difference ± 1.96 * SD.

*For the Pearson correlation pooled analysis, data of the ProMo cohort were excluded because KES was presented in a different unit (Newton) than the other cohorts (Newton meters).

†P < .05.

Fig. 1. Bland-Altman plots of z scores of HGS and z scores KES. Results are stratified by cohort and age: (A and B) MyoAge cohort (A: healthy young, B: healthy old), (C and D) MMU cohort (C: healthy young, D: healthy old), (E) DHEAge cohort, (F) geriatric outpatients, (G) ProMo cohort, and the (H) pooled analysis. The solid line represents the mean difference in HGS and KES, while the dashed lines represent the upper and lower 95% LOA (mean difference $\pm$ 1.96 SD). Gray dots represent males and black dots represent females.

Discussion

This study showed a low to moderate agreement between HGS and KES at population level and individual level for 5 cohorts differing in age and health status.

Among healthy individuals, the present study showed a low correlation between HGS and KES from Pearson correlation analysis. Previous studies showed strong correlations among 155 individuals aged 20–90 years (males 0.70, female: 0.82)¹² and among 164 individuals aged 18–85 years (0.77–0.96).¹³ This discrepancy may be explained by

the different inclusion criteria because the aforementioned studies required individuals to be able to walk unaided, whereas the cohorts encompassing healthy individuals in our study included individuals who were able to walk more than 250 m with walking aid permitted¹⁶ or no specific criteria regarding the use of walking aid and walking distance.^{17,18} Another explanation for the discrepancy in correlations is the varied physical activity level of the study population. Studies have shown that a higher daily physical activity level was significantly associated with higher KES but not with HGS in community-dwelling older adults.^{27,28} Another study included only a limited number of individuals and found a moderate to strong correlation in 20 healthy young individuals aged 20–32 years (male [$n = 10$]: 0.63, female [$n = 10$]: 0.83) and a low correlation in 18 healthy older individuals aged 62–82 years (male [$n = 9$]: 0.35, female [$n = 9$]: 0.05).¹⁴ For geriatric outpatients, the moderate correlation between HGS and KES is in discrepancy with the low correlation (male: 0.35, female: 0.37) in a previous study, which included community-dwelling older individuals with health problems in 3 or 4 domains in functional, somatic, mental, and social domains and resulted in larger population variance.¹⁵

As a result of different rates of decline between HGS and KES across aging,^{9–11} it was hypothesized that the agreement between HGS and KES would be weaker in healthy older individuals compared with healthy young individuals. This hypothesis was supported by ICC values being lower and the range of 95% LOA being wider in healthy older individuals compared with healthy young. This is consistent with a cross-sectional study in healthy young and healthy older men with the same level of daily physical activity which revealed that lower limb muscles strength was significantly lower in older men than in young men while upper limbs muscles strength was similar between the age groups.²⁹ Differences may be further accelerated by using compensation strategies (ie, extensive use of arm muscles when rising from a chair).³⁰

It was expected that the agreement of HGS and KES would be lower as a function of health status. However, ICC values showed higher agreement and Bland-Altman analysis showed a smaller range of 95% LOA in geriatric outpatients and older individuals post hip fracture compared with healthy older individuals. Apart from higher population variance, which results in higher ICC values, HGS weakness may increasingly link to KES weakness in lower health status; physiological “floor” effects may further contribute as both HGS and KES may approach their low limits.³¹ The result might also be explained by the potentially higher variance in physical activity among healthy older individuals compared with geriatric outpatients and older individuals post-hip fracture.

Our findings suggested that measure of a single muscle group should not be regarded as a proxy for overall muscle strength. Even within the same population of age and health status, Bland-Altman analysis showed that the agreement between HGS and KES were lower in some individuals compared with the others. Therefore, it may pose a challenge in using one single muscle group strength measurement as a surrogate of overall muscle strength on an individual basis or in clinical practices.³² Some feasibility issues such as the availability of standardized protocol and the need for special equipment pose a challenge in measuring KES in clinical practice. However, instrumented KES measurement such as hand-held dynamometry³³ and isokinetic dynamometry³⁴ should be used instead of manual muscle testing because of its subjectivity and the lack of sensitivity.³⁵

Our findings showed a low agreement between HGS and KES, however, whether HGS, KES, or both are associated with clinical outcomes was not investigated. A population-based cohort study ($n = 1755$) showed that lower KES in female individuals was associated with increased mortality and hospitalization whereas lower HGS in male individuals was associated with increased risk of mortality alone.³² Another study in community-dwelling older females showed that a faster rate of decline in HGS but not KES was predicted of mortality.³⁶ These results suggest that there were sex-specific

differences in the association between HGS and KES, mortality, and hospitalization. Another point to be noted is that the reliability and accuracy of measuring HGS and especially KES is not known in our study. Therefore, it remains questionable of whether it is worthwhile to measure both HGS and KES.

A strength of this study is the inclusion of different cohorts representing different age and health status, thereby making the results generalizable to a wider population. However, HGS and KES was measured using different types of devices and protocols in the cohorts, resulting in the use of different units (Newton meters/Newton or kilograms), which made it necessary to use z scores in ICC and Bland-Altman analysis. It is recommended that in future studies the measurement of HGS and KES be conducted according to the same standardized operation procedure to ensure reproducibility and consistency across different studies.

One limitation of this study is that the reliability and accuracy of HGS and KES is unknown. It is difficult to know whether individuals truly gave a maximal voluntary effort in each trial. Different conditions of individuals including pain in joints and osteoarthritis were not registered and could have influenced the muscle strength. In addition, HGS and especially KES measurement are not gold standard to quantify muscle strength.

Conclusions

A low to moderate agreement between HGS and KES was found as a function of age and health status at population level. Within the same population of age and health status, agreement between HGS and KES also varied on individual level. The use of 1 muscle group strength measure seems unjustified as an indicator of overall limb muscle strength.

Acknowledgments

We thank Marjon Stijntjes, Jantsje Pasma, Astrid Bijlsma, Yoann Barnouin, Thomas Maden-Wilkinson, Alex Ireland, and Thomas Maden-Wilkinson for their contribution to collect the data.

Supplementary Data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.jamda.2018.04.019>.

References

- Welsh TJ, Gordon AL, Gladman JR. Comprehensive geriatric assessment—a guide for the nonspecialist. *Int J Clin Pract* 2014;68:290–293.
- Rijk JM, Roos PR, Deckx L, et al. Prognostic value of handgrip strength in people aged 60 years and older: A systematic review and meta-analysis. *Geriatr Gerontol Int* 2016;16:5–20.
- Taekema DG, Ling CH, Kurrle SE, et al. Temporal relationship between handgrip strength and cognitive performance in oldest old people. *Age Ageing* 2012;41:506–512.
- Bijlsma AY, Pasma JH, Lambers D, et al. Muscle strength rather than muscle mass is associated with standing balance in elderly outpatients. *J Am Med Dir Assoc* 2013;14:493–498.
- Bohannon RW. Hand-grip dynamometry predicts future outcomes in aging adults. *J Geriatr Phys Ther* 2008;31:3–10.
- Ling CH, Taekema D, de Craen AJ, et al. Handgrip strength and mortality in the oldest old population: The Leiden 85-plus study. *CMAJ* 2010;182:429–435.
- Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 2010;39:412–423.
- Martin HJ, Yule V, Syddall HE, et al. Is hand-held dynamometry useful for the measurement of quadriceps strength in older people? A comparison with the gold standard Biodex dynamometry. *Gerontology* 2006;52:154–159.
- Frontera WR, Hughes VA, Fielding RA, et al. Aging of skeletal muscle: A 12-year longitudinal study. *J Appl Physiol* (1985) 2000;88:1321–1326.
- Lynch NA, Metter EJ, Lindle RS, et al. Muscle quality. I. Age-associated differences between arm and leg muscle groups. *J Appl Physiol* 1999;86:188–194.

11. Candow DG, Chilibeck PD. Differences in size, strength, and power of upper and lower body muscle groups in young and older men. *J Gerontol A Biol Sci Med Sci* 2005;60:148–156.
12. Samson MM, Meeuwssen IB, Crowe A, et al. Relationships between physical performance measures, age, height and body weight in healthy adults. *Age Ageing* 2000;29:235–242.
13. Bohannon RW, Magasi SR, Bubela DJ, et al. Grip and knee extension muscle strength reflect a common construct among adults. *Muscle Nerve* 2012;46:555–558.
14. Samuel D, Wilson K, Martin HJ, et al. Age-associated changes in hand grip and quadriceps muscle strength ratios in healthy adults. *Aging Clin Exp Res* 2012;24:245–250.
15. Chan OYA, van Houwelingen AH, Gussekloo J, et al. Comparison of quadriceps strength and handgrip strength in their association with health outcomes in older adults in primary care. *Age* 2014;36:9714.
16. McPhee JS, Hogrel JY, Maier AB, et al. Physiological and functional evaluation of healthy young and older men and women: Design of the European MyoAge study. *Biogerontology* 2013;14:325–337.
17. Piasecki M, Ireland A, Stashuk D, et al. Age-related neuromuscular changes affecting human vastus lateralis. *J Physiol* 2016;594:4525–4536.
18. Baulieu EE, Thomas G, Legrain S, et al. Dehydroepiandrosterone (DHEA), DHEA sulfate, and aging: Contribution of the DHEAge Study to a sociobiomedical issue. *Proc Natl Acad Sci U S A* 2000;97:4279–4284.
19. Reijnierse EM, Trappenburg MC, Leter MJ, et al. The association between parameters of malnutrition and diagnostic measures of sarcopenia in geriatric outpatients. *Plos One* 2015;10:e0135933.
20. Salpakoski A, Tormakangas T, Edgren J, et al. Effects of a multicomponent home-based physical rehabilitation program on mobility recovery after hip fracture: A randomized controlled trial. *J Am Med Dir Assoc* 2014;15:361–368.
21. Fess EE. *Clinical Assessment recommendations*. 2 ed. Chicago: American Society of Hand Therapists; 1992.
22. Reijnierse EM, de Jong N, Trappenburg MC, et al. Assessment of maximal handgrip strength: How many attempts are needed? *J Cachexia Sarcopenia Muscle* 2017;8:466–474.
23. Mukaka MM. Statistics corner: A guide to appropriate use of correlation coefficient in medical research. *Malawi Med J* 2012;24:69–71.
24. Shrout PE, Fleiss JL. Intraclass correlations: Uses in assessing rater reliability. *Psychol Bull* 1979;86:420–428.
25. Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *J Chiropr Med* 2016;15:155–163.
26. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;1:307–310.
27. Ikenaga M, Yamada Y, Takeda N, et al. Dynapenia, gait speed and daily physical activity measured using triaxial accelerometer in older Japanese men. *J Sports Med Phys Fitness* 2014;3:147–154.
28. Aoyagi Y, Park H, Watanabe E, et al. Habitual physical activity and physical fitness in older Japanese adults: The Nakanojo Study. *Gerontology* 2009;55:523–531.
29. Nogueira FRD, Libardi CA, Vechin FC, et al. Comparison of maximal muscle strength of elbow flexors and knee extensors between younger and older men with the same level of daily activity. *Clin Interv Aging* 2013;8:401–407.
30. Macaluso A, De Vito G. Muscle strength, power and adaptations to resistance training in older people. *Eur J Appl Physiol* 2004;91:450–472.
31. Kallman DA, Plato CC, Tobin JD. The Role of Muscle Loss in the Age-Related Decline of Grip Strength - Cross-Sectional and Longitudinal Perspectives. *J Gerontol* 1990;45:M82–M88.
32. Guadalupe-Grau A, Carnicero JA, Gomez-Cabello A, et al. Association of regional muscle strength with mortality and hospitalisation in older people. *Age Ageing* 2015;44:790–795.
33. Stark T, Walker B, Phillips JK, et al. Hand-held dynamometry correlation with the gold standard isokinetic dynamometry: A systematic review. *PM R* 2011;3:472–479.
34. Hartmann A, Knols R, Murer K, et al. Reproducibility of an isokinetic strength-testing protocol of the knee and ankle in older adults. *Gerontology* 2009;55:259–268.
35. Bohannon RW. Manual muscle testing: Does it meet the standards of an adequate screening test? *Clin Rehabil* 2005;19:662–667.
36. Xue QL, Beamer BA, Chaves PH, et al. Heterogeneity in rate of decline in grip, hip, and knee strength and the risk of all-cause mortality: The Women's Health and Aging Study II. *J Am Geriatr Soc* 2010;58:2076–2084.