VU Research Portal

The assessment of diabetic polyneuropathy in daily clinical practice: reproducibility and validity of Semmes Weinstein monofilaments examination and clinical neurological examination

Valk, G.D.; de Sonnaville, J.J.J.; van Houtum, W.H.; Heine, R.J.; van Eijk, J.T.M.; Bouter, L.M.; Bertelsman, F.W.

published in

Muscle and Nerve

1997

DOI (link to publisher)

[10.1002/\(SICI\)1097-4598\(199701\)20:1<116::AID-MUS19>3.0.CO;2-2](https://doi.org/10.1002/(SICI)1097-4598(199701)20:1<116::AID-MUS19>3.0.CO;2-2)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Valk, G. D., de Sonnaville, J. J. J., van Houtum, W. H., Heine, R. J., van Eijk, J. T. M., Bouter, L. M., & Bertelsman, F. W. (1997). The assessment of diabetic polyneuropathy in daily clinical practice: reproducibility and validity of Semmes Weinstein monofilaments examination and clinical neurological examination. *Muscle and Nerve*, 20, 116-118. [https://doi.org/10.1002/\(SICI\)1097-4598\(199701\)20:1<116::AID-MUS19>3.0.CO;2-2](https://doi.org/10.1002/(SICI)1097-4598(199701)20:1<116::AID-MUS19>3.0.CO;2-2)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

THE ASSESSMENT OF DIABETIC POLYNEUROPATHY IN DAILY CLINICAL PRACTICE: REPRODUCIBILITY AND VALIDITY OF SEMMES WEINSTEIN MONOFILAMENTS EXAMINATION AND CLINICAL NEUROLOGICAL EXAMINATION

GERLOF D. VALK, MD, JEROEN J.J. DE SONNAVILLE, MD, WILLIAM H. VAN HOUTUM, MD, ROBERT J. HEINE, MD, PhD, JACQUES T.M. VAN EIJK, PhD, LEX M. BOUTER, PhD, and FREDERIK W. BERTELSMANN, MD, PhD

Polyneuropathy is present in over 80% of both type 1 and 2 diabetic patients with foot lesions.⁶ A reduction in the incidence of foot ulcerations can be achieved if polyneuropathy is identified at an early stage.^{1,5,17}

For the detection of diabetic polyneuropathy in daily clinical practice a simple, sensitive, and inexpensive screening method is required. Previous studies have indicated that both the Semmes Weinstein monofilaments examination (SWME) and the clinical neurological examination (CNE) can possibly be useful for diagnosing diabetic polyneuropathy in daily clinical practice and, moreover, may even help in identifying patients at risk for amputations.^{4,11,14,16}

The aim of the present study was to assess the reproducibility of both the SWME and the CNE in

type 1 and type 2 diabetic patients. Moreover, the sensitivity and specificity of both modes of examination were quantified, using vibration perception thresholds (VPT) as “gold standard.”

MATERIALS AND METHODS

Subjects. Sixty-eight diabetic patients participated in the study, 36 males and 32 females, with a mean age of 51.6 years (range 20–71) and a mean diabetes duration of 17.6 years (range 1–50). The neuropathy status at the time of recruitment of the patients was unknown. Individuals with polyneuropathy due to other causes were excluded from the study. The participants were asked to visit the research ward on two separate days with an interval of 2–4 weeks. Both the SWME and the CNE were performed by two different observers on one of the days, and repeated by one of these two observers on the other day. At the first visit, VPT was assessed by a third independent observer. On both occasions the observers were blinded to the results of the tests carried out by the other observers. Approval was granted by the ethical committee of the Free University Hospital.

SWME. During the SWME, the monofilaments were applied on both feet at three different sites (first toe, medial surface, and base of the third metatarsal bone). Three sizes of monofilaments were used for this study; 4.17 (1 g force), 5.07 (10 g force),

From the Department of Neurology (Drs. Valk and Bertelsmann), Institute for Research in Extramural Medicine (Drs. Valk and Bouter), Department of Internal Medicine (Drs. de Sonnaville, van Houtum, and Heine), Department of General Practice, Nursing Home Medicine and Social Medicine (Dr. van Eijk), and Department of Epidemiology and Biostatistics (Dr. Bouter), Free University Hospital, Amsterdam, The Netherlands.

Acknowledgments: We thank Ms. E. Dekker and C. Blok for help in performing the study and Dr. J.N.D. de Neeling and W. Devillee for statistical advice.

Address reprint requests to Dr. F.W. Bertelsmann, Department of Neurology, Free University Hospital, P.O. Box 7057, 1007 MB Amsterdam, The Netherlands.

Accepted for publication July 16, 1996.

CCC 0148-639X/97/010116-03
© 1997 John Wiley & Sons, Inc.

and 6.10 (75 g force).^{10,11} The SWME threshold was defined as the total number of times the application of the monofilaments was not felt, and could vary between 0 and 18.

CNE. The CNE was quantified on the basis of a previously developed scoring system, which has been described in an earlier report.¹⁶ The CNE involves clinical testing of (1) sensory modalities (pinprick, light touch, vibration, and position sense) of the feet; (2) anatomic level below which light touch sensation is impaired; (3) muscle strength of the feet; and (4) ankle reflexes. The total score of the CNE could vary between 0 and 37.

VPT. The VPT of the foot was assessed with a Vibrometer (Somedic, Stockholm, Type 4). Details of this examination, and age-related reference values, have been described previously.^{8,13} Values $\geq$ mean of age-related reference values + 2 standard deviations (SD) were considered to indicate the presence of polyneuropathy.

Statistical Analysis. The mean value and standard deviation of all measurements were calculated for the tests compared. The within-subject standard deviation (within-subject SD), the coefficient of variation (ratio of within-subject SD and mean of measurements), and the Pearson correlation coefficient with corresponding 95% confidence intervals (95% CI) were computed as measures of reproducibility.²

For selecting the best diagnostic cutoff point on the scale of measurement, the receiver operating characteristics (ROC) curve method was used.^{3,15} Sensitivity and specificity figures were compared by

calculating the area under the curves (AUC) and the significance of the difference between these areas.⁹

The Pearson correlation coefficient (95% CI) was used to measure the degree of association between the results of the SWME and the CNE.

RESULTS

Reproducibility. The SWME and the CNE were assessed in 64 patients on two separate occasions. The measures of reproducibility of the SWME and the CNE assessments are presented in Table 1.

ROC Curves, Sensitivity, and Specificity. For calculating sensitivity and specificity figures, the VPT was used as "gold standard." The best diagnostic cutoff point, at which maximum sensitivity and reasonable specificity figures were achieved, was >0 for the SWME and >4 for the CNE. When both examinations were performed by the same observer, the AUC $\pm$ standard error (SE) of the CNE (0.86 ± 0.05) was significantly larger than that of the SWME (0.77 ± 0.06) ($P = 0.01$).

Using the best cutoff point, sensitivity figures of the SWME were 87.5% and 95.8%, respectively, and specificity figures were 61.4% and 45.5%, respectively, for the two different observers. Sensitivity figures of the CNE were 91.3% and 82.6%, respectively, and specificity figures were 63.4% and 60.9%, respectively.

Association between Measurements. Correlations between the SWME and the CNE were strong, and highly significant (correlation coefficients ranging from 0.69 to 0.83).

Table 1. Test-retest reproducibility of Semmes Weinstein monofilaments examination and clinical neurological examination assessments of diabetic polyneuropathy.

	Pooled data [mean (SD)]	Measures of reproducibility		
		Within-subject SD	Coefficient of variation	Correlation coefficient (95% CI)
SWME ($\log_{10}$)				
Intraobserver	0.39 (0.35)	0.16	0.41	0.80
Test-retest*				(0.69–0.87)
Interobserver	0.40 (0.37)	0.18	0.45	0.77
Test-retest†				(0.65–0.85)
CNE				
Intraobserver	7.15 (6.29)	2.93	0.41	0.79
Test-retest*				(0.68–0.87)
Interobserver	7.15 (5.96)	2.76	0.39	0.79
Test-retest†				(0.68–0.87)

SWME, Semmes Weinstein monofilaments examination; CNE, clinical neurological examination. Outcomes of SWME: $\log_{10}$ -transformed (range from 0 to 1.3).

*One observer on two separate days.

†Two observers on the same day.

DISCUSSION

In this study the reproducibility of both the SWME and the CNE was assessed, because these modes of examination are currently used for the detection of diabetic polyneuropathy in daily clinical practice. We calculated the within-subject SD, because it quantifies the scatter of the measured values around the mean of the subjects. Because the within-subject SD does not permit a comparison of methods with different scales of measurement, we also computed the coefficient of variation and, in addition, the test-retest correlation coefficients.

The results show that both modes of examination are sufficiently reproducible for the assessment of diabetic polyneuropathy in daily clinical practice. This finding is in accordance with the small number of reproducibility studies that have been reported previously.^{7,10,12} To our knowledge, no other data on the reproducibility of the SWME in diabetic patients are available.

By calculating the AUC, the ROC method allows comparison between competitive methods of examination.⁹ In the present study, the AUC of the CNE was significantly larger than the AUC of the SWME, indicating a greater discriminating ability of the CNE in comparison with the SWME.

Our results also show that the outcomes of the SWME and the CNE are strongly correlated, which indicates construct validity of both methods of examination.

In conclusion, the results of the present study show that the SWME and the CNE are both sufficiently reproducible and, in addition, reasonably sensitive and specific for diagnosing diabetic polyneuropathy in daily clinical practice.

REFERENCES

1. Bild DE, Selby JV, Sincock P, Browner WS, Braveman P, Showstack JA: Lower-extremity amputation in people with diabetes. Epidemiology and prevention. *Diabetes Care* 1989;12:24-31.
2. Bland JM, Altman DG: Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;i:307-310.
3. Borthey AL, Malerbi DA, Franco LJ: The ROC curve in the evaluation of fasting capillary blood glucose as a screening test for diabetes and IGT. *Diabetes Care* 1994;11:1269-1272.
4. Boucher P, Teasdale N, Courtemanche R, Bard C, Fleury M: Postural stability in diabetic polyneuropathy. *Diabetes Care* 1995;18:638-645.
5. Boulton AJM: Clinical presentation and management of diabetic neuropathy and foot ulceration. *Diabet Med* 1991;8(Symposium):S52-S57.
6. Caputo GM, Cavanagh PR, Ulbrecht JS, Gibbons GW, Karchmer AW: Assessment and management of foot disease in patients with diabetes. *N Engl J Med* 1994;331:854-860.
7. Dyck PJ, Kratz KM, Lehman KA, Karnes JL, Melton LJ III, O'Brien PC, Fitchy WJ, Windebank AJ, Smith BE, Fow PA, Service FJ, Rizza RA, Zimmerman BR: The Rochester diabetic neuropathy study: design, criteria for types of neuropathy, selection bias, and reproducibility of neuropathic tests. *Neurology* 1991;41:799-807.
8. Goldberg JM, Lindblom U: Standardised method of determining vibratory perception thresholds for diagnosing and screening in neurological investigation. *J Neurol Neurosurg Psychiatry* 1979;42:793-803.
9. Hanley JA, McNeil BJ: A method of comparing the areas under receiver operating characteristic curves derived from the same cases. *Radiology* 1983;148:839-843.
10. Holewski JJ, Stess RM, Graf PM, Grunfeld K: Aesthesiometry: quantification of cutaneous pressure sensation in diabetic peripheral neuropathy. *J Rehabil Res Dev* 1988;25:1-10.
11. Kumar S, Fernando DJS, Veves A, Knowles EA, Young MJ, Boulton AJM: Semmes-Weinstein monofilaments: a simple, effective and inexpensive screening device for identifying diabetic patients at risk for foot ulceration. *Diabetes Res Clin Pract* 1991;13:63-68.
12. Maser RE, Nielsen VK, Bass EB, Manjoo Q, Dorman JS, Kelsey SF, Becker DJ, Orchard TJ: Measuring diabetic neuropathy: assessment and comparison of clinical examination and quantitative sensory testing. *Diabetes Care* 1989;12:270-275.
13. de Neeling JND, Beks PJ, Bertelsmann FW, Heine RJ, Bouter LM: Sensory thresholds in older adults: reproducibility and reference values. *Muscle Nerve* 1994;17:454-461.
14. Rith-Najarian SJ, Stolusky T, Gohdes DM: Identifying diabetic patients at high risk for lower-extremity amputation in a primary health care setting. *Diabetes Care* 1992;15:1386-1389.
15. Tsuji I, Nakamoto K, Hasegawa T, Hisashige A, Inawashiro H, Fukao A, Hisamichi S: Receiver operating characteristic analysis on fasting plasma glucose, HbA_{1c}, and fructosamine on diabetes screening. *Diabetes Care* 1991;14:1075-1077.
16. Valk GD, Nauta JJP, Strijers RLM, Bertelsmann FW: Clinical examination versus neurophysiological examination in the diagnosis of diabetic polyneuropathy. *Diabet Med* 1992;9:716-721.
17. Ward JD: The diabetic leg. *Diabetologia* 1982;22:141-147.