VU Research Portal

Relevant outcomes for nutrition interventions to treat and prevent malnutrition in older people

Correa-Pérez, Andrea; Lozano-Montoya, Isabel; Volkert, Dorothee; Visser, Marjolein; Cruz-Jentoft, Alfonso J.

published in

European Geriatric Medicine
2018

DOI (link to publisher)

[10.1007/s41999-018-0024-8](https://doi.org/10.1007/s41999-018-0024-8)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Correa-Pérez, A., Lozano-Montoya, I., Volkert, D., Visser, M., & Cruz-Jentoft, A. J. (2018). Relevant outcomes for nutrition interventions to treat and prevent malnutrition in older people: a collaborative senator-ontop and manuel delphi study. *European Geriatric Medicine*, 9(2), 243-248. <https://doi.org/10.1007/s41999-018-0024-8>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Relevant outcomes for nutrition interventions to treat and prevent malnutrition in older people: a collaborative senator-ontop and manuel delphi study

Andrea Correa-Pérez¹ · Isabel Lozano-Montoya^{1,2} · Dorothee Volkert³ · Marjolein Visser^{4,5} · Alfonso J. Cruz-Jentoft¹

Received: 20 October 2017 / Accepted: 3 January 2018 / Published online: 19 January 2018
© European Geriatric Medicine Society 2018

Abstract

Background and aims Research in malnutrition in older people is limited by the lack of consensus on relevant outcomes. Researchers of two European initiatives, the ‘Malnutrition in the Elderly (MaNuEL) Knowledge Hub’ (mostly experts in nutrition) and the Optimal Evidence-Based Non-drug Therapies in Older People (ONTOP) project (geriatricians) agreed to merge forces performing a systematic review of the effectiveness of nutritional interventions for the prevention and treatment of malnutrition in older persons. In a first step, we aimed to identify relevant outcomes for this review using a systematic approach and to explore if the rating of relevant outcomes differed depending on the researchers’ professional background.

Methods Following the ONTOP protocol, we searched all outcomes used in research of nutritional interventions for the prevention and treatment of malnutrition in older people. We carried out a web-based Delphi survey using a standardized list of 13 potentially relevant outcomes among 41 experts in geriatrics and nutrition who were asked to rate each outcome from 1 to 9 points: low importance (score 1–3), important but non-critical (score 4–6), and critical (score 7–9). Participants were informed that only those outcomes rated as critical (7–9 points) would be used in the literature review. After two rounds consultation, we compared the results from each group of experts: the ONTOP group formed by 13 geriatricians and the MaNuEL group formed by 28 experts in nutrition. Mean values were used for overall rating and the Mann–Whitney *U* test was used to see the differences on ratings between both groups.

Results Mortality, morbidity, functional status, nutritional status and quality of life were considered critical outcomes by the whole group of experts. However, by analysing the ratings by the experts’ professional background, geriatricians only rated mortality, morbidity and functional status as critical, while experts in nutrition (MaNuEL group) rated nutritional status, changes in dietary intake, compliance with the intervention, quality of life, and frailty status outcomes as critical too. Two outcomes, changes in dietary intake and compliance with the intervention, were rated with a significant different between the two professional groups ($p < 0.05$).

Conclusions Five outcomes were considered critical for research in nutritional interventions for the prevention and treatment of malnutrition in older persons: mortality, morbidity, functional status, nutritional status and quality of life by the whole panel of experts. However, more consensus is needed on the relevance of specific outcomes of nutritional interventions. Consensus processes within but also between relevant organizations are required to reach consensus and to contribute to this aim.

Keywords Malnutrition · Elderly · Geriatrics · Critical outcomes · Nutritional interventions

Introduction

Malnutrition is considered one of the “geriatric giants”, being a common problem in older people in any setting [1]. Malnutrition is associated with many negative health outcomes, including mortality [2, 3], morbidity, functional

✉ Andrea Correa-Pérez
andrea.correa@salud.madrid.org

Extended author information available on the last page of the article

impairment, disability [4], reduced quality of life, and increased health care cost (hospitalizations, institutionalization) [5–7]. The relevance of these health outcomes may vary across the lifetime. The 2006 ESPEN Guidelines in Geriatrics acknowledged that, while reducing morbidity and mortality is a priority in younger patients, in geriatric patients maintenance of function and quality of life is often the most important aim [8].

Despite increasing scientific interest in malnutrition in older persons over the last decades, many uncertainties remain [9]: definition of malnutrition and the optimal screening tools [10] and the effectiveness of nutritional interventions [11, 12], for example. In 2016, within the European Joint Program Initiative “A healthy diet for a healthy life”, the ‘Malnutrition in the Elderly (MaNuEL) Knowledge Hub’ was launched to support networking activities in this field [9]. One of the objectives of MaNuEL is to review the effectiveness of nutritional and other non-pharmacological interventions for the prevention and treatment of malnutrition in older persons. In parallel, a second European initiative, the Optimal Evidence-Based Non-drug Therapies in Older People (ONTOP) project, within the SENATOR trial (Software ENgine for the Assessment and optimization of drug and non-drug Therapy in Older persons; <https://www.senator-project.eu/>) [13, 14] is performing formal reviews of systematic reviews (SRs) on different geriatric syndromes, one of them being malnutrition [15]. Researchers of both groups decided to merge forces to perform this task. When planning the clinical questions to be used for the literature review using the PICO methodology [16], it became clear that there were different views on what were relevant outcomes of nutrition interventions for the prevention and treatment of malnutrition in older persons between researchers coming from the nutrition field and those working in clinical geriatric medicine. This fact could have a relevant impact in the selection of articles to be included in systematic reviews. Thus, in a first step, we aimed to identify relevant outcomes for this review using a systematic approach. In addition, we decided to explore if the rating of relevant outcomes differed depending on researchers’ professional background.

Methods

To identify relevant outcomes for non-pharmacological nutrition interventions, following the ONTOP protocol, a web-based Delphi survey was carried out [17, 18].

First, a search of the most frequently used outcomes reported in clinical trials, systematic reviews and clinical guidelines of non-pharmacological interventions in older people with malnutrition or at risk of malnutrition was

made in different databases (Pubmed, Embase, Cochrane library). A list of 13 outcomes was drafted (Table 1) and a survey spreadsheet was made. Experts were invited to complete the survey through an e-mail which contained a web-link to the survey. Two parallel surveys were sent to two different expert advisory panels. The first (ONTOP panel) was formed by 13 geriatricians from nine European countries (ES, FR, AT, IS, SE, BE, IE, GB, IT), who had been advising using the same methodology on relevant outcomes for other geriatric syndromes [19–22]. The second panel mostly formed by experts in geriatric nutrition (MaNuEL group) was formed by 28 researchers from seven European countries (ES, FR, AT, DE, CH, IE, NL) and New Zealand [9]. The experts who were invited to join the survey are listed in the acknowledgements. Free commercial software was used to create the online survey [23]. Stata v.12 was used to analyse and compare the answers from participants.

The same list of outcomes was sent to all participants (Table 1), asking them to rate each outcome from 1 to 9 points as follows: low importance (score 1–3), important but non-critical (score 4–6), and critical (score 7–9). An open box allowed proposing additional outcomes and giving comments. Participants were informed that only those outcomes rated as critical would be used in the literature review. A second round was launched with the same list of outcomes and the overall rating obtained for each outcome in the first round, so raters could hold or change their rating based on this feedback. The participants answered the survey anonymously.

Following the ONTOP protocol, overall rating from all participants was calculated by the mean and standard deviation, as it was previously assessed for other geriatric syndromes [19–22]. Mann–Whitney *U* test was used for comparison of the ratings between professional groups (geriatricians versus nutrition experts).

Results

From 13 experts of the ONTOP group and 28 from the MaNuEL group, 6 (46.1%) and 19 (67.9%) participated in the first round, and 5 (38.5%) and 14 (50.0%) in the second round. As the surveys were anonymous, it was not possible to know if participants from each group were the same between rounds. After first round consultation, morbidity and functional status were identified as critical (Table 1); no additional relevant outcomes were proposed, although some raters asked for clarifications of some of the terms used (e.g., nutritional status, difference between functional and frailty status, and difference between morbidity and adverse events). These clarifications were included in the outcome definition at the second round.

Table 1 Results of a first round Delphi survey to define critical outcomes of nutrition interventions for the prevention and treatment of malnutrition in older people

Potential outcomes	Geriatricians (<i>n</i> = 6) (mean ± SD)	Experts on nutrition (<i>n</i> = 19) (mean ± SD)	Total (<i>n</i> = 25) (mean ± SD)	Overall rate
Mortality ^a	7.0 ± 1.4	6.9 ± 1.8	6.9 ± 1.7	Important
Morbidity (hospital complications, infections, pressure sores...)	7.0 ± 1.1	7.7 ± 1.4	7.6 ± 1.4	Critical
Functional status	5.8 ± 2.8	8.0 ± 1.3	7.7 ± 1.5	Critical
Cognitive status	4.5 ± 2.2	5.8 ± 1.9	5.5 ± 2.0	Important
Nutritional status ^b	5.8 ± 2.5	8.3 ± 1.1	7.8 ± 1.3	Critical
Changes in dietary intake	5.3 ± 1.6	6.8 ± 2.1	6.5 ± 2.0	Important
Blood biomarkers (albumin, transferrin...)	4.3 ± 1.2	5.0 ± 1.8	4.8 ± 1.7	Important
Compliance with the intervention	4.2 ± 1.9	6.9 ± 2.1	6.3 ± 2.4	Important
Health care cost (LOS, admission to hospital/nursing home...)	5.3 ± 1.0	6.0 ± 1.8	5.8 ± 1.6	Important
Falls	5.2 ± 1.2	5.7 ± 2.0	5.6 ± 1.9	Important
Quality of life ^b	5.7 ± 1.4	7.3 ± 1.6	6.9 ± 1.6	Important
Frailty status	4.8 ± 1.7	6.7 ± 1.2	6.3 ± 1.6	Important
Adverse events	4.8 ± 2.1	6.7 ± 1.5	6.2 ± 1.8	Important

n number of raters^aOnly critical outcomes for geriatricians (ONTOP group)^bOnly critical outcomes for the experts on nutrition (MaNuEL group)**Table 2** Results of a second round Delphi survey to define critical outcomes of nutrition interventions for the prevention and treatment of malnutrition in older people

Potential outcomes	Geriatricians (<i>n</i> = 5) (mean ± SD)	Experts on nutrition (<i>n</i> = 14) (mean ± SD)	Total (<i>n</i> = 19) (mean ± SD)	Overall rate
Mortality	7.4 ± 1.5	7.3 ± 1.1	7.3 ± 1.2	Critical
Morbidity (hospital complications, infections, pressure sores...)	7.4 ± 0.9	7.8 ± 1.1	7.7 ± 1.0	Critical
Functional status (mobility, ADL, physical performance...)	7.8 ± 0.4	7.4 ± 1.7	7.5 ± 1.4	Critical
Cognitive status (including dementia and delirium)	4.6 ± 1.8	5.7 ± 2.1	5.4 ± 2.0	Important
Nutritional status ^a (weight change, BMI, skin folds, muscle mass...)	6.8 ± 2.3	8.3 ± 0.9	7.9 ± 1.5	Critical
Changes in dietary intake ^a	5.0 ± 2.2	7.5 ± 1.8	6.8 ± 2.2	Important
Blood biomarkers (albumin, transferrin...)	5.8 ± 2.3	4.9 ± 2.0	5.1 ± 2.1	Important
Compliance with the intervention ^a	4.6 ± 2.5	7.3 ± 1.1	6.6 ± 1.9	Important
Health care cost (LOS, admission to hospital/nursing home...)	6.2 ± 1.3	6.2 ± 1.6	6.2 ± 1.5	Important
Falls	6.0 ± 2.0	5.9 ± 2.0	5.9 ± 1.9	Important
Quality of Life ^a	6.2 ± 2.4	7.5 ± 0.9	7.2 ± 1.5	Critical
Frailty status ^a (changes in frailty scores)	5.0 ± 2.8	7.1 ± 1.5	6.5 ± 2.1	Important
Adverse events (diarrhoea, nausea...)	5.2 ± 2.4	6.9 ± 1.5	6.5 ± 1.9	Important

n number of raters^aOnly critical outcomes for the experts on nutrition (MaNuEL group)

Table 2 shows the final rating of each outcome after the second round for the whole group and the two professional subgroups. More outcomes were rated as critical after the second round consultation. Mortality, morbidity, functional status, nutritional status and quality of life were considered critical by 15 (79%) experts and thus were selected to be used in the systematic review.

Ratings were different depending on the group of experts. Geriatricians only rated mortality, morbidity and functional status as critical, while experts on nutrition also rated nutritional status, changes in dietary intake, compliance with the intervention, quality of life, and frailty status as critical (Table 2).

Finally, nutritional status and quality of life obtained an overall mean enough to be rated as critical.

The sample size was too small to calculate kappa coefficients, so final rating dispersion was also calculated by median, 25th and 75th percentiles (Table 3). Mann–Whitney *U* tests showed significant differences between expert groups. Dietary intake and compliance with the intervention were considered as critical outcomes only by the nutritional experts from the MaNuEL group (Table 3).

Discussion

To make informed clinical decisions, patients and health care providers need to be aware of what outcomes are expected to change with any given intervention. When research or systematic reviews of research are planned,

choosing relevant outcomes is key to make sound evidence-based recommendations.

Unfortunately, there is still little consensus on what are the relevant outcomes to be used in nutritional interventions for the prevention and treatment of malnutrition in older people. The 2006 ESPEN guidelines on geriatric nutrition suggested a list of outcomes that start in the provision of sufficient amounts of nutrients, move to maintenance or improvement of nutritional status, and finish in improving function, quality of life, morbidity and mortality [8].

Geriatricians (clinicians in care of complex older people) seem to focus on clinical outcomes. Possibly, they expect nutrition interventions to merge with other pharmacological and non-pharmacological interventions in improving the most relevant clinical outcomes (such as function, morbidity and mortality) and seem to be less interested in intermediate outcomes (such as nutritional intake and nutritional status) and thus in the mechanisms of action. On the other hand, the experts on nutrition while agreeing on those major clinical outcomes as relevant would also like to have nutrition-related intermediate outcomes confirmed (compliance with the nutritional intervention, changes in dietary intake, and changes in nutritional status) and also emphasize quality of life.

The main limitation of our study is the limited number of professionals included in this Delphi survey, and the fact that all of them are highly skilled professionals with research activities in older patients. They may not represent general geriatricians or non-geriatric focused nutritionists.

Table 3 Differences on second round outcome rating between geriatricians and experts on nutrition

Outcomes	Geriatricians (<i>n</i> = 5) P50 (P25, P75)	Experts on nutrition (<i>n</i> = 14) P50 (P25, P75)	<i>p</i> value
Mortality	8 (7, 8)	7 (6, 8)	0.70
Morbidity (hospital complications, infections, pressure sores...)	8 (7, 8)	8 (7, 9)	0.44
Functional status (mobility, ADL, physical performance...)	8 (8, 8)	8 (6, 9)	1.00
Cognitive status (including dementia and delirium)	4 (3, 6)	5 (5, 6)	0.32
Nutritional status (weight change, BMI, skin folds, muscle mass...)	7 (7, 8)	8,5 (8, 9)	0.10
Changes in dietary intake	6 (6, 6)	8 (7, 9)	0.02*
Blood biomarkers (albumin, transferrin...)	6 (6, 7)	5 (3, 6)	0.30
Compliance with the intervention	6 (3, 6)	7 (7, 8)	0.01*
Health care cost (LOS, admission to hospital/nursing home...)	6 (5, 7)	6 (6, 7)	0.74
Falls	7 (5, 7)	6 (4, 7)	0.78
Quality of life	6 (5, 8)	7,5 (7, 8)	0.27
Frailty status (changes in frailty scores)	6 (2, 7)	7 (6, 9)	0.19
Adverse events (diarrhoea, nausea...)	6 (6, 6)	7 (6, 8)	0.18

n number of raters

*Significant differences between geriatricians and experts on nutrition based on *U* Mann–Whitney test

However, they do represent a wide range of European countries and different work environments and expertise.

Perhaps the most relevant conclusion is that five outcomes were consistently considered critical by the whole group of experts: mortality, morbidity, functional status, nutritional status and quality of life. These outcomes were also assessed in recent systematic reviews [24, 25]. Clear guidelines on which research outcomes to include and how to assess these specific outcomes, if issued by competent professional organizations, may foster nutrition research, comparability of research results and the ability to merge data in meta-analysis. This may be a crucial step in building solid evidence in the effectiveness of nutritional interventions in the future. Consensus processes within but also between relevant organizations are required to reach consensus and to contribute to this aim.

Acknowledgements The authors particularly thank the members who participated in the Delphi process to identify relevant outcomes. List of experts who were invited to join the survey: Experts from the SENATOR-ONTOP project: Dr. Hubert Blain, Montpellier, France; Dr. Regina Roller-Wirnsberger, Graz, Austria; Dr. Adalsteinn Gudmundsson, Reikiavik, Iceland; Dr. Gunnar Akner, Örebro, Sweden; Dr. Mirko Petrovic, Ghent, Belgium; Dr. Denis O'Mahony, Cork, Ireland; Dr. Roy L. Soiza, Aberdeen, UK; Dr. Fabiana Trotta, Ancona, Italy; Dr. Giuseppina dell'Aquila, Ancona, Italy; Dr. Iosief Abraha, Ancona, Italy; Dr. Antonio Cherubini, Ancona, Italy; Dr. Andrea Corsonello, Ancona, Italy; Dr. Fabio Salvi, Ancona, Italy. Experts from the MaNuEL Knowledge Hub project: Dr. Dorothee Volkert, Nuremberg, Germany; Dr. Karin Schindler, Vienna, Austria; Dr. Claire Sulmont Rosse, Dijon, France; Dr. Dominique Dardevet, Clermont-Ferrand, France; Dr. Isabelle Maître, Angers, France. Dr. Yves Rolland, Toulouse, France; Dr. Heiner Boeing, Nuthetal, Germany; Dr. Marta Stelmach, Nuthetal, Germany; Dr. Christine Brombach, Wädenswil, Switzerland; Dr. Gabriele Nagel, Ulm, Germany; Dr. Marion Flechtner-Mors, Ulm, Germany; Dr. Antje Hebestreit, Bremen, Germany; Dr. Maike Wolters, Bremen, Germany; Dr. Corinna Geisler, Kiel, Germany; Dr. Eva Kiesswetter, Nürnberg, Germany; Dr. Clare Corish, Dublin, Ireland; Dr. Eibhlís O'Connor, Limerick, Ireland; Dr. Eileen O'Herlihy, Cork, Ireland; Dr. Eileen Gibney, Dublin, Ireland; Dr. Ilse Reinders, Amsterdam, The Netherlands; Dr. Lisette de Groot, Wageningen, The Netherlands; Dr. Elke Naumann, Nijmegen, the Netherlands; Dr. Eva Leistra, Amsterdam, The Netherlands; Dr. Ruth Teh, Auckland, New Zealand; Dr. Marjolein Visser, Amsterdam, The Netherlands; Dr. Christa Lohrmann, Graz, Austria; Dr. Virginie Van Wymelbeke, Dijon, France; Dr. Marian de van der Schueren, Nijmegen, The Netherlands.

Author contributions All authors have contributed actively to this work and agreed with the final version to be submitted.

Funding sources The preparation of this paper was supported by the MAINutrition in the ELderly (MaNuEL) knowledge hub. This work is supported by the Joint Programming Initiative 'Healthy Diet for a Healthy Life'. The funding agencies supporting this work are (in alphabetical order of participating Member State): Austria: Federal Ministry of Science, Research and Economy (BMWFW); France: Ecole Supérieure d'Agricultures (ESA); Germany: Federal Ministry of Food and Agriculture (BMEL) represented by Federal Office for Agriculture and Food (BLE); Ireland: Department of Agriculture, Food and the Marine (DAFM), and the Health Research Board

(HRB); Spain: Instituto de Salud Carlos III, and the SENATOR trial (FP7-HEALTH-2012-305930); The Netherlands: The Netherlands Organisation for Health Research and Development (ZonMw). The authors from Spain have only received Grants from the SENATOR trial funded by the European Union Seventh Framework Programme (FP7/2007–2013) under Grant agreement no. 305930.

Compliance with ethical standards

Conflict of interest Dr. Correa-Pérez reports Grants from EUROPEAN UNION FP7 PROGRAM (FP7/2007–2013) under Grant agreement no. 305930, during the conduct of the study; Dr. Cruz-Jentoft reports Grants from EUROPEAN UNION FP7 PROGRAM, during the conduct of the study; Dr. Volkert reports grants from Medical Nutrition Industrie (MNI), Grants from Nestlé Nutrition Institute, outside the submitted work; Dr. Lozano-Montoya has nothing to disclose; Dr. Visser has nothing to disclose.

Ethical standards The ONTOP-SENATOR project has been approved by the Ethics Committee from the coordinator site University College Cork (Cork, Ireland), and by the local Ethics Committee from the Hospital Universitario Ramón y Cajal (IRYCIS) (Madrid, Spain) where this work was conducted.

Informed consent For this type of study, informed consent is not required.

References

1. Rojer AGM, Kruijenga HM, Trappenburg MC, Reijnierse EM, Sipilä S, Narici MV et al (2016) The prevalence of malnutrition according to the new ESPEN definition in four diverse populations. *Clin Nutr* 35:758–762. <https://doi.org/10.1016/j.clnu.2015.06.005>
2. Söderström L, Rosenblad A, Thors Adolfsson E, Bergkvist L (2017) Malnutrition is associated with increased mortality in older adults regardless of the cause of death. *Br J Nutr* 117:532–540. <https://doi.org/10.1017/S0007114517000435>
3. Agarwal E, Ferguson M, Banks M, Batterham M, Bauer J, Capra S et al (2013) Malnutrition and poor food intake are associated with prolonged hospital stay, frequent readmissions, and greater in-hospital mortality: results from the Nutrition Care Day Survey 2010. *Clin Nutr* 32:737–745. <https://doi.org/10.1016/j.clnu.2012.11.021>
4. Marshall S, Bauer J, Isenring E (2014) The consequences of malnutrition following discharge from rehabilitation to the community: a systematic review of current evidence in older adults. *J Hum Nutr Diet* 27:133–141. <https://doi.org/10.1111/jhn.12167>
5. Abizanda P, Sinclair A, Barcons N, Lizán L, Rodríguez-Mañas L (2016) Costs of malnutrition in institutionalized and community-dwelling older adults: a systematic review. *J Am Med Dir Assoc* 17:17–23. <https://doi.org/10.1016/j.jamda.2015.07.005>
6. Lim SL, Ong KCB, Chan YH, Loke WC, Ferguson M, Daniels L (2012) Malnutrition and its impact on cost of hospitalization, length of stay, readmission and 3-year mortality. *Clin Nutr* 31:345–350. <https://doi.org/10.1016/j.clnu.2011.11.001>
7. Muscaritoli M, Krznarić Z, Barazzoni R, Cederholm T, Golay A, Van Gossum A et al (2016) Effectiveness and efficacy of nutritional therapy—a cochrane systematic review. *Clin Nutr*. <https://doi.org/10.1016/j.clnu.2016.06.022>
8. Volkert D, Chourdakis M, Faxen-Irving G, Frühwald T, Landi F, Suominen MH et al (2015) ESPEN guidelines on nutrition in

- dementia. *Clin Nutr* 34:1052–1073. <https://doi.org/10.1016/j.clnu.2015.09.004>
9. Visser M, Volkert D, Corish C, Geisler C, de Groot LC, Cruz-Jentoft AJ et al (2017) Tackling the increasing problem of malnutrition in older persons: the Malnutrition in the Elderly (MaNuEL) Knowledge Hub. *Nutr Bull* 42:178–186. <https://doi.org/10.1111/mbu.12268>
 10. Rojer AGM, Kruijenga HM, Trappenburg MC, Reijnders EM, Sipilä S, Narici MV et al (2016) The prevalence of malnutrition according to the new ESPEN definition in four diverse populations. *Clin Nutr* 35:758–762. <https://doi.org/10.1016/j.clnu.2015.06.005>
 11. Health Council of the Netherlands (2011) Undernutrition in the elderly. Health Council Netherlands, Hague
 12. de van der Schueren MAE, Wijnhoven HAH, Kruijenga HM, Visser M (2016) A critical appraisal of nutritional intervention studies in malnourished, community dwelling older persons. *Clin Nutr* 35:1008–1014. <https://doi.org/10.1016/j.clnu.2015.12.013>
 13. SENATOR-project.eu n.d. <https://www.senator-project.eu/> (accessed May 11, 2017)
 14. Soiza RL, Subbarayan S, Antonio C, Cruz-Jentoft AJ, Petrovic M, Gudmundsson A et al (2017) The SENATOR project: developing and trialling a novel software engine to optimize medications and nonpharmacological therapy in older people with multimorbidity and polypharmacy. *Ther Adv Drug Saf* 8:81–85. <https://doi.org/10.1177/2042098616675851>
 15. Abraha I, Cruz-Jentoft A, Soiza RL, O'Mahony D, Cherubini A (2015) Evidence of and recommendations for non-pharmacological interventions for common geriatric conditions: the SENATOR-ONTOP systematic review protocol. *BMJ Open* 5:e007488. <https://doi.org/10.1136/bmjopen-2014-007488>
 16. Guyatt GH, Oxman AD, Kunz R, Atkins D, Brozek J, Vist G et al (2011) GRADE guidelines: 2. Framing the question and deciding on important outcomes. *J Clin Epidemiol* 64:395–400. <https://doi.org/10.1016/j.jclinepi.2010.09.012>
 17. Hsu C-C (2007) The Delphi technique: making sense of consensus. *Pract Assess Res Eval* 12:1–8
 18. Diamond IR, Grant RC, Feldman BM, Pencharz PB, Ling SC, Moore AM et al (2014) Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 67:401–409. <https://doi.org/10.1016/j.jclinepi.2013.12.002>
 19. Abraha I, Trotta F, Rimland JM, Cruz-Jentoft A, Lozano-Montoya I, Soiza RL et al (2015) Efficacy of non-pharmacological interventions to prevent and treat delirium in older patients: a systematic overview. The SENATOR project ONTOP series. *PLoS One* 10:1–31. <https://doi.org/10.1371/journal.pone.0123090>
 20. Vélez-Díaz-Pallarés M, Lozano-Montoya I, Abraha I, Cherubini A, Soiza RL, O'Mahony D et al (2015) Nonpharmacologic interventions to heal pressure ulcers in older patients: an overview of systematic reviews (The SENATOR-ONTOP Series). *J Am Med Dir Assoc* 16:448–469. <https://doi.org/10.1016/j.jamda.2015.01.083>
 21. Lozano-Montoya I, Vélez-Díaz-Pallarés M, Abraha I, Cherubini A, Soiza RL, O'Mahony D et al (2016) Nonpharmacologic interventions to prevent pressure ulcers in older patients: an overview of systematic reviews (The Software ENgine for the Assessment and optimization of drug and non-drug therapy in older persons [SENATOR] definition of optimal evidence. *J Am Med Dir Assoc* 17:370.e1–370.e10. <https://doi.org/10.1016/j.jamda.2015.12.091>
 22. Lozano-Montoya I, Correa-Pérez A, Abraha I, Soiza RL, Cherubini A, O'Mahony D et al (2017) Nonpharmacological interventions to treat physical frailty and sarcopenia in older patients: a systematic overview—the SENATOR Project ONTOP Series. *Clin Interv Aging* 12:721–740. <https://doi.org/10.2147/CIA.S132496>
 23. SurveyMonkey Inc. San Mateo, California, USA n.d. <https://www.surveymonkey.com> (accessed September 1, 2016)
 24. Avenell A, Smith TO, Curtain JP, Mak JC, Myint PK (2016) Nutritional supplementation for hip fracture aftercare in older people. *Cochrane Database Syst Rev* 11:CD001880. <https://doi.org/10.1002/14651858.cd001880.pub6>
 25. Munk T, Tolstrup U, Beck AM, Holst M, Rasmussen HH, Hovhannisyan K et al (2016) Individualised dietary counselling for nutritionally at-risk older patients following discharge from acute hospital to home: a systematic review and meta-analysis. *J Hum Nutr Diet* 29:196–208. <https://doi.org/10.1111/jhn.12307>

Affiliations

Andrea Correa-Pérez¹ · Isabel Lozano-Montoya^{1,2} · Dorothee Volkert³ · Marjolein Visser^{4,5} · Alfonso J. Cruz-Jentoft¹

¹ Servicio de Geriátría, Hospital Universitario Ramón y Cajal (IRYCIS), Crta. Colmenar Viejo Km 9.1, 28034 Madrid, Spain

² Present Address: Hospital Central de la Cruz Roja San José y Santa Adela, Avenida Reina Victoria 22–26, 28003 Madrid, Spain

³ Institute for Biomedicine of Aging, Friedrich-Alexander-Universität Erlangen-Nürnberg, Kobergerstr. 60, 90408 Nuremberg, Germany

⁴ Department of Internal Medicine, VU University Medical Centre, P.O. Box 7057, 1007 MB Amsterdam, The Netherlands

⁵ Department of Health Sciences, Vrije Universiteit, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands