VU Research Portal

Routine Follow-Up Radiographs for Ankle Fractures Seldom Add Value to Clinical Decision-Making

van Gerven, Pieter; Weil, Nikki L.; Termaat, Marco F.; Rubinstein, Sidney M.; El Moumni, Mostafa; Zuidema, Wietse P.; Hoogendoorn, Jochem M.; van der Meulen, Hub G.W.M.; van Tulder, Maurits W.; Schipper, Inger B.

published in

Journal of Foot and Ankle Surgery
2018

DOI (link to publisher)

[10.1053/j.jfas.2018.03.035](https://doi.org/10.1053/j.jfas.2018.03.035)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

van Gerven, P., Weil, N. L., Termaat, M. F., Rubinstein, S. M., El Moumni, M., Zuidema, W. P., Hoogendoorn, J. M., van der Meulen, H. G. W. M., van Tulder, M. W., & Schipper, I. B. (2018). Routine Follow-Up Radiographs for Ankle Fractures Seldom Add Value to Clinical Decision-Making: A Retrospective, Observational Study. *Journal of Foot and Ankle Surgery*, 57(5), 957-960. <https://doi.org/10.1053/j.jfas.2018.03.035>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Contents lists available at ScienceDirect

The Journal of Foot & Ankle Surgery

journal homepage: www.jfas.org

Routine Follow-Up Radiographs for Ankle Fractures Seldom Add Value to Clinical Decision-Making: A Retrospective, Observational Study

Pieter van Gerven, MD ¹, Nikki L. Weil, MD ², Marco F. Termaat, MD, PhD ³, Sidney M. Rubinstein, PhD ⁴, Mostafa El Moumni, MD, PhD ⁵, Wietse P. Zuidema, MD ⁶, Jochem M. Hoogendoorn, MD, PhD ⁷, Hub G. W. M. van der Meulen, MD ⁸, Maurits W. van Tulder, PhD ⁹, Inger B. Schipper, MD, PhD, FACS ¹⁰

¹PhD Candidate, Department of Trauma Surgery, Leiden University Medical Center, Leiden, The Netherlands

²Researcher, Department of Trauma Surgery, Leiden University Medical Center, Leiden, The Netherlands

³Trauma Surgeon, Department of Trauma Surgery, Leiden University Medical Center, Leiden, The Netherlands

⁴Senior Researcher, Department of Health Sciences, Faculty of Science, Amsterdam Public Health Research Institute, Vrije Universiteit, Amsterdam, The Netherlands

⁵Trauma Surgeon, Department of Surgery, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

⁶Trauma Surgeon, Department of Surgery, VU Medical Centre, Amsterdam, The Netherlands

⁷Trauma Surgeon, Department of Surgery, Haaglanden Medical Center, The Hague, The Netherlands

⁸Trauma Surgeon, Department of Surgery, HAGA Hospital, The Hague, The Netherlands

⁹Professor, Department of Health Sciences, Faculty of Science, Amsterdam Public Health Research Institute, Vrije Universiteit, Amsterdam, The Netherlands

¹⁰Professor of Trauma Surgery and Trauma Surgeon, Department of Trauma Surgery, Leiden University Medical Center, Leiden, The Netherlands

ARTICLE INFO

Level of Clinical Evidence: 3

Keywords:

ankle trauma
clinical relevance
Lauge-Hansen
radiology
surgery
Weber

ABSTRACT

Currently, the routine use of radiographs for uncomplicated ankle fractures represents good clinical practice. However, radiographs are associated with waiting time, radiation exposure, and costs. Studies have suggested that radiographs seldom alter the treatment strategy if no clinical indication for the imaging study was present. The objective of the present study was to evaluate the effect of routine radiographs on the treatment strategy during the follow-up period of ankle fractures. All patients aged ≥18 years, who had visited 1 of the participating clinics with an eligible ankle fracture in 2012 and with complete follow-up data were included. The data were retrospectively analyzed. The sociodemographic and clinical characteristics and the number of, and indications for, the radiographs taken were collected from the medical records of the participating clinics. We assessed the changes in treatment strategy according to the radiographic findings. In 528 patients with an ankle fracture, 1174 radiographs were performed during the follow-up period. Of these radiographs, 936 (79.7%) were considered routine. Of the routine radiographs taken during the follow-up period, only 11 (1.2%) resulted in changes to the treatment strategy. Although it is common practice to take radiographs routinely during the follow-up period for ankle fractures, the results from the present study suggest that routine radiographs seldom alter the treatment strategy. This limited clinical relevance should be weighed against the health care costs and radiation exposure associated with the use of routine radiographs. For a definitive recommendation, however, the results of our study should be confirmed by a prospective trial, which we are currently conducting.

© 2018 by the American College of Foot and Ankle Surgeons. All rights reserved.

Routine radiography during outpatient fracture treatment is known to contribute to the increasing costs of health care (1). The cost-effectiveness of diagnostic imaging has become an increasingly

important factor in clinical decision-making with health care costs increasing globally (2). Despite this, routine radiographs performed during outpatient clinical visits of patients with an ankle fracture are a common worldwide practice (3,4). The arguments for routine radiography include monitoring of bone healing, identification of complications, resident education, reassurance for the physician and patient, and medicolegal motives (5). Currently, the added value of routine radiographs is under discussion. Several studies examining the value of radiographs immediately after splinting and radiographs taken at the first postoperative outpatient clinic visit have suggested that

Financial Disclosure: None reported.

Conflicts of interest: None reported.

Address correspondence to: P. van Gerven, MD, Department of Surgery, Leiden University Medical Center, P.O. Box 9600, Postzone K6-R, Leiden 2300 RC, The Netherlands.

E-mail address: p.van_gerven@lumc.nl (P. van Gerven).

radiographs without a clear clinical indication (e.g., pain, loss of mobility, or subsequent trauma to the ankle) will not lead to changes in treatment strategies (1,6–11). These radiographs did, however, contribute to additional radiation exposure and unnecessary costs. In the Netherlands, with a population of 17 million people, the costs of radiographs during the follow-up period for ankle fractures has been ~3 million Euros annually, based on an incidence of 15,000/y and 4 occasions per patient when a radiographic assessment is performed, costing €50 each (12). Considering that the incidence of ankle fractures is expected to increase worldwide in the coming decades owing to an aging population (13), the clinical value of routine radiographs for monitoring fracture healing and delivering good quality care must be established.

We undertook a retrospective cohort study to identify cases in which an outpatient clinic visit during the follow-up period of ankle fractures, which included a routine radiograph that led to a change in treatment strategy. The objective of the present study was to evaluate whether routine radiographs performed during the follow-up period for patients with an ankle fracture altered the treatment strategy. We hypothesized that routine radiographs during the follow-up period of uncomplicated ankle fractures would not alter the treatment strategy.

Patients and Methods

Study Population

We retrospectively analyzed the information from consecutive patients with complete follow-up data available from 4 level 1 trauma centers in the Netherlands, 2 university hospitals and 2 large teaching hospitals. Patients aged ≥18 years with non-Weber type A ankle fractures (unimalleolar, bimalleolar, or trimalleolar fractures with a Lauge-Hansen classification of supination adduction II, supination eversion II-IV, pronation eversion I-IV, or pronation abduction I-III) (14) that had occurred from January 1, 2012 to December 31, 2012 were eligible for inclusion. Distortions and isolated Danis-Weber classification type A fractures (15) were not included. The exclusion criteria were pathologic fractures, open fractures, multiple fractures, and severe injuries (injury severity score ≥16). The follow-up period consisted of the time the patient was receiving treatment at 1 of our affiliated hospitals. No active monitoring was pursued after this period.

Study Procedure

The present investigation was performed in compliance with the current laws and ethical standards in the Netherlands. All data were stored in accordance with Dutch privacy legislation. All participating centers used a follow-up protocol that recommends radiographs at follow-up consultations 1, 2, 6, and 12 weeks after trauma or surgical fixation. The following data were extracted from the medical records: baseline patient characteristics, including age, sex, and American Society of Anesthesiologists score; fracture type according to Lauge-Hansen (14) and Danis-Weber (15) classification schemes; treatment strategies; the date of trauma and date of discharge from monitoring; the dates and number of, and indications for, the radiographic assessments; and whether the initial treatment strategy was changed by the information obtained from the radiographs.

In the present study, the standard set of anteroposterior, lateral, and mortise views was counted as 1 radiographic assessment. The fracture type was classified according to the radiographs taken at the emergency department or, when the patient had first been treated at a different emergency department, during the first consultation visit. A radiograph was considered routine if the physician had not documented the clinical indication for performing the radiograph in the medical record.

A distinction was made between radiographs taken during the first 3 weeks after trauma (defined as the treatment period, during which a treatment strategy was drafted and surgical fixation might be performed) and radiographs taken after the first 3 weeks (defined as the follow-up period, in which the main reasons for taking radiographs were to monitor bone healing and assess for complications). In the present study, we focused solely on the radiographs taken during the follow-up period. The patients were stratified into 2 groups according to the treatment strategy (i.e., surgical or conservative treatment).

Statistical Analysis

Descriptive statistics are reported for the baseline characteristics, fracture type, and radiographic characteristics. The outcome values are reported separately for conser-

Table 1

Baseline characteristics

Characteristic	Total Cohort (n = 528)	Conservative Treatment (n = 261)	Surgical Treatment (n = 267)	p Value
Male sex	238 (45)	121 (46)	117 (44)	.56
Age (y)	49.9 ± 19.5	53.5 ± 20.5	46.5 ± 18.0	< .05*
ASA score				
1	281 (53)	135 (52)	146 (55)	.50
2	166 (32)	72 (28)	94 (35)	.06
3	71 (13)	48 (18)	23 (9)	< .05*
unknown	10 (2)	6 (2)	4 (1)	.50
Fracture type				
Lauge Hansen SA	7 (1)	7 (3)	0 (0)	< .05*
Lauge Hansen SE	360 (68)	198 (76)	162 (61)	< .05*
Lauge Hansen PE	135 (26)	40 (15)	95 (36)	< .05*
Lauge Hansen PA	15 (3)	7 (3)	8 (3)	.87
Posterior malleolar only	10 (2)	8 (3)	2 (0.7)	.51
Weber C stress fracture only	1 (0.1)	1 (0.3)	0 (0)	.311

Abbreviations: ASA, American Society of Anesthesiologists; PA, pronation abduction; PE, pronation exorotation; SA, supination adduction; SE, supination exorotation.

* Statistically significant ($p < .05$).

Data presented as n (%) or mean ± standard deviation.

vatively and surgically treated patients. Categorical data were compared using a χ^2 test. Continuous data were compared using an unpaired t test. Statistical significance was defined at the 5% level ($p \leq .05$). All analyses were performed using SPSS statistics, version 23 (IBM Corp., Armonk, NY).

Results

In the cohort of 601 consecutive patients with an ankle fracture, 73 were excluded by the exclusion criteria. The study group included 528 patients, 238 (45%) males and 290 females (55%). The mean age of all patients was 49.9 ± 19.5 years (standard deviation). Of the 528 patients, 261 (49%) were treated conservatively and 267 (51%) were treated surgically. The baseline characteristics are listed in Table 1. The median follow-up period was 14.1 (range 1.1 to 133) weeks for all patients.

The details regarding the use of radiographs and the influence of the radiographic findings on the treatment strategy are listed in Table 2. In the conservatively treated patients, 257 radiographs were performed during the treatment period (median per patient of 1; range 0 to 3), and 415 radiographs were performed during the follow-up

Table 2

Usage of routine radiographs in the follow-up protocol of ankle fractures

Variable	Patients (n = 528)	Conservative Treatment (n = 261)	Surgical Treatment (n = 267)
Radiographs taken during treatment period			
Total	621	257	364
Median	1	1	1
Range	0 to 4	0 to 3	0 to 4
Radiograph taken during follow-up period (n)			
Total	1174	415	759
Median	2	2	3
Range	0 to 11	0 to 6	0 to 11
Routine radiographs	936 (80%)	373 (90%)	563 (74.2%)
Radiographs for clinical indication	238 (20%)	42 (10%)	196 (25.8%)
Radiographs leading to a change in treatment strategy	23 (2.0%*)	8 (1.9%*)	15 (2.0%*)
Routine radiographs leading to a change in treatment strategy	11 (1.2%†)	6 (1.6%†)	5 (0.9%†)

* Radiographs leading to a change in treatment strategy per number of radiographs taken during the follow-up period.

† Routine radiographs leading to a change in treatment strategy per number of routine radiographs.

Table 3

Routine radiographs leading to a change in treatment strategy (n = 11)

Variable	n (%)
Change in treatment strategy	11 (1.2)
Prolonged cast immobilization (2 wk)	6 (0.6)
Changed to surgical treatment 3 wk after trauma	2 (0.2)
Changed to surgical treatment 6 wk after trauma	1 (0.1)
Changed to surgical treatment 5 mo after trauma	1 (0.1)
Cancellation of planned implant removal	1 (0.1)

* Of all routine radiographs.

period (median 2, range 0 to 6). Of the 415 radiographs taken during the follow-up period, 337 (90%) were scored as routine radiographs. In the surgically treated patients, 364 radiographs (median 1, range 0 to 4) were performed during the treatment period, and 759 radiographs (median 3, range 0 to 11) were performed during the follow-up period. Of the 759 radiographs taken during the follow-up period, 563 (74%) were scored as routine radiographs. In the conservatively and surgically treated patients, 6 of 337 and 5 of 563 routinely scored radiographs, respectively, resulted in a change in the treatment strategy (Table 3). Cast immobilization was prolonged by 2 weeks for 6 patients, an initially conservative treatment plan was changed to surgical treatment for 4 patients, and a planned implant removal was canceled for 1 patient because no radiologic consolidation was visible. Of the 4 patients who were scheduled for surgery because of findings from routine radiographs, 2 were assigned surgical treatment during their second outpatient clinic visit, which was 21 days after the initial trauma. The third patient complained of pain during the first 3 months after the trauma and was referred for physiotherapy. During the next outpatient clinic visit 5 months after the trauma, no complaints were documented; however, the patient was assigned surgical treatment because no signs of consolidation were seen on the radiographs. The fracture of the fourth patient scheduled for surgery was 2 weeks old before presentation at the emergency department and was initially deemed suitable for a conservative treatment strategy. The patient was assigned to surgery during the first outpatient visit 4 weeks later owing to secondary loss of reduction.

In the present cohort, 1174 (65.4%) of the total of 1795 radiographs were taken during the follow-up period. Of these 1174 radiographs, 936 (79.7%) were considered routine. For the general Dutch population, this could mean that 65.4% (€1,962,000) of the total annual radiographic costs of €3 million is spent within the follow-up period. Of these costs, 79.7% (€1,563,714) can be attributed to routine radiography. This indicates that using the data found in the present cohort, 52% of all the costs involved in radiography of ankle fractures could potentially be saved by omitting routine radiographs during the follow-up period.

Discussion

We assessed the effect of conducting routine radiographs during the follow-up period on clinical decision-making in a large cohort of patients with ankle fractures. Our results suggest that only a small percentage (1.2%) of routine radiographs performed during the follow-up period will lead to changes in patient management, with effort and cost involved in generating these radiographs. Just 2 of 936 radiographs taken during the follow-up period (0.2%) led to surgical fixation based on radiologic findings (i.e., secondary dislocation in 1 patient and nonunion in 1 patient scheduled for surgery). These findings should be considered in light of the increasing health care costs and unnecessary exposure to radiation. Although the quantified radiation dose of a single ankle radiograph is low (16), it is difficult to defend administering even small amounts of ionizing radiation, if the indication

to do so is lacking. In addition, each radiograph requires an investment in time from the patient, their companion, and the health care professionals involved.

We divided the therapy of our patients into a treatment period and a follow-up period and focused solely on the latter. We did this to diminish any bias that might arise due to differences in fracture-specific, surgeon-specific, or hospital-specific preferences in the early phases of ankle fracture treatment. Previous studies have also focused on routine radiographs taken in later stages of treatment, when protocols are more standardized or have a greater level of adherence (1,9). The present results are consistent with previous studies (1,5–7,9). For example, Ghattas et al (1), Miniaci-Coxhead et al (10), Ovaska et al (11), and Harish et al (7) demonstrated that radiographs taken at the first postoperative clinic visit of patients with various fracture types did not provide any additional clinically relevant information. Eastley et al (9) studied the effect of radiographs late in the follow-up period of distal radius fractures. To the best of our knowledge, to date, no studies have evaluated the use of routine radiographs in the follow-up period of patients with ankle fractures. The present study explored the use of routine radiographs in a large cohort of patients with a non-Weber type A ankle fracture. We choose not to include isolated Danis Weber type A fractures (Lauge-Hansen supination adduction I), because these mainly represent ligamentous injuries, and no radiologic follow-up is recommended for this type of trauma (3). All types of ankle fractures requiring radiologic follow-up (Lauge-Hansen supination eversion, pronation eversion, supination adduction II, pronation adduction) and all treatment strategies (surgically and conservatively treated patients) were included in the present evaluation.

However, the present study had some important limitations. Given its retrospective character, clinically relevant information that might affect fracture healing (e.g., smoking habits) (17) could not be retrieved from the medical records for many patients. Subsequently, the observed number of changes in treatment strategy might be an underestimation of the assumed effects of these radiographs, because the radiographs can also confirm a correct treatment strategy and acknowledge its continuation. This effect could not be measured using our study design, because this is often not noted in the medical records.

Perhaps even more important is that the clinical indications to generate a radiograph might not always have been properly documented. If no clinical indication was noted in the medical records, a radiograph was considered “routine,” potentially leading to an underestimation of the number of radiographs performed for a clinical indication. We undertook a crude estimation of the costs of routine radiographs during the follow-up period of ankle fractures. Given the potential underestimation of the number of radiographs performed for a clinical indication, these results should be interpreted with care. Second, our analysis does not represent either a cost-effectiveness analysis or a cost-benefit analysis, because the data on the cost associated with a possible gain of health in terms of quality-adjusted life-years or incremental cost differences could not be retrieved from the medical records in the retrospective study design. Similarly, documentation on the continuation of the preset treatment strategy based on the radiographic findings was probably also lacking in many cases. We only considered the documented reasons for a change in treatment, which created a bias such that the total influence of radiographs on the continuation of treatment strategy would have been underestimated. Nevertheless, even if we included a certain range of cases in which continuation of treatment was influenced by routine radiographs, the overall added value of these radiographs would seem overestimated.

In conclusion, although it is common practice to routinely take radiographs during the follow-up period for ankle fractures, the current results suggest that these radiographs seldom influence clinical decision-making and can possibly be omitted. Because of the study limitations, the results of these analyses and the clinical

consequences of a reduced imaging protocol should be confirmed in a prospective trial. Our research group is currently conducting a randomized controlled trial in which a group receiving routine radiographs is compared with a group in which radiographs in the follow-up period are performed only when deemed necessary. These results could help in weighing the clinical importance of routine radiographs and help establish guidelines for their use in the treatment of patients with uncomplicated ankle fractures.

Acknowledgments

We would like to thank Pieta Krijnen, PhD, for her help in the conception of the research plan and critical revisions of the early manuscript and Meltem Akin, MD, for her help in extracting the data from the patients' medical records.

References

- Ghattas TN, Dart BR, Pollock AG, Hinkin S, Pham A, Jones TL. Effect of initial postoperative visit radiographs on treatment plans. *J Bone Joint Surg Am* 95:e57, 2013.
- Mushlin AI. Challenges and opportunities in economic evaluations of diagnostic tests and procedures. *Acad Radiol* 6(suppl 1):S128–S131, 1999.
- Schipper IB, Termaat MF, Rhemrev S, Meylaerts SAG, Bartlema K, Stichter W, Embden D, van de Bremer J, Krug E, Hoogendoorn JM, Alleman J, van der Meulen H, van der Elst M, Greeven A, Niggebrugge A, Ritchie E, van der Linden F, de Vries M, Nagels J, Turkcan K, Cheragwandi A, Dijkstra S. *Richtlijnen Voor Behandeling Van Letsels Van Het Steun En Bewegingsapparaat (Clinical Guideline for the Treatment of Trauma to the Musculoskeletal System)*, pp. 216–222, Optima Grafische Communicatie, Rotterdam, The Netherlands, 2016.
- Chaudhry S, DelSole EM, Egol KA. Post-splinting radiographs of minimally displaced fractures: good medicine or medicolegal protection? *J Bone Joint Surg Am* 94:e128, 2012.
- Chaudhry S, DelSole EM, Egol KA. Post-splinting radiographs of minimally displaced fractures: good medicine or medicolegal protection? *J Bone Joint Surg Am* 94:e128, 2012.
- Chakravarthy J, Mangat K, Qureshi A, Porter K. Postoperative radiographs following hip fracture surgery. do they influence patient management? *Int J Clin Pract* 61:421–424, 2007.
- Harish S, Vince AS, Patel AD. Routine radiography following ankle fracture fixation: a case for limiting its use. *Injury* 30:699–701, 1999.
- Huffaker S, Earp BE, Blazar PE. The value of post-operative radiographs in clinical management of AO type A distal radius fractures. *J Hand Surg Eur* 40:790–795, 2015.
- Eastley N, Aujla R, Khan Z. Radiographs late in the follow up of uncomplicated distal radius fractures: are they worth it? Clinical outcome and financial implications. *Orthop Rev (Pavia)* 4:88–90, 2012.
- Miniaci-Coxhead SL, Martin EA, Ketz JP. Quality and utility of immediate formal postoperative radiographs in ankle fractures. *Foot Ankle Int* 36:1196–1201, 2015.
- Ovaska MT, Nuutinen T, Madanat R, Makinen TJ, Soderlund T. The role of outpatient visit after operative treatment of ankle fractures. *Injury* 47:2575–2578, 2016.
- Nederlandse Zorgautoriteit. Tarievenlijst Eerstelijnsdiagnostiek; 2015. Available at <https://zorgproducten.nza.nl/ZoekZorgproduct.aspx>. Accessed April 28, 2017.
- Centraal Bureau voor de Statistiek. Population Prognosis in the Netherlands, 2015. Available at <https://opendata.cbs.nl/statline/#/CBS/nl/dataset/83597NED/table?ts=1525167532486>. Accessed May 1, 2017.
- Lauge-Hansen N. Fractures of the ankle. II. Combined experimental-surgical and experimental-roentgenologic investigations. *Arch Surg* 60:957–985, 1950.
- Weber BG. *Die Verletzungen Des Oberen Sprunggelenkes*, Huber Verlag, Bern, 1966.
- Lin EC. Radiation risk from medical imaging. *Mayo Clin Proc* 85:1142–1146, 2010.
- Lee JJ, Patel R, Biermann JS, Dougherty PJ. The musculoskeletal effects of cigarette smoking. *J Bone Joint Surg Am* 95:850–859, 2013.