VU Research Portal

Toxicokinetics of Zn and Cd in the earthworm *Eisenia andrei* exposed to metal-contaminated soils under different combinations of air temperature and soil moisture content

González-Alcaraz, M. Nazaret; Loureiro, Susana; van Gestel, Cornelis A.M.

published in

Chemosphere

2018

DOI (link to publisher)

[10.1016/j.chemosphere.2018.01.019](https://doi.org/10.1016/j.chemosphere.2018.01.019)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

González-Alcaraz, M. N., Loureiro, S., & van Gestel, C. A. M. (2018). Toxicokinetics of Zn and Cd in the earthworm *Eisenia andrei* exposed to metal-contaminated soils under different combinations of air temperature and soil moisture content. *Chemosphere*, 197, 26-32. <https://doi.org/10.1016/j.chemosphere.2018.01.019>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Toxicokinetics of Zn and Cd in the earthworm *Eisenia andrei* exposed to metal-contaminated soils under different combinations of air temperature and soil moisture content

M. Nazaret González-Alcaraz ^{a, *}, Susana Loureiro ^b, Cornelis A.M. van Gestel ^a

^a Department of Ecological Science, Faculty of Science, Vrije Universiteit, De Boelelaan 1085, 1081 HV, Amsterdam, The Netherlands

^b Department of Biology & CESAM, Campus Universitário de Santiago, University of Aveiro, 3810-193, Aveiro, Portugal

HIGHLIGHTS

- Climate change simulated by higher air temperature and lower soil moisture content.
- Zn toxicokinetics in *Eisenia andrei* not affected by climate conditions.
- Faster Cd kinetics in earthworms at higher air temperature and soil moisture content.
- Cd kinetics at higher air temperature slowed down with decreasing soil moisture.
- Higher Cd-BAFs in earthworms incubated under warmer and drier conditions.

ARTICLE INFO

Article history:

Received 23 October 2017

Received in revised form

15 December 2017

Accepted 5 January 2018

Available online 8 January 2018

Handling Editor: Jim Lazorchak

Keywords:

Bioaccumulation

Bioavailability

Climate change

Heavy metals

Mining wastes

Soil invertebrates

ABSTRACT

This study evaluated how different combinations of air temperature (20 °C and 25 °C) and soil moisture content (50% and 30% of the soil water holding capacity, WHC), reflecting realistic climate change scenarios, affect the bioaccumulation kinetics of Zn and Cd in the earthworm *Eisenia andrei*. Earthworms were exposed for 21 d to two metal-contaminated soils (uptake phase), followed by 21 d incubation in non-contaminated soil (elimination phase). Body Zn and Cd concentrations were checked in time and metal uptake (k_1) and elimination (k_2) rate constants determined; metal bioaccumulation factor (BAF) was calculated as k_1/k_2 . Earthworms showed extremely fast uptake and elimination of Zn, regardless of the exposure level. Climate conditions had no major impacts on the bioaccumulation kinetics of Zn, although a tendency towards lower k_1 and k_2 values was observed at 25 °C + 30% WHC. Earthworm Cd concentrations gradually increased with time upon exposure to metal-contaminated soils, especially at 50% WHC, and remained constant or slowly decreased following transfer to non-contaminated soil. Different combinations of air temperature and soil moisture content changed the bioaccumulation kinetics of Cd, leading to higher k_1 and k_2 values for earthworms incubated at 25 °C + 50% WHC and slower Cd kinetics at 25 °C + 30% WHC. This resulted in greater BAFs for Cd at warmer and drier environments which could imply higher toxicity risks but also of transfer of Cd within the food chain under the current global warming perspective.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

Metal soil contamination by anthropogenic activities (e.g. mining, smelting, agriculture, waste disposal) is an environmental problem worldwide (COM, 2006; FAO and ITPS, 2015; He et al.,

* Corresponding author. Present address: Department of Biology & CESAM, Campus Universitário de Santiago, University of Aveiro, 3810-193, Aveiro, Portugal.
E-mail address: nazaret.gonzalez@ua.pt (M.N. González-Alcaraz).

2015). Metals exert toxic effects on soil living organisms (van Straalen, 2004; Stankovic et al., 2014), affecting the sustainability of terrestrial ecosystems and, in some cases, human health (Naveed et al., 2014; Zhou et al., 2016; Morgado et al., 2017). Toxicity is known to be related to the metal fraction that can be taken up by organisms and subsequently interact with biological targets (i.e. metal bioavailability; Peijnenburg et al., 2007) rather than to the total metal concentration in the soil. Numerous studies have considered metal body concentrations as estimation of bioavailable

fractions (Heikens et al., 2001). However, metal uptake rates are considered better predictors of their bioavailability (van Straalen et al., 2005). Metal uptake and elimination might occur simultaneously in organisms. To cope with this issue, more accurate uptake rates are estimated when toxicokinetics studies include uptake phases (organisms exposed to contaminated soil) followed by elimination phases without uptake (organisms transferred to non-contaminated soil) (van Straalen et al., 2005).

Metal bioavailability depends on multiple factors such as the considered species, the properties of the soil matrix (e.g. pH, organic matter and texture) and exposure time (Heikens et al., 2001; Allen, 2002; Nahmani et al., 2007; Peijnenburg et al., 2007). Climate conditions, especially air temperature and soil moisture content, also play an important role since they can influence the performance of soil organisms as well as the speciation and therefore the bioavailability of the metals present in the system (Holmstrup et al., 2010; Augustsson et al., 2011; González-Alcaraz and van Gestel, 2015). In the actual context of global warming, studies concerning how climate factors may affect metal bioavailability and thus toxicity to soil organisms are gaining more interest (Løkke et al., 2013; Stahl et al., 2013; Noyes and Lema, 2015). This climatic approach is essential for the future risk assessment of metal-contaminated soils and will help developing adequate remediation strategies (Landis et al., 2013; Rohr et al., 2013).

Earthworms are major components of the soil community (Lavelle and Spain, 2001; Lavelle et al., 2006). They are good bio-indicators of soil health and quality and of the biological impact of metal contamination (Spurgeon et al., 2003). Earthworms have been widely used to evaluate metal bioaccumulation (Heikens et al., 2001; Nahmani et al., 2007) although not many studies have been performed considering future climate predictions. A previous work showed that climate conditions differently affected the bioaccumulation of metals in earthworms depending on the element considered, although in that study no elimination phase in non-contaminated soil was considered after metal exposure (González-Alcaraz and van Gestel, 2016b). The present study is a further attempt to better predict metal bioaccumulation in earthworms under future climate change scenarios, considering both uptake and elimination phases. Therefore, the aim was to evaluate if variations in air temperature and soil moisture content affect the uptake and elimination kinetics of Zn and Cd in the earthworm *Eisenia andrei* exposed to a metal-contaminated soil, tested at two dilution rates with non-contaminated soil. To achieve this goal a toxicokinetics approach was followed under different combinations of air temperature (20 °C and 25 °C) and soil moisture content (50% and 30% of the soil water holding capacity, WHC), earthworms being exposed for 21 d to metal-contaminated soils (uptake phase) followed by 21 d incubation in non-contaminated soil (elimination phase). We hypothesize that different climate conditions would lead to changes in metal bioaccumulation kinetics in earthworms.

2. Materials and methods

2.1. Metal-contaminated test soil

An agricultural field located inside the Campo de Cartagena plain, one of the main intensive irrigated agricultural areas in southern Europe (IMIDA, 2005), and in the vicinity of the former mining district of La Unión-Sierra de Cartagena (Murcia, SE Spain; Figure S1, Supplementary material) was selected to collect the test soil. The area is characterized by a Mediterranean semiarid climate with an annual average temperature of ~18 °C, an annual average precipitation of ~250–300 mm (most falling in spring and autumn in form of short intensive rainfall events) and an average evapotranspiration rate of ~850 mm year⁻¹. The abandonment of the old

tailings has continued leading to the dispersion of great volumes of metal mining wastes via water and/or wind erosion, affecting a wide variety of surrounding ecosystems (Conesa and Jiménez-Cárceles, 2007; Conesa and Schulin, 2010). Numerous studies have pointed at metal contamination problems existing in the area and the urgent need of restoration programs (Jiménez-Cárceles et al., 2008; Párraga-Aguado et al., 2013; Bes et al., 2014; González-Alcaraz and van Gestel, 2016a).

Soil samples were collected (top 20 cm) from three randomly distributed points inside the agricultural field, air dried, sieved through a 2 mm mesh and homogenized before being characterized. No earthworms were found in the agricultural field during soil sampling. The test soil showed clay texture, neutral pH in 0.01 M CaCl₂ (~7), high electrical conductivity (EC ~3 dS m⁻¹), moderate organic matter content determined as loss on ignition (LOI ~5%), high cation exchange capacity (CEC ~16 cmol_c kg⁻¹) and ~47% of WHC (Table 1). Total metal concentrations were high (Cd ~26 mg kg⁻¹, Cu ~80 mg kg⁻¹, Pb ~8733 mg kg⁻¹ and Zn ~8835 mg kg⁻¹; Table 1), compared to the geochemical background levels established for the zone (Cd ~0.3 mg kg⁻¹, Cu ~15 mg kg⁻¹, Pb ~9 mg kg⁻¹ and Zn ~42 mg kg⁻¹; Hernández Bastida et al., 2005; Martínez-Sánchez and Pérez-Sirvent, 2007; Pérez-Sirvent et al., 2009) and the intervention values set for agricultural soils by the nearby Andalusia Region (Cd ~25 mg kg⁻¹, Cu ~595 mg kg⁻¹, Pb ~275 mg kg⁻¹ and Zn ~10,000 mg kg⁻¹; BOJA, 2015). Porewater

Table 1

General characterization of the metal-contaminated test soil from SE Spain and the Lufa 2.2 control soil used for the toxicokinetics study with the earthworm *Eisenia andrei* under different combinations of air temperature and soil moisture content. Values are average ± SD (n = 3). EC (electrical conductivity). LOI (total organic matter determined as loss on ignition). CEC (cation exchange capacity). WHC (water holding capacity). d.l. (detection limit).

Parameter	Test soil	Lufa 2.2 soil
pH 0.01 M CaCl ₂ ^a	7.01 ± 0.05	5.21 ± 0.04
EC (dS m ⁻¹) ^b	2.95 ± 0.09	0.10 ± 0.002
LOI (%) ^c	5.30 ± 0.10	3.12 ± 0.05
CEC (cmol _c kg ⁻¹) ^d	16.3 ± 0.6	7.8 ± 1.9
WHC (%) ^e	46.5 ± 0.5	44.4 ± 0.7
Texture ^f	Clay	Sandy loam
Porewater metals ^g		
Cd (μg L ⁻¹)	28.7 ± 2.1	<d.l. (3)
Cu (μg L ⁻¹)	43.3 ± 1.2	59.0 ± 17.1
Pb (μg L ⁻¹)	67.3 ± 13.1	32.3 ± 17.5
Zn (μg L ⁻¹)	383 ± 37	16.0 ± 17.5
0.01 M CaCl ₂ -extractable metals ^h		
Cd (μg kg ⁻¹)	81.6 ± 2.9	<d.l. (15)
Cu (μg kg ⁻¹)	<d.l. (30)	<d.l. (30)
Pb (μg kg ⁻¹)	<d.l. (225)	<d.l. (225)
Zn (μg kg ⁻¹)	989 ± 87	246 ± 3
Total metals ⁱ		
Cd (mg kg ⁻¹)	25.6 ± 0.1	<d.l. (0.2)
Cu (mg kg ⁻¹)	80.3 ± 4.4	3.1 ± 0.1
Pb (mg kg ⁻¹)	8733 ± 2479	15.0 ± 2.1
Zn (mg kg ⁻¹)	8835 ± 96	23.6 ± 2.5

^a 1:5 (w:v) soil:0.01 M CaCl₂ suspensions after 2 h shaking at 200 rpm.

^b 1:5 (w:v) soil:H₂O suspensions after 2 h shaking at 200 rpm.

^c Combustion following a heating ramp from 200 °C to 500 °C for 8 h.

^d Saturation of soil exchange complex with 1 M CH₃COONa pH 8.2 and displacement of adsorbed sodium with 1 M CH₃COONH₄ pH 7.0 (Chapman, 1965). Sodium concentration determination by flame atomic absorption spectroscopy (AAS; Perkin-Elmer Analyst 100).

^e Sandbox method after soil saturation with water for 3 h (ISO, 1999).

^f Laser grain size HELOS-QUIXEL analyzer (Konert and Vandenberghe, 1997).

^g Soil saturation with water at 100% WHC for 7 d, centrifugation for 45 min at 2000 rcf over a 0.45 μm membrane filter and metal concentrations determined by flame AAS.

^h Metal concentrations determined in 0.01M CaCl₂ extracts by flame AAS.

ⁱ Acid digestion in 4:1(v:v) HNO₃ 65%:HCl 37% at 140 °C for 7 h. Metal concentrations determined by flame AAS.

metal concentrations were $\sim 29 \mu\text{g L}^{-1}$ for Cd, $\sim 43 \mu\text{g L}^{-1}$ for Cu, $\sim 67 \mu\text{g L}^{-1}$ for Pb and $\sim 383 \mu\text{g L}^{-1}$ for Zn (Table 1). Exchangeable metals (extracted with 0.01 M CaCl_2) showed low concentrations except for Cd ($\sim 82 \mu\text{g kg}^{-1}$) and Zn ($\sim 989 \mu\text{g kg}^{-1}$) (Table 1).

2.2. Experimental set-up

2.2.1. Test species

Eisenia andrei Bouché 1972 (Class Oligochaeta, Family Lumbricidae) was cultured at the Vrije Universiteit (Amsterdam, The Netherlands) for >10 years in clean horse manure free of any pharmaceuticals at 20 °C, 75% relative humidity and complete darkness (OECD, 2010). Earthworms were originally obtained from ECT Oekotoxikologie in Flörsheim (Germany) where they were genotyped to confirm their species identity (Römbke et al., 2016).

Before starting the toxicokinetics experiment, synchronized sexually mature earthworms (well-developed clitella and ~ 300 – 700 mg fresh weight) were transferred to clean soil (Lufa 2.2; Speyer, Germany) and kept for several hours (~ 6) for acclimation to soil conditions and to replace the gut content of horse manure by soil (Vijver et al., 2005; OECD, 2010). This acclimatization phase was performed in complete darkness at 20 °C and 75% relative humidity.

2.2.2. Soil preparation

The metal-contaminated test soil was mixed with the standard reference soil Lufa 2.2 (Table 1) at ratios (w:w) of 1:1 (50% metal-contaminated soil, hereafter named test soil 1:1) and 1:3 (25% metal-contaminated soil, hereafter named test soil 1:3). Soil mixtures were prepared with dry soils. This dilution approach allowed earthworms to burrow in the soil since the clay texture of the original study soil limited their movement (authors' visual observation from pilot tests performed with the metal-contaminated test soil). To prevent changes in metal availability in the mixing process, the pH (in 0.01 M CaCl_2) of the Lufa 2.2 soil was adjusted with CaCO_3 to approximately 7 (by adding $4 \text{ mg CaCO}_3 \text{ g}^{-1}$ dry soil) to mimic the pH of the metal-contaminated test soil (Table 1). The WHC of each soil mixture ($\sim 42\%$ for soil 1:1 and $\sim 39\%$ for soil 1:3) was determined using the sandbox method after saturation of the soil with water for 3 h (ISO, 1999).

2.2.3. Toxicokinetics

Toxicokinetics tests with *E. andrei* were performed according to the standardized test guideline OECD 317 (OECD, 2010). The climate conditions recommended by the guideline are 20 °C of air temperature and a soil moisture content of approximately 50% of the soil WHC (standard climate conditions; OECD, 2010). From these standard conditions and in order to recreate future climate predictions for southern parts of Europe (~ 4 °C of temperature increase and ~ 10 – 20% of soil moisture content decrease; Bates et al., 2008; Forzieri et al., 2014), an increase of 5 °C in air temperature and a decrease of 20% in soil WHC were chosen. Toxicokinetics tests were performed for both soil mixtures (soil 1:1 and soil 1:3) under four different climate conditions: 1) 20 °C + 50% WHC (standard climate conditions), 2) 20 °C + 30% WHC, 3) 25 °C + 50% WHC and 4) 25 °C + 30% WHC (climate conditions simulating warming and drier environments).

Toxicokinetics tests consisted of two phases (uptake and elimination), each one lasting 21 d. Before each phase earthworms were rinsed with demineralized water, dried on filter paper and weighed. In the uptake phase earthworms were exposed to both soil mixtures (soil 1:1 and soil 1:3), and then transferred to pH-adjusted Lufa 2.2 soil for the elimination phase. In both phases earthworms were kept individually in 100 mL glass jars containing 30 g of soil previously moistened and 2 g (dry weight) of moistened

horse dung for food. Soil moistening was done just before starting the experiment. Tests were run under the different climate conditions established in controlled climate chambers with 75% relative humidity and a 12:12 h light:dark photoperiod (OECD, 2010). Soil moisture content was checked twice a week by weighing the test jars and water loss replenished with demineralized water to keep the initial soil moisture content. At time points 0 (background body metal concentrations), 1, 3, 7, 10, 14 and 21 d during the uptake phase and 22, 24, 28, 31, 35 and 42 d during the elimination phase three earthworms were sacrificed for the determination of the body metal concentrations (three replicates per soil mixture/climate condition/time point). Sampled earthworms were depurated on moist filter paper for 24 h in a petri dish to fully purge their gut content (OECD, 2010), rinsed with demineralized water, dried on filter paper, weighted (to evaluate weight change throughout the experiment) and frozen at -20 °C.

Two control sets were performed, one with the original Lufa 2.2 soil (pH in 0.01 M CaCl_2 ~ 5.2 ; Table 1) and another one with the pH-adjusted Lufa 2.2 soil used for soil mixture preparation (pH in 0.01 M CaCl_2 ~ 7.0). The first control allowed checking for earthworm performance in non-contaminated soil (OECD, 2010), the second control if soil pH was causing differences in earthworm performance. Control tests were performed under the four climate conditions established following the methodology described above. Earthworm survival, weight change and body metal concentrations were checked at the end of the uptake (after 21 d) and elimination (after 42 d) phases (six replicates per control soil/climate condition/time point).

2.2.4. Chemical analysis

Frozen earthworms were freeze-dried for 48 h, weighted and digested in 4:1 (v:v) HNO_3 65%:HCl 37% in Teflon bombs heated for 7 h at 140 °C in a destruction oven (Binder). The concentrations of Zn and Cd were measured by flame atomic absorption spectroscopy (Perkin-Elmer AAnalyst 100; detection limit 3 mg L^{-1}). Body metal concentrations are expressed on a dry weight (d.w.) basis. Quality control was checked with the certified reference materials DOLT4 (Dogfish liver, LGCS Standards) and Bovine Liver (BCR-185R); recoveries were 110–117% for Zn and 113–119% for Cd.

2.2.5. Kinetic modelling

For each soil mixture (soil 1:1 and soil 1:3) a first-order one-compartment kinetic model was applied to describe metal uptake and elimination rates in the earthworms. Eqs. (1) and (2) were used to describe the uptake and elimination phases, respectively:

$$C_t = C_0 + (k_1/k_2) * C_{\text{exp}} * (1 - e^{-k_2 * t}) \quad (1)$$

$$C_t = C_0 + (k_1/k_2) * C_{\text{exp}} * (e^{-k_2 * (t-t_c)} - e^{-k_2 * t}) \quad (2)$$

where C_t = body metal concentration in earthworms ($\mu\text{g g}^{-1}$ d.w.) at time t (d); C_0 = background body metal concentration in earthworms ($\mu\text{g g}^{-1}$ d.w.); k_1 = uptake rate constant ($\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$); k_2 = elimination rate constant (d^{-1}); C_{exp} = total metal concentration in soil calculated from mixture proportion ($\mu\text{g g}^{-1}$ dry soil); t_c = time at which the earthworms were transferred to non-contaminated soil (21 d). Uptake and elimination equations were fitted simultaneously. A growth rate constant (k_g) was included in the kinetic model to consider changes in earthworm body weight throughout the experiment, but this did not affected k_1 and k_2 values. Results shown therefore are those derived using Eqs. (1) and (2).

A kinetic metal bioaccumulation factor (BAF) in earthworms was calculated as k_1/k_2 (Peijnenburg et al., 1999). Half-life for the elimination of the metals from the earthworms after exposure to

the soil mixtures was calculated as $\ln(2)/k_2$.

2.2.6. Statistical analyses

Statistical analyses were performed with IBM SPSS Statistics 22 and differences were considered significant at $p < 0.05$. For each soil mixture (soil 1:1 and soil 1:3), differences in k_1 and k_2 values among the climate conditions tested were evaluated by generalized likelihood ratio tests (Sokal and Rohlf, 1969; van Gestel and Hensbergen, 1997). No statistical analyses could be performed for earthworm fresh weight due to the fact that organisms from the same soil/climate condition/time point were pooled together for cleaning the gut content before being weighed. This made it difficult to distinguish earthworms based on their initial fresh weight. Therefore the data from the different replicates were pooled.

3. Results and discussion

3.1. Earthworm performance under different climate conditions

The validity of the tests performed with *E. andrei* was evaluated according to the following criteria (OECD, 2010): 1) mortality at the end of the test $\leq 10\%$; 2) weight loss at the end of the uptake and elimination phases compared to the initial fresh weight for each phase $\leq 20\%$. These criteria apply both for controls (original Lufa 2.2 soil and pH-adjusted Lufa 2.2 soil) and soil mixtures (soil 1:1 and soil 1:3) under standard climate conditions ($20^\circ\text{C} + 50\%$ WHC). No mortality was registered in controls and soil 1:3 (25% metal-contaminated soil). In soil 1:1 (50% metal-contaminated soil) one earthworm died (3% mortality). When exposed to standard climate conditions, the earthworms tended to lose weight throughout the experiment (average weight loss at the end of the uptake and elimination phases, respectively): original Lufa 2.2 soil ($\sim 13\%$ and $\sim 20\%$); pH-adjusted Lufa 2.2 soil ($\sim 13\%$ and $\sim 16\%$); soil 1:1 ($\sim 8\%$ and $\sim 6\%$); soil 1:3 ($\sim 10\%$ and $\sim -2\%$) (data not shown). Therefore, the validity criteria established by OECD were met.

Earthworm body weight was affected by changing air temperature and soil moisture content compared to the standard climate conditions. In both control soils earthworm weight loss at the end of the uptake and elimination phases was most pronounced at 25°C . At $20^\circ\text{C} + 30\%$ WHC earthworm weight loss was $\sim 9\text{--}11\%$ for the original Lufa 2.2 soil and $\sim 16\text{--}19\%$ for the pH-adjusted Lufa 2.2 soil (data not shown). At 25°C , regardless of the soil moisture content (50% and 30% WHC), earthworm weight loss was $\sim 16\text{--}33\%$ for the original Lufa 2.2 soil and $\sim 20\text{--}29\%$ for the pH-adjusted Lufa 2.2 soil (data not shown). This trend agrees with a previous study where earthworms showed higher weight loss at 25°C compared to 20°C , and no influence was found of the pH of the Lufa 2.2 soil (González-Alcaraz and van Gestel, 2016b). Lima et al. (2011, 2015) also found greater weight loss for *E. andrei* in Lufa 2.2 soil with increasing air temperature (20°C vs. 26°C) and no effect of soil moisture content (60%, 40%, 20% and 10% of soil WHC). However, our results do not agree with other studies showing decreasing body weight with lowered soil moisture content in the earthworm species *Eisenia fetida* (Diehl and Williams, 1992) and *Aporrectodea caliginosa* (Holmstrup, 2001).

In both soil mixtures (soil 1:1 and soil 1:3) earthworms incubated at $25^\circ\text{C} + 30\%$ WHC reached the highest weight loss values at the end of the uptake phase ($\sim 49\%$ and $\sim 44\%$ after 21 d exposure, respectively; Figure S2, Supplementary material), showing a synergistic interaction between metal contamination and warmer and drier conditions (Friis et al., 2004; Holmstrup et al., 2010; González-Alcaraz and van Gestel, 2016b). When transferred from metal-contaminated to non-contaminated soil, however, earthworms tended to gain weight, especially those incubated at $25^\circ\text{C} + 30\%$ WHC (weight gain $\sim 22\%$ and $\sim 14\%$ after 21 d in clean soil for

organisms earlier exposed to soil 1:1 and soil 1:3, respectively; Figure S2, Supplementary material).

3.2. Metal toxicokinetics in earthworms under different climate conditions

Background body metal concentrations in earthworms were $\sim 100\text{--}120\ \mu\text{g g}^{-1}$ d.w. for Zn (Fig. 1) and $\sim 2\text{--}3\ \mu\text{g g}^{-1}$ d.w. for Cd (Fig. 2), normal levels for earthworms from non-contaminated soils (Zn $\sim 90\text{--}120\ \mu\text{g g}^{-1}$ d.w. and Cd $\sim 3\text{--}6\ \mu\text{g g}^{-1}$ d.w.; Janssen et al., 1997; van Gestel et al., 2002). Similar body metal concentrations were found in earthworms exposed to control soils for 42 d under the different climate conditions tested (Zn $\sim 80\text{--}160\ \mu\text{g g}^{-1}$ d.w. and Cd $\sim 2\text{--}11\ \mu\text{g g}^{-1}$ d.w.; data not shown). When earthworms were exposed to metal contamination different bioaccumulation patterns were observed for Zn (essential element) and Cd (non-essential element) (Figs. 1 and 2).

Body Zn concentrations increased rapidly after few days of exposure to both soil mixtures (soil 1:1 and soil 1:3), reaching a steady state at body Zn concentrations $\sim 240\text{--}420\ \mu\text{g g}^{-1}$ d.w. (Fig. 1). When transferred to non-contaminated soil, body Zn concentrations rapidly decreased to background levels ($\sim 110\text{--}140\ \mu\text{g g}^{-1}$ d.w.; Fig. 1). This bioaccumulation pattern seems typical for Zn as it has previously been shown also in other studies (Spurgeon and Hopkin, 1999; Świątek et al., 2017), and may be explained from the presence of efficient regulation mechanisms. Zinc regulation in earthworms occurs via excretion (Spurgeon and Hopkin, 1999), leading to high k_2 values (15–42 fold higher than k_1 values; Table 2) and short half-lives (< 1 d; Table 2). Changing air temperature and soil moisture content had no major effects on the bioaccumulation pattern of Zn (Fig. 1). This agrees with González-

Fig. 1. Uptake and elimination kinetics of Zn in the earthworm *Eisenia andrei* exposed to the mixtures of the metal-contaminated test soil with the pH-adjusted Lufa 2.2 soil under the different climate conditions tested: (A) soil 1:1 (50% metal-contaminated soil); (B) soil 1:3 (25% metal-contaminated soil). Uptake and elimination phases lasted 21 d each. Dots represent average body concentrations (on a dry weight basis, d.w.) $\pm$ SE ($n = 3$). Lines represent modelled Zn body concentrations using Eqs. (1) and (2). WHC (water holding capacity).

Fig. 2. Uptake and elimination kinetics of Cd in the earthworm *Eisenia andrei* exposed to the mixtures of the metal-contaminated test soil with the pH-adjusted Lufa 2.2 soil under the different climate conditions tested: (A) soil 1:1 (50% metal-contaminated soil); (B) soil 1:3 (25% metal-contaminated soil). Uptake and elimination phases lasted 21 d each. Dots represent average body concentrations (on a dry weight basis, d.w.) $\pm$ SE ($n = 3$). Lines represent modelled Cd body concentrations using Eqs. (1) and (2). WHC (water holding capacity).

Alcaraz and van Gestel (2016b) who found no impact of climate conditions on Zn bioaccumulation in *E. andrei* exposed for 21 d to metal-contaminated soils of different properties. Despite this, for both soil mixtures (soil 1:1 and soil 1:3), the treatment at 25 °C + 30% showed lower k_1 and k_2 values compared to the other climate conditions tested (Table 2).

Unlike Zn, body Cd concentrations in earthworms tended to increase with exposure time throughout the uptake phase and stayed more or less constant or slowly decreased upon transfer to non-contaminated soil (Fig. 2). This is a typical pattern for non-essential elements, with earthworms generally showing very slow or no elimination of Cd (Spurgeon and Hopkin, 1999; Lock and Janssen, 2001; Smith et al., 2010; Giska et al., 2014). Cadmium detoxification in earthworms occurs via its sequestration by metallothioneins (Stürzenbaum et al., 2001, 2004; Conder et al., 2002; Vijver et al., 2006). This agrees with the low k_2 values obtained

(1.5–21 fold lower than k_1 values; Table 3). At the end of the uptake phase (21 d of exposure), higher body Cd concentrations were found in earthworms incubated at 50% of the soil WHC, regardless of the air temperature (2.4–3.6 and 1.2–3.1 fold higher in soil 1:1 and soil 1:3, respectively; Fig. 2). The treatment at 25 °C + 50% WHC showed the highest k_1 (~ 0.16 vs. ~ 0.01 – 0.06 $\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$ in soil 1:1; ~ 0.31 vs. ~ 0.04 – 0.19 $\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$ in soil 1:3) and k_2 (~ 0.11 vs. ~ 0 – 0.01 d^{-1} in soil 1:1; ~ 0.12 vs. ~ 0.002 – 0.04 d^{-1} in soil 1:3) values (Table 3). This could be related to a higher metabolic activity when earthworms (poikilothermic organisms) were incubated at higher temperature, enhancing Cd uptake and elimination, which resulted in shorter half-lives (~ 7 vs. ~ 75 – 81 d in soil 1:1; ~ 6 vs. ~ 16 – 377 d in soil 1:3; Table 3). However, this was not the case for earthworms incubated at 25 °C + 30% WHC which showed lower k_1 and k_2 values (Table 3), similar to what happened for Zn bioaccumulation (Table 2). This difference was more marked compared to the treatments moistened at 50% of the soil WHC, especially in soil 1:3 (25% metal-contaminated soil): k_1 values were 5–9 fold lower and k_2 values 24–62 fold lower at 25 °C + 30% WHC (significant, $p < 0.05$; Table 3). A warmer and drier environment could have hindered earthworm performance, as shown by the greater weight loss upon exposure to metal-contaminated soils (Figure S2, Supplementary material), slowing down metal uptake and elimination. Therefore, the bioaccumulation pattern of Cd in earthworms changed when changing climate conditions. This agrees with the results of González-Alcaraz and van Gestel (2016b), although they found increasing k_1 and k_2 values at higher air temperature and/or lower soil moisture content. Differences in the properties of the test soils as well as not including an elimination phase in non-contaminated soil in the toxicokinetic study could be responsible of the different results obtained.

BAF values can be used as indicators of soil metal bioavailability (Fründ et al., 2011) and to predict risks of trophic transfer (Smith et al., 2010); $\text{BAF} > 1$ indicates metal accumulation within organisms. For Zn, due to its fast elimination, BAFs were below 1 both in soil 1:1 and soil 1:3 and under the different climate conditions tested (~ 0.02 – 0.07 $\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$; Table 2). But for Cd, BAFs were above 1 (~ 1.50 – 21.3 $\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$; Table 3), indicating that earthworms concentrated Cd within their body (Smith et al., 2010). BAFs for Cd differed among exposure concentrations and climate conditions. Higher BAFs were found when earthworms were exposed to soil 1:3 (25% metal-contaminated soil) (Table 3), in agreement with increasing BAFs for metals at lower exposure levels (McGeer et al., 2003). Moreover, in soil 1:3, the treatment at 25 °C + 30% WHC showed the highest BAF value compared to the other climate conditions tested (~ 21.3 vs. ~ 3.0 – 5.3 $\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$; Table 3). Therefore the bioaccumulation potential of Cd in earthworms not only depended on the exposure level but also on the climate conditions, with greater Cd bioaccumulation at warmer and drier environments.

Table 2

Uptake rate constant (k_1), elimination rate constant (k_2), bioaccumulation factor (BAF) and half-life for the bioaccumulation of Zn in the earthworm *Eisenia andrei* exposed to the mixtures of the metal-contaminated test soil with the pH-adjusted Lufa 2.2 soil under the different climate conditions tested: soil 1:1 (50% metal-contaminated soil) and soil 1:3 (25% metal-contaminated soil). No 95% confidence intervals could be calculated for the k_1 and k_2 values. WHC (water holding capacity).

Contaminated soil	Climate condition	k_1 ($\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$)	k_2 (d^{-1})	BAF ($\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$)	Half-life (d)
1:1	20 °C + 50% WHC	0.47	16.9	0.03	0.04
	20 °C + 30% WHC	0.37	14.0	0.02	0.05
	25 °C + 50% WHC	0.50	13.0	0.04	0.05
	25 °C + 30% WHC	0.37	11.1	0.03	0.06
1:3	20 °C + 50% WHC	1.00	15.0	0.07	0.05
	20 °C + 30% WHC	0.86	14.4	0.06	0.05
	25 °C + 50% WHC	0.61	12.9	0.05	0.05
	25 °C + 30% WHC	0.51	12.6	0.04	0.06

Table 3

Uptake rate constant (k_1), elimination rate constant (k_2), bioaccumulation factor (BAF) and half-life for the bioaccumulation of Cd in the earthworm *Eisenia andrei* exposed to the mixtures of the metal-contaminated test soil with the pH-adjusted Lufa 2.2 soil under the different climate conditions tested: soil 1:1 (50% metal-contaminated soil) and soil 1:3 (25% metal-contaminated soil). 95% confidence intervals are given in between brackets. For each percentage of contaminated soil, different letters at the same column indicate significant differences among climate conditions (likelihood ratio test, $p < 0.05$). WHC (water holding capacity).

Contaminated soil	Climate condition	k_1 ($\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$)	k_2 (d^{-1})	BAF ($\text{g}_{\text{soil}} \text{g}^{-1} \text{earthworm d}^{-1}$)	Half-life (d)
1:1	20 °C + 50% WHC	0.055 b (0.035–0.074)	0.009 b (0 ^a –0.029)	6.42	81.3
	20 °C + 30% WHC	0.011 c (0.004–0.017)	0 ^a	–	–
	25 °C + 50% WHC	0.159 a (0.103–0.216)	0.106 a (0.062–0.150)	1.50	6.5
	25 °C + 30% WHC	0.020 c (0.011–0.030)	0.009 b (0 ^a –0.036)	2.19	75.2
1:3	20 °C + 50% WHC	0.194 a (0.113–0.276)	0.044 ac (0.012–0.077)	4.38	15.6
	20 °C + 30% WHC	0.087 b (0.057–0.117)	0.016 b (0 ^a –0.037)	5.30	42.3
	25 °C + 50% WHC	0.306 a (0.183–0.431)	0.115 a (0.061–0.169)	3.02	6.1
	25 °C + 30% WHC	0.039 c (0.023–0.055)	0.002 bc (0 ^a –0.024)	21.3	377

^a Each zero comes from a negative k_2 value generated by the mathematical model (Eqs. (1) and (2)). It means that there was no elimination.

4. Conclusions

The earthworm *E. andrei* rapidly accumulated Zn to a steady state level when exposed to metal-contaminated soils, but also rapidly eliminated Zn to reach background levels upon transfer to non-contaminated soil. This suggests efficient regulation of Zn body concentrations. Air temperature (20 °C and 25 °C) and soil moisture content (50% and 30% of the soil WHC) had no major impacts on the bioaccumulation kinetics of Zn, although a tendency to lower uptake and elimination rates was observed at 25 °C + 30% WHC. On the contrary, different combinations of air temperature and soil moisture content changed the bioaccumulation kinetics of Cd. Earthworms incubated at high soil moisture content had higher body Cd concentrations upon exposure to metal contamination. When high temperature was combined with high soil moisture content earthworms showed faster uptake and elimination rates for Cd. However, when high temperature was combined with low soil moisture content, slower Cd kinetics was found (lower uptake and elimination rates at 25 °C and 30% of the soil WHC). This resulted in higher BAFs for Cd when earthworms were incubated under warmer and drier environments. These findings could not only imply higher toxicity risks for earthworms in metal-contaminated soils under the actual global warming perspective, but also of transfer/biomagnification of Cd within the food chain. The latter is of major concern if we take into account that earthworms are at the lower levels of most wildlife food chains. Therefore, and considering future climate predictions, more studies concerning the influence of climate factors on metal bioavailability to soil invertebrates are needed to properly predict and manage their potential risks.

Conflicts of interest

There is no conflict of interest.

Acknowledgements

The authors acknowledge funding to the GLOBALTOX project through the Research Executive Agency (REA-European Commission) under the Marie Skłodowska-Curie actions (H2020-MSCA-IF-2015/H2020-MSCA-IF-2015, Project ID 704332). We thank Rudo A. Verweij from the Vrije Universiteit (Amsterdam, The Netherlands) for his valuable contribution to the laboratory work.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.chemosphere.2018.01.019>.

References

- Allen, H.E., 2002. Bioavailability of Metals in Terrestrial Ecosystems: Importance of Partitioning for Bioavailability to Invertebrates, Microbes, and Plants. Society of Environmental Toxicology and Chemistry (SETAC), Pensacola, Florida.
- Augustsson, A., Filipsson, M., Öberg, T., Bergbäck, B., 2011. Climate change - an uncertainty factor in risk analysis of contaminated land. *Sci. Total Environ.* 409, 4693–4700.
- Bates, B.C., Kundzewicz, Z.W., Wu, S., Palutikof, J.P., 2008. Climate Change and Water. Technical Paper of the Intergovernmental Panel on Climate Change. IPCC Secretariat, Geneva, Switzerland.
- Bes, C.M., Pardo, T., Bernal, M.P., Clemente, R., 2014. Assessment of the environmental risks associated with two mine tailing soils from the La Unión-Cartagena (Spain) mining district. *J. Geochem. Explor.* 147, 98–106.
- BOJA, 2015. Decreto 18/2015, de 27 de enero, por el que se aprueba el reglamento que regula el régimen aplicable a los suelos contaminados. Boletín Oficial de la Junta de Andalucía. Consejería de Medio Ambiente de la Junta de Andalucía, Spain.
- Chapman, H.D., 1965. Cation-exchange capacity. In: Black, C.A. (Ed.), *Methods of Soil Analysis, Part 2: Chemical and Microbiological Properties*. American Society of Agronomy, Madison, pp. 891–900.
- COM, 2006. Communication from the Commission to the Council, the European Parliament, the European Economic and Social Committee and the Committee of the Regions - Thematic Strategy for Soil Protection [SEC(2006)620] [SEC(2006)1165].
- Conder, J.M., Seals, L.D., Lanno, R.P., 2002. Method for determining toxicologically relevant cadmium residues in the earthworm *Eisenia fetida*. *Chemosphere* 49, 1–7.
- Conesa, H.M., Jiménez-Cárceles, F.J., 2007. The Mar Menor lagoon (SE Spain): a singular natural ecosystem threatened by human activities. *Mar. Pollut. Bull.* 54, 839–849.
- Conesa, H.M., Schulin, R., 2010. The Cartagena-La Unión mining district (SE Spain): a review of environmental problems and emerging phytoremediation solutions after fifteen years research. *J. Environ. Monit.* 12, 1225–1233.
- Diehl, W., Williams, D., 1992. Interactive effects of soil moisture and food on growth and aerobic metabolism in *Eisenia fetida* (Oligochaeta). *Comp. Biochem. Physiol.* A 102, 179–184.
- FAO and ITPS, 2015. Status of the World's Soil Resources (SWRS) – Main Report. Food and Agriculture Organization of the United Nations and Intergovernmental Technical Panel on Soils, Rome, Italy.
- Forzieri, G., Feyen, L., Rojas, R., Flörke, M., Wimmer, F., Biamchi, A., 2014. Ensemble projections of future streamflow droughts in Europe. *Hydrol. Earth Syst. Sci.* 18, 85–108.
- Friis, K., Damgaard, C., Holmstrup, M., 2004. Sublethal soil copper concentrations increase mortality in the earthworm *Aporrectodea caliginosa* during drought. *Ecotoxicol. Environ. Saf.* 57, 65–73.
- Fründ, H.C., Graefe, U., Tischer, S., 2011. Earthworms as bioindicators of soil quality. In: Karaca, A. (Ed.), *Biology of Earthworms, Soil Biology*, vol. 24. Springer-Verlag Berlin Heidelberg, Germany, pp. 261–278.
- Giska, I., van Gestel, C.A.M., Skip, B., Laskowski, R., 2014. Toxicokinetics of metals in the earthworm *Lumbricus rubellus* exposed to natural polluted soils e relevance of laboratory tests to the field situation. *Environ. Pollut.* 190, 123–132.
- González-Alcaraz, M.N., van Gestel, C.A.M., 2015. Climate change effects on enchytraeid performance in metal-polluted soils explained from changes in metal bioavailability and bioaccumulation. *Environ. Res.* 142, 177–184.
- González-Alcaraz, M.N., van Gestel, C.A.M., 2016a. Toxicity of a metal(loid)-polluted agricultural soil to *Enchytraeus crypticus* changes under a global warming perspective: variations in air temperature and soil moisture content. *Sci. Total Environ.* 573, 203–211.
- González-Alcaraz, M.N., van Gestel, C.A.M., 2016b. Metal/metalloid (As, Cd and Zn) bioaccumulation in the earthworm *Eisenia andrei* under different scenarios of climate change. *Environ. Pollut.* 215, 178–186.
- He, Z., Shentu, J., Yang, X., Baligar, V.C., Zhang, T., Stoffella, P.J., 2015. Heavy metal contamination of soils: sources, indicators, and assessment. *J. Environ. Indic.*

- 9, 17–18.
- Heikens, A., Peijnenburg, W.J.G.M., Hendriks, A.J., 2001. Bioaccumulation of heavy metals in terrestrial invertebrates. *Environ. Pollut.* 113, 385–393.
- Hernández Bastida, J.A., Fernández Tapia, M.T., Alarcón Bernal, M.A., 2005. Valores de fondo y valores genéricos de referencia para Cd, Co, Cr, Cu, Mn, Ni, Pb y Zn en suelos del Campo de Cartagena, Murcia (SE Spain). *Edafología* 12, 105–114.
- Holmstrup, M., 2001. Sensitivity of life history parameters in the earthworm *Aporrectodea caliginosa* to small changes in soil water potential. *Soil Biol. Biochem.* 33, 1217–1223.
- Holmstrup, M., Bindesbøl, A.M., Oostingh, G.J., Duschl, A., Scheil, V., Köhler, H.R., Loureiro, S., Soares, A.M.V.M., Ferreira, A.L.G., Kienle, C., Gerhardt, A., Laskowski, R., Kramarz, P.E., Bayley, M., Svendsen, C., Spurgeon, D.J., 2010. Interactions between effects of environmental chemicals and natural stressors: a review. *Sci. Total Environ.* 408, 3746–3762.
- IMIDA, 2005. Gestión integrada del agua en la Región de Murcia: El caso del Campo de Cartagena. IMIDA (Instituto Murciano de Investigación y Desarrollo Agrario y Alimentario), Murcia, Spain.
- ISO, 1999. ISO 11267: Soil Quality – Inhibition of Reproduction of Collembola (*Folsomia candida*) by Soil Pollutants. ISO (International Organization for Standardization Organization), Geneva, Switzerland.
- Janssen, R.P.T., Posthuma, L., Baerselman, R., Den Hollander, H.A., van Veen, R.P.M., Peijnenburg, W.J., 1997. Equilibrium partitioning of heavy metals in Dutch field soils. II. Prediction of metal accumulation in earthworms. *Environ. Toxicol. Chem.* 16, 2479–2488.
- Jiménez-Cárceles, F.J., Álvarez-Rogel, J., Conesa Alcaraz, H.M., 2008. Trace element concentrations in saltmarsh soils strongly affected by wastes from metal sulphide mining areas. *Water Air Soil Pollut.* 188, 283–295.
- Konert, M., Vandenbergh, J., 1997. Comparison of laser grain size analysis with pipette and sieve analysis: a solution for the underestimation of the clay fraction. *Sedimentology* 44, 523–535.
- Landis, W.G., Dorda, J.L., Brooks, M.L., Chapman, P.M., Menzie, C.A., Stahl Jr., R.G., Stauber, J.L., 2013. Ecological risk assessment in the context of global climate change. *Environ. Toxicol. Chem.* 32, 79–92.
- Lavelle, P., Decaëns, T., Aubert, M., Barot, S., Blouin, M., Bureau, F., Margerie, P., Mora, P., Rossi, J.P., 2006. Soil invertebrates and ecosystem services. *Eur. J. Soil Biol.* 42, 3–15.
- Lavelle, P., Spain, A.V., 2001. *Soil Ecology*. Kluwer Scientific Publications, Amsterdam, The Netherlands.
- Lima, M.P.R., Cardoso, D.N., Soares, A.M.V.M., Loureiro, S., 2015. Carbaryl toxicity prediction to soil organisms under high and low temperature regimes. *Ecotoxicol. Environ. Saf.* 114, 263–272.
- Lima, M.P.R., Soares, A.M.V.M., Loureiro, S., 2011. Combined effects of soil moisture and carbaryl to earthworms and plants: simulation of flood and drought events. *Environ. Pollut.* 159, 1844–1851.
- Lock, K., Janssen, C.R., 2001. Zinc and cadmium body burdens in terrestrial oligochaetes: use and significance in environmental risk assessment. *Environ. Toxicol. Chem.* 20, 2067–2072.
- Løkke, H., Rags, A.M.J., Holmstrup, M., 2013. Tools and perspectives for assessing chemical mixtures and multiple stressors. *Toxicology* 313, 73–82.
- Martínez-Sánchez, M.J., Pérez-Sirvent, C., 2007. Niveles de fondo y niveles genéricos de referencia de metales pesados en suelos de la Región de Murcia. *Consejería de Desarrollo Sostenible y Ordenación del Territorio*, Murcia, España.
- McGeer, J.C., Brix, K.V., Skeaff, J.M., DeForest, D.K., Brigham, S.I., Adams, W.J., Green, A., 2003. Inverse relationship between bioconcentration factor and exposure concentration for metals: implications for hazard assessment of metals in the aquatic environment. *Environ. Toxicol. Chem.* 22, 1017–1037.
- Morgado, R.G., Loureiro, S., González-Alcaraz, M.N., 2017. Changes in ecosystem structure and soil functions due to soil contamination. In: Duarte, A., Cachada, A., Rocha-Santos, T. (Eds.), *Soil Pollution: from Monitoring to Remediation*, first ed. Academic Press, Elsevier, pp. 59–87.
- Nahmani, J., Hodson, M.E., Black, S., 2007. A review of studies performed to assess metal uptake by earthworms. *Environ. Pollut.* 145, 402–424.
- Naveed, M., Moldrup, P., Arthur, E., Holmstrup, M., Nicolaisen, M., Tuller, M., Herath, L., Hamamoto, S., Kawamoto, K., Komatsui, T., Vogel, H.J., Wollensen de Jonge, L., 2014. Simultaneous loss of soil biodiversity and functions along a copper contamination gradient: when soil goes to sleep. *Soil Sci. Soc. Am. J. Abstr. Soil Phys.* 78, 1239–1250.
- Noyes, P.D., Lema, S.C., 2015. Forecasting the impacts of chemical pollution and climate change interactions on the health of wildlife. *Curr. Zool.* 61, 669–689.
- OECD, 2010. Guidelines for Testing Chemicals No. 317-Bioaccumulation in Terrestrial Oligochaetes. Organization for Economic Co-operation and Development, Paris, France.
- Párraga-Aguado, I., Álvarez-Rogel, J., González-Alcaraz, M.N., Jiménez-Cárceles, F.J., Conesa, H.M., 2013. Assessment of metal(loid)s availability and their uptake by *Pinus halepensis* in a Mediterranean forest impacted by abandoned tailings. *Ecol. Eng.* 58, 84–90.
- Peijnenburg, W.J.G.M., Baerselman, R., de Groot, A.C., Jager, T., Posthuma, L., van Veen, R.P., 1999. Relating environmental availability to bioavailability: soil type-dependent metal accumulation in the Oligochaete *Eisenia andrei*. *Ecotoxicol. Environ. Saf.* 44, 294–310.
- Peijnenburg, W.J.G.M., Zablotzka, M., Vijver, M.G., 2007. Monitoring metals in terrestrial environments within a bioavailability framework and a focus on soil extraction. *Ecotoxicol. Environ. Saf.* 67, 163–179.
- Pérez-Sirvent, C., Martínez-Sánchez, M.J., García-Lorenzo, M.L., Molina, J., Tudela, M.L., 2009. Geochemical background levels of zinc, cadmium and mercury in anthropically influenced soils located in a semi-arid zone (SE, Spain). *Geoderma* 148, 307–317.
- Rohr, J.R., Johnson, P., Hickey, C.W., Helm, R.C., Fritz, A., Brasfield, S., 2013. Implications of global climate change for natural resources damage assessment, restoration and rehabilitation. *Environ. Toxicol. Chem.* 32, 93–101.
- Römbke, J., Aira, M., Backeljau, T., Breugelmans, K., Domínguez, J., Funke, E., Graf, N., Hajibabaei, M., Pérez-Losada, M., Porto, P.G., Schmelz, R.M., Vierna, J., Vizcaíno, A., Pfenniger, M., 2016. DNA barcoding of earthworms (*Eisenia fetida/andrei* complex) from 28 ecotoxicological test laboratories. *Appl. Soil Ecol.* 104, 3–11.
- Smith, B.A., Egeler, P., Gilberg, D., Hendershot, W., Stephenson, G.L., 2010. Uptake and elimination of cadmium and zinc by *Eisenia andrei* during exposure to low concentrations in artificial soil. *Arch. Environ. Contam. Toxicol.* 59, 264–273.
- Sokal, R.R., Rohlf, F.J., 1969. *Biometry*. W.H. Freeman, San Francisco, USA.
- Spurgeon, D.J., Hopkin, S.P., 1999. Comparisons of metal accumulation and excretion kinetics in earthworm (*Eisenia fetida*) exposed to contaminated field and laboratory soils. *Appl. Soil Ecol.* 11, 227–243.
- Spurgeon, D.J., Weeks, J.M., van Gestel, C.A.M., 2003. A summary of eleven years progress in earthworm ecotoxicology: the 7th international symposium on earthworm ecology, Cardiff, Wales, 2002. *Pedobiologia* 47, 588–606.
- Stahl Jr., R.G., Hooper, M.J., Balbus, J.M., Clements, W., Fritz, A., Gouin, T., Helm, R., Hickey, C., Landis, W., Moe, J., 2013. The influence of global climate change on the scientific foundations and applications of environmental toxicology and chemistry: introduction to a SETAC international workshop. *Environ. Toxicol. Chem.* 32, 13–19.
- Stankovic, S., Kalaba, P., Stankovic, A.R., 2014. Biota as toxic metal indicators. *Environ. Chem. Lett.* 12, 63–84.
- Stürzenbaum, S.R., Georgiev, O., Morgan, A.J., Kille, P., 2004. Cadmium detoxification in earthworms: from genes to cells. *Environ. Sci. Technol.* 38, 6283–6289.
- Stürzenbaum, S.R., Winters, C., Galay, M., Morgan, A.J., Kille, P., 2001. Metal ion trafficking in earthworms - identification of cadmium specific metallothionein. *J. Biol. Chem.* 276, 34013–34018.
- Świątek, Z.M., van Gestel, C.A.M., Bednarska, A.J., 2017. Toxicokinetics of zinc-oxide nanoparticles and zinc ions in the earthworm *Eisenia andrei*. *Ecotoxicol. Environ. Saf.* 143, 151–158.
- van Gestel, C.A.M., Hensbergen, P.J., 1997. Interaction of Cd and Zn toxicity for *Folsomia candida* Willem (Collembola: Isotomidae) in relation to bioavailability in soil. *Environ. Toxicol. Chem.* 16, 1177–1186.
- van Gestel, C.A.M., Henzen, L., Dirven-van Breemen, E.M., Kamerman, J.W., 2002. Influence of soil remediation techniques on the bioavailability of heavy metals. In: Sunahara, G.I., Renoux, A.Y., Thellen, C., Gaudet, C.I., Pilon, A. (Eds.), *Environmental Analysis of Contaminated Sites*. John Wiley & Sons Ltd., Chichester, UK, pp. 361–388.
- van Straalen, N.M., 2004. The use of soil invertebrates in ecological surveys of contaminated soils. In: Doelman, P., Eijssackers, H.J.P. (Eds.), *Vital Soil – Function, Value and Properties*. Elsevier, The Netherlands, pp. 159–195.
- van Straalen, N.M., Donker, M.H., Vijver, M.G., van Gestel, C.A.M., 2005. Bioavailability of contaminants estimated from uptake rates into soil invertebrates. *Environ. Pollut.* 136, 409–417.
- Vijver, M.G., van Gestel, C.A.M., van Straalen, N.M., Lanno, R.P., Peijnenburg, W.J.G.M., 2006. Biological significance of metals partitioned to subcellular fractions within earthworms (*Aporrectodea caliginosa*). *Environ. Toxicol. Chem.* 25, 807–814.
- Vijver, M.G., Vink, J.P.M., Jager, T., Wolterbeek, H.Th., van Straalen, N.M., van Gestel, C.A.M., 2005. Biphasic elimination and uptake kinetics of Zn and Cd in the earthworm *Lumbricus rubellus* exposed to contaminated floodplain soil. *Soil Biol. Biochem.* 37, 1843–1851.
- Zhou, T., Li, L., Zhang, X., Zheng, J., Joseph, S., Pan, G., 2016. Changes in organic carbon and nitrogen in soil with metal pollution by Cd, Cu, Pb and Zn: a meta-analysis. *Eur. J. Soil Sci.* 67, 237–246.