VU Research Portal

An expansion formula for type A and Kronecker quantum cluster algebras

Çanakç, Ike; Lampe, Philipp

published in

Journal of Combinatorial Theory. Series A
2020

DOI (link to publisher)

[10.1016/j.jcta.2019.105132](https://doi.org/10.1016/j.jcta.2019.105132)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Çanakç, ., & Lampe, P. (2020). An expansion formula for type A and Kronecker quantum cluster algebras. *Journal of Combinatorial Theory. Series A*, 171, 1-30. [105132]. <https://doi.org/10.1016/j.jcta.2019.105132>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

ELSEVIER

Contents lists available at ScienceDirect

Journal of Combinatorial Theory,
Series A

www.elsevier.com/locate/jcta

An expansion formula for type A and Kronecker
quantum cluster algebras[☆] İlke Çanakçı^a, Philipp Lampe^b^a Department of Mathematics, Vrije Universiteit, Amsterdam 1081 HV,
the Netherlands^b School of Mathematics, Statistics and Actuarial Science (SMSAS), Canterbury
CT2 7FS, United Kingdom

A R T I C L E I N F O

Article history:

Received 5 November 2018

Accepted 27 August 2019

Available online 31 October 2019

Keywords:

Quantum cluster algebras

Non-commutative rings

Surface cluster algebras

Triangulations

Arcs

Snake graphs

Perfect matchings

Lattices

Stembridge phenomenon

Mathematical physics

A B S T R A C T

We introduce an expansion formula for elements in quantum cluster algebras associated to type A and Kronecker quivers with principal quantization. Our formula is parametrized by perfect matchings of snake graphs as in the classical case. In the Kronecker type, the coefficients are q -powers whose exponents are given by a weight function induced by the lattice of perfect matchings. As an application, we prove that a reflectional symmetry on the set of perfect matchings satisfies Stembridge's $q = -1$ phenomenon with respect to the weight function. Furthermore, we discuss a relation of our expansion formula to generating functions of BPS states.

© 2019 Elsevier Inc. All rights reserved.

Contents

1. Introduction	2
2. Background on (quantum) cluster algebras and snake graphs	4
3. An expansion formula for quantum cluster algebras of type A	12
4. An expansion formula for the Kronecker quantum cluster algebra	14

[☆] The first author was supported by EPSRC grant EP/P016014/1 and the second author was supported by EPSRC grants EP/N005457/1 and EP/M004333/1.

E-mail addresses: i.canakci@vu.nl (İ. Çanakçı), P.B.Lampe@kent.ac.uk (P. Lampe).

5. The Stembridge phenomenon	25
References	28

1. Introduction

Cluster algebras were introduced by Fomin–Zelevinsky [29–31] and by Berenstein–Fomin–Zelevinsky [7] in a series of four articles to give an algebraic framework for the study of total positivity and dual canonical bases in Lie theory. Cluster algebras occur naturally in many areas of mathematics such as representation theory, geometry, combinatorics and mathematical physics.

An important class of cluster algebras, namely *surface cluster algebras*, was introduced by Fomin–Shapiro–Thurston [27] using combinatorics of (oriented) surfaces with marked points. Subsequently, Fomin–Thurston [28] gave an intrinsic formulation of surface cluster algebras, building on earlier works of Gekhtman–Shapiro–Vainshtein [34], Fock–Goncharov [25,26] and Penner [52], considering hyperbolic structures on the surface. Surface cluster algebras are also important from a classification point of view since all but finitely many cluster algebras of finite mutation type are those associated to marked surfaces or of rank 2 by Felikson–Shapiro–Tumarkin [23,24].

Quantum deformations of cluster algebras were introduced by Berenstein–Zelevinsky [9] to develop a setting for a general notion of canonical bases. Their construction also builds on Fock–Goncharov [25,26] but is given in a more systematic way and, in particular, they show most structural results of cluster algebras in the quantum setting.

Combinatorial, geometric and representation theoretic aspects of surface cluster algebras have been studied significantly in the classical case. An expansion formula for elements in surface cluster algebras was given by [47] extending on [64,62,46] and further combinatorial aspects of surface cluster algebras were explored in [14,15,17,16,13]. For some results on representation theoretic aspects of surface cluster algebras, see [6,10,18,59,69]. In the quantum setting most of research goes into understanding cluster variables and a ‘good’ basis in representation theoretic terms (see for instance [40,58,57,56,55,60]), into identifying quantum cluster algebra structures in Lie theory (see [33,35–37]) or into the so-called positivity conjecture which is shown to be true in [21] for all quantum cluster algebras. Representation theoretic expressions for quantum cluster variables were given by [60,61]; combinatorial expressions for quantum cluster variables were given by [41,42]; geometric descriptions for quantum cluster variables were given by [44]. Also see [8] for the non-commutative version of cluster algebras.

In the quantum setting, there is not much known solely for surface type quantum cluster algebras. The aim of this article is to take this direction and extend the expansion formula to give explicit Laurent polynomial expressions for quantum cluster variables in the quantum cluster algebra of type A_n and for quantum cluster variables and certain other elements in the quantum cluster algebra of Kronecker type considered with principal quantization. More precisely, for surface type cluster algebras and for their quantum

analogues, cluster variables (and quantum cluster variables) are in bijection with (isotopy classes) of arcs in the surface. Furthermore, there is a planar graph, called *snake graph*, associated to every arc in the surface. The explicit formula of [47] for elements in the cluster algebra is parameterized by perfect matchings of snake graphs. In this article we show that they can also be used to give Laurent polynomial expansions for elements in the quantum cluster algebra.

To state our main result, we introduce some notation. The quantum deformation of a cluster algebra is obtained by making each cluster into a *quasi-commuting* family of variables. Then the quantum cluster algebra $\mathcal{A}_q$ is a $\mathbb{Z}[q^{\pm 1/2}]$ -subalgebra of the skew-field of fractions of the *based quantum torus* (see Section 2.3). The based quantum torus admits a $\mathbb{Z}[q^{\pm 1/2}]$ -basis indexed by $\mathbb{Z}^m$, denoted $\{M[a] \mid a \in \mathbb{Z}^m\}$.

Let $\mathcal{A}_q$ be the quantum cluster algebra with principal quantization associated with an adjacency quiver Q of a triangulation of a surface $\mathcal{S}$ which is either a disc with (finitely) many marked points on the boundary or an annulus with exactly one marked point in each boundary component.

Theorem A (Theorem 3.1, Theorem 4.20). *With the notation above, let γ be a (generalized) arc in $\mathcal{S}$, x_γ be the corresponding element in $\mathcal{A}_q$ and $\mathcal{G}$ be the snake graph associated with γ . Then*

$$x_\gamma = \sum_{P \models \mathcal{G}} q^{\Omega(P)} M[\nu(P)]$$

where $\nu(P) \in \mathbb{Z}^m$ is a vector associated with a perfect matching P of $\mathcal{G}$ and where $\Omega(P) = 0$ if $\mathcal{S}$ is a disc and $\Omega(P)$ is a half-integer given explicitly by the position of the perfect matching in the perfect matching lattice if $\mathcal{S}$ is the annulus with exactly one marked point in each of the boundary components.

We note that our formula coincides with the classical case [47,54] in the limit $q \rightarrow 1$. For type A, the proof of the theorem uses work of Tran [65] on quantum F -polynomials.

We then get a curious application of Theorem A by comparing our expansion formula for self-crossing arcs in the Kronecker type with generating functions of BPS states. Relations between BPS states and cluster theory have been studied for example by [1,2,4,19,20,32,66]. Córdova–Neitzke [20] consider a supersymmetric quantum field theory of quiver type whose line defects W_n are parametrized by natural numbers n and compute the generating function $F(W_n)$ of BPS states recursively using a Coulomb branch formula for small n .

Question A (Question 4.23). Is the following statement true? For $n \in \mathbb{N}$, the generating function $F(W_n)$ coincides with the element associated to a self-crossing arc in the quantum cluster algebra of Kronecker type.

We answer the question affirmatively for $n \in \{0, 1, 2, 3, 4\}$, see Proposition 4.22.

In an independent work, Huang [39] provides an expansion formula for quantum cluster algebras with *boundary* coefficients generalizing [47]; here the q -powers of the terms are given by recursions in the perfect matching lattice.

If we plug in $q = 1$ in Theorem A, we obtain the classical expansion formula. The substitution $q = -1$ is related to Stembridge's $q = -1$ phenomenon. The snake graph $\mathcal{G}$ associated with a (non-crossing) arc in the annulus with exactly one marked point on each boundary component admits an axis of symmetry. Reflection across this axis induces an involution σ on the set of perfect matchings $\text{Match}(\mathcal{G})$. Stembridge's $q = -1$ phenomenon [64] asserts that the number of fixed points under this involution is equal to the evaluation of coefficients at $q = -1$. We define a map w as a slight variation of the map Ω . We then get the following result.

Theorem B (Corollary 5.6). *Let $p, r \in \mathbb{N}$. There exists $X(q) \in \mathbb{Z}[q]$ such that the following conditions hold.*

- (1) *Up to a power of $q^{\pm 1/2}$, the polynomial $X(q)$ is equal to $\sum_P q^{\Omega(P)}$ where the sum runs over all P in the graded component $\text{Match}(\mathcal{G})_{p,r}$ of the perfect matching lattice.*
- (2) *The value $X(-1)$ is equal to the number of fixed points of the restriction of σ to $\text{Match}(\mathcal{G})_{p,r}$.*

In other words, the quadruple $(\text{Match}(\mathcal{G})_{p,r}, \sigma, X(q), w)$ satisfies Stembridge's $q = -1$ phenomenon.

The paper is organized as follows. Section 2 is devoted to background on surface cluster algebras, snake graphs and general introduction to quantum cluster algebras. Section 3 concerns with the expansion formula for quantum cluster algebras of type A. Section 4 gives the formula for the Kronecker type and links it to BPS states. Finally in Section 5, we establish a connection to Stembridge's $q = -1$ phenomenon.

Acknowledgments: We would like to thank Hugh Thomas for stimulating discussions and for bringing Córdova–Neitzke's paper to our attention. We would like to thank Dylan Allegretti for insightful comments on a previous version of the manuscript.

2. Background on (quantum) cluster algebras and snake graphs

2.1. Cluster algebras

Let us recall some basic features of cluster algebras. For more details we refer the reader to the literature.

Let $m \geq n$ be natural numbers. In this article we use the notations $[r] = \{k \mid 1 \leq k \leq r\}$ and $[r, s] = \{k \mid r \leq k \leq s\}$ for $r, s \in \mathbb{N}$. Moreover, we denote by I the $n \times n$ identity matrix and by 0 the $n \times n$ zero matrix.

An integer $m \times n$ matrix B is called an *exchange matrix* if the submatrix of B on rows and columns $[n]$ is skew-symmetrizable. We denote by $\text{Ex}(m, n)$ the set of all $m \times n$ exchange matrices. A central notion in cluster theory is the mutation $\mu_k: \text{Ex}(m, n) \rightarrow \text{Ex}(m, n)$ for $k \in [n]$, defined by $B = (b_{ij}) \mapsto \mu_k(B)$ with

$$\mu_k(B)_{i,j} = \begin{cases} -b_{ij}, & \text{if } i = k \text{ or } j = k; \\ b_{ij} + (b_{ik}|b_{kj}| + |b_{ik}|b_{kj})/2, & \text{otherwise.} \end{cases}$$

A *cluster* is a tuple $(\mathbf{x}, \mathbf{y}) = (x_1, \dots, x_n, y_1, \dots, y_{m-n})$ of m algebraically independent variables over $\mathbb{Q}$. The elements x_i with $i \in [n]$ are called *cluster variables* and the elements y_i with $i \in [1, m-n]$ are called *frozen variables*. A *seed* is a triple $(B, \mathbf{x}, \mathbf{y})$ formed by a cluster and an exchange matrix. We extend the notion of mutation to seeds in the following way. The *mutation* of a seed $(B, \mathbf{x}, \mathbf{y})$ at $k \in [n]$ is the seed $(\mu_k(B), \mathbf{x}', \mathbf{y})$ where $\mathbf{x}'$ is obtained from $\mathbf{x}$ by replacing x_k with

$$x'_k = \frac{1}{x_k} \left(\prod_{\substack{1 \leq i \leq n \\ b_{ik} > 0}} x_i^{b_{ik}} \prod_{\substack{n+1 \leq i \leq m \\ b_{ik} > 0}} y_{i-n}^{b_{ik}} + \prod_{\substack{1 \leq i \leq n \\ b_{ik} < 0}} x_i^{-b_{ik}} \prod_{\substack{n+1 \leq i \leq m \\ b_{ik} < 0}} y_{i-n}^{-b_{ik}} \right).$$

The *cluster algebra* $\mathcal{A}(B, \mathbf{x}, \mathbf{y})$ is the subalgebra of $\mathbb{Q}(\mathbf{x}, \mathbf{y})$ generated by all cluster variables in all seeds obtained from $(B, \mathbf{x}, \mathbf{y})$ by sequences of mutations.

Definition 2.1 (*Principal coefficients and C-matrices*).

(a) Let $\tilde{B}$ be an $n \times n$ exchange matrix. The $(2n) \times n$ block matrix

$$B = \begin{pmatrix} \tilde{B} \\ I \end{pmatrix}$$

is said to be the extension of $\tilde{B}$ with *principal coefficients*.

(b) Suppose we mutate B along a sequence $\mathbf{i} = (i_r, \dots, i_2, i_1)$ of mutable indices to obtain a matrix

$$\mu_{\mathbf{i}}(B) = B_{\mathbf{i}} = \begin{pmatrix} \tilde{B}_{\mathbf{i}} \\ C_{\mathbf{i}} \end{pmatrix}.$$

Then we call $C_{\mathbf{i}}$ the *C-matrix* (or *coefficient matrix*) of $B_{\mathbf{i}}$. The column vectors of $C_{\mathbf{i}}$ are called *c-vectors*.

Let $\tilde{B}$ be an $n \times n$ skew-symmetrizable matrix and let B be its extension with principal coefficients as above. We choose an initial cluster with cluster variables $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and frozen variables $\mathbf{y} = (y_1, y_2, \dots, y_n)$.

Suppose we mutate $(B, \mathbf{x}, \mathbf{y})$ along a sequence $\mathbf{i} = (i_r, \dots, i_2, i_1)$ of mutable indices to obtain a seed $(B_{\mathbf{i}}, \mathbf{x}_{\mathbf{i}}, \mathbf{y}_{\mathbf{i}})$. Let $x' \in \mathbf{x}_{\mathbf{i}}$. Then we denote the c -vector corresponding to the column of C with the same label as x' by $c(x')$.

Fomin–Zelevinsky [31, Proposition 6.1] prove that the assignments $x_i \mapsto e_i$ and $y_i \mapsto -b_i$, where e_i is the i -th standard basis vector and b_i is the i -th column vector of $\tilde{B}$, define a $\mathbb{Z}^n$ -grading of the Laurent polynomial ring $\mathbb{Z}[x_i^{\pm 1}, y_i^{\pm 1} \mid i \in [n]]$ such that each cluster variable $x' \in \mathcal{A}(B, \mathbf{x}, \mathbf{y})$ is homogeneous.

Definition 2.2 (*G-matrices*). The g -vector $g(x')$ of a cluster variable $x' \in \mathcal{A}(B, \mathbf{x}, \mathbf{y})$ is given by the degree of x' with respect to the $\mathbb{Z}^n$ -grading. The G -matrix of a cluster $\mathbf{x}' = (x'_1, x'_2, \dots, x'_n)$ is defined to be the matrix with columns $g(x'_1), g(x'_2), \dots, g(x'_n)$.

Suppose we mutate $(B, \mathbf{x}, \mathbf{y})$ along a sequence $\mathbf{i} = (i_r, \dots, i_2, i_1)$ of mutable indices to obtain a seed $(B_{\mathbf{i}}, \mathbf{x}_{\mathbf{i}}, \mathbf{y}_{\mathbf{i}})$. Nakanishi [50, Theorem 4.1] and Nakanishi–Zelevinsky [51, Theorem 1.2] prove that the C -matrix of $B_{\mathbf{i}}$ and the G -matrix of $\mathbf{x}_{\mathbf{i}}$ possess the tropical duality $C_{\mathbf{i}} G_{\mathbf{i}}^T = I$.

2.2. Surface cluster algebras and snake graphs

The focus of this article is on cluster algebras of type A_n and of Kronecker type each of which can be realized as a cluster algebra of surface type [27, 28]. We will recall surface cluster algebras in this section following the exposition in [14, 47] in order to give a self-contained exposition of the paper.

Let $\mathcal{S}$ be a connected oriented 2-dimensional Riemann surface (with possibly empty) boundary and $\mathcal{M}$ be a finite set of marked points. We further assume that $\mathcal{M}$ is contained in the boundary of $\mathcal{S}$ (though this is not necessary for the general construction) and that each boundary component contains at least one marked point. If $\mathcal{S}$ is a disc, we let $\mathcal{M}$ to have at least 4 marked points. We call a pair $(\mathcal{S}, \mathcal{M})$ *bordered surface with marked points*.

Definition 2.3 (*Arcs, compatible arcs, triangulation*). Let $\mathcal{S}$ be a bordered surface with marked points. An *arc* of $\mathcal{S}$ is a non-self-intersecting curve, considered up to isotopy, with endpoints at marked points of $\mathcal{S}$. We further assume that an arc is disjoint from the boundary of $\mathcal{S}$ except the endpoints, which might coincide, and it does not cut out a monogon or digon. Two arcs are *compatible* if they do not intersect in the interior of $\mathcal{S}$, that is, there are representatives in the corresponding isotopy classes that do not intersect except the endpoints may coincide. A *triangulation* $\mathcal{T}$ is a maximal collection of pairwise distinct compatible arcs. The connected components in the complement $\mathcal{S} \setminus \mathcal{T}$ are called *triangles*.

Let $\mathcal{S}$ be a bordered surface with marked points and let $\mathcal{T} = \{\tau_1, \dots, \tau_n\}$ be a triangulation of $\mathcal{S}$. Every triangle Δ is bounded by three arcs τ_r, τ_s, τ_t . Without loss of

Fig. 1. The domain of an arc (left) and the construction of its snake graph (right).

generality we may assume that, when going through the boundary of Δ in positive direction, we pass the arcs in order $\tau_r \rightarrow \tau_s \rightarrow \tau_t \rightarrow \tau_r$. To every triangle Δ we associate the $n \times n$ matrix $B^\Delta = (b_{ij}^\Delta)$ with entries $b_{rs} = b_{st} = b_{tr} = 1$, $b_{sr} = b_{ts} = b_{rt} = -1$, and $b_{ij} = 0$ otherwise.

Definition 2.4 (*Surface cluster algebras*). Let $\mathcal{S}$ be a bordered surface with marked points and $\mathcal{T} = \{\tau_1, \dots, \tau_n\}$ be a triangulation of $\mathcal{S}$.

- For each $\tau_i \in \mathcal{T}$ with $i \in [n]$ we associate formal variables x_i and y_i . Then $(\mathbf{x}, \mathbf{y}) = (x_1, \dots, x_n, y_1, \dots, y_n)$ is an initial cluster.
- The *signed adjacency matrix* of $\mathcal{T}$ is $\tilde{B} = \sum_{\Delta} B^\Delta$ where the sum runs over all triangles. The *adjacency quiver* $Q_{\mathcal{T}}$ of $(\mathcal{S}, \mathcal{T})$ is a quiver with vertex set $[n]$ such that the number of arrows $i \rightarrow j$ is equal to b_{ij} if $b_{ij} > 0$ and zero otherwise.
- The *surface cluster algebra* $\mathcal{A}(\mathcal{S}, \mathcal{T})$ with principal coefficients is the cluster algebra generated by the seed $(B, \mathbf{x}, \mathbf{y})$ where B is the principal extension of $\tilde{B}$.

2.2.1. Snake graphs

In this section, we recall the construction of snake graphs of [47]. We fix a surface $\mathcal{S}$ and a triangulation $\mathcal{T} = \{\tau_1, \tau_2, \dots, \tau_n\}$.

Let γ be a (generalized) arc in $(\mathcal{S}, \mathcal{T})$. We consider the domain $\text{dom}(\gamma)$ of γ in $(\mathcal{S}, \mathcal{T})$, that is, $\text{dom}(\gamma)$ is the union of all the triangles γ crosses (counted with multiplicity), see Fig. 1.

Fix an orientation of γ . For each crossing of γ with $\mathcal{T}$, we will associate a weighted tile, i.e. a square of fixed side length drawn in the plane with sides aligned horizontally and vertically, as follows:

- if γ crosses an arc τ_i , the associated weighted tile G_i has face weight $w(G_i) = \tau_i$ and edge weights induced by the quadrilateral Q_i in $\mathcal{T}$ which contains τ_i as diagonal, see Fig. 1;
- glue successive tiles $G_{i_1}, G_{i_2}, \dots, G_{i_k}$ as follows: set the diagonals τ_k and τ_{k+1} from the top-left corner to the bottom-right corner of G_k and G_{k+1} , respectively and glue the tiles G_k and G_{k+1} along the edge with weight induced from the third side of the triangle bounded by τ_k and τ_{k+1} in $\mathcal{T}$, see Fig. 1.

Remark 2.5. The snake graph $\mathcal{G}_\gamma$ associated to an arc γ in a triangulated surface is simply an unfolding of the domain of γ where zig-zag pieces in the triangulation correspond to straight subgraphs in G_γ and fans in the triangulation correspond to corner subgraphs in G_γ . The edge and face weights on $\mathcal{G}_\gamma$ are induced by the triangulation $\mathcal{T}$.

Definition 2.6. A *perfect matching* of a graph $\mathcal{G}$ is a set of edges of $\mathcal{G}$ such that each vertex v in $\mathcal{G}$ is incident to precisely one edge e in P . We denote by $\text{Match}(\mathcal{G})$ the set of perfect matchings of the graph $\mathcal{G}$, and for brevity we write $P \models \mathcal{G}$ if P is a perfect matching of $\mathcal{G}$.

The set $\text{Match}(\mathcal{G}_\gamma)$ of a snake graph $\mathcal{G}_\gamma$ admits a rich combinatorial structure which we will discuss next.

Definition 2.7 (*Perfect matching lattice*). The *perfect matching graph* $L(\mathcal{G})$ of a snake graph $\mathcal{G}$ is defined as follows: its vertices are the perfect matchings of $\mathcal{G}$, and two vertices are connected by an edge if the perfect matchings are obtained from each other by a single twist.

By [47], the perfect matching graph admits a lattice structure with maximal and minimal perfect matchings given by those containing only boundary edges of $\mathcal{G}$. We refer to [47] for a precise definition of the minimum and maximum perfect matchings and explain the lattice later in more detail.

Example 2.8. The minimal and maximal perfect matchings of the snake graph are and and its perfect matching lattice is given in Fig. 2.

The twist admits a nice combinatorial interpretation. For every perfect matching P of a snake graph $\mathcal{G}$, we consider the symmetric difference $\text{Sym}(P) = (P \cup P_{\min}) \setminus (P \cap P_{\min})$. The set $\text{Sym}(P)$ defines a union of cycles in the planar realization of $\mathcal{G}$. We say that a tile G of $\mathcal{G}$ is *twisted* if it lies in the interior of one of these cycles. The set of twisted tiles of P is denoted by $\text{Twist}(P)$. Furthermore, for an arc τ we define $\text{Twist}(P)_\tau$ to be the subset of $\text{Twist}(P)$ consisting of the tiles G with face weight τ . The name is justified by a theorem of Musiker–Schiffler–Williams according to which the set $\text{Twist}(P)$ is equal to the set of tiles twisted along any shortest path from $P_{\min}$ to P in $L(\mathcal{G})$.

Musiker–Schiffler–Williams [48, Theorem 5.2] show that the graph $L(\mathcal{G})$ is the Hasse diagram of the poset $\text{Match}(\mathcal{G}, \leq)$ when we define $P_1 \leq P_2$ if and only if $\text{Twist}(P_1) \subseteq \text{Twist}(P_2)$. In fact, it is a distributive lattice with minimal element $P_{\min}$ and maximal element $P_{\max}$. Their theorem is a consequence of work by Propp [54, Theorem 2].

Fig. 2. The lattice of perfect matchings of a snake graph.

2.2.2. Laurent polynomials associated to snake graphs

Let $\mathcal{S}$ be a surface with triangulation $\mathcal{T} = \{\tau_1, \tau_2, \dots, \tau_n\}$. Let γ be an arc in $\mathcal{S}$ and $\mathcal{G}$ its associated snake graph. Let P be a perfect matching of $\mathcal{G}$. Then

- we assign formal variables x_{τ_i}, y_{τ_i} to each arc τ_i in $\mathcal{T}$, and often use the abbreviations $x_i = x_{\tau_i}$ and $y_i = y_{\tau_i}$;
- the *weight monomial* $x(P)$ of P is given by $x(P) = \prod_{e \in P} x_{w(e)}$;
- the *height monomial* $y(P)$ of P is given by $y(P) = \prod_{G \in \text{Twist}(P)} y_{w(G)}$;
- the *crossing monomial* of γ with respect to $\mathcal{T}$ is given by $\text{cross}(\gamma, \mathcal{T}) = \prod_{j \in J} x_j$ where J is the index set of the arcs in $\mathcal{S}$ that γ crosses;
- the *Laurent polynomial* associated to $\mathcal{G}$ with respect to $\mathcal{T}$ is defined as

$$x_{\mathcal{G}} = \frac{1}{\text{cross}(\gamma, \mathcal{T})} \sum_{P \models \mathcal{G}} x(P)y(P).$$

Theorem 2.9 ([47]). Let $\mathcal{S}$ be a surface with triangulation $\mathcal{T}$ and let $\mathcal{A}(\mathcal{S})$ be the cluster algebra associated to $\mathcal{S}$. Let γ be a (generalized) arc of $\mathcal{S}$ and $\mathcal{G}$ be its associated snake graph with respect to $\mathcal{T}$ and $x_{\mathcal{G}}$ be the Laurent polynomial associated to γ . Then $x_{\mathcal{G}} = x_{\gamma}$.

Remark 2.10. Let $x_{\gamma} \in \mathcal{A}(\mathcal{S}, \mathcal{T})$ be the cluster variable associated to an arc γ . The weight monomial from Section 2.2.2 is related to the g -vector of x_{γ} with respect to the initial

seed associated with $\mathcal{T}$, see [48, Proposition 6.2]; namely

$$g(x_\gamma) = \deg(x(P_{\min})) - \deg(\text{cross}(\mathcal{T}, \gamma)) \in \mathbb{Z}^n. \quad (1)$$

In the above situation, for every perfect matching P of $\mathcal{G}_\gamma$ we put

$$\nu(P) = (\deg(x(P)) - \deg(\text{cross}(\mathcal{T}, \gamma)), y(P)) \in \mathbb{Z}^{2n}.$$

2.3. Quantum cluster algebras

In this section we give a brief introduction to quantum cluster algebras. The section follows Berenstein–Zelevinsky [9]. Let $q^{1/2}$ be an indeterminate and let $q^{-1/2}$ be its formal inverse.

Definition 2.11 (*Principal quantization*). Let $\tilde{B}$ be an $n \times n$ skew-symmetric matrix. We consider the extension with principal coefficients B of size $(2n) \times n$ and a $(2n) \times (2n)$ matrix Λ as follows:

$$B = \begin{pmatrix} \tilde{B} \\ I \end{pmatrix}, \quad \Lambda = \begin{pmatrix} 0 & -I \\ I & -\tilde{B} \end{pmatrix}.$$

We call the pair (B, Λ) the *principal quantization pair* of $\tilde{B}$ and call Λ the *principal quasi-commutation matrix* of $\tilde{B}$.

Zelevinsky [67, Example 0.5] proves that the principal quantization pair (B, Λ) attached to a skew-symmetric $n \times n$ matrix $\tilde{B}$ is always *compatible*, that is, the matrices obey the relation $B^T \Lambda = (I \ 0)$.

Fomin–Zelevinsky’s *sign coherence conjecture* asserts that the entries in a c -vector are either all non-negative or all non-positive. Fomin–Zelevinsky [31, Proposition 5.6] prove that the sign coherence conjecture is equivalent to the *constant term conjecture* for F -polynomials. This conjecture holds true by a result of Derksen–Weyman–Zelevinsky [22, Theorem 1.7]. Later different proofs of sign-coherence were given by Nagao [49, Theorem 9.9], Plamondon [53, Theorem 3.13] and Gross–Hacking–Keel–Kontsevich [38, Corollary 5.5]. We use sign-coherence in the proof of the following proposition.

Proposition 2.12. *Let $\tilde{B}$ be an $n \times n$ skew-symmetric matrix. If we mutate the principal quantization pair (B, Λ) along a sequence $\mathbf{i} = (i_r, \dots, i_2, i_1)$ of mutable indices, then*

$$\mu_{\mathbf{i}}(B, \Lambda) = (B_{\mathbf{i}}, \Lambda_{\mathbf{i}}) = \left(\begin{pmatrix} \tilde{B}_{\mathbf{i}} \\ C_{\mathbf{i}} \end{pmatrix}, \begin{pmatrix} 0 & -G_{\mathbf{i}}^T \\ G_{\mathbf{i}} & G_{\mathbf{i}} \tilde{B}_{\mathbf{i}}^T G_{\mathbf{i}}^T \end{pmatrix} \right).$$

Proof. By construction $B_{\mathbf{i}} = \mu_{\mathbf{i}}(B)$ is obtained from B by ordinary matrix multiplication, hence the form of $B_{\mathbf{i}}$ follows from Definition 2.1. We can show by induction that

the upper left part of $\Lambda_{\mathbf{i}}$ is zero. Let $\mathbf{i}' = (i_{r-1}, \dots, i_1)$. By definition $\Lambda_{\mathbf{i}} = E_{\varepsilon}^T \Lambda_{\mathbf{i}'} E_{\varepsilon}$, where we use the same notation as Berenstein–Zelevinsky [9, Equation (3.2)]. The sign coherence of the c vectors implies there is always a sign ε such that E_{ε} is block diagonal, which proves that the upper left part of $\Lambda_{\mathbf{i}}$ is zero. Moreover, the pair $(B_{\mathbf{i}}, \Lambda_{\mathbf{i}})$ must satisfy the compatibility condition $B_{\mathbf{i}}^T \Lambda_{\mathbf{i}} = (I \ 0)$ so that the lower left part of $\Lambda_{\mathbf{i}}$ is equal to $G_{\mathbf{i}}$ due to the relation $C_{\mathbf{i}} G_{\mathbf{i}}^T = I$. The upper right part of $\Lambda_{\mathbf{i}}$ must be equal to $-G_{\mathbf{i}}^T$ due to skew-symmetry. The structure of the lower right part can be read off from the compatibility condition. $\square$

Suppose that Λ is a skew-symmetric integer $m \times m$ matrix. The *based quantum torus* $T(\Lambda)$ is a $\mathbb{Z}[q^{\pm 1/2}]$ -algebra with $\mathbb{Z}[q^{\pm 1/2}]$ -basis $\{M[a] \mid \mathbb{Z}^m\}$ indexed by $\mathbb{Z}^m$; on basis elements the multiplication is defined as

$$M[a]M[b] = q^{\frac{1}{2}a^T \Lambda b} M[a+b] \quad (2)$$

for all $a, b \in \mathbb{Z}^m$.

The based quantum torus $T(\Lambda)$ is an Ore domain and it is contained in its skew field of fractions $\mathcal{F}$. We refer the reader to [9] for details about this construction. Note that $\mathcal{F}$ is an algebra over $\mathbb{Q}(q^{1/2})$. We say that two elements $f_1, f_2 \in \mathcal{F}$ are *q-commuting* if there exists an integer k such that $f_i f_j = q^{k/2} f_j f_i$. Examples of *q-commuting* elements include the basis elements of the based quantum torus: for all $a, b \in \mathbb{Z}^m$ we have $M[a]M[b] = q^{a^T \Lambda b} M[b]M[a]$.

A *quantum cluster* is a tuple $(\mathbf{x}', \mathbf{y}') = (x'_1, \dots, x'_n, y'_1, \dots, y'_{m-n})$ of m pairwise *q-commuting* elements in $\mathcal{F}$. The elements x'_i with $i \in [n]$ are called *quantum cluster variables* and the elements y'_i with $i \in [1, m-n]$ are called *frozen variables*. The quantum cluster $(\mathbf{x}, \mathbf{y}) = (M[e_1], \dots, M[e_m])$, where e_i denotes the i -th standard basis vector of $\mathbb{Z}^m$, is called the *initial quantum cluster*. The exponents arising in the *q-commutativity* relations among the quantum cluster and frozen variables in a quantum cluster $(\mathbf{x}', \mathbf{y}')$ assemble a skew-symmetric matrix Λ' . We call Λ' the *q-commutativity matrix* of the quantum cluster. We denote the basis elements in the based quantum torus $T(\Lambda')$ by $M_{\Lambda'}[a]$ with $a \in \mathbb{Z}^m$.

A *quantum seed* is a tuple $(B', \Lambda', \mathbf{x}', \mathbf{y}')$ such that $(\mathbf{x}', \mathbf{y}')$ is a quantum cluster with *q-commutativity* matrix Λ' and there exists a positive integer k such that $(B')^T \Lambda = (kI \ 0)$. The last property is called the *compatibility condition*.

The *mutation* $\mu_k(B', \Lambda', \mathbf{x}', \mathbf{y}') = (B'', \Lambda'', \mathbf{x}'', \mathbf{y}'')$ of the quantum seed $(B', \Lambda', \mathbf{x}', \mathbf{y}')$ at $k \in [n]$ is constructed as follows. The quantum cluster $(\mathbf{x}'', \mathbf{y}'')$ is obtained from $(\mathbf{x}', \mathbf{y}')$ by replacing x'_k with

$$x''_k = M_{\Lambda'} \left(-e_k + \sum_{b_{ki} > 0} b_{ki} e_i \right) + M_{\Lambda'} \left(-e_k - \sum_{b_{ik} < 0} b_{ik} e_i \right).$$

This cluster consists again of pairwise q -commuting elements by the compatibility condition. Then Λ'' is defined as the q -commutativity matrix of $(\mathbf{x}'', \mathbf{y}'')$. Lastly, $B'' = \mu_k(B')$ is given by the usual mutation of exchange matrices.

The *quantum cluster algebra* $\mathcal{A}_q(B, \Lambda, \mathbf{x}, \mathbf{y})$ is the subalgebra of $\mathcal{F}$ generated by all quantum cluster variables and frozen variables in all quantum seeds that can be obtained from $(B, \Lambda, \mathbf{x}, \mathbf{y})$ by sequences of mutations. We also use the shorthand notation $\mathcal{A}_q(B, \Lambda)$ for $\mathcal{A}_q(B, \Lambda, \mathbf{x}, \mathbf{y})$.

3. An expansion formula for quantum cluster algebras of type A

3.1. A quantum expansion formula for type A

Let $n \geq 1$ be a natural number. We consider the marked oriented surface $(\mathcal{S}, M)$ which is formed by a disc $\mathcal{S}$ with a set M of $n + 3$ marked points on the boundary. The cluster algebra $\mathcal{A}(\mathcal{S}, M)$ has finite type A_n . We fix a triangulation $\mathcal{T}$ of $(\mathcal{S}, M)$, and we denote by Q the quiver of $\mathcal{T}$. Without loss of generality we may assume that Q is a quiver of type A_n .

Let (B, Λ) be the principal quantization pair of the signed adjacency matrix of Q . We are interested in the quantum cluster algebra $\mathcal{A}_q(B, \Lambda)$. We denote the initial quantum cluster by $(\mathbf{x}, \mathbf{y}) = (x_i, y_i)_{i \in [1, n]}$.

Our goal is to prove the following theorem about the expansion of quantum cluster variables as $\mathbb{Z}[q^{\pm 1/2}]$ -linear combinations of quantum Laurent monomials in the initial quantum cluster variables. As it turns out, the coefficients are all equal to 1.

Theorem 3.1. *If γ is any arc in $(\mathcal{S}, M)$, then*

$$x_\gamma = \sum_{P \models \mathcal{G}_\gamma} M[\nu(P)] \in \mathcal{A}_q(B, \Lambda).$$

The proof of Theorem 3.1 will be given at the end of Section 3.

3.2. Quantum F -polynomials

The main tool in the proof of the expansion formula is quantum F -polynomials. These polynomials are a quantized version of the F -polynomials from Fomin–Zelevinsky’s fourth cluster algebra article [31]. Quantum F -polynomials have been introduced and studied by Tran [65]. Let us recall some notions and results from Tran’s article.

Definition 3.2 (*$\widehat{y}$ -variables*). Suppose $i \in [1, n]$. Let b_i be the i -th column of the exchange matrix B . We put $\widehat{y}_i = M[b_i]$. Moreover, for every subset $S = \{s_1, \dots, s_r\} \subseteq [1, n]$ with $s_1 < \dots < s_r$ we set

$$\widehat{y}_S = q^{\frac{1}{2} \sum_{1 \leq i < j \leq r} b_{s_j, s_i}} \widehat{y}_{s_1} \cdots \widehat{y}_{s_r}.$$

Fig. 3. A successor-closed subset and the corresponding perfect matching.

For two indices $i, j \in [1, n]$ we have $b_i^T \Lambda b_j = -b_{ji}$ so that

$$\widehat{y}_i \widehat{y}_j = M[b_i] M[b_j] = q^{\frac{1}{2} b_i^T \Lambda b_j} M[b_i + b_j] = q^{-\frac{1}{2} b_{ji}} M[b_i + b_j] = q^{-b_{ji}} \widehat{y}_j \widehat{y}_i.$$

We conclude that $\widehat{y}_S$ is a standard basis element in Berenstein–Zelevinsky’s quantum torus. More precisely, if we denote by $b_S = \sum_{s \in S} b_s$ the sum of the column vectors of B indexed by elements of S , then $\widehat{y}_S = M[b_S]$.

We consider an arc γ and the associated snake graph $\mathcal{G}_\gamma$. The weights of the tiles of $\mathcal{G}_\gamma$ are given by a subset $T \subseteq Q_0$ such that the full subquiver of Q on the vertex set T is connected.

Definition 3.3 (*Properties of subsets*). Suppose that $S \subseteq T$.

- (1) We say that S is *successor-closed* if the following condition holds. If $j \in S$, $i \in T$ and there exists an arrow $j \rightarrow i$ in Q , then $i \in S$.
- (2) We denote by $\Phi(S)$ the number of connected components of the full subquiver of Q on the vertex set S .

Note that successor-closed subsets of T are in natural bijection with perfect matchings of $\mathcal{G}_\gamma$. More precisely, for every successor-closed subset $S \subseteq T$ there is a unique perfect matching P such that the weights of the twisted tiles of P are equal to S .

Example 3.4. Suppose that $T = \{1, 2, 3, 4, 5\}$ and that the full subquiver of Q on the vertex set T is given by the quiver on the left of Fig. 3. Then the set $S = \{1, 3, 4\}$ is successor-closed. In the middle of the figure we display the snake graph and its minimal perfect matching. Twisting the tiles labelled 1, 3, 4 yields the perfect matching on the right of the figure.

In this example we have $|S| = 3$ and $\Phi(S) = 2$. Note that the number of arrows inside S is equal to 1, which is equal to the difference of the previous two numbers.

We can generalize the observation in the following way.

Proposition 3.5. Suppose that $S \subseteq T$ is successor-closed. Then

$$|S| - |\{(k, j) \in S \times T \mid (k, j) \in Q_1\}| = \Phi(S).$$

Proof. The full subquiver of Q on the vertex set S is a (directed) forest, i.e. a disjoint union of (directed) trees. In every connected component the difference between the number of vertices and the number of arrows is equal to 1. $\square$

Recall that g_γ denotes the g -vector of the cluster variable x_γ . Tran, see [65, Proposition 7.3], shows that the entry of the g -vector at an index $s \in Q_0$ is equal to

$$(g_\gamma)_s = |\{j \in T \mid \exists(s \rightarrow j) \in Q_1\}| - 1. \quad (3)$$

In the next statement we view the g -vector as an element in $\mathbb{Z}^{2n}$ via the canonical inclusion $\mathbb{Z}^n \hookrightarrow \mathbb{Z}^{2n}$.

Theorem 3.6 (Tran [65], Theorems 5.3, 7.4). *If we put*

$$F_\gamma = \sum_{S \subseteq T} q^{\frac{1}{2}\Phi(S)} \hat{y}_S$$

where the sum runs over all successor-closed subsets S of T , then

$$x_\gamma = F_\gamma M[g_\gamma].$$

The element F_γ is called the *quantum F -polynomial*.

Proof of Theorem 3.1. We apply Tran's Theorem 3.6. Using $\hat{y}_S = M[b_S]$, where b_S is the sum of the column vectors of B indexed by S , we obtain

$$x_\gamma = \sum_S q^{\frac{1}{2}\Phi(S)} M[b_S] M[g_\gamma] = \sum_S q^{\frac{1}{2}\Phi(S) + \frac{1}{2}b_S^T \Lambda g_\gamma} M[b_S + g_\gamma].$$

We show that $\Phi(S) + b_S^T \Lambda g_\gamma$ is zero for every successor-closed subset $S \subseteq T$. This claim completes the proof because it implies that every coefficient in the expansion of x_γ as a linear combination of standard basis elements is equal to 1. Recall that the last n entries of g_γ are 0 by convention. This implies $b_i^T \Lambda g_\gamma = (g_\gamma)_i$ for every $i \in S$. We use equation (3), sum the expression over all $i \in S$ and apply Proposition 3.5 to obtain the result. $\square$

4. An expansion formula for the Kronecker quantum cluster algebra

4.1. The setup

We consider the marked oriented surface $(\mathcal{S}, M)$ which is formed by an annulus $\mathcal{S}$ and a set $M \subseteq \partial\mathcal{S}$ of two marked points, one on each boundary component. Fig. 4 shows a triangulation $\mathcal{T}$ of $(\mathcal{S}, M)$ by two arcs τ_1, τ_2 . The quiver of $\mathcal{T}$ is the Kronecker quiver $1 \rightrightarrows 2$. The quiver of its principal extension, denoted Q , is also shown in Fig. 4. Note that Q is an ice quiver with two mutable vertices $1, 2$ and two frozen vertices $1', 2'$.

Fig. 4. A triangulation of an annulus and its associated quiver.

Fig. 5. The exchange graph of the Kronecker cluster algebra.

The triangulation gives rise to a seed of the associated cluster algebra $\mathcal{A}(\mathbf{x}, \mathbf{y}, B)$. The set of cluster variables of this cluster algebra admits a natural parametrization by the set of integers $\mathbb{Z}$, see Fig. 5. Hence the set of cluster variables of $\mathcal{A}(\mathbf{x}, \mathbf{y}, B)$ can be written as $\{x_n \mid n \in \mathbb{Z}\}$; in this notation the exchange relations become $x_{n-1}x_{n+1} = x_n^2 + y_1^{n-1}y_2^{n-2}$ for $n \geq 2$ and equations for $n \leq 1$ can be written down similarly.

Fomin–Zelevinsky’s Laurent phenomenon asserts that $x_n \in \mathbb{Z}[x_1^{\pm 1}, x_2^{\pm 1}, y_1, y_2]$ for every $n \in \mathbb{Z}$. There are two different explicit formulae to write x_n as a Laurent polynomial in $\mathbf{x}$. The first formula we state is a variation of a theorem by Caldero–Zelevinsky [12, Theorem 4.1]. The authors establish their formula by computing Euler characteristics of quiver Grassmannians and by using Caldero–Chapoton’s formula [11]. Later Zelevinsky [68] gave a simple inductive proof of the formula.

Theorem 4.1 (cf. Caldero–Zelevinsky). *For every $n \geq 0$ we have*

$$x_{n+3} = x_1^{-n-1}x_2^{n+2} + \sum_{p+r \leq n} \binom{n-r}{p} \binom{n+1-p}{r} x_1^{2p-n-1} x_2^{2r-n} y_1^{n+1-r} y_2^p.$$

It is easy to see that the cluster variable x_{n+3} is attached to an arc γ in $(\mathcal{S}, M)$ that crosses τ_1 exactly $n+1$ times and crosses τ_2 exactly n times. Moreover, it can be seen that the snake graph $\mathcal{G}_n$ associated to γ is given as follows.

Definition 4.2 (Snake graphs of Kronecker type). The snake graph $\mathcal{G}_n$ is the following straight snake graph consisting of $2n+1$ tiles of alternating face weight 1 and 2 and with exactly $n+1$ tiles of weight 1 and exactly n tiles of weight 2; the top and bottom edges of the tile of face weight 1 have edge weights 2 and the tile of face weight 2 has edge weights 1:

For some discussion which will become clear later, we also need the snake graph $\mathcal{H}_n$, which is obtained from $\mathcal{G}_n$ by removing the last tile (with weight 1). Note that $\mathcal{H}_n$ contains exactly n tiles with face weight 1 and exactly n tiles with face weight 2.

The perfect matching P_{min} of $\mathcal{G}_n$ is formed by all the edges of weight 1 and its weight monomial is equal to $x(P_{min}) = x_2^{2n+2}$. The following statement is a precedent of Theorem 2.9 due to [45,46].

Theorem 4.3 (*Musiker–Propp, Musiker–Schiffler*). *Let $n \geq 0$. Then we have*

$$x_{n+3} = \frac{1}{x_1^{n+1}x_2^n} \left(\sum_{P \models \mathcal{G}_n} x(P)y(P) \right).$$

Remark 4.4. Suppose that $n \geq 0$. The cluster variable x_{-n} corresponds to an arc in $(\mathcal{S}, M)$ that crosses τ_1 exactly n times and crosses τ_2 exactly $n+1$ times. Its snake graph is obtained from $\mathcal{G}_n$ by reversing the roles of the indices 1 and 2. There are formulae for x_{-n} analogous to Theorem 4.1 and 4.3 and all other statements made later in the text. For better readability we refrain from writing down statements for both x_{-n} and x_{n+3} and focus on (quantum) cluster variables with positive indices throughout the text.

In the following we abbreviate $\text{Twist}(P)_{\tau_i}$ by $\text{Twist}(P)_i$ for $i \in \{1, 2\}$.

Definition 4.5 (*Level sets*). Let p, r, n be natural numbers. The *level set* $\text{Match}(\mathcal{G}_n)_{p,r}$ is the set of perfect matchings P of $\mathcal{G}_n$ with $|\text{Twist}(P)_1| = n+1-r$ and $|\text{Twist}(P)_2| = p$. (In other words, $\text{Match}(\mathcal{G}_n)_{p,r}$ is the set of perfect matchings obtained from P_{min} by twisting $n+1-r$ tiles of weight 1 and p tiles of weight 2.) In this case we also write $P \models_p^r \mathcal{G}_n$. Similarly, we let $\text{Match}(\mathcal{H}_n)_{p,r}$ be the set of perfect matchings P of $\mathcal{H}_n$ with $|\text{Twist}(P)_1| = n-r$ and $|\text{Twist}(P)_2| = p$. In this case we also write $P \models_p^r \mathcal{H}_n$.

Remark 4.6. Let p, r, n be natural numbers.

- Let $P \in \text{Match}(\mathcal{G}_n) \setminus \{P_{min}\}$. If a tile $G \in \text{Twist}(P)$ has face weight 2, then its neighboring tiles G' and G'' must belong to $\text{Twist}(P)$ as well. In particular, we must have $|\text{Twist}(P)_1| \geq |\text{Twist}(P)_2| + 1$. It follows that $\text{Match}(\mathcal{G}_n)_{p,r} = \emptyset$ unless $p+r \leq n$ or $(p, r) = (0, n+1)$.
- A perfect matching satisfies $P \models_p^r \mathcal{G}_n$ if and only if $y(P) = y_1^{n+1-r} y_2^p$.
- Combining Theorems 4.1 and 4.3 we see that $|\text{Match}(\mathcal{G}_n)_{p,r}| = \binom{n-r}{p} \binom{n+1-p}{r}$ if $p+r \leq n$. Note that there is a direct combinatorial proof of this identity by

Musiker–Propp [45, Section 2.1]. The same authors also show that $|\text{Match}(\mathcal{H}_n)_{p,r}| = \binom{n-r}{p} \binom{n-p}{r}$ if $p+r \leq n$.

4.2. The quantum cluster algebra of Kronecker type

We consider the quantum cluster algebra $\mathcal{A}_q(B, \Lambda)$ constructed from the principal quantization pair

$$B = \begin{pmatrix} \tilde{B} \\ I \end{pmatrix} = \begin{pmatrix} 0 & 2 \\ -2 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \Lambda = \begin{pmatrix} 0 & -I \\ I & -\tilde{B} \end{pmatrix} = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & -2 \\ 0 & 1 & 2 & 0 \end{pmatrix}.$$

The initial seed of $\mathcal{A}_q(B, \Lambda)$ is denoted by $(\mathbf{x}, \mathbf{y}, B, \Lambda)$. In particular, we have $x_1x_2 = x_2x_1$, $y_1y_2 = q^{-2}y_2y_1$ and $x_iy_i = q^{-1}y_ix_i$ for $i \in \{1, 2\}$.

By a slight abuse of notation we denote the quantum cluster variables of $\mathcal{A}_q(B, \Lambda)$ again by x_n with $n \in \mathbb{Z}$. By equation (1) the g -vector of x_{n+3} is

$$g(x_{n+3}) = \deg(x(P_{\min})) - \deg(\text{cross}(\mathcal{T}, \gamma)) = \binom{0}{2n+2} - \binom{n+1}{n} = \binom{-n-1}{n+2} \quad (4)$$

where γ is the arc corresponding to x_{n+3} . From this we can deduce that the G - and C -matrix for the cluster $(x_{n+3}, x_{n+2}, y_1, y_2)$ are

$$G = \begin{pmatrix} -n-1 & -n \\ n+2 & n+1 \end{pmatrix}, \quad C = G^{-T} = \begin{pmatrix} -n-1 & n+2 \\ -n & n+1 \end{pmatrix}.$$

So according to Proposition 2.12 we obtain the q -commutativity relations $x_{n+3}y_1 = q^{n+1}y_1x_{n+3}$, $x_{n+3}y_2 = q^{-n-2}y_2x_{n+3}$, and $y_1y_2 = q^{-2}y_2y_1$. These relations imply that

$$x_{n+2}^{-1}y_1^{n+2}y_2^{n+1} = q^{1-2(n+2)(n+1)}y_2^{n+1}y_1^{n+2}x_{n+2}^{-1}.$$

Hence the exchange relations become

$$\begin{aligned} x_{n+2}x_{n+4} &= x_{n+3}^2 + q^{-1/2+(n+2)(n+1)}y_1^{n+2}y_2^{n+1} \quad (n \geq 0), \\ x_{n+4}x_{n+2} &= x_{n+3}^2 + q^{1/2+(n+2)(n+1)}y_1^{n+2}y_2^{n+1} \quad (n \geq 0). \end{aligned}$$

4.3. An expansion formula via quantum binomial coefficients

Let us recall some definitions from quantum algebra. Given two natural numbers k, n with $k \leq n$. The *quantum integer*, the *quantum factorial* and the *quantum binomial coefficient* are defined as

(b) The element belongs to the quantum cluster algebra. More precisely,

$$\begin{aligned} s_1 &= M[1, -1, 1, 1] + M[-1, -1, 1, 0] + M[-1, 1, 0, 0] + qM[1, 1, 0, 1] - qM[1, 1, 0, 1] \\ &= (M[2, -1, 0, 1] + M[0, -1, 0, 0]) (M[-1, 0, 1, 0] + M[-1, 2, 0, 0]) - qM[1, 1, 0, 1] \\ &= x_0x_3 - q^{1/2}x_1x_2y_2 \in \mathcal{A}_q(B, \Lambda). \end{aligned}$$

(c) The name s_1 is chosen in accordance with the name of the corresponding non-quantized element in the classical cluster algebra $\mathcal{A}(\mathbf{x}, \mathbf{y}, B)$, see Zelevinsky [68]. This element plays a crucial role in the construction of bases of $\mathcal{A}(\mathbf{x}, \mathbf{y}, B)$. It is associated with the loop inside the annulus.

Lemma 4.10. *The equation $x_ns_1 = x_{n+1} + qx_{n-1}y_1y_2$ holds for every $n \geq 2$.*

Proof. We prove the lemma by induction on n . Suppose that $n = 2$. By definition we have

$$\begin{aligned} x_2s_1 &= x_2 (M[-1, 1, 0, 0] + M[-1, -1, 1, 0] + M[1, -1, 1, 1]) \\ &= x_1^{-1}x_2^2 + q^{-1/2}x_1^{-1}y_1 + qx_1y_1y_2 = x_3 + qx_1y_1y_2. \end{aligned}$$

For the induction step it is enough to show that for every $n \geq 3$,

$$\begin{aligned} x_n^{-1}x_{n+1} + qx_n^{-1}x_{n-1}y_1y_2 &= x_{n-1}^{-1}x_n + qx_{n-1}^{-1}x_{n-2}y_1y_2 \\ \Leftrightarrow x_{n-1}x_{n+1} + qx_{n-1}^2y_1y_2 &= x_n^2 + qx_nx_{n-2}y_1y_2. \end{aligned}$$

By the exchange relations the last equation is equivalent to

$$\begin{aligned} x_n^2 + q^{-1/2+(n-1)(n-2)}y_1^{n-1}y_2^{n-2} + qx_{n-1}^2y_1y_2 \\ &= x_n^2 + q \left(x_{n-1}^2 + q^{1/2+(n-2)(n-3)}y_1^{n-2}y_2^{n-3} \right) y_1y_2 \\ \Leftrightarrow q^{-1/2+(n-1)(n-2)}y_1^{n-1}y_2^{n-2} \\ &= qq^{1/2+(n-2)(n-3)}q^{2(n-3)}y_1^{n-1}y_2^{n-2}. \end{aligned}$$

This equation is true so that we have proved the lemma. $\square$

Sketch of the proof of Theorem 4.7. Using Lemma 4.10 we can proceed in a similar way as in the proof of the closely related formula from [41, Theorem 5.3]. $\square$

Definition 4.11 (*Quantum coefficients*). For natural numbers p, r, n with $p + r \leq n$ we denote the coefficients in Theorem 4.7 by

$$c_{p,r,n} = \begin{bmatrix} n-r \\ p \end{bmatrix}_q \begin{bmatrix} n+1-p \\ r \end{bmatrix}_q.$$

Moreover we set $c_{0,n+1,n} = 1$ and $c_{p,r,n} = 0$ otherwise. Furthermore, for $p + r \leq n$ we put

$$d_{p,r,n} = \begin{bmatrix} n-r \\ p \end{bmatrix}_q \begin{bmatrix} n-p \\ r \end{bmatrix}_q$$

and set $d_{p,r,n} = 0$ for $p + r > n$.

Remark 4.12. The ring homomorphism $\mathbb{Z}[q^{\pm 1/2}] \rightarrow \mathbb{Z}[q^{\pm 1/2}]$ defined by $q^{1/2} \mapsto q^{-1/2}$ is called bar involution. It is usually denoted by $\overline{(\cdot)}$. Theorem 4.7 implies that the coefficients in the quantum Laurent expansion with respect to Berenstein–Zelevinsky’s basis elements are bar invariant, that is, $\overline{c_{p,r,n}} = c_{p,r,n}$ for all p, r, n . For a related result about F -polynomials see Tran [65, Corollary 6.5]. For related results about quantum cluster varieties see Allegretti–Kim [5, Theorem 1.2 (4)] and Allegretti [3, Theorem 4.7 (3)].

4.4. Expansion formulae via perfect matchings

Notation 4.13. Let us denote the $2n + 1$ tiles of the snake graph $\mathcal{G}_n$ in Definition 4.2 by

$$G_{-n}, G_{-(n-1)}, \dots, G_{n-1}, G_n$$

from left to right. Similarly, denote the $2n$ tiles of the snake graph $\mathcal{H}_n$ by

$$H_{-n}, H_{-(n-1)}, \dots, H_{n-2}, H_{n-1}$$

from left to right.

Definition 4.14 (*Exponents of tiles*). Let $n \geq 0$.

- (1) The function α assigns every tile of $\mathcal{G}_n$ the half-integer $\alpha(G_i) = i/2$ if G_i has weight 1, and the half-integer $\alpha(G_i) = -i/2$ if G_i has weight 2.
- (2) The function α assigns every tile of $\mathcal{H}_n$ the half-integer $\alpha(H_i) = (i-1)/2$ if H_i has weight 1, and the half-integer $\alpha(H_i) = -i/2$ if H_i has weight 2.

An example is shown in Fig. 7. For every Kronecker snake graph $\mathcal{G}$ (e.g. $\mathcal{G} = \mathcal{G}_n$ or $\mathcal{G} = \mathcal{H}_n$ for some $n \geq 1$) define a map

$$\alpha: \text{Match}(\mathcal{G}) \rightarrow \frac{1}{2}\mathbb{Z}, \quad P \mapsto \sum_{G \in \text{Twist}(P)} \alpha(G).$$

Definition 4.15 (*Expansions from matchings*). For $n \geq 0$ we put

$$r_n = \sum_{P \models \mathcal{G}_n} q^{\alpha(P)} M[\nu(P)], \quad s_n = \sum_{P \models \mathcal{H}_n} q^{\alpha(P)} M[\nu(P)].$$

Fig. 7. The map α for $\mathcal{G}_3$ (left) and $\mathcal{H}_3$ (right).

Notice that $\alpha(P) = 0$ for every perfect matching P of $\mathcal{H}_1$. Hence the formula for s_1 in Definition 4.15 agrees with the formula for s_1 given in Definition 4.8. Note that $s_0 = 1$ since $\mathcal{H}_0$ (the snake graph with no tiles consisting of a single vertical unweighted edge) has exactly 1 perfect matching P with $\nu(P) = 0$ and $\alpha(P) = 0$.

Lemma 4.16. *The following recursive formulae are true.*

$$(a) \quad r_n x_1 = q^{1/2} s_n y_1 + r_{n-1} x_2, \quad (b) \quad x_2 s_n = r_{n-1} + q^{-1/2} M[1, 0, 1, 1] s_{n-1}.$$

Proof. By definition we have

$$r_n x_1 = \left(\sum_{P \models \mathcal{G}_n} q^{\alpha(P)} M[\nu(P)] \right) M[1, 0, 0, 0] = \sum_{P \models \mathcal{G}_n} q^{\alpha(P)} M[\nu(P)] M[1, 0, 0, 0]. \quad (6)$$

Notice that the minimum perfect matching of $\mathcal{G}_n$ satisfies $\nu(P_{\min}) = (-n-1, n+2, 0, 0)^T$. Here, the first two entries are given by the g -vector $(-n-1, n+2)$ of the cluster variable, see equations (1) and (4), and the last two entries are zero because $P_{\min}$ does not admit twisted tiles with weights 1 or 2. Whenever we twist a tile of weight 1 we loose two edges of weight 2. Whenever we twist a tile of weight 2 we gain two edges of weight 1. Hence every perfect matching satisfies

$$\nu(P) = (-n-1 + 2y_2(P), n+2 - 2y_1(P), y_1(P), y_2(P))^T.$$

Here we use $y_i(P)$ as a shorthand notation for $|\text{Twist}(P)_i|$ for $i \in \{1, 2\}$. From this we can conclude that

$$\frac{1}{2} \nu(P)^T \Lambda \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \frac{1}{2} \nu(P)^T \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & -2 \\ 0 & 1 & 2 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \frac{1}{2} y_1(P).$$

This term arises as the exponent in the multiplication of the two basis elements in equation (6). It follows that

$$r_n x_1 = \sum_{P \models \mathcal{G}_n} q^{\alpha(P) + 1/2 y_1(P)} M[\nu(P) + (1, 0, 0, 0)^T].$$

We split the sum into two parts. Let e be the rightmost vertical edge of $\mathcal{G}_n$. The map $P \mapsto P' = P \setminus \{e\}$ induces a bijection between perfect matchings of $\mathcal{G}_n$ that contain e and perfect matchings of the graph $\mathcal{H}_n$. For such a perfect matching P the rightmost tile of weight 1 is necessarily twisted. We obtain

$$\begin{aligned} & \sum_{\substack{P \models \mathcal{G}_n \\ e \in P}} q^{\alpha(P) + \frac{1}{2}y_1(P)} M[\nu(P) + (1, 0, 0, 0)^T] \\ &= \sum_{P' \models \mathcal{H}_n} q^{\alpha(P' \cup \{e\}) + \frac{1}{2}|\text{Twist}_1(P' \cup \{e\})|} M[\nu(P) + (1, 0, 0, 0)^T] \\ &= \sum_{P' \models \mathcal{H}_n} q^{\alpha(P' \cup \{e\}) + \frac{1}{2}|\text{Twist}_1(P')| + \frac{1}{2}} M[\nu(P') + (0, 0, 1, 0)^T] \\ &= \left(\sum_{P' \models \mathcal{H}_n} q^{\alpha(P' \cup \{e\}) + \frac{1}{2}|\text{Twist}_1(P')| + \frac{1}{2}x_1(P') - y_2(P')} M[\nu(P')] \right) M[0, 0, 1, 0] \end{aligned}$$

when we denote $\nu(P') = (x_1(P'), x_2(P'), y_1(P'), y_2(P'))^T$. We claim that this term is equal to $q^{1/2}s_n y_1$. To prove the claim it is enough to show that the exponent in the term in the last line of the previous equation equals $\alpha(P') + 1/2$. The term $x_1(P') - 2y_2(P')$ is invariant under twisting since we gain two edges of weight 1 whenever we twist a tile of weight 2. This expression becomes $-n$ for the minimum matching. Hence every perfect matching P' of $\mathcal{H}_n$ satisfies $x_1(P') - 2y_2(P') = -n$. We have $\alpha(P) - \alpha(P') = -\frac{1}{2}y_1(P') + \frac{1}{2}(n+1)$ because the map α gets shifted by $-1/2$ for every tile with label 1 when passing from P to P' , see Fig. 7, and P contains the twisted rightmost tile with exponent $(n+1)/2$. This establishes the claim.

Next we investigate the second part of the sum. If e is not contained in a perfect matching P of $\mathcal{G}_n$, then P contains the rightmost horizontal edges e_1 and e_2 of $\mathcal{G}_n$. The map $P \mapsto P'' = P \setminus \{e_1, e_2\}$ induces a bijection between perfect matchings of $\mathcal{G}_n$ that contain e_1 and e_2 and perfect matchings of the graph $\mathcal{G}_{n-1}$. For such a perfect matching P the two rightmost tiles with weights 1 and 2 are not twisted. We obtain

$$\begin{aligned} & \sum_{\substack{P \models \mathcal{G}_n \\ e_1, e_2 \in P}} q^{\alpha(P) + \frac{1}{2}y_1(P)} M[\nu(P) + (1, 0, 0, 0)^T] \\ &= \sum_{P'' \models \mathcal{G}_{n-1}} q^{\alpha(P'' \cup \{e_1, e_2\}) + \frac{1}{2}|\text{Twist}_1(P'' \cup \{e_1, e_2\})|} M[\nu(P) + (1, 0, 0, 0)^T] \\ &= \sum_{P'' \models \mathcal{G}_{n-1}} q^{\alpha(P'' \cup \{e_1, e_2\}) + \frac{1}{2}|\text{Twist}_1(P'')|} M[\nu(P'') + (0, 1, 0, 0)^T] \\ &= \left(\sum_{P'' \models \mathcal{G}_{n-1}} q^{\alpha(P'' \cup \{e_1, e_2\}) + \frac{1}{2}|\text{Twist}_1(P'')| - \frac{1}{2}y_2(P'')} M[\nu(P'')] \right) M[0, 1, 0, 0]. \end{aligned}$$

We claim that this term is equal to $r_{n-1}x_2$. We show that the exponent of the term in the last line of the previous equation equals $\alpha(P'')$. By construction $y_2(P) = |\text{Twist}_2(P'')|$. We have $\alpha(P) - \alpha(P'') = -\frac{1}{2}|\text{Twist}_1(P'')| + \frac{1}{2}|\text{Twist}_1(P'')|$ because the map α gets shifted by $\pm 1/2$ (depending on the weight) when passing from P to P'' .

This completes the proof of part (a). Part (b) is shown in a similar way. $\square$

Definition 4.17 (*Coefficients from perfect matchings*). Let p, r, n be natural numbers such that $p \leq n+1$ and $r \leq n$. We put

$$\tilde{c}_{p,r,n} = \sum_{\substack{P \models \mathcal{G}_n \\ y_1(P)=n+1-r \\ y_2(P)=p}} q^{\alpha(P)}, \quad \tilde{d}_{p,r,n} = \sum_{\substack{P \models \mathcal{H}_n \\ y_1(P)=n-r \\ y_2(P)=p}} q^{\alpha(P)}.$$

Remark 4.18. Let $n \geq 0$. A combination of Definitions 4.15 and 4.17 yields

$$r_n = \sum_{p,r \geq 0} \tilde{c}_{p,n+1-r,n} M[2p-n-1, 2r-n, n+1-r, p],$$

$$s_n = \sum_{p,r \geq 0} \tilde{d}_{p,n-r,n} M[2p-n, 2r-n, n-r, p].$$

Lemma 4.19. *The following recursions hold for all p, r, n .*

$$(a) \quad q^{\frac{n+1-r}{2}} \tilde{c}_{p,n+1-r,n} = q^{\frac{n+1}{2}} \tilde{d}_{p,n-r,n} + q^{\frac{p}{2}} \tilde{c}_{p,n+1-r,n-1}$$

$$(b) \quad q^{-\frac{p}{2}} \tilde{d}_{p,n-r,n} = \tilde{c}_{p,n-r,n-1} + q^{-\frac{n-r}{2}} \tilde{d}_{p-1,n-r-1,n-1}$$

$$(c) \quad q^{\frac{n+1-r}{2}} c_{p,r,n} = q^{\frac{n+1}{2}} d_{p,r,n} + q^{\frac{p}{2}} c_{p,n+1-r,n-1}$$

$$(d) \quad q^{-\frac{p}{2}} d_{p,r,n} = c_{p,r,n-1} + q^{-\frac{n-r}{2}} d_{p-1,r,n-1}$$

Proof. We substitute the expressions from Remark 4.18 in Lemma 4.16 and compare coefficients. Evaluation of Lemma 4.16 (a) at $M[2p-n, 2r-n, n+1-r, p]$ yields part (a) of this lemma, evaluation of Lemma 4.16 (b) at $M[2p-n, 2r-n+1, n-r, p]$ yields part (b) of this lemma.

For part (c) we use the quantum Pascal rule to get

$$c_{p,r,n} = \begin{bmatrix} n-r \\ p \end{bmatrix}_q \left(q^{\frac{r}{2}} \begin{bmatrix} n-p \\ r \end{bmatrix}_q + q^{-\frac{n+1-p-r}{2}} \begin{bmatrix} n-p \\ r-1 \end{bmatrix}_q \right)$$

$$= q^{\frac{r}{2}} d_{p,r,n} + q^{-\frac{n+1-p-r}{2}} c_{p,r-1,n-1}.$$

Part (d) is proved in the same way as part (c). $\square$

The following theorem is a generalization of Theorem 4.3 to the quantum cluster algebra of Kronecker type $\mathcal{A}_q(B, \Lambda)$.

Theorem 4.20. *For every $n \geq 0$ the following equality holds true:*

$$x_{n+3} = \sum_{p,r \geq 0} \tilde{c}_{p,n+1-r,n} M[2p-n-1, 2r-n, n+1-r, p] = \sum_{P \models \mathcal{G}_n} q^{\alpha(P)} M[v(P)];$$

$$s_n = \sum_{p,r \geq 0} \tilde{d}_{p,n-r,n} M[2p-n, 2r-n, n-r, p] = \sum_{P \models \mathcal{H}_n} q^{\alpha(P)} M[\nu(P)].$$

Moreover, $s_n \in \mathcal{A}_q(B, \Lambda)$.

Proof. We have to show that the equalities $c_{p,r,n} = \tilde{c}_{p,n+1-r,n}$ and $d_{p,r,n} = \tilde{d}_{p,n-r,n}$ are both true for all natural numbers p, r, n . We prove these statements by induction on n .

For the base case suppose that $n = 0$. In this case the statement for the coefficients d and $\tilde{d}$ is true because by construction $d_{0,0,0} = \tilde{d}_{0,0,0} = 1$ is the only non-zero coefficient. For the statement for the coefficients c and $\tilde{c}$, note that the single tile G in the snake graph $\mathcal{G}_1$ satisfies $\alpha(G) = 0$. Hence $\alpha(P) = 0$ for both perfect matchings $P \models \mathcal{G}_1$. This implies $\tilde{c}_{0,0,0} = \tilde{c}_{0,1,0} = 1$ and $\tilde{c}_{p,r,0} = 0$ otherwise. On the other hand, using $[0]_q = 1$ and $[1]_q = 1$ in Definition 4.11 we see that $c_{0,0,0} = c_{0,1,0} = 1$ and $c_{p,r,0} = 0$ otherwise.

The induction step follows from Lemma 4.19. If the statements are true for $n-1$, then parts (b) and (d) of the lemma imply the statement for the coefficients d and $\tilde{d}$, and using this equality together with parts (a) and (c) of the lemma we establish the equality of coefficients c and $\tilde{c}$.

Lemma 4.16 (a) implies $s_n y_1 \in \mathcal{A}_q(B, \Lambda)$. Since the frozen variable y_1 is invertible we can conclude that s_n lies in $\mathcal{A}_q(B, \Lambda)$. $\square$

Remark 4.21. Some authors do not invert the frozen variables in the definition of a (quantum) cluster algebra. Using the recursions from Lemmas 4.10 and 4.16, one can prove by induction that s_n belongs to the quantum cluster algebra without inverted frozen variables.

4.5. BPS states

There is a connection between the elements

$$s_n = \sum_{P \models \mathcal{H}_n} q^{\alpha(P)} M[\nu(P)]$$

associated to self-crossing arcs and Bogomol'nyi–Prasad–Sommerfield states (BPS states) in supersymmetric 4-dimensional quantum field theories. Córdova–Neitzke, see [20], consider a supersymmetric quantum field theory of quiver type whose line defects W_n are parametrized by natural numbers n . The authors consider a generating function $F(W_n)$ of BPS states such that the coefficients are counting BPS states of a given electromagnetic central charge and a given angular momentum (in the x_3 direction).

Córdova–Neitzke use a Coulomb branch formula, see [20, Equation (3.22)], to compute $F(W_n)$ recursively. The Coulomb branch formula is based on work of Manschot–Pioline–Sen, see [43], which relates the Coulomb branch to moduli spaces of quiver representations. Córdova–Neitzke compute $F(W_n)$ explicitly for $n \in \{0, 1, 2, 3, 4\}$. A direct comparison with our expansion formula establishes the following proposition.

Proposition 4.22. *For $n \in \{0, 1, 2, 3, 4\}$ we have $s_n = F(W_n)$.*

Question 4.23. Does $s_n = F(W_n)$ hold for all $n \in \mathbb{N}$?

Several authors have studied variations of the question in the classical commutative setup. Gaiotto–Moore–Neitzke [32, Section 5] conjectured that if $q = 1$, then $F(W_n)$ is equal to a Fock–Goncharov canonical basis element in a cluster variety. Córdova–Neitzke give a second method to calculate $F(W_n)$ using the Higgs branch. Allegretti [3] relates generating functions of BPS states calculated via the Higgs branch to elements in cluster algebras.

5. The Stembridge phenomenon

Stembridge’s $q = -1$ phenomenon [63, Section 0] asserts the following: Suppose that we have a finite set B of combinatorial origin together with a natural weight function $w \in B \rightarrow \mathbb{Z}$. We consider the Laurent polynomial $X(q) = \sum_{b \in B} q^{w(b)} \in \mathbb{Z}[q^{\pm 1}]$ with $X(1) = |B|$. Moreover suppose that B admits a natural involution $\sigma: B \rightarrow B$. Stembridge’s phenomenon asserts that often $X(-1) = |B^\sigma|$ is equal to the number of fixed points in B under the action of $\sigma: B \rightarrow B$.

We consider the quantum cluster algebra attached to the Kronecker quiver from Section 4. Suppose that p, r, n are natural numbers with $p + r \leq n$. The graph $\mathcal{G}_n$ admits a horizontal and a vertical axis of symmetry. We consider the reflection across the vertical axis of symmetry. The reflection induces an involution $\sigma: \text{Match}(\mathcal{G}_n) \rightarrow \text{Match}(\mathcal{G}_n)$. Note that σ leaves the height monomial $y(P)$ of a perfect matching invariant. By restriction we obtain an involution $\sigma: \text{Match}(\mathcal{G}_n)_{p,r} \rightarrow \text{Match}(\mathcal{G}_n)_{p,r}$. To construct the polynomial X let us introduce the following variation of quantum numbers and quantum binomial coefficients.

Definition 5.1 (*Gaussian integers and binomial coefficients and evaluations at $q = -1$*). Let k, n be natural numbers with $k \leq n$.

- The polynomial $(n)_q = (q^n - 1)/(q - 1) = 1 + q + \dots + q^{n-1} \in \mathbb{Z}[q]$ is called *Gaussian integer*. The *Gaussian factorial* is defined as $(n)_q! = (n)_q(n-1)_q \dots (1)_q \in \mathbb{Z}[q]$. The *Gaussian binomial coefficient* is defined as $\binom{n}{k}_q = (n)_q! / [(k)_q!(n-k)_q!] \in \mathbb{Z}[q]$.
- The natural numbers $(n)_{-1}$, $(n)_{-1}!$ and $\binom{n}{k}_{-1}$ are defined to be the evaluations of the polynomials in part (a) at $q = -1$.

Remark 5.2. The Gaussian integers and quantum integers agree up to a power of $q^{1/2}$. In particular,

$$[n]_q = q^{\frac{1-n}{2}} \frac{q^n - 1}{q - 1} = (n)_q, \quad [n]_{q!} = q^{-n(n-1)/4} (n)_q!, \quad \left[\begin{matrix} n \\ k \end{matrix} \right]_q = q^{-k(n-k)/2} \binom{n}{k}_q,$$

$$c_{p,r,n} = q^{-[r+(n-r-p)(p+r)]/2} \binom{n-r}{p}_q \binom{n+1-p}{r}_q.$$

Let us describe the evaluation of Gaussian binomial coefficients at $q = -1$.

Lemma 5.3. For all natural numbers n, k with $k \leq n$ the following equation holds:

$$\binom{n}{k}_{-1} = \begin{cases} 0 & \text{if } k \equiv 1 \pmod{2} \text{ and } n \equiv 0 \pmod{2}; \\ \binom{\lfloor n/2 \rfloor}{\lfloor k/2 \rfloor} & \text{otherwise.} \end{cases}$$

Proof. In this proof, all congruences are read modulo 2. The polynomial $(n)_q \in \mathbb{Z}[q]$ is divisible by $q + 1$ if and only if n is even. In this case we may write $(n)_q = (q + 1)h_n$ for some polynomial $h_n \in \mathbb{Z}[q]$ with $h_n(-1) = n/2$. If n is odd, then $(n)_{-1} = 1$.

Thus, when we write the rational function $\binom{n}{k}_q$ in lowest terms, the irreducible polynomial $q + 1$ occurs with multiplicity $\lfloor n/2 \rfloor - \lfloor k/2 \rfloor - \lfloor (n - k)/2 \rfloor$. This multiplicity is equal to 1 if $k \equiv 1$ and $n \equiv 0$, and it is 0 otherwise. In particular, $\binom{n}{k}_{-1} = 0$ if $(k, n) \equiv (1, 0) \pmod{2}$. For $(k, n) \not\equiv (1, 0) \pmod{2}$ we may evaluate $\binom{n}{k}_q = (n)_q! / [(k)_q! (n - k)_q!]$ at $q = -1$ by replacing every $(l)_q$ with 1 for every odd l and every $(l)_q$ with $l/2$ for every even l . The desired equality follows. $\square$

The following corollary is a direct consequence of Lemma 5.3.

Corollary 5.4. Suppose that p, r, n are natural numbers with $p + r \leq n$. If (p, r, n) is congruent to $(1, 0, 0)$, $(1, 1, 1)$, $(0, 1, 1)$, or $(1, 1, 0)$ modulo 2, then

$$\binom{n-r}{p}_{-1} \binom{n+1-p}{r}_{-1} = 0.$$

If (p, r, n) is congruent to $(0, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$, or $(1, 0, 1)$ modulo 2, then

$$\binom{n-r}{p}_{-1} \binom{n+1-p}{r}_{-1} = \binom{\lfloor (n-r)/2 \rfloor}{\lfloor p/2 \rfloor} \binom{\lfloor (n+1-p)/2 \rfloor}{\lfloor r/2 \rfloor}.$$

Theorem 5.5. Assume that p, r, n are natural numbers such that $p + r \leq n$. Then

$$\binom{n-r}{p}_{-1} \binom{n+1-p}{r}_{-1} = |\{P \in \text{Match}(\mathcal{G}_n)_{p,r} \mid \sigma(P) = P\}|.$$

Proof. We distinguish the following cases:

(1) Assume that n is even. In this case the central tile of $\mathcal{G}_n$ has weight 1.

(1.1) Assume that p is even.

(1.1.1) Assume that r is even. Then for every $P \models_p^r \mathcal{G}_n$ the number of tiles $G \in \text{Twist}(G)_1$ of weight 1, namely $n+1-r$, is odd. Hence the central tile must be twisted.

A perfect matching $P \models_p^r \mathcal{G}_n$ with $\sigma(P) = P$ is uniquely determined by a perfect matching of the full subgraph isomorphic to $\mathcal{H}_{n/2}$ on the left of the central tile (because this determines the structure of P on the right of the central tile). Hence the number of matchings $P \models_p^r \mathcal{G}_n$ with $\sigma(P) = P$ is equal to the cardinality of the set $\text{Match}(\mathcal{H}_{n/2})_{p',r'}$ where $p' = p/2$ and $r' = r/2$. Remark 4.6 and Lemma 5.3 imply that $|\{P \in \text{Match}(\mathcal{G}_n)_{p,r} \mid \sigma(P) = P\}|$ is equal to

$$\binom{(n-r)/2}{p/2} \binom{(n-p)/2}{r/2} = \binom{n-r}{p}_{-1} \binom{n+1-p}{r}_{-1}.$$

(1.1.2) Assume that r is odd. Then for every $P \models_p^r \mathcal{G}_n$ the number of tiles $G \in \text{Twist}(G)_1$ of weight 1, namely $n+1-r$, is even. Hence the central tile cannot be twisted, and locally at the central tile the perfect matching contains the top and the bottom edges.

A perfect matching $P \models_p^r \mathcal{G}_n$ with $\sigma(P) = P$ is uniquely determined by a perfect matching of the full subgraph isomorphic to $\mathcal{G}_{(n-2)/2}$ on the left of the central tile (because this determines the structure of P on the right of the central tile). Hence the number of matchings $P \models_p^r \mathcal{G}_n$ with $\sigma(P) = P$ is equal to the cardinality of the set $\text{Match}(\mathcal{G}_{(n-2)/2})_{p',r'}$ where $p' = p/2$ and $r' = (r+1)/2$. Remark 4.6 and Lemma 5.3 imply that $|\{P \in \text{Match}(\mathcal{G}_n)_{p,r} \mid \sigma(P) = P\}|$ is equal to

$$\binom{(n-r-1)/2}{p/2} \binom{(n-p)/2}{(r-1)/2} = \binom{n-r}{p}_{-1} \binom{n+1-p}{r}_{-1}.$$

(1.2) Assume that p is odd. Any perfect matching $P \models_p^r \mathcal{G}_n$ with $\sigma(P) = P$ arises from $P_{\min}$ by twisting the same number of tiles labelled 2 on each side of the central tile. Hence $|\{P \in \text{Match}(\mathcal{G}_n)_{p,r} \mid \sigma(P) = P\}| = 0$. The desired equality follows from Corollary 5.4.

(2) The case when n is odd can be dealt with in an analogous way. $\square$

Corollary 5.6. *Let p, r, n be natural numbers with $p+r \leq n$. Put*

$$X(q) = \binom{n-r}{p}_q \binom{n+1-p}{r}_q = q^{[r+(n-r-p)(p+r)]/2} \sum_{P \models_p^r \mathcal{G}_n} q^{\alpha(P)}$$

and define a map $w: \text{Match}(\mathcal{G}_n)_{p,r} \rightarrow \mathbb{N}$ by

$$P \mapsto \frac{r + (n - r - p)(p + r)}{2} + \alpha(P).$$

Then the quadruple $(\text{Match}(\mathcal{G}_n)_{p,r}, \sigma, X, w)$ satisfies Stembridge's $q = -1$ phenomenon.

References

- [1] M. Alim, S. Cecotti, C. Córdova, S. Espahbodi, A. Rastogi, C. Vafa, BPS quivers and spectra of complete $\mathcal{N} = 2$ quantum field theories, *Comm. Math. Phys.* 323 (3) (2013) 1185–1227.
- [2] M. Alim, S. Cecotti, C. Córdova, S. Espahbodi, A. Rastogi, C. Vafa, $\mathcal{N} = 2$ quantum field theories and their BPS quivers, *Adv. Theor. Math. Phys.* 18 (1) (2014) 27–127.
- [3] D. Allegretti, A duality map for the quantum symplectic double, preprint, arXiv:1605.01599, 2016.
- [4] D. Allegretti, Categorified canonical bases and framed BPS states, preprint, arXiv:1806.10394, 2018.
- [5] D. Allegretti, H.K. Kim, A duality map for quantum cluster varieties from surfaces, *Adv. Math.* 306 (2017) 1164–1208.
- [6] C. Amiot, P.-G. Plamondon, The cluster category of a surface with punctures via group actions, preprint, arXiv:1707.01834, 2017.
- [7] A. Berenstein, S. Fomin, A. Zelevinsky, Cluster algebras III. Upper bounds and double Bruhat cells, *Duke Math. J.* 126 (1) (2005) 1–52.
- [8] A. Berenstein, V. Retakh, Noncommutative marked surfaces, *Adv. Math.* 328 (2018) 1010–1087.
- [9] A. Berenstein, A. Zelevinsky, Quantum cluster algebras, *Adv. Math.* 195 (2005) 405–455.
- [10] T. Brüstle, J. Zhang, On the cluster category of a marked surface without punctures, *Algebra Number Theory* 5 (4) (2011) 529–566.
- [11] Ph. Caldero, F. Chapoton, Cluster algebras as Hall algebras of quiver representations, *Comment. Math. Helv.* 81 (3) (2006) 595–616.
- [12] Ph. Caldero, A. Zelevinsky, Laurent expansions in cluster algebras via quiver representations, *Mosc. Math. J.* 6 (3) (2006) 411–429.
- [13] İ. Çanakçı, K. Lee, R. Schiffler, On cluster algebras for surfaces without punctures and one marked point, *Proc. Amer. Math. Soc., Ser. B* 2 (2015) 35–49.
- [14] İ. Çanakçı, R. Schiffler, Snake graph calculus and cluster algebras from surfaces, *J. Algebra* 382 (2013) 240–281.
- [15] İ. Çanakçı, R. Schiffler, Snake graph calculus and cluster algebras from surfaces II: self-crossing snake graphs, *Math. Z.* 281 (1) (2015) 55–102.
- [16] İ. Çanakçı, R. Schiffler, Cluster algebras and continued fractions, *Compos. Math.* 154 (3) (2018) 565–593.
- [17] İ. Çanakçı, R. Schiffler, Snake graph calculus and cluster algebras from surfaces III: band graphs and snake rings, *Int. Math. Res. Not. IMRN* 2019 (4) (2019) 1145–1226, <https://doi.org/10.1093/imrn/rnx157>.
- [18] İ. Çanakçı, S. Schroll, With an appendix by C. Amiot: extensions in Jacobian algebras and cluster categories of unpunctured surfaces, *Adv. Math.* 313 (2017) 1–49.
- [19] M. Cirařici, Line defects and (framed) BPS quivers, *J. High Energy Phys.* 2013 (11) (2013) 141.
- [20] C. Córdova, A. Neitzke, Line defects, tropicalization, and multi-centered quiver quantum mechanics, *J. High Energy Phys.* 2014 (9) (2014) 099.
- [21] B. Davison, Positivity for quantum cluster algebras, *Ann. of Math.* (2) 187 (1) (2018) 157–219.
- [22] H. Derksen, J. Weyman, A. Zelevinsky, Quivers with potentials and their representations II: applications to cluster algebras, *J. Amer. Math. Soc.* 23 (3) (2010) 749–790.
- [23] A. Felikson, M. Shapiro, P. Tumarkin, Skew-symmetric cluster algebras of finite mutation type, *J. Eur. Math. Soc.* 14 (4) (2012) 1135–1180.
- [24] A. Felikson, M. Shapiro, P. Tumarkin, Cluster algebras of finite mutation type via unfoldings, *Int. Math. Res. Not.* 2012 (8) (2012) 1768–1804.
- [25] V. Fock, A. Goncharov, Moduli spaces of local systems and higher Teichmüller theory, *Publ. Math. Inst. Hautes Études Sci.* 103 (1) (2006) 1–211.
- [26] V. Fock, A. Goncharov, Cluster ensembles, quantization and the dilogarithm, *Ann. Sci. Éc. Norm. Supér.* 42 (6) (2009) 865–930.
- [27] S. Fomin, M. Shapiro, D. Thurston, Cluster algebras and triangulated surfaces. Part I: cluster complexes, *Acta Math.* 201 (1) (2008) 83–146.
- [28] S. Fomin, D. Thurston, Cluster algebras and triangulated surfaces. Part II: lambda lengths, *Mem. Amer. Math. Soc.* 255 (2018) 1223.

- [29] S. Fomin, A. Zelevinsky, Cluster algebras I: foundations, *J. Amer. Math. Soc.* 15 (2) (2002) 497–529.
- [30] S. Fomin, A. Zelevinsky, Cluster algebras II: finite type classification, *Invent. Math.* 154 (1) (2003) 63–121.
- [31] S. Fomin, A. Zelevinsky, Cluster algebras IV: coefficients, *Compos. Math.* 143 (1) (2007) 112–164.
- [32] D. Gaiotto, G. Moore, A. Neitzke, Framed BPS states, *Adv. Theor. Math. Phys.* 17 (2) (2013) 241–397.
- [33] Ch. Geiß, B. Leclerc, J. Schröer, Cluster structures on quantum coordinate rings, *Selecta Math.* 19 (2) (2013) 337–397.
- [34] M. Gekhtman, M. Shapiro, A. Vainshtein, Cluster algebras and Weil–Petersson forms, *Duke Math. J.* 127 (2) (2005) 291–311.
- [35] K.R. Goodearl, M. Yakimov, The Berenstein–Zelevinsky quantum cluster algebra conjecture, preprint, arXiv:1602.00498, 2016.
- [36] J. Grabowski, Examples of quantum cluster algebras associated to partial flag varieties, *J. Pure Appl. Algebra* 215 (7) (2011) 1582–1595.
- [37] J. Grabowski, S. Launois, Graded quantum cluster algebras and an application to quantum Grassmannians, *Proc. Lond. Math. Soc.* 109 (3) (2014) 697–732.
- [38] M. Gross, P. Hacking, S. Keel, M. Kontsevich, Canonical bases for cluster algebras, *J. Amer. Math. Soc.* 31 (2018) 497–608.
- [39] M. Huang, Quantum cluster algebras from unpunctured triangulated surfaces, preprint, arXiv:1807.06910 [math], 2018.
- [40] Y. Kimura, F. Qin, Graded quiver varieties, quantum cluster algebras and dual canonical basis, *Adv. Math.* 262 (2014) 261–312.
- [41] Ph. Lampe, A quantum cluster algebra of Kronecker type and the dual canonical basis, *Int. Math. Res. Not.* 2011 (12) (2011) 2970–3005.
- [42] Ph. Lampe, Quantum cluster algebras of type A and the dual canonical basis, *Proc. Lond. Math. Soc.* (3) 108 (1) (2014) 1–43.
- [43] J. Manschot, B. Pioline, A. Sen, On the Coulomb branch and Higgs branch formulae for multi-centered black holes and quiver invariants, *J. High Energy Phys.* 2013 (5) (2013) 166.
- [44] G. Muller, Skein and cluster algebras of marked surfaces, *Quantum Topol.* 7 (3) (2016) 435–503.
- [45] G. Musiker, J. Propp, Combinatorial interpretations for rank-two cluster algebras of affine type, *Electron. J. Combin.* 14 (1) (2007) 15.
- [46] G. Musiker, R. Schiffler, Cluster expansion formulas and perfect matchings, *J. Algebraic Combin.* 32 (2) (2010) 187–209.
- [47] G. Musiker, R. Schiffler, L. Williams, Positivity for cluster algebras from surfaces, *Adv. Math.* 227 (6) (2011) 2241–2308.
- [48] G. Musiker, R. Schiffler, L. Williams, Bases for cluster algebras from surfaces, *Compos. Math.* 149 (2) (2013) 217–263.
- [49] K. Nagao, Donaldson–Thomas theory and cluster algebras, *Duke Math. J.* 162 (7) (2013) 1313–1367.
- [50] T. Nakanishi, Periodicities in cluster algebras and dilogarithm identities, in: *Representations of Algebras and Related Topics*, EMS Series of Congress Reports, 2011, pp. 407–443.
- [51] T. Nakanishi, A. Zelevinsky, On tropical dualities in cluster algebras, *Contemp. Math.* 565 (2012) 217–226.
- [52] R.C. Penner, The decorated Teichmüller space of punctured surfaces, *Comm. Math. Phys.* 113 (1987) 299–339.
- [53] P.-G. Plamondon, Cluster algebras via cluster categories with infinite-dimensional morphism spaces, *Compos. Math.* 147 (6) (2011) 1921–1954.
- [54] J. Propp, Lattice structure for orientations of graphs, preprint, arXiv:math/0209005, 2002.
- [55] F. Qin, Quantum cluster variables via Serre polynomials, with an appendix by Bernhard Keller, *J. Reine Angew. Math.* 668 (2012) 149–190.
- [56] F. Qin, t -analog of q -characters, bases of quantum cluster algebras, and a correction technique, *Int. Math. Res. Not. IMRN* 2014 (22) (2014) 6175–6232.
- [57] F. Qin, Quantum groups via cyclic quiver varieties I, *Compos. Math.* 152 (2) (2016) 299–326.
- [58] F. Qin, Triangular bases in quantum cluster algebras and monoidal categorification conjectures, *Duke Math. J.* 166 (12) (2017) 2337–2442.
- [59] Y. Qiu, Y. Zhou, Cluster categories for marked surfaces: punctured case, *Compos. Math.* 153 (9) (2017) 1779–1819.
- [60] D. Rupel, On a quantum analog of the Caldero–Chapoton formula, *Int. Math. Res. Not. IMRN* 2011 (14) (2011) 3207–3236.

- [61] D. Rupel, Quantum cluster characters for valued quivers, *Trans. Amer. Math. Soc.* 367 (2015) 7061–7102.
- [62] R. Schiffler, H. Thomas, On cluster algebras arising from unpunctured surfaces, *Int. Math. Res. Not. IMRN* 2019 (17) (2009) 3160–3189.
- [63] J. Stembridge, Canonical bases and self-evacuating tableaux, *Duke Math. J.* 82 (3) (1996) 585–606.
- [64] C. Szántó, On the cardinalities of Kronecker quiver Grassmannians, *Math. Z.* 269 (2011) 833–846.
- [65] T. Tran, F-polynomials in quantum cluster algebras, *Algebr. Represent. Theory* 14 (2011) 1025–1061.
- [66] H. Williams, Toda systems, cluster characters, and spectral networks, preprint, arXiv:1411.3692.
- [67] A. Zelevinsky, Quantum cluster algebras: Oberwolfach talk, February 2005, preprint, arXiv:math/0502260, 2005.
- [68] A. Zelevinsky, Semicanonical basis generators of the cluster algebra of type $A_1^{(1)}$, *Electron. J. Combin.* 14 (1) (2007) 4.
- [69] J. Zhang, Y. Zhou, B. Zhu, Cotorsion pairs in the cluster category of a marked surface, *J. Algebra* 391 (2013) 209–226.