VU Research Portal

Associations Between Nutrient Intake and Corresponding Nutritional Biomarker Levels in Blood in a Memory Clinic Cohort

de Leeuw, Francisca A.; van de Rest, Ondine; Doorduijn, Astrid S.; Fieldhouse, Jay L.P.; de van der Schueren, Marian A.E.; Visser, Marjolein; van den Heuvel, Ellen G.H.M.; Scheltens, Philip; Teunissen, Charlotte E.; Kester, Maartje I.; van der Flier, Wiesje M.

published in

Journal of the American Medical Directors Association
2020

DOI (link to publisher)

[10.1016/j.jamda.2020.04.031](https://doi.org/10.1016/j.jamda.2020.04.031)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

de Leeuw, F. A., van de Rest, O., Doorduijn, A. S., Fieldhouse, J. L. P., de van der Schueren, M. A. E., Visser, M., van den Heuvel, E. G. H. M., Scheltens, P., Teunissen, C. E., Kester, M. I., & van der Flier, W. M. (2020). Associations Between Nutrient Intake and Corresponding Nutritional Biomarker Levels in Blood in a Memory Clinic Cohort: The NUDAD Project. *Journal of the American Medical Directors Association*, 21(10), 1436-1438. <https://doi.org/10.1016/j.jamda.2020.04.031>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

JAMDA

journal homepage: www.jamda.com

Research Letter

Associations Between Nutrient Intake and Corresponding Nutritional Biomarker Levels in Blood in a Memory Clinic Cohort: The NUDAD Project

To the Editor:

Diet is a promising intervention target to prevent or slow Alzheimer's disease (AD).^{1,2} Early (predementia) stages of AD offer a unique opportunity for dietary interventions.² Nutritional assessment methods to estimate nutrient intake have, however, not been validated in clinical populations. Hence, we assessed the association between nutrient intake assessed by food frequency questionnaire (FFQ) and nutrient status measured by nutritional biomarkers in blood in a clinical sample of controls, mild cognitive impairment (MCI), and patients with AD.

Methods

From the Nutrition, the Unrecognized Determinant in Alzheimer's Disease (NUDAD) study we included 90 participants: 29 patients with AD, 22 patients with MCI, and 39 individuals with subjective cognitive decline (SCD) who served as controls.³ All participants completed a FFQ and had at least 1 nutritional biomarker measurement. Written informed consent was obtained from all participants and the local Medical Ethics Committee approved the study.

Usual dietary intake was measured using the semiquantitative 238-item HELIUS FFQ with a reference period of 4 weeks.⁴ In addition, participants reported their nutritional supplement use. If the nutrient dosage could not be obtained from participants or the (online) instructions leaflet ($n = 11$), it was imputed with the nutrient dosage of the most frequently used comparable supplement in this study.

All nutritional biomarker were measured in fasting plasma, serum, or whole blood samples, except for folic acid and vitamin B12, which were measured in nonfasting samples, and vitamin A, which was measured in both fasting (89%) and nonfasting (11%) samples.

To investigate associations of nutrient intake with nutritional biomarker levels, we used linear regression analyses. Furthermore, we assessed whether associations differed per diagnostic group (controls/MCI/AD). Lastly, we performed sensitivity analyses excluding supplement users for each nutrient separately.

Results

Overall, mean age was 67 ± 9 years, 42 (47%) were female, and vitamin and mineral supplement use was observed in 40 (44%) participants.

Table 1 shows the associations between nutrient (diet + supplement) intake and their corresponding nutritional biomarker levels in blood. Intake and blood levels were associated for vitamin B1, vitamin B6, vitamin B12, vitamin C, folate, and eicosapentaenoic acid, and trends were observed for retinol activity equivalents, total omega-3 fatty acids, and docosahexaenoic acid (DHA). No associations were found for zinc.

Subsequently, we evaluated if associations differed per diagnostic group. Interactions ($P < .10$) were found between diagnostic group and nutrient intake for vitamin B6, vitamin C, total omega-3 fatty acids, and DHA. Stratification by diagnostic group showed associations for vitamin B6 in all diagnostic groups [B(SE) controls; 0.78 (0.06), MCI; 1.05 (0.20), AD; 0.45 (0.14), $P < .01$], for omega-3 fatty acids and DHA in controls [B(SE) controls; 0.50 (0.20), $P = .021$, 0.38 (0.13), $P = .009$, MCI; -0.38 (0.30), $P = .230$, -0.13 (0.27), $P = .646$, AD; 0.25 (0.17), $P = .166$, -0.27 (0.25), $P = .285$] and for vitamin C in AD [B(SE) controls; 0.22 (0.15), $P = .152$, MCI; 0.03 (0.27), $P = .927$, AD; 0.75 (0.11), $P < .001$].

Finally, we re-analyzed the data in the total cohort excluding supplement users for each nutrient separately. Associations remained comparable in effect size for most nutrients, but attenuated for DHA, vitamin B1, B6, and B12 [B(SE) 0.07(0.15), -0.41 (0.60), 0.31(0.23), -0.22 (0.47), $P > .05$].

Discussion

The main finding of this study is that nutrient intake of omega-3 fatty acids, A, B and C vitamins, as assessed by the HELIUS FFQ and self-reported supplement use, is moderately associated with their corresponding nutritional biomarker levels in a memory-clinic cohort of controls and patients with MCI or AD. Associations were largely similar across diagnostic groups, except for total omega-3 fatty acids and DHA that were only associated in controls, and vitamin C that was only associated in AD. Almost one-half of the participants used nutritional supplements.

In the subgroup of nonsupplement users, associations between nutrient intake and nutritional biomarker levels remained largely comparable to our findings in the total cohort.

Among the strengths of this study is that we used a well characterized, clinical population with different stages of cognitive decline. A potential limitation is that the FFQ used in this study has a reference period of 4 weeks, which is relatively short to assess habitual diet. The burden of memory, however, may be less in comparison to other FFQs that use a reference period up to a year,

Table 1
Associations between Nutrient Intake and Nutritional Biomarker Levels in Blood

Determinants	Outcome	All		
		B(SE)	P	n
Nutrient intake	Nutritional biomarker			
Retinol activity equivalents, mcg/d	Vitamin A, $\mu\text{mol/L}$	0.23 (0.12)	.057	87
Vitamin B1*, mg/d	Vitamin B1, nmol/L	0.48 (0.09)	<.001	79
Vitamin B6*, mg/d	Vitamin B6, nmol/L	0.71 (0.07)	<.001	79
Vitamin B12, mcg/d	Vitamin B12, pmol/L	0.35 (0.10)	<.001	77
Vitamin C, mg/d	Vitamin C, $\mu\text{mol/L}$	0.49 (0.09)	<.001	78
Folate equivalents, mcg/d	Folate, nmol/L	0.49 (0.09)	<.001	77
Zinc, mg/d	Zinc, $\mu\text{mol/L}$	−0.10 (0.11)	.373	79
Omega-3 fatty acids, % of total fatty acids intake	Omega-3 fatty acids, % of plasma total fatty acids	0.21 (0.12)	.101	79
DHA, % of total fatty acids intake	DHA, % of plasma total fatty acids	0.19 (0.12)	.102	79
EPA, % of total fatty acids intake	EPA, % of plasma total fatty acids	0.41 (0.11)	<.001	79

EPA, eicosapentaenoic acid; SD, standard deviation.

The HELIUS FFQ was not designed to measure these nutrients.

Linear regression analyses adjusted for sex, age, diagnosis, and total energy intake were used, using separate models for each nutritional biomarker. Data were log-transformed and converted to z scores before data analysis. Effect size is change of SD nutritional biomarker blood level per 1 SD increase in nutrient intake.

which could be an advantage in a cohort of cognitively impaired individuals. Moreover, most plasma and serum biomarkers reflect relatively short-term intake. Future studies should include nutritional biomarkers in adipose tissue or erythrocytes that assess longer-term intake.⁵

In conclusion, nutrient intake and nutrient status were moderately associated and largely similar across diagnostic groups. Our findings indicate that the HELIUS FFQ provides valid estimates of nutrient intake in a memory-clinic cohort.

Acknowledgments

The Alzheimer Center Amsterdam is supported by Stichting Alzheimer Nederland and Stichting VUmc fonds. The clinical database structure was developed with funding from Stichting Dioraphte. Research of the Alzheimer Center Amsterdam is part of the neurodegeneration research program of Amsterdam Neuroscience. W.F. is recipient of a donation by stichting Equilibrio, and of a ZonMW Memorabel grant (#733050814). F.L., A.D., M.K., and J.F. are appointed at the NUDAD project, which is funded by NWO-FCB (project number 057-14-004). W.F. holds the Pasman chair.

We acknowledge members of the NUDAD project team: Amsterdam University Medical Center location VUmc: Wiesje van der Flier, Maartje Kester, Philip Scheltens, Charlotte Teunissen, Marian de van der Schueren, Francien de Leeuw, Astrid Doorduijn, Jay Fieldhouse, Heleen Hendriksen, José Overbeek, and Els Dekkers; Vrije Universiteit Amsterdam: Marjolein Visser; Wageningen University and Research: Ondine van de Rest and Sanne Boesveldt; DSM: Peter van-Dael and Manfred Eggersdorfer; Nutricia Research: John Sijben, Nick van Wijk, Amos Attali, and Martin Verkuil; FrieslandCampina: Rolf Bos, Cecile Singh-Povel, Ellen van den Heuvel, and Martijn Veltkamp.

The authors would like to thank Eva Anne Hartman for her help with the administration and cleaning of the nutritional supplement data. F.L., O.R., A.D., J.F., M.S., M.V. and M.K. report no disclosures. E.H. is an employee of FrieslandCampina, P.S. has received consultancy/speaker fees (paid to the institution) from Novartis, Vivoryon, Genentech and EIP Pharma. C.T. has a collaboration contract with ADx Neurosciences, performed contract research or

received grants from Probiobdrug, Biogen, Esai, Toyama, Janssen prevention center, Boehringer, AxonNeurosciences, Fujirebio, EIP farma, PeopleBio, and Roche. Research programs of W.F. have been funded by ZonMW, NWO, EU-FP7, EU-JPND, Alzheimer Nederland, CardioVascular Onderzoek Nederland, Health-Holland, Topsector Life Sciences & Health, stichting Dioraphte, Gieskes-Strijbis fonds, stichting Equilibrio, Pasman stichting, Biogen MA Inc, Boehringer Ingelheim, Life-MI, AVID, Roche BV, Janssen Stellar, Combinostics. W.F. has performed contract research for Biogen MA Inc and Boehringer Ingelheim. W.F. has been an invited speaker at Boehringer Ingelheim and Biogen MA Inc. All funding is paid to her institution.

References

1. Gustafson DR, Clare Morris M, Scarmeas N, et al. New perspectives on Alzheimer's disease and nutrition. *J Alzheimers Dis* 2015;46:1111–1127.
2. Livingston G, Sommerlad A, Orgeta V, et al. Dementia prevention, intervention, and care. *Lancet* 2017;390:2673–2734.
3. Doorduijn AS, de van der Schueren MAE, van de Rest O, et al. Olfactory and gustatory functioning and food preferences of patients with Alzheimer's disease and mild cognitive impairment compared to controls: The NUDAD project. *J Neurol* 2020;267:144–152.
4. Beukers MH, Dekker LH, de Boer EJ, et al. Development of the HELIUS food frequency questionnaires: Ethnic-specific questionnaires to assess the diet of a multiethnic population in The Netherlands. *Eur J Clin Nutr* 2015;69:579–584.
5. Corella D, Ordovas JM. Biomarkers: Background, classification and guidelines for applications in nutritional epidemiology. *Nutr Hosp* 2015;31:177–188.

Francisca A. de Leeuw, MD
Alzheimer Center Amsterdam, Department of Neurology
Amsterdam Neuroscience
Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands

Ondine van de Rest, PhD
Division of Human Nutrition and Health
Wageningen University and Research
Wageningen, The Netherlands

Astrid S. Doorduijn, MSC
Alzheimer Center Amsterdam
Department of Neurology
Amsterdam Neuroscience
Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands

Department of Nutrition and Dietetics
Public Health research institute
Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands

Jay L.P. Fieldhouse, MSC
Alzheimer Center Amsterdam
Department of Neurology
Amsterdam Neuroscience
Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands

Marian A.E. de van der Schueren, PhD
Department of Nutrition and Health
HAN University of Applied Sciences
Nijmegen, The Netherlands

Marjolein Visser, PhD
Department of Health Sciences
Faculty of Science

*Vrije Universiteit Amsterdam and the Amsterdam Public Health
Research Institute
Amsterdam, The Netherlands*

*Ellen G.H.M. van den Heuvel, PhD
Friesland Campina
Amersfoort, The Netherlands*

*Philip Scheltens, MD, PhD
Alzheimer Center Amsterdam
Department of Neurology
Amsterdam Neuroscience
Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands*

*Charlotte E. Teunissen, PhD
Neurochemistry Laboratory and Biobank
Department of Clinical Chemistry
Amsterdam Neuroscience*

*Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands*

*Maartje I. Kester, MD, PhD
Department of Neurology
Flevoziekenhuis
Almere, The Netherlands*

*Wiesje M. van der Flier, PhD
Alzheimer Center Amsterdam
Department of Neurology
Amsterdam Neuroscience
Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands*

*Department of Epidemiology and Biostatistics
Amsterdam Neuroscience
Vrije Universiteit Amsterdam
Amsterdam UMC, Amsterdam, The Netherlands*