VU Research Portal

Overwintering fires in boreal forests

Scholten, Rebecca C.; Jandt, Randi; Miller, Eric A.; Rogers, Brendan M.; Veraverbeke, Sander

published in

Nature

2021

DOI (link to publisher)

[10.1038/s41586-021-03437-y](https://doi.org/10.1038/s41586-021-03437-y)

document version

Peer reviewed version

document license

Other

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Scholten, R. C., Jandt, R., Miller, E. A., Rogers, B. M., & Veraverbeke, S. (2021). Overwintering fires in boreal forests. *Nature*, 593(7859), 399-404. <https://doi.org/10.1038/s41586-021-03437-y>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Article citation: Scholten, R.C., Jandt, R., Miller, E.A. *et al.* Overwintering fires in boreal forests. *Nature* **593**, 399–404 (2021). <https://doi.org/10.1038/s41586-021-03437-y>

Overwintering fires in boreal forests

Rebecca C. Scholten^{1*}, Randi Jandt², Eric A. Miller³, Brendan M. Rogers⁴ and Sander Veraverbeke¹

¹ Faculty of Science, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, North Holland, The Netherlands.

² Alaska Fire Science Consortium, University of Alaska, Fairbanks, AK, USA.

³ Bureau of Land Management, Alaska Fire Service, Fort Wainwright, AK 99703, USA.

⁴ Woodwell Climate Research Center, Falmouth, MA, USA.

*Correspondence to: r.c.scholten@vu.nl.

Forest fires are usually viewed within the context of a single fire season, in which weather conditions and fuel supply can combine to create conditions favourable for fire ignition – usually from lightning or human activity – and spread^{1–3}. But some fires exhibit overwintering behaviour, in which fires smoulder through the non-fire season and flare-up in the subsequent spring^{4,5}. Boreal forests, with deep organic soils favourable for smouldering⁶ and accelerated climate warming⁷ may present unusually favourable conditions for overwintering. Still, the extent of overwintering in boreal forests and the

underlying factors influencing overwintering behaviour remain unclear. Here we show that overwintering fires in boreal forests are associated with hot summers generating large fire years and deep burning into organic soils, conditions that have become more frequent in our study areas in recent decades. Our results are based on an algorithm to detect overwintering fires in Alaska, USA, and Northwest Territories, Canada, using field and remote sensing datasets. Between 2002 and 2018, overwintering fires were responsible for 0.8 % of the total burned area; however, in one year this amounted to 38 %. The spatiotemporal predictability of overwintering fires could be leveraged by fire management agencies to facilitate early detection, which may result in reduced carbon emissions and firefighting costs.

Arctic-boreal regions are warming faster than the global average^{7,8}, and are estimated to store more than twice as much carbon as the Earth's atmosphere in their organic soils⁹. Fires are a natural disturbance in boreal forests and release carbon from above- and belowground carbon pools into the atmosphere. A large fraction of the carbon emissions from fires in northern high latitudes originates from belowground carbon pools as these fires often burn deep into organic soils^{10–12}. In a warming climate, boreal fire regimes are intensifying and fires may burn deeper into organic soils and thereby threaten soil carbon reservoirs^{10,13}. Moreover, increasing summer temperatures in northern high latitudes lead to more severe fire weather¹⁴ and more lightning ignitions² that enable fires to burn more area, although regional differences in decadal burned area trends exist and the relatively short length of consistent burned area time series influences the interpretation¹⁵. Containment expenses increase exponentially with fire size, and therefore large fires constitute the majority of the budget allocated to fire management agencies in the USA and Canada^{16,17}. The stagnant fire management budgets¹⁸ in the USA and Canada are under pressure because of the increasing threats of climate warming and continued expansion of dwellings into the wildland

urban interface. Prevention and aggressive initial attack on undesired fires may be a viable way to lower firefighting costs^{16,18}.

Traditionally, the fire season in high latitudes begins with the lightning season in June or early-season human activities like debris burning^{2,3,19}. Once ignited, boreal fires can smoulder in organic soils during periods when weather does not favour flaming spread, and, after days or months, re-emerge under weather conditions that favour flaming^{1,20,21}. Smouldering boreal fires often remain undetected, especially in remote areas, which poses challenges for fire managers¹. Recently, fire managers in Alaska, USA, and the Northwest Territories, Canada, started reporting an increasing number of extreme manifestations of this holdover phenomenon. In such cases, some fires hibernate in deep organic soil layers for seven to eight months during the winter and re-emerge early the next fire season, in what can appear to be a new ignition. Limited and often anecdotal evidence of these overwintering fires exists in recent Alaskan fire management reports and operating plans^{22,23}, as well as news reports²⁴. Overwintering, or “zombie” fires, are an understudied phenomenon in boreal forests and may have severe implications for fire management, human health, and climate^{4,5}.

Detection of large overwintering fires

Overwintering fires typically undergo four temporal stages. Towards the end of a fire season, the fire seemingly stops burning as flaming spread ceases (Fig. 1A). Unnoticed, it smoulders during winter under the snow cover (Fig. 1B). As soon as fire weather facilitates fire spread the following fire season, the fire flames up (Fig. 1C, Extended Data Fig. 1), thereby burning additional area (Fig. 1D). This sequence of events allows for the identification of spatial and temporal characteristics of overwintering fires. First, newly burned areas of an overwintering fire are found

near the original burn scar, and second, they require no additional ignition source and can therefore re-emerge early in the fire season, before the main lightning season.

We developed an algorithm to retrospectively identify and map overwintering fires based on these two spatiotemporal characteristics (Extended Data Fig. 2). We analysed the locations of 45 small overwintering fires reported by fire managers in Alaska and the Northwest Territories between 2005 and 2017 to determine a suitable threshold for the distance between a holdover and its fire of origin. With small fires we refer to fires whose re-emergence remained undetected in the Moderate Resolution Imaging Spectroradiometer (MODIS) active fire product²⁵. Small overwintering fire sizes ranged from 0.04 to 42.5 ha, and 78 % of the fires burned less than 1 ha. We found that 89 % of these small overwintering fires started within the fire perimeter from the year before, and 93 % travelled less than 500 m over the winter (Extended Data Fig. 3). Our results are consistent with laboratory experiments on boreal peat, which have shown that smouldering fires spread around 100 - 250 m/yr (10 - 30 mm/h) depending on oxygen supply and water and mineral contents of the peat⁶. We therefore adopted a threshold of 1000 m to search for overwintering flare-ups in the vicinity of burned area from the year before. The 1000 m threshold was chosen because it is the approximate nadir pixel size of the MODIS active fire product that we used to detect fires.

In Alaska and Northwest Territories, fine fuels are conducive to flaming spread as early as mid- to late May, even before convective thunderstorm activity starts²⁶. The timing of dry fuel availability is dependent on the onset of snowmelt in spring, which varies in time and space²³. Spring snow melt is dependent on winter and spring temperatures and precipitation and therefore a suitable proxy of spring and winter weather conditions, and has been shown to be an effective predictor of fire activity²⁷. To capture the annual variability in dry fuel availability, we based the temporal constraint of our algorithm on the regional yearly snowmelt day, which we calculated from the

MODIS daily fractional snow cover product (MODSCAG)²⁸. We analysed the difference between the regional snowmelt day and the detection dates of the 45 reported overwintering fires, and found that overwintering fires re-emerged on average 27 days (standard deviation: 18.5 days) after the regional snowmelt onset. Regionally, overwintering fires on average re-emerged at the end of May (Julian day: 150, standard deviation: 17.9 days). We therefore used the 90th percentile of 48 days after the regional snowmelt onset as the temporal threshold within the detection algorithm for overwintering fires. Since the satellite product detects fires on average two days later than fire managers, we increased the temporal threshold to a total of 50 days. Lastly, in addition to the spatial and temporal constraints, our algorithm eliminated fires that started close to human infrastructure or close in space and time to a recorded lightning strike (Extended Data Fig. 4). Excluding areas close to infrastructure means some overwintering fires may be missed, but eliminates a larger number of false positives from spring pile burning and other anthropogenic influences around settlements.

In addition to the 45 small reported overwintering fires, which we used for algorithm development, we used a subset of nine larger overwintering fires (mean size: 20312, standard deviation: 24185 ha), which were large enough to be detected by the MODIS active fire product, as validation data for our algorithm (Supplementary Table 1). We extracted the re-emergence date and distance to burn scars of the antecedent year for all ignitions from the Alaskan Fire Emissions Database (AKFED)²⁹, and applied our detection algorithm to them. We detected seven out of nine of the large field-verified overwintering fires.

Furthermore, we identified 20 previously unreported large overwintering fires in Alaska and the Northwest Territories between 2002 and 2018 (Supplementary Table 2). Large overwintering fires constitute 0.8 % of the total burned area and 0.5 % of the total carbon emissions, yet their relative

contribution can be substantial in individual years and amounted to more than 5 % in three years in Alaska (2007, 2008 and 2010) and two years in the Northwest Territories (2002 and 2015). For example, in Alaska in 2008, the contribution of a single overwintering fire that burned 13700 ha amounted to 38 % of the annual burned area.

Temporal drivers of overwintering fires

Years with large annual burned area more frequently produced overwintering fires (Fig. 2). Fire season temperature and annual burned area were strongly correlated for both Interior Alaska (Extended Data Fig. 5, Spearman's $\rho = 0.59$, $p < 0.001$) and the Northwest Territories ($\rho = 0.41$, $p = 0.097$), and we found increasing temperature trends in both areas, and in burned area for Interior Alaska (Fig. 2). Temperature trends differed within regions and the largest warming was observed in western Interior Alaska and central Northwest Territories (Extended Data Fig. 5). Fire season temperature and burned area correlated strongly with the number of overwintering fires in both regions (Fig. 2, Extended Data Fig. 6). Several fires of the large fire years 2009 and 2015 in Alaska and the extreme 2014 fire season in the Northwest Territories overwintered. While burned area in the antecedent year is prerequisite for the occurrence of overwintering fires, we found that fires survived winter following the six hottest summers in the Northwest Territories, whereas overwintering was not observed after the seven coolest summers. Our results based on average fire season temperature are further supported by an analysis that focused on extreme temperatures (Extended Data Figure 7). We found that the number of hot days that surpassed the longer-term 90th percentile of daily maximum temperature during the fire season correlated strongly with annual burned area (Alaska: $\rho = 0.71$, $p < 0.001$; Northwest Territories: $\rho = 0.50$, $p = 0.001$), and with the number of fires that overwintered. Large scale climatic drivers thus govern the survival and growth of overwintering fires. In autumn, fires in boreal regions are

usually extinguished by substantial rain events¹. Extended fire seasons and droughts associated with climate warming^{30,31} may counteract the natural fire extinction in autumn and instead increase the chances of fires entering a smouldering phase. An important driver modulating the emergence of large overwintering fires may therefore be warm and extreme summers that facilitate long and large fire seasons^{31,32}. Within our time series, we found no evidence that winter and spring meteorology or the snowmelt timing influence the survival of large overwintering fires (Extended Data Tables 1 and 2).

Our reference data on overwintering fires contained five times as many small fires that were undetected by satellite imagery, as large, detected fires, suggesting that, when re-emerging in spring, these fires usually remain relatively small and undetected, and only occasionally grow large when fire weather conditions favour fire spread. Large overwintering fires on average experienced more severe fire weather at the time of the flare-up than small overwintering fires (Extended Data Table 3), yet this relationship may partly be confounded by interacting effects of fire spread direction and limited fuel availability in the burned area of the antecedent year.

Spatial drivers of overwintering fires

For a fire to overwinter it needs to burn deep into the organic soil or underneath tree roots so that the organic soil can protect and insulate from adverse winter conditions³³. Severe fires burn deep into the soil organic layers³⁴ and may thus help sustain the smouldering phase of overwintering fires during winter. We analysed burn depth data from AKFED, and found that, on average, fires that promoted overwintering had burned deeper into the organic soil layer than those that stopped burning at the end of the fire season for both Alaska (14.0 cm vs. 12.6 cm, $p = 0.07$) and the Northwest Territories (16.7 cm vs. 14.1 cm, $p = 0.02$) (Fig. 3, Table 1), indicating that deep burning may facilitate overwintering of fires. Regionally, burn depth is correlated with the 90th

percentile of daily maximum temperature in summer in both regions (Extended Data Fig. 7, Alaska: $\rho = 0.56$, $p = 0.02$, Northwest Territories: $\rho = 0.48$, $p = 0.047$). Extreme temperatures have increased since 1979 in western Interior Alaska and central and southern Northwest Territories (Extended Data Fig. 5).

Burn depth in organic layers is co-influenced by fire weather, topographic landscape position and vegetation and soil characteristics^{11,12,35,36}. We compared topographic indicators, pre-fire tree cover and tree species dominance, and carbon in the organic soil layer of fires that produced overwintering fires to those that did not facilitate overwintering (Table 1). Overwintering fires were associated with flat, low-elevation areas, both in Alaska and the Northwest Territories (Fig. 4, Table 1). Lowland terrain in Alaska and the Northwest Territories typically features thick organic soil. Indeed, overwintering fires occur more often in areas with higher carbon contents in the upper soil layer (0 - 30 cm) in Alaska ($p = 0.003$), however, this driver was not significant in the Northwest Territories. Tree cover and species modulate fire severity by their influence on fuel availability and connectivity³⁷. We found that fires that produced overwintering fires have a higher tree cover ($p = 0.001$) and a larger fraction of black spruce ($p = 0.09$) in Alaska, yet these drivers were not significant in the Northwest Territories. Fires occur in more varied landscapes with regard to soil carbon content and forest composition in Alaska compared to the Northwest Territories (Table 1), which may explain why some of these drivers were significant in Alaska but not in the Northwest Territories.

Climate change and fire management

We identified three main drivers of overwintering fires that are influenced directly by climate warming: summer temperature extremes, large annual fire extent and deep burning. Higher temperatures in boreal regions lead to intensified drought and elongated fire seasons³². Longer

fire seasons allow fires to spread faster and grow larger, thereby leading to large area burned³⁸. Summer heat and drought induce deep drying of surface organic fuels, and are thus associated with higher fire severity and deep burning¹². Increasing summer temperatures associated with climate warming may thus promote the survival of overwintering fires in the future. Likewise, earlier onset of spring fire weather conditions may lead to a larger fraction of these fires growing large. At the same time, ecosystem shifts towards a dominance of deciduous vegetation due to increasing fire severity³⁹ and higher temperatures⁴⁰ may constrain the occurrence of overwintering fires in the future. Hence, the fate of overwintering fires in the changing boreal biome will depend on counteracting processes that facilitate or constrain their occurrence.

Overwintering fires are currently a relatively rare phenomenon in boreal forests. Yet, because of their long duration and extended smouldering phase, overwintering fires may substantially influence soil functioning and post-fire recovery trajectories³⁴. We estimated that large overwintering fires in Alaska and the Northwest Territories emitted 3.5 (standard deviation: 1.1) Tg carbon between 2002 and 2018, 64 % of which occurred during the 2015 NWT and 2010 Alaska fire seasons. The contribution of smouldering combustion is generally underestimated in carbon emission estimates from boreal fires²⁰. Thus, our estimate is likely conservative, since overwintering fires exhibit a substantial smouldering phase and may burn deeper than our emissions model currently predicts. In addition, smouldering fires emit relatively more methane and less carbon dioxide in comparison to flaming fires⁴¹, yet methane has a much larger global warming potential⁴².

Carbon emissions from overwintering fires currently contribute 0.5 % of the total carbon emissions from fires in Alaska and Northwest Territories, yet this fraction may grow larger with climate warming. We have shown that overwintering fires have temporal and spatial

predictability. Space- and airborne monitoring of the edges of burn scars from the preceding year in lowland forested peatlands early in the fire season, and especially after a year with large burned area, may prove beneficial for detecting and suppressing flare-ups from overwintering fires while they are small. Fire suppression has shown to be most successful and cost-effective when applied early and on small fires^{16,17}. Out of the 26 overwintering fires for which we had suppression cost data in Alaska, the single largest fire caused 80 % (\$2.2 million) of the total costs incurred by all overwintering fires (Supplementary Tables 1 and 2). Early detection and attack on overwintering fires could thus contribute to savings in the fire management budget that is under increasing pressure¹⁸. In addition, targeted monitoring and early suppression of overwintering fires could help fire managers preserve terrestrial carbon stores when suppression is part of a climate change mitigation strategy.

References:

1. Sedano, F. & Randerson, J. T. Multi-scale influence of vapor pressure deficit on fire ignition and spread in boreal forest ecosystems. *Biogeosciences* **11**, 3739–3755 (2014).
2. Veraverbeke, S. *et al.* Lightning as a major driver of recent large fire years in North American boreal forests. *Nat. Clim. Chang.* **7**, 529–534 (2017).
3. Calef, M. P., McGuire, A. D. & Chapin, F. S. Human influences on wildfire in alaska from 1988 through 2005: An analysis of the spatial patterns of human impacts. *Earth Interact.* **12**, (2008).
4. McCarty, J. L., Smith, T. E. L. & Turetsky, M. R. Arctic fires re-emerging. *Nat. Geosci.* **13**, 656–658 (2020).
5. Irannezhad, M., Liu, J., Ahmadi, B. & Chen, D. The dangers of Arctic zombie wildfires. *Science* (80-.). **369**, 1171 (2020).

- 227 6. Rein, G. Smouldering Fires and Natural Fuels. in *Fire Phenomena and the Earth System:*
228 *An Interdisciplinary Guide to Fire Science* (ed. Belcher, C. M.) 15–34 (2013).
229 doi:10.1002/9781118529539.
- 230 7. Post, E. *et al.* The polar regions in a 2°C warmer world. *Sci. Adv.* **5**, (2019).
- 231 8. Overland, J. E., Wang, M., Walsh, J. E. & Stroeve, J. C. Future Arctic climate changes:
232 Adaptation and mitigation time scales. *Earth's Futur.* **2**, 68–74 (2014).
- 233 9. Tarnocai, C. *et al.* Soil organic carbon pools in the northern circumpolar permafrost
234 region. *Global Biogeochem. Cycles* **23**, 1–11 (2009).
- 235 10. Walker, X. J. *et al.* Increasing wildfires threaten historic carbon sink of boreal forest soils.
236 *Nature* **572**, 520–523 (2019).
- 237 11. Turetsky, M. R. *et al.* Recent acceleration of biomass burning and carbon losses in
238 Alaskan forests and peatlands. *Nat. Geosci.* **4**, 27–31 (2011).
- 239 12. Walker, X. J. *et al.* Soil organic layer combustion in boreal black spruce and jack pine
240 stands of the Northwest Territories, Canada. *Int. J. Wildl. Fire* **27**, 125–134 (2018).
- 241 13. Turetsky, M. R. *et al.* Global vulnerability of peatlands to fire and carbon loss. *Nat.*
242 *Geosci.* **8**, 11–14 (2015).
- 243 14. Flannigan, M. D. *et al.* Fuel moisture sensitivity to temperature and precipitation: climate
244 change implications. *Clim. Change* **134**, 59–71 (2016).
- 245 15. Coops, N. C., Hermosilla, T., Wulder, M. A., White, J. C. & Bolton, D. K. A thirty year,
246 fine-scale, characterization of area burned in Canadian forests shows evidence of
247 regionally increasing trends in the last decade. *PLoS One* **13**, 1–12 (2018).
- 248 16. USDA Forest Service USDI NASF. *Large fire cost reduction action plan.* (2003).
- 249 17. Podur, J. & Wotton, M. Will climate change overwhelm fire management capacity? *Ecol.*

- 250 *Modell*. **221**, 1301–1309 (2010).
- 251 18. Tymstra, C., Stocks, B. J., Cai, X. & Flannigan, M. D. Wildfire management in Canada:
252 Review, challenges and opportunities. *Prog. Disaster Sci.* **5**, 100045 (2020).
- 253 19. Stocks, B. J. *et al.* Large forest fires in Canada, 1959–1997. *J. Geophys. Res.* **108**, (2002).
- 254 20. Wiggins, E. B. *et al.* Evidence for a larger contribution of smoldering combustion to
255 boreal forest fire emissions from tower observations in Alaska. *Atmos. Chem. Phys.*
256 **Preprint**, (2020).
- 257 21. Rein, G., Garcia, J., Simeoni, A., Tihay, V. & Ferrat, L. Smouldering natural fires:
258 Comparison of burning dynamics in boreal peat and Mediterranean humus. *WIT Trans.*
259 *Ecol. Environ.* **119**, 183–192 (2008).
- 260 22. Baber, C. & McMaster, R. *2019 Alaska Statewide Annual Operating Plan. Alaska Master*
261 *Cooperative Wildland Fire Management and Stafford Act Response Agreement Exhibit C*
262 (2019) doi:10.1201/9781315585796-14.
- 263 23. AICC. Wildland Fire Summary and Statistics Annual Report. 1–35 (2010).
- 264 24. Alaska Division of Forestry. State Forestry monitoring hot spots that overwintered from
265 Deshka Landing Fire. [https://akfireinfo.com/2020/04/10/state-forestry-monitoring-hot-](https://akfireinfo.com/2020/04/10/state-forestry-monitoring-hot-spots-that-overwintered-from-deshka-landing-fire/)
266 spots-that-overwintered-from-deshka-landing-fire/ (2020).
- 267 25. Giglio, L., Schroeder, W. & Justice, C. O. The collection 6 MODIS active fire detection
268 algorithm and fire products. *Remote Sens. Environ.* **178**, 31–41 (2016).
- 269 26. Kasischke, E. S., Rupp, T. S. & Verbyla, D. L. Fire trends in the Alaskan boreal forest. in
270 *Alaska’s Changing Boreal Forest* (ed. F.S. Chapin, III, M. Oswood, K.V. Cleve, L. A. V.
271 and D. L. V.) 285–301 (Oxford University Press, 2006).
- 272 27. Westerling, A. L., Hidalgo, H. G., Cayan, D. R. & Swetnam, T. W. Warming and earlier

- spring increase Western U.S. forest wildfire activity. *Science* (80-.). **313**, 940–943 (2006).
28. Painter, T. H. *et al.* Retrieval of subpixel snow covered area, grain size, and albedo from MODIS. *Remote Sens. Environ.* **113**, 868–879 (2009).
29. Scholten, R. C., Jandt, R. R., Miller, E. A., Rogers, B. M. & Veraverbeke, S. ABoVE: Ignitions, burned area and emissions of fires in AK, YT, and NWT, 2001-2018. (2020) doi:<https://doi.org/10.3334/ORNLDAAAC/1812>.
30. Xiao, J. & Zhuang, Q. Drought effects on large fire activity in Canadian and Alaskan forests. *Environ. Res. Lett.* **2**, (2007).
31. Flannigan, M. D. *et al.* Global wildland fire season severity in the 21st century. *For. Ecol. Manag.* **294**, 51–64 (2013).
32. Jolly, W. M. *et al.* Climate-induced variations in global wildfire danger from 1979 to 2013. *Nat. Commun.* **6**, 1–11 (2015).
33. Adams, W. H. *The role of fire in the Alaska taiga. An unsolved problem. Technical report, Unpublished report of the Bureau of Land Management, State Office, Anchorage, Alaska, USA.* (1974).
34. Certini, G. Effects of fire on properties of forest soils: A review. *Oecologia* **143**, 1–10 (2005).
35. Kane, E. S., Kasischke, E. S., Valentine, D. W., Turetsky, M. R. & McGuire, A. D. Topographic influences on wildfire consumption of soil organic carbon in interior Alaska: Implications for black carbon accumulation. *J. Geophys. Res. Biogeosciences* **112**, 1–11 (2007).
36. Hoy, E. E., Turetsky, M. R. & Kasischke, E. S. More frequent burning increases vulnerability of Alaskan boreal black spruce forests. *Environ. Res. Lett.* **11**, 1–11 (2016).

37. Miyanishi, K. & Johnson, E. A. Process and patterns of duff consumption in the mixedwood boreal forest. *Can. J. For. Res.* **32**, 1285–1295 (2002).
38. Kasischke, E. S. & Turetsky, M. R. Recent changes in the fire regime across the North American boreal region - Spatial and temporal patterns of burning across Canada and Alaska. *Geophys. Res. Lett.* **33**, (2006).
39. Johnstone, J. F. *et al.* Factors shaping alternate successional trajectories in burned black spruce forests of Alaska. *Ecosphere* **11**, (2020).
40. Mekonnen, Z. A., Riley, W. J., Randerson, J. T., Grant, R. F. & Rogers, B. M. Expansion of high-latitude deciduous forests driven by interactions between climate warming and fire. *Nat. Plants* **5**, (2019).
41. Andreae, M. O. & Merlet, P. Emission of trace gases and aerosols from biomass burning. *Global Biogeochem. Cycles* **15**, 955–966 (2001).
42. Dean, J. F. *et al.* Methane Feedbacks to the Global Climate System in a Warmer World. *Rev. Geophys.* **56**, 207–250 (2018).
43. Beaudoin, A., Bernier, P. Y., Villemaire, P., Guindon, L. & Guo, X. J. Tracking forest attributes across Canada between 2001 and 2011 using a k nearest neighbors mapping approach applied to MODIS imagery. *Can. J. For. Res.* **48**, 85–93 (2018).

Table 1. Spatial variables differ for fires that produced overwintering fires compared to fires that did not produce overwintering fires. P-values are based on Welch t-test. Analysis is based on all (small and large, reported and newly identified) overwintering fires. References for data sources are given in the methods.

Region	Variable	Source	$\mu_{\text{overwinter}} (\pm \text{S.d.})$	$\mu_{\text{other}} (\pm \text{S.d.})$	p
Alaska	Burn depth	Alaska Fire Emissions Database v3	14.0 cm ($\pm$ 3.6)	12.6 cm ($\pm$ 3.3)	0.067
NWT			16.7 cm ($\pm$ 4.4)	14.1 cm ($\pm$ 3.3)	0.019
Alaska	Elevation	ArcticDEM	214.5 m ($\pm$ 149.5)	402.7 m ($\pm$ 358.7)	< 0.001
NWT			270.5m ($\pm$ 95.4)	356.9 m ($\pm$ 224.0)	0.001
Alaska	Slope	ArcticDEM	2.51° ($\pm$ 2.94)	6.71° ($\pm$ 6.95)	< 0.001
NWT			1.86° ($\pm$ 1.20)	2.90° ($\pm$ 4.08)	0.001
Alaska	Fraction tree cover	MODIS vegetation continuous fields product (MOD44B)	0.37 ($\pm$ 0.16)	0.27 ($\pm$ 0.17)	0.001
NWT			0.23 ($\pm$ 0.09)	0.24 ($\pm$ 0.12)	0.87
Alaska	Fraction black spruce	Fuel Characteristic Classification System	0.35 ($\pm$ 0.25)	0.25 ($\pm$ 0.23)	0.09
NWT		Beaudoin et al. (2018) ⁴³	0.28 ($\pm$ 0.17)	0.23 ($\pm$ 0.16)	0.34
Alaska	Organic carbon content in upper (0-30 cm) soil layer	Northern Circumpolar Soil Carbon Database	14.87 kg/m ² ($\pm$ 7.4)	9.5 ($\pm$ 5.1)	0.003
NWT			8.3 kg/m ² ($\pm$ 5.0)	9.3 ($\pm$ 6.1)	0.42

Fig. 1. Landsat 8 false-colour time series of a 2015 fire in Alaska that generated an overwintering fire in 2016. A burn scar at the end of the fire season (white perimeter, A) had seemingly extinguished but was smouldering under the snow layer (B) until favourable conditions enabled the fire to re-emerge (C) thereby creating additional burned area (blue perimeter, D). Fire perimeters were taken from the Alaska Large Fire Database. The Landsat composites used the spectral bands centred at 2.20 μm (red), 0.86 μm (green) and 0.65 μm (blue). Imagery was plotted in R.

Fig. 2. Temporal drivers of overwintering fires and their long-term trends. Burned area is correlated with the average daily maximum temperature of May – September for Alaska (a, e, Spearman’s $\rho = 0.59$, $p < 0.001$) and Northwest Territories (b, f, $\rho = 0.41$, $p < 0.01$), and is increasing in Alaska ($p = 0.10$). May – September maximum temperatures are increasing in Alaska ($p = 0.04$) and Northwest Territories ($p = 0.01$). The number of large overwintering fires correlates with the May – September temperatures in Alaska (e, $\rho = 0.54$, $p = 0.03$) and Northwest Territories (f, $\rho = 0.77$, $p < 0.001$), and with burned area from the year in Alaska (c, e, $\rho = 0.49$, $p = 0.05$) and Northwest Territories (d, f, $\rho = 0.44$, $p = 0.08$). Large overwintering fires include flare-ups from official reports and additional fires identified by our algorithm. Dashed lines represent significant trends, shaded areas their 95% confidence interval. White areas in a and b refer to the period from 2001 to 2018. Extended Data Figure 6 offers scatterplots of all correlations for visual inspection.

Fig. 3. Burn scars that generate overwintering fires (light gray) had burned deeper into the soil organic layer compared to fire scars that did not generate overwintering fires (dark gray) in Alaska (a, $p = 0.07$) and Northwest Territories (b, $p = 0.02$). Vertical lines represents the median, plus (+) signs the mean, and lower and upper hinges correspond to 25th and 75th percentiles. Whiskers extend up to 1.5 times the interquartile range, sample points beyond that are represented as dots. We included overwintering fires from government reports and our algorithm.

Fig. 4. Overwintering flare-ups (blue dots) predominantly occur in lowland areas of Alaska (a) and the Northwest Territories (b). Small overwintering fires that were not detected by the Moderate Resolution Imaging Spectroradiometer active fire products are represented as small dots. Large fires (reported and identified by our algorithm) are represented by large dots. White areas represent data gaps in the ArcticDEM. Maps were plotted in R.

Methods

Verified overwintering fires. Fire managers in Alaska and Canada routinely document information on all fires detected in their territory. These data are assembled in the Alaska Wildland Fire Maps (AWFP; <https://fire.ak.blm.gov/predsvcs/maps.php>) and the Canadian National Fire Database (CNFD)¹⁹. These databases contain the discovery date and location of fires as well as numerous fire attributes such as the size, end date, estimated costs and fire cause. Fire managers attribute ignition causes based on expert knowledge, ground truth or helicopter data, and other sources such as satellite imagery and lightning data. Causes in the fire databases only include human and lightning sources. With rising awareness of overwintering fires, however, some fire managers sparsely started documenting these re-emerging fires in a separate database. We assembled the timing and location of re-emergence of 54 overwintering fires, 42 in Alaska (AK), USA, and 12 in Northwest Territories (NWT), Canada, from these fire management reports. The key characteristics of the overwintering fires used in our study can be found in Supplementary Table 1. Cost data for the fires in Supplementary Tables 1 and 2 were taken from interagency Incident Status Summaries (209 reports) and provided by the Bureau of Land Management, where available, and supplemented by the estimated costs listed in the AWFP database.

Burned area, ignition locations. burn depth and carbon emissions. We derived the burned area and day of burning for Alaska and Northwest Territories between 2001 to 2018 at 500 m spatial resolution using the Alaskan Fire Emissions Database (AKFED) version 3²⁹, which was updated with input from the Moderate Resolution Imaging Spectroradiometer (MODIS) Collection 6^{2,25}. Daily burned area was retrieved by combining fire perimeter data from the AWFP and the CNFD, and remotely sensed surface reflectance and active fire data from MODIS. After integration of

MODIS Collection 6, total burned area and carbon emissions remained within 5 % of previous estimates^{2,44}.

The location of the first day of burning of a fire marks the ignition point. We therefore extracted the location and timing of ignitions from local minima within the day of burning variable, denoting the earliest burn date, within each fire perimeter. Fires originating from multiple separate ignitions sometimes grow together in a multi-ignition fire complex. Our algorithm therefore allowed for several ignition points per fire perimeter by using a local minimum search radius of 5 km. Although MODIS provides daily coverage of active fires and burned area, the actual ignition location can be obscured if clouds are present, or if a fire starts several hours before the satellite overpass and spreads fast. In these cases, the local minimum contained multiple pixels with the same day of burning. When multiple neighbouring pixels burned at the same day, we estimated the ignition location as the centroid of these pixels and calculated the spatial uncertainty of the ignition locations from the standard deviation in the x and y coordinates of these burned pixels. The spatial standard deviation of the ignition location is as a measure for the ignition location uncertainty. Since the native resolution of the MODIS active fire data is 926.6 m, we added a buffer of 1 km to all ignition locations. For fires with multiple burned pixels on the start date, we extended this 1 km buffer with the spatial standard deviation of the ignition location.

Burn depth and emissions were also derived from AKFED version 3, which predicts carbon consumption and burn depth based on remotely sensed pre-fire tree cover, the differenced Normalized Burn Ratio, and temperature and the drought code at the day of burning using a nonlinear multiplicative regression model ($R^2 = 0.39$)². The model was developed using field observations from black spruce (*Picea mariana*) ecosystems. Primary sources of uncertainties that influence the carbon consumption estimate include the unexplained variance in the regression

model, the underlying land cover classifications, and consumption scaling for non-black spruce ecosystems. To eliminate uncertainties from consumption scaling in our spatial analysis, we excluded burn scars with high dominance (more than 90 %) of ecosystems other than black spruce. Pixel-level uncertainties in carbon consumption were within 20-25 % of the pixel-level predictions⁴⁴.

We used fire perimeter data from CLFD and AAFP to calculate burned area for 1975 to 2000. Since these fire perimeters do not account for unburned islands in the mapped area, we normalised their burned area with AKFED burned area. As in Veraverbeke et al. (2017)², we assumed that uncertainties in fire perimeter mapping have reduced since the integration of Landsat imagery in fire mapping around 1975⁴⁵. The minimum mapping unit (MMU) for the CNFD was 200 ha, and changed over time for AAFP, from 405 ha before 1987 to 40.5 ha between 1987 and 2014, and finally to 4.5 ha starting from 2015. We calculated the ratio of AKFED burned area, which has a MMU of 25 ha, over the ratio of burned area retrieved from the fire perimeters between 2001 and 2018. To remove uncertainties due to the MMU, we calculated separate ratios excluding all perimeters smaller than 405 ha and 40.5 for Alaska, and all fires smaller than 200 ha for the Northwest Territories. We multiplied the burned area estimates with the according ratio per region and, in the case for Alaska, time frame. The derived ratios were 0.971 for the Northwest Territories and 0.829 for fires larger than 405 ha, and 0.825 for fires larger than 40.5 ha in Alaska.

Lightning data and lightning ignition attribution. We acquired data on lightning strikes between 2001 and 2018 detected by the Alaskan Lightning Detection Network (ALDN)⁴⁶ and the Canadian Lightning Detection Network (CLDN)⁴⁷, which contained information on location and timing of cloud-to-ground lightning strikes. The ALDN was started by the Bureau of Land Management Alaska Fire Service (BLM-AFS) in 1976 and has since gradually increased in

detection accuracy, efficiency and coverage. The detection accuracy is highest for interior Alaska and decreases towards the coast. A significant upgrade to the system in 2000 led to an increased detection accuracy and efficiency of 0.5 - 2 km and 80 – 90 %, respectively⁴⁶. The replacement of the Impact lightning system with a Time of Arrival (TOA) system in 2012 resulted in a further 1.5-fold increase in the detection efficiency, and an increased accuracy stemming from the counting of strokes per flash instead of lightning flashes⁴⁸.

Lightning data from the CLDN is available since 1998 and provided by Environment and Climate Change Canada. The CLDN was upgraded gradually, with the largest changes in Northern Canada comprising the addition of two sites in northern Yukon in 2003 and sensor upgrades in NWT and Yukon in 2008 and 2010^{47,49}. For southern NWT, where most of the lightning activity takes place, the CLDN detects approximately 80-90 % of the lightning flashes with a positional accuracy of 500 m⁵⁰. At the periphery of the sensor network, the efficiency decreases to about 70 % with positional accuracies between 12 and 22 km. Lightning detection and accuracy decreases to approximately 30 % 300 km beyond the sensor network. Towards north-eastern NWT, accuracy and efficiency markedly decline due to a lack of sensor sites.

Between 2001 and 2018, the positional accuracies of the ALDN and CLDN vary substantially in time and space. We therefore adapted a conservative estimate of 2 km for the overall accuracy of the sensor networks and buffered all detected lightning strikes in AK and NWT using this 2 km buffer.

We used a spatial and a temporal constraint to assess if an ignition may have been caused by lightning. Both, the ignition locations as well as the lightning locations contain location uncertainties. For the spatial constraint, we thus overlaid all ignition locations including their spatial uncertainty buffer with the buffered lightning strikes of the same year. Subsequently, we

compared the date of the lightning strike with the ignition date. Fires often smoulder for several days after a lightning strike before they are detected, yet the lag time between a lightning strike and the ignition detection in boreal forests of North America is not known. Lag times of two or three days have been inferred for fires in Australia, Finland and Florida, U.S.^{51–53}. We extended the lag time threshold to six days to account for longer holdover times that may occur due to the prolonged smouldering in organic soils in boreal North America. Thus, we classified fires with a lightning strike up to six days before the ignition date as started by lightning. We also accounted for a temporal uncertainty of one day in ignition timing⁵⁴. We thereby identified 85 % of the lightning ignitions as reported in the AWFP and CNFD (Extended Data Figure 4A).

Infrastructure data. We used vector data on roads and other infrastructure elements to assess if an ignition may have been caused by anthropogenic activity. For Alaska, the Alaska Infrastructure 1:63,360 shapefile (2006) provided by the Department of Natural Resources comprises all roads and trails, and power and electrical lines. The same infrastructure elements are available from various infrastructure datasets of NWT including the 2010 Road Network File by Statistics Canada⁵⁵ and the Roads 1M dataset by the Government of Yukon⁵⁶, which we combined here.

In Alaska, 99 % of ignitions up to 5 km from settlements are human-induced³. We used all fires classified as human-ignited by the AWFP and CNFD to derive a distance threshold from our data. First, we calculated the distance between each ignition point including its uncertainty buffer and its nearest infrastructure element for all ignition points that fell within a distance of 5 km of an infrastructure element. Then, we derived a statistical distribution of these distances for all fires that were classified as human-ignited in the official fire databases. 75 % of the ignitions were within 1 km of an infrastructure element (Extended Data Figure 4B). We therefore classified fires that fell within 1 km of an infrastructure element as human-ignited.

Snow cover data. We determined the regional first snow-free day of spring between 2001 and 2018 from the MODIS daily fractional snow cover product (MODSCAG)²⁸. MODSCAG computes the snow fraction of each 500 m pixel using spectral mixture analysis and has shown to outperform normalized difference snow index-based methods, especially during periods of accumulation and melt⁵⁷. We flagged a pixel as snow-free when its fractional snow cover dropped below 15 %. We determined the period between March 21 (Julian day 80) and July 1 (Julian day 182) as spring season and selected the first snow-free day of each pixel during this period. Pixels that were still snow covered by July 1 were flagged as permanent snow cover and excluded from the analysis. We also excluded pixels with persistent missing data due to cloud cover on four or more days preceding the first snow-free day detection. The resulting retrieval contained data for 98 % of interior Alaska and 87 % of interior Northwest Territories.

For a regional estimate, we calculated the yearly mean of the first snow-free day within the interior boreal regions of Alaska and the Northwest Territories. For Alaska this refers to the intermontane boreal ecoregions between the Brooks Range and the Alaska Range, excluding the coastal Bering ecoregions⁵⁸. For NWT we selected the taiga plains and taiga shield ecozones in Northwest Territories⁵⁹. Only pixels that contained data for all years and had not burned during the 18-year timeframe were included in the regional mean.

Climate and fire weather data. We extracted meteorological data from North America Regional Reanalysis⁶⁰ (NARR) for our climate analysis. NARR provides climate reanalysis data since 1979 at a 32 km resolution based on the NCEP Eta atmospheric model and the Regional Climate Data Assimilation System. We extracted 3-hourly air temperature at 2m, relative humidity, wind speed and precipitation over Alaska and the Northwest Territories, and calculated monthly means of the 3-hour period that included local solar noon. We derived vapour pressure deficit (VPD) and fire

weather variables following the Canadian Fire Weather Index System (CFWIS)⁶¹ from meteorological variables.

Detection of large overwintering fires. 45 of the ground-truthed overwintering fires (10 from NWT and 35 from AK, in the following referred to as ‘small fires’) were too small to be detected from the MODIS active fire product that was used within AKFED²⁵ (Supplementary Table 1). We used the spatial and temporal characteristics of these 45 small fires to derive spatial and temporal thresholds for a detection algorithm for larger overwintering fires that can be detected from MODIS imagery. The nine remaining overwintering fires from the fire management reports were large enough to be detected by MODIS and were used as reference data for validation of the detection algorithm (Extended Data Fig. 2).

Overwintering fires re-emerge within or in close proximity to burned area from the year before and earlier in the year than the majority of lightning- and human-ignited fires. We calculated the shortest distance between each of the 45 small overwintering fire locations reported by fire managers and any area burned in the previous year based on our burned area product and derived a threshold of 1 km based on the statistical distribution of these distances (Extended Data Figure 3) and the spatial resolution of our satellite product. Distributions of the difference between the detection date of the small overwintering fires and the regional snowmelt served for a temporal threshold. We chose a threshold of 48 days, which comprises the 90 % quantile of the distribution. On average, fires are detected by our satellite product within 1.7 days of the discovery date of the fire agencies. We therefore extended the threshold by two days to account for the differences in data sources. We applied both thresholds to all ignitions detected by MODIS between 2002 and 2018 to identify potential overwintering fires. From these, we further excluded ignitions that were likely caused by lightning by filtering out all ignitions in spatiotemporal vicinity of a lightning

strike. We intersected ignitions and lightning strikes including their spatial uncertainties (2 km for all lightning strikes and the individual positional inaccuracy of each ignition) and allowed for a lag time of six days between lightning strikes and ignition in combination with an uncertainty of one day in the ignition timing. We also excluded ignitions with a likely human origin when these occurred with 1 km of infrastructure, thereby accounting for the spatial uncertainty of the ignitions location.

Uncertainty of our algorithm. Our estimate of the number of overwintering fires based on these four constraints and moderate resolution satellite data is likely conservative. For the Northwest Territories, for example, some estimates suggest that about one third of all fires in 2015 were caused by overwintering flare-ups⁶². Our algorithm however only classified 4 % of the ignitions to be overwintering fires, although 17.5 % of the ignitions were within a 1 km distance from a previous year fire. Many of these ignitions occurred close to a human infrastructure element or late in the season and were therefore removed by our algorithm to avoid false positives. However, our reference data on overwintering fires suggest that 35 % of the small fires are indeed found within our infrastructure buffer, and emergence dates as late as July have been reported by fire managers.

Furthermore, many overwintering fires occur in unburned islands or stay relatively small and are therefore not detected by the MODIS active fire product. The Visible Infrared Imaging Radiometer Suite (VIIRS) active fire detection data product⁶³ has a higher spatial resolution of 375 m and is therefore capable of detecting smaller fires. Indeed, using VIIRS data we could detect a further 8 of the 31 overwintering fires that were too small to be detected by MODIS. However, VIIRS data are only available from 2012 onward, which renders it less useful than MODIS for the analysis of longer time periods.

Spatial drivers of overwintering fires. We extracted burn depth from AKFED for all burn scars. We excluded burn scars with high dominance (more than 90 %) of white spruce, pine and deciduous ecosystems because the burn depth model was developed for black spruce ecosystems. We tested the statistical difference in mean burn depth between burn scars that produced overwintering fires and those that did not using Welch's t-test^{64,65}. We thereby assumed that overwintering fires were caused by the closest fire of the previous year.

A variety of datasets were used to analyse additional spatial drivers. The analysis was carried out analogous to the burn depth analysis by comparing the mean over the entire burn scar between fires that produced overwintering fires and those that did not using Welch's t-test. We extracted the elevation and slope for all burn scars from the 100 m resolution ArcticDEM v3.0^{66,67}. ArcticDEM provides high-resolution (up to 2 m) digital surface models of the Arctic from 0.32 to 0.5 m resolution panchromatic satellite imagery of the DigitalGlobe collection including WorldView-1 (2007), WorldView-2 (2009), WorldView-3 (2014), and GeoEye-1 (2008)⁶⁸. Annual Terra MODIS Vegetation Continuous Fields Collection 6 data at 250 m resolution (MOD44B)⁶⁹ for the years 2000-2017 were used to derive pre-fire tree cover for each burn scar. Tree species fractions were taken from the Fuel Characteristic Classification System layer of the year 2001⁷⁰⁻⁷² for Alaska, and from^{43,73} for the Northwest Territories. We aggregated the tree species into black spruce (*Picea mariana*), white spruce (*Picea glauca*), deciduous, tundra-grass-shrub and non-vegetated ecosystems, and pine (only present in the Northwest Territories) as described in². Organic carbon content in the upper organic soil layer (0-30 cm depth) was extracted from the Northern Circumpolar Soil Carbon Database⁷⁴.

Temporal drivers of overwintering fires. We analysed the relationship between the number of overwintering fires, annual burned area and daily maximum temperatures from NARR on a

regional scale. Based on scatterplots between all three variables, we chose Spearman correlations because of non-linearity (Extended Data Fig. 6a, b, e), the presence of outliers (Extended Data Fig. 6d) and small sample sizes (Extended Data Fig. 6 c-f, n = 17). P-values were computed for all correlations.

To analyse the influence of winter and spring weather, we computed Spearman's correlations between overwintering fires and the regional snowmelt, as well as winter and spring temperature, vapour pressure deficit, precipitation and relative humidity. Analogous to the spatial drivers analyses, we also tested for differences in the snowmelt date and fire weather variables in spring between fire scars that facilitated overwintering, and those that did not using Welsh's t-test. For the fire weather variables, we hereby took the average of the 50 days after the average snowmelt of each fire. We further compared vapour pressure deficit and fire weather variables at the day of detection for small and large overwintering fires using Welsh's t-test to assess the influence of spring fire weather on the growth of overwintering fires.

References:

44. Veraverbeke, S., Rogers, B. M. & Randerson, J. T. Daily burned area and carbon emissions from boreal fires in Alaska. *Biogeosciences Discuss.* **11**, 17579–17629 (2015).
45. Kasischke, E. S. *et al.* Quantifying burned area for North American forests: Implications for direct reduction of carbon stocks. *J. Geophys. Res. Biogeosciences* **116**, 1–17 (2011).
46. Farukh, M. A. & Hayasaka, H. Active Forest Fire Occurrences in Severe Lightning Years in Alaska. *J. Nat. Disaster Sci.* **33**, 71–84 (2012).
47. Burrows, W. R. & Kochtubajda, B. A decade of cloud-to-ground lightning in Canada: 1999-2008. Part 1: Flash density and occurrence. *Atmos. - Ocean* **48**, 177–194 (2010).

48. Bieniek, P. A. *et al.* Lightning Variability in Dynamically Downscaled Simulations of Alaska's Present and Future Summer Climate. *J. Appl. Meteorol. Climatol.* **59**, 1139–1152 (2020).
49. Kochtubajda, B. *et al.* Exceptional cloud-to-ground lightning during an unusually warm summer in Yukon, Canada. *J. Geophys. Res. Atmos.* **116**, 1–20 (2011).
50. Kochtubajda, B., Stewart, R. & Tropea, B. Lightning and Weather Associated with the Extreme 2014 Wildfire Season in Canada's Northwest Territories. in *24th International Lightning Detection Conference* 1–4 (2016).
51. Dowdy, A. J. & Mills, G. A. Atmospheric and fuel moisture characteristics associated with lightning-attributed fires. *J. Appl. Meteorol. Climatol.* **51**, 2025–2037 (2012).
52. Larjavaara, M., Pennanen, J. & Tuomi, T. J. Lightning that ignites forest fires in Finland. *Agric. For. Meteorol.* **132**, 171–180 (2005).
53. Duncan, B. W., Adrian, F. W. & Stolen, E. D. Isolating the lightning ignition regime from a contemporary background fire regime in east-central Florida, USA. *Can. J. For. Res.* **40**, 286–297 (2010).
54. Veraverbeke, S. *et al.* Mapping the daily progression of large wildland fires using MODIS active fire data. *Int. J. Wildl. Fire* **23**, 655–667 (2014).
55. Statistics Canada. Road Network File 2010.
<https://www150.statcan.gc.ca/n1/en/catalogue/92-500-X> (2016).
56. Government of Yukon. Corporate Spatial Warehouse.
ftp://ftp.geomaticsyukon.ca/GeoYukon/Transportation/Roads_1M/ (2018).
57. Rittger, K., Painter, T. H. & Dozier, J. Assessment of methods for mapping snow cover from MODIS. *Adv. Water Resour.* **51**, 367–380 (2013).

58. Gallant, A. L., Binnian, E. F., Omernik, J. M. & Shasby, M. B. *Ecoregions of Alaska. USGS Professional Paper 1567*. (1995) doi:0160482909\r0607010010.
59. Canadian Council on Ecological Areas (CCEA). Canada Ecozones. <https://www.ccea.org/ecozones-downloads> (2016).
60. Mesinger, F. *et al.* North American regional reanalysis. *Bull. Am. Meteorol. Soc.* **87**, 343–360 (2006).
61. Van Wagner, C. E. *Development and structure of the Canadian Fire Weather Index System. Forestry Technical Report* vol. 35 (1987).
62. Alaska Fire Science Consortium. *Opportunities to Apply Remote Sensing in Boreal/Arctic Wildfire Management & Science: A Workshop*. (2019).
63. Schroeder, W., Oliva, P., Giglio, L. & Csiszar, I. A. The New VIIRS 375m active fire detection data product: Algorithm description and initial assessment. *Remote Sens. Environ.* **143**, 85–96 (2014).
64. Welch, B. L. The Significance of the Difference Between Two Means when the Population Variances are Unequal. *Biometrika* **29**, 350–362 (1938).
65. Welch, B. L. The generalization of ‘Student’s’ problem when several different population variances are involved. *Biometrika* vol. 34 28–35 (1947).
66. Morin, P. *et al.* ArcticDEM; A Publically Available, High Resolution Elevation Model of the Arctic. *Geophys. Res. Abstr.* **18**, 2016–8396 (2016).
67. Porter, C. *et al.* ArcticDEM. *Harvard Dataverse* (2018) doi:<https://doi.org/10.7910/DVN/OHHUKH>.
68. Dai, C., Durand, M., Howat, I. M., Altenau, E. H. & Pavelsky, T. M. Estimating River Surface Elevation From ArcticDEM. *Geophys. Res. Lett.* **45**, 3107–3114 (2018).

69. Hansen, M. C. *et al.* Global Percent Tree Cover at a Spatial Resolution of 500 Meters: First Results of the MODIS Vegetation Continuous Fields Algorithm. *Earth Interact.* **7**, 1–15 (2003).
70. Pettinari, M. L. & Chuvieco, E. Generation of a global fuel data set using the Fuel Characteristic Classification System. *Biogeosciences* **13**, 2061–2076 (2016).
71. Ottmar, R. D., Sandberg, D. V, Riccardi, C. L. & Prichard, S. J. An overview of the Fuel Characteristic Classification System - Quantifyin, classifying, and creating fuelbeds for resource planning. *Can. J. For. Res.* **37**, 2383–2393 (2007).
72. Riccardi, C. L. *et al.* The fuelbed: a key element of the Fuel Characteristic Classification System. *Can. J. For. Res.* **37**, 2394–2412 (2007).
73. Beaudoin, A., Bernier, P. Y., Villemaire, P., Guindon, L. & Guo, X. . Species composition, forest properties and land cover types across Canada’s forests at 250m resolution for 2001 and 2011. *Nat. Resour. Canada, Can. For. Serv. Laurentian For. Centre, Quebec, Canada* (2017) doi:10.23687/ec9e2659-1c29-4ddb-87a2-6aced147a990.
74. Hugelius, G. *et al.* The northern circumpolar soil carbon database: Spatially distributed datasets of soil coverage and soil carbon storage in the northern permafrost regions. *Earth Syst. Sci. Data* **5**, 3–13 (2013).

Acknowledgments. We thank the three anonymous reviewers and the editor for their suggestions on the initial manuscript. We would like to thank Cheryl Van Der Horn and Gabriella Branson (Alaska Interagency Coordination Center), and Matthew Coyle (Forest Management Division, Department of Environment and Natural Resources, Government of the Northwest Territories) for providing ground truth data on overwintering fires. We wish to thank Environment and Climate Change Canada for their generous permission to use Canadian Lightning Detection Network data,

and the Bureau of Land Management, Alaska Fire Service, for providing cost information. We thank NASA JPL's Snow Data Center for making their MODSCAG data available. This work was funded by the Netherlands Organization for Scientific Research (NWO) through a Vidi grant (Fires Pushing Trees North) awarded to S.V. B.M.R. acknowledges support from the National Aeronautics and Space Administration (NASA) Arctic-Boreal Vulnerability Experiment (ABOVE; NNX15AU56A).

Author contributions. S.V. and R.C.S. designed the research. R.C.S. performed the analysis with input from S.V.. B.M.R. contributed to the interpretation of cost data. R.R.J. and E.A.M. contributed to the interpretation of field data. R.C.S. drafted the manuscript. All authors participated in manuscript editing.

Competing interests. The authors declare no competing interests.

Data availability. The location and timing of ignition of the overwintering fires used in this study are found in the Supplementary material. Daily burned area, emissions and ignitions data for Alaska, and the Northwest Territories are archived at the Oak Ridge National Laboratory Distributed Active Archive Center for biogeochemical dynamics (<https://doi.org/10.3334/ORNLDAAAC/1812>). Lightning data is available from the Alaska Interagency Coordination Center (<https://fire.ak.blm.gov/predsvcs/maps.php>) and from Environment and Climate Change Canada. Infrastructure data is available for Alaska from the Department of Natural Resources (<https://catalog.data.gov/dataset/alaska-infrastructure-1-63360>), and for the Northwest Territories from Statistics Canada (<https://www150.statcan.gc.ca/n1/en/catalogue/92-500-X>) and the Government of Yukon (https://hub.arcgis.com/datasets/322b6cf3fa1444c289a1d611a4778ead_42/data). MODSCAG snow fraction data is freely available from the JPL Snow Data Server

(<http://snow.jpl.nasa.gov/portal/>). All climate data used in this study is available from the North America Regional Reanalysis (<https://psl.noaa.gov/data/gridded/data.narr.html>). All data used for the analysis of spatial drivers is freely available, including the ArcticDEM (<https://doi.org/10.7910/DVN/OHHUKH>), Northern Circumpolar Soil Carbon Database v2 (<https://doi.org/10.5879/ECDS/000000002>) and Fuel Characteristic Classification System (<https://www.landfire.gov/fccs.php>).

Code Availability. Code used to analyse the data is available from <https://github.com/screbec/Overwintering-fires> or <https://doi.org/10.5281/zenodo.4549321>.

Extended Data:

Extended Data Fig. 1. Aerial view of the Seven Mile Slough Fire in Alaska on 9 May, 2011.

Smouldering hotspots (a) had overwintered and burned in the duff layer below the spruces of an unburned island. Green tree crowns of the fallen trees (b) in the original unburned island (perimeter approximated in black) suggest that tree roots were damaged due to subsurface burning. (Photo by Eric Miller)

Extended Data Fig. 2. Workflow used to detect large overwintering fires. First, ignition locations, dates, and causes according to official fire databases were extracted. In four steps, the algorithm filters these ignitions by date, distance to an old fire scar, and co-occurrence of lightning strikes and infrastructure elements. Small overwintering fires that were not detected by satellite products were used to derive thresholds for the algorithm.

Extended Data Fig. 3. Overwintering fires emerge earlier after the seasonal snowmelt (a) and closer to a fire scar from the year before (b) than other fires. Other fires refer to all fires not classified as overwintering in official fire databases. Day since regional snowmelt was calculated from the timing of the ignition points from the Alaskan Fire Emissions Database when possible, complemented with data from government sources for small fires.

Extended Data Fig. 4. Histograms of (a) lag time between lightning strikes and ignition detections and (b) distance to road for human ignitions. Human and lightning ignitions were characterized based on the Alaskan Wildland Fire Maps (AK) and Canadian National Fire Database (NWT). The black lines indicate the thresholds used to eliminate potential overwintering fires due to spatial proximity to infrastructure and spatiotemporal proximity to lightning strikes.

Extended Data Fig. 5: Average and extreme temperature trends for interior Alaska and the Northwest Territories a, b, Average of the daily maximum temperature of the summer months May – September. c, d, Its 90th percentile. e, f, Number of hot days surpassing the 90th percentile. Panels a, c, e show data for interior Alaska, and panels b, d, f for the taiga plains and taiga shield of the Northwest Territories.

Extended Data Fig. 6: Scatter plots and Spearman correlations of summer temperature, burned area and overwintering flare-ups. a, b, Daily mean maximum temperature between May and September (MJJAS) and annual burned area. c, d, Antecedent year's burned area and the number of overwintering flare-ups. e, f, MJJAS maximum temperature and the number of overwintering flare-ups. Panels a, c, e show data for Alaska, and panels b, d, f for the Northwest Territories.

Extended Data Fig. 7: Scatter plots and Spearman correlations of temperature extremes and burned area, overwintering flare-ups and burn depth. a, b, Number of MJJAS hot days (days with a maximum temperature hotter than the 1979-2020 90th percentile) and burned area. c, d, Number of MJJAS hot days and overwintering ignitions. e, f, 90th percentile of MJJAS temperature and average burn depth. Panels a, c, e show data for Alaska, and panels b, d, f for the Northwest Territories.

Extended Data Tab. 1: Correlation of winter and spring meteorology with the number of overwintering flare-ups in Alaska and Northwest Territories. Results that are significant on a 0.1 level are shaded light grey, and those on a 0.05 level dark grey.

Region	Variable	-	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May
Alaska	Regional snow melt	-0.22								
Northwest Territories		-0.06								
Alaska	Average temperature		0.04	0.17	0.05	-0.05	0.16	0.22	0.18	0.08
Northwest Territories			0.37	-0.06	0.08	0.24	0.11	0.29	-0.14	0.22
Alaska	Vapour Pressure Deficit (VPD)		-0.24	-0.07	0.23	-0.08	0.06	0.06	0.1	0.3
Northwest Territories			0.19	-0.07	-0.03	0.21	0.22	0	-0.41	0.37
Alaska	Total precipitation		-0.38	0.06	-0.23	0.13	-0.07	0.08	-0.2	-0.02
Northwest Territories			-0.16	0.28	0.01	-0.09	0.05	0.06	0	-0.08
Alaska	Relative humidity		0.48	0.12	-0.07	0.22	0.06	0.18	-0.31	0.12
Northwest Territories			-0.11	-0.47	0.11	0.21	-0.11	0.04	0.16	0.04

Extended Data Tab. 2: Average first snow-free day, vapour pressure deficit (VPD) and moisture codes and fire danger indices for days 0 to 50 after the snowmelt did not differ significantly for burn scars that produced overwintering fires and those that did not. P-values are based on Welch t-test. Analysis is based on all (small and large, reported and newly identified) overwintering fires.

Region	Variable	$\mu_{\text{overwinter}} (\pm \text{s.d.})$	$\mu_{\text{other}} (\pm \text{s.d.})$	<i>p</i>
Alaska	First snow-free day	117.9 ($\pm$ 13.6)	122 ($\pm$ 12.3)	0.23
Northwest Territories		128.7 ($\pm$ 9.3)	130.8 ($\pm$ 12.5)	0.35
Alaska	Vapour Pressure Deficit (VPD)	836.8 ($\pm$ 176.2)	804.5 ($\pm$ 210)	0.45
Northwest Territories		1125.8 ($\pm$ 295.3)	1115.2 ($\pm$ 246.8)	0.88
Alaska	Fine Fuel Moisture Code (FFMC)	77.7 ($\pm$ 5)	76.5 ($\pm$ 5.6)	0.31
Northwest Territories		81.9 ($\pm$ 3.8)	80.7 ($\pm$ 4.7)	0.19
Alaska	Duff Moisture Code (DMC)	21.8 ($\pm$ 8.2)	21.9 ($\pm$ 10)	0.97
Northwest Territories		34 ($\pm$ 13.4)	33.6 ($\pm$ 13.9)	0.9
Alaska	Drought Code (DC)	121.9 ($\pm$ 36.6)	134.9 ($\pm$ 39.7)	0.16
Northwest Territories		166.1 ($\pm$ 35.7)	172.1 ($\pm$ 46.5)	0.48
Alaska	Initial Spread Index (ISI)	3.8 ($\pm$ 1.2)	3.8 ($\pm$ 1.3)	0.92
Northwest Territories		6.1 ($\pm$ 1.9)	6 ($\pm$ 1.8)	0.93
Alaska	Buildup Index (BUI)	14.6 ($\pm$ 5.2)	14.8 ($\pm$ 6.6)	0.84
Northwest Territories		22.6 ($\pm$ 8.4)	23 ($\pm$ 10.3)	0.84
Alaska	Fire Weather Index (FWI)	5.2 ($\pm$ 2.3)	5.2 ($\pm$ 2.7)	0.95
Northwest Territories		9.7 ($\pm$ 4.1)	9.6 ($\pm$ 4.2)	0.9
Alaska	Daily Severity Rating (DSR)	0.9 ($\pm$ 0.6)	0.9 ($\pm$ 0.7)	0.89
Northwest Territories		2.4 ($\pm$ 1.4)	2.3 ($\pm$ 1.6)	0.97

738 **Extended Data Tab. 3: Moisture codes and fire danger indices at the day of detection by**
739 **the AKFED product for overwintering fires smaller and larger than 1 km².** Bold numbers
740 represent significant differences at $p < 0.1$.

Region	Variable	$\mu_{\text{small}} (\leq 1 \text{ km}^2) (\pm \text{S.d.})$	$\mu_{\text{large}} (> 1 \text{ km}^2) (\pm \text{S.d.})$	p
Alaska	Vapour Pressure Deficit (VPD)	928.7 ($\pm$ 483.6)	1138.4 ($\pm$ 515.3)	0.43
Northwest Territories		1634.2 ($\pm$ 654.7)	1610.7 ($\pm$ 528.8)	0.95
Alaska	Fine Fuel Moisture Code (FFMC)	83.5 ($\pm$ 7.3)	88.2 ($\pm$ 3.8)	0.14
Northwest Territories		84.8 ($\pm$ 6.5)	82.9 ($\pm$ 19.4)	0.68
Alaska	Duff Moisture Code (DMC)	25.3 ($\pm$ 20.0)	34.8 ($\pm$ 19.4)	0.36
Northwest Territories		62.0 ($\pm$ 44.3)	62.8 ($\pm$ 35.7)	0.97
Alaska	Drought Code (DC)	127.4 ($\pm$ 54.4)	156.6 ($\pm$ 41.3)	0.26
Northwest Territories		277.2 ($\pm$ 87.7)	271.3 ($\pm$ 68.7)	0.91
Alaska	Initial Spread Index (ISI)	4.6 ($\pm$ 3.0)	8.7 ($\pm$ 4.1)	0.05
Northwest Territories		7.1 ($\pm$ 5.4)	7.5 ($\pm$ 5.7)	0.90
Alaska	Buildup Index (BUI)	16.4 ($\pm$ 11.6)	22.4 ($\pm$ 11.3)	0.32
Northwest Territories		39.6 ($\pm$ 26.2)	52.5 ($\pm$ 38.6)	0.49
Alaska	Fire Weather Index (FWI)	6.1 ($\pm$ 4.6)	13.2 ($\pm$ 7.3)	0.05
Northwest Territories		15.1 ($\pm$ 14.1)	18.4 ($\pm$ 16.7)	0.71
Alaska	Daily Severity Rating (DSR)	0.9 ($\pm$ 1.0)	3.1 ($\pm$ 2.7)	0.08
Northwest Territories		4.8 ($\pm$ 7.0)	7.1 ($\pm$ 8.6)	0.61

741