VU Research Portal

Auxiliary subunits of the AMPA: The Shisa family of proteins

Abdollahi Nejat, Mazyar; Klaassen, Remco V.; Spijker, Sabine; Smit, August B.

published in

Current Opinion in Pharmacology
2021

DOI (link to publisher)

[10.1016/j.coph.2021.03.001](https://doi.org/10.1016/j.coph.2021.03.001)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Abdollahi Nejat, M., Klaassen, R. V., Spijker, S., & Smit, A. B. (2021). Auxiliary subunits of the AMPA: The Shisa family of proteins. *Current Opinion in Pharmacology*, 58, 52-61. <https://doi.org/10.1016/j.coph.2021.03.001>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Auxiliary subunits of the AMPA receptor: The Shisa family of proteins

Mazyar Abdollahi Nejat, Remco V. Klaassen,
Sabine Spijker and August B. Smit

Abstract

AMPA receptors mediate fast synaptic transmission in the CNS and can assemble with several types of auxiliary proteins in a spatio-temporal manner, from newly synthesized AMPA receptor tetramers to mature AMPA receptors in the cell membrane. As such, the interaction of auxiliary subunits with the AMPA receptor plays a major role in the regulation of AMPA receptor biogenesis, trafficking, and biophysical properties. Throughout the years, various ‘families’ of proteins have been identified and today the approximate full complement of AMPAR auxiliary proteins is known. This review presents the current knowledge on the most prominent AMPA-receptor-interacting auxiliary proteins, highlights recent results regarding the Shisa protein family, and provides a discussion on future research that might contribute to the discovery of novel pharmacological targets of auxiliary subunits.

Addresses

Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive Research, Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

Corresponding author: Smit, August B. (guus.smit@vu.nl)

Current Opinion in Pharmacology 2021, 58:52–61

This review comes from a themed issue on **Neurosciences: Ion channel accessory proteins**

Edited by **Daniel Bertrand** and **Guss Smith**

For complete overview about the section, refer [Neurosciences: Ion channel accessory proteins](#)

Available online 21 April 2021

<https://doi.org/10.1016/j.coph.2021.03.001>

1471-4892/© 2021 Published by Elsevier Ltd.

Keywords

Ionotropic glutamate receptor, Auxiliary proteins, Modulation of channel function, Receptor targeting.

Introduction

α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors (AMPA) are ionotropic glutamate receptors expressed at the neuronal postsynaptic membrane and mediate fast synaptic transmission in the CNS [1]. AMPAR subunit composition, phosphorylation state, and changes in the number are well known to directly affect

synaptic transmission and plasticity [2]. In addition, AMPARs have been found to associate with a wide range of auxiliary proteins, which play a critical role in regulating intracellular trafficking, expression at the synaptic membrane, and AMPAR functional properties [3]. Differential expression of AMPAR core subunits and their co-assembly with auxiliary proteins dramatically increases receptor diversity and modulatory potential in the CNS [4,5]. The first family of AMPAR auxiliary subunits identified were the transmembrane AMPAR regulatory proteins (TARPs), of which TARP $\gamma 2$ (stargazin) was found to be absent in the severely ataxic stargazer mouse mutant [6,7]. Since then, the advent of interaction proteomics has enabled the identification of a full complement of AMPAR associated proteins [8,9].

In 2010, interaction proteomics led to the identification of Shisa9/CKAMP44 as a novel AMPAR-interacting protein [10], whose structure and function were distinct from other established auxiliary subunits [11,12]. More specifically, Shisa9 was characterized as a single-pass transmembrane protein that induced fast and strong desensitization of AMPAR currents within the hippocampus. In the past few years, several additional members of the Shisa family have been reported to exert a role as AMPAR modulatory proteins, namely Shisa6/CKAMP52 [13–15], Shisa7/CKAMP59 [13,16], and Shisa8/CKAMP39 [13]. Although the Shisa family members share a high structural similarity, they are very distinct in both their (brain) regional expression profile and their modulation of AMPAR function [13,17]. This review provides a short overview of the known AMPAR auxiliary subunits including their function and effect on AMPARs while highlighting the Shisa protein family.

Auxiliary subunits

AMPA receptors assemble with a wide range of auxiliary subunits during their life cycle [3,5,18], from the moment of synthesis in the endoplasmic reticulum (ER) to full integration at postsynaptic sites [5]. Multiple studies using electrophysiology, proteomics, imaging, and mutagenesis have elucidated the many important functions of these proteins [5,18]. Apart from AMPAR assembly, trafficking, and expression at the synapse, auxiliary subunits are directly involved in the regulation of the receptor’s biophysical properties [3,5].

Stargazin was identified as the first transmembrane protein to interact with the AMPAR [19]. Stargazin (Transmembrane AMPAR Regulatory Protein (TARP) $\gamma 2$) acts during AMPAR biogenesis between tetramerization and ER exit, is essential for receptor maturation, and affects AMPAR surface expression [20–23]. Other members of the TARP family have subsequently been identified as AMPAR-interacting proteins: Type-I TARPs (subunits $\gamma 2$ - $\gamma 4$ and $\gamma 8$) and type-II TARPs (subunits $\gamma 5$ and $\gamma 7$) [21,24–26]. In addition to AMPAR maturation and trafficking, TARPs affect the pharmacological and gating properties of the receptor, as observed in both *in vitro* (Table 1) and *ex/in vivo* (Table 2) studies [25,27]. The effects of the TARP proteins on AMPAR conductance properties are specific for each family member and are subunit-dependent [28,29]. Importantly, TARPs have been demonstrated to directly associate with PSD-95 [19,30,31], one of the key organizers of the postsynaptic density (PSD) scaffold. This interaction regulates AMPAR surface trafficking and is essential for both long-term potentiation (LTP) and long-term depression (LTD) in the hippocampus [30,32]. In particular, TARP $\gamma 2$ phosphorylation facilitates the interaction with PSD-95 and regulates lateral diffusion of AMPARs [32–35]. In addition, TARP $\gamma 8$ phosphorylation was found to be crucial for CaMKII-dependent LTP [36].

In recent years, major efforts have been undertaken to identify the full complement of AMPAR-interacting proteins, using immunopurification of brain-derived AMPAR complexes and mass spectrometric analysis [8,9,37]. From these efforts stem the establishment of Cornichon homolog proteins (CNIHs) as an additional auxiliary subunit family [37]. These four-transmembrane domain proteins are expressed widely throughout the brain, and play a role in controlling the export of AMPARs from ER [38–40]. Moreover, CNIHs modulate AMPAR activation, deactivation, desensitization, and synaptic transmission (Tables 1 and 2) [3,37,41–43]. Remarkably, studies have found the association of either TARP $\gamma 2$ or $\gamma 8$ with CNIH-2 at hippocampal AMPARs, suggesting an interplay between TARPs and CNIHs [39,41,44].

Using interaction proteomics analysis of the germline-specific gene 1 like (GSG1L), a distant homolog of TARPs was identified as auxiliary subunits of the AMPAR [8,45]. GSG1L directly interacts with AMPARs at synapses and modulates AMPAR trafficking, surface expression, and gating properties (Tables 1 and 2; Figure 1) [8,45–49]. Synapse differentiation induced gene 1 (SynDIG1) was first identified as an AMPAR-interacting transmembrane protein that regulates the development, strength, and number of excitatory synapses [50]. However, since SynDIG1 does not directly affect biophysical properties or AMPAR surface trafficking it was later suggested to serve as a regulatory protein for excitatory synaptogenesis [51]. SynDIG4,

also known as proline-rich transmembrane protein 1 (Prpt1), was identified as an AMPAR-associated protein [8,9,45]. SynDIG4 (Prpt1) was shown to be a component of the PSD [52], although the major fraction of Prpt1 colocalizes with GluA1 subunits at extrasynaptic sites in neurons [53]. It was shown to control levels of AMPARs at the cell surface and influence the gating properties of AMPARs (Tables 1 and 2) [54,55]. Ferric-chelate reductase 1 like (FRRS1L), also known as C9orf4 [8], is a single-pass transmembrane protein that effectively associates with the ER-resident CPT1c protein and plays an important role in the regulation of excitatory synaptic strength [56–58]. FRRS1L-CPT1c complexes co-assemble with the AMPAR during early biogenesis in the ER and FRRS1L-CPT1c-containing AMPARs exclusively localize at ER membranes (Figure 1). Brechet et al. have suggested that FRRS1L-CPT1c assemblies act as a ‘catalyst’ of AMPAR biogenesis and delivery of synaptic AMPARs specifically in the ER and are not present in AMPAR complexes at the synapse [56]. Additional protein quantifications and correlation analyses led to the model that all AMPARs expressed at the cell surface emerge from FRRS1L-containing AMPAR complexes in the ER [56]. In addition, secreted proteins acting at the AMPAR have been found [8,9]. Noelin1 (Olfactomedin-1) was shown to affect the membrane diffusion properties of the receptor while leaving kinetic and conductance properties of the AMPAR unaffected (Tables 1 and 2) [59].

In conclusion, a broad range of proteins can interact with AMPARs at varying stages of its life cycle, from newly synthesized AMPAR tetramers to mature AMPARs in the cell membrane (Figure 1), affecting biogenesis, trafficking, and biophysical properties of the receptor. The modulatory properties of the abovementioned auxiliary proteins on AMPAR surface expression and gating properties are summarized in Table 1 (*in vitro*) and Table 2 (*in vivo*).

Shisa9

The type-I transmembrane protein Shisa9 (CKAMP44) was discovered by interaction proteomics in 2010 and described as the first Shisa family member interacting with AMPARs [10]. Shisa9 specifically associates with AMPARs independent of subunit-type [10]. The 44 kDa protein Shisa9 is widely expressed throughout the brain and is most abundant in the dentate gyrus (DG) of the hippocampus [10]. Recent large-scale single-cell RNA sequencing studies of the hippocampus CA1 revealed the highest expression of Shisa9 in GABAergic interneurons, specifically in CCK⁺ and Reln⁺ subclasses [60]. Shisa9 function in inhibitory cells has not been studied to date and it will be of interest to reveal its role in these neurons. Shisa9 localizes at postsynaptic sites in primary neurons and interacts with multiple scaffold proteins, which are vital for the regulation of synaptic plasticity [61]. For instance, the C-terminal region of the

Table 1

Summary of AMPAR-interacting auxiliary proteins and their modulation of AMPAR surface expression and gating properties *in vitro*.

	AMPA surface expression	Current amplitude	Deactivation rate	Desensitization rate	Recovery from desensitization rate	Methods used
<i>In vitro</i>						
TARP $\gamma 2$	↑ [20,28,45,74]	↑ [20,28,75,76]	↓ [8,20,28,37,43,68,75,77]	↓ [8,20,28,43,68,75–77]	↔ [8,44,68,76,77]	[8,28,37]: Oocytes; [20,44,68]: HEK293; [45,74]: HEK; [75]: Oocytes/HEK293; [43,76,77]: HEK293 outside-out patch
TARP $\gamma 8$	↑ [17]	↑ [17,69]	↓ [29,39,42,44,69,74,77]	↓ [17,29,44,69,77]	↑ [44,69]	[17,29,42,44,69]: HEK293; [39,74]: HEK; [77]: tsA201 outside-out patch
CNIHs	↑ [37,58,74]	↑ [39,74]	↓ [8,37,39,42–44]	↓ [8,37,43,44]	↔ [8,37,42,44]	[8,58]: Oocytes (CNIH-2); [37]: Cultured cells (HeLa)/Oocytes (CNIH-2/3); [39]: HEK (CNIH-2); [42,44,74]: HEK293 (CNIH-2); [43]: HEK 293 outside-out patch (CNIH-3)
GSG1L	↑ [45]	N/A	↓ [8,45,47]	↓ [8,45,47]	↓ [8,46,47]	[8]: Oocytes; [45]: HEK outside-out patch; [46,47]: HEK
SynDIG4 (Prnt1)	N/A	↓ [54]	↓ [54]	↓ [54]	- [54]	[54]: Oocytes, outside-out patch
FRRS1L	- [57,58]	- [57]	N/A	N/A	↓ [56]	[56]: tsA201 outside-out patch; [57]: HEK ; [58]: Oocytes
CPT1c	↔ [58,78]	↑ [78]	N/A	- [78]	- [56]	[56]: tsA201 outside-out patch; [58]: Oocytes; [78]: tsA201 outside-out patch
Noelin1 (Olfacto-medin-1)	N/A	N/A	- [59]	- [59]	- [59]	[59]: HEK293
Shisa9 (CKAMP44)	- [10]	↓ [10,17]	N/A	- [17]	↓ [17]	[10]: Oocytes; [17]: HEK293
Shisa6 (CKAMP52)	↑ [13]	-↓* [13]	↔* [13,14]	↔* [13,14]	↔* [13]	[13]: HEK293/Oocytes; [14]: HEK293
Shisa7 (CKAMP59)	-↓* [13]	-↓* [13]	- [13,16]	↔ [13,16]	↔ [13,16]	[13]: HEK293/Oocytes; [16]: HEK293
Shisa8 (CKAMP39)	-↓* [13]	↓ [13]	-↓* [13]	-↑* [13]	↓ [13]	[13]: HEK293/Oocytes

↑ depicts increase, ↓ decrease, ↔ variable effects/conflicting results, – no effect. *Effect depends on type of GluA subunit. N/A, not available; HEK, human embryonic kidney. The table represents our best approximation of data in the literature.

Table 2

Summary of AMPAR-interacting auxiliary proteins and their modulation of AMPAR surface expression and gating properties *ex/in vivo*.

	AMPA surface expression	Current amplitude	Deactivation rate	Desensitization rate	Recovery from desensitization rate	Methods used
<i>Ex/in vivo</i>						
TARP $\gamma 2$	$\leftrightarrow$ [21,29,42,77,79,80]	$\leftrightarrow$ [21,29]	- [77]	$\downarrow$ [80]	$\uparrow$ [80]	[21]: CN, KO (CB); [29]: CN outside-out patch, KO (CB); [42]: Acute slices (HC); [77]: CN (CB); [79]: IB (CB); [80]: CN (HC)
TARP $\gamma 8$	$\uparrow$ [17,29,39,81]	$\leftrightarrow$ [17,21,29,39,42,81]	$\leftrightarrow$ [17,39]	$\downarrow$ [17]	$\uparrow$ [17,42]	[17]: CN outside-out patch, KO (HC (DG)); [21]: CN, KO (CB); [29]: CN outside-out patch, KO (CB); [39]: IB (HC)/CN, KO (HC (CA1)); [42]: CN (CB, HC); [81]: CN (HC)
CNIHs	$\uparrow$ [39,51,74]	$\uparrow$ [39]	$\leftrightarrow$ [39,74]	$\leftrightarrow$ [39,74]	- [39,42]	[39]: Acute slices (HC)/CN, KO (HC) (CNIH-2); [42]: CN (CB) (CNIH-2); [51]: CN, KD (HC); [74]: CN (CB, HC)
GSG1L	$\downarrow$ [46]	$\downarrow$ [46,48]	$\leftrightarrow$ [46,48]	$\leftrightarrow$ [46,48]	$\uparrow$ [46]	[46]: CN, KO (HC (CA1)) [48]: CN, KO (HC (DG))
SynDIG4 (Prnt1)	$\uparrow$ [55]	$\leftrightarrow$ [54,55]	- [54]	N/A	N/A	[54]: Acute slices, KO (HC (CA1)); [55]: Acute slices (HC (CA1))
FRRS1L	$\leftrightarrow$ [56,57]	$\uparrow$ [56–58]	- [56,57]	N/A	N/A	[56]: CN, KD (HC); [57]: CN, single-cell KO (HC (CA1)); [58]: Acute slices outside-out patch, KO (HC (CA1))
CPT1c	$\uparrow$ [56,78]	$\uparrow$ [56,82]	- [56,82]	N/A	N/A	[56]: CN, KD ((HC); [78]: CN (cortex); [82]: CN, KO (HC)
Noelin1 (Olfacto-medin-1)	N/A	- [59]	N/A	N/A	N/A	[59]: CN (HC)
Shisa9 (CKAMP44)	$\uparrow$ [17,83]	$\leftrightarrow^*$ [10,17,83]	$\leftrightarrow$ [10,17,83]	$\leftrightarrow^*$ [10,17,83]	$\downarrow$ [10,17,83]	[10]: Acute slices outside-out patch, KO (HC (DG))/ nucleated patch, KO, KD (HC); [17]: CN outside-out patch, KO (HC (DG)/nucleated patch (HC (DG)); [83]: Acute slices, nucleated patch, KO (dLGN relay neurons)
Shisa6 (CKAMP52)	$\leftrightarrow$ [14,15]	- [14,15]	$\leftrightarrow^*$ [14,15]	$\downarrow$ [14]	N/A	[14]: IB, KO (HC)/Acute slices, KO (HC (CA1))/ Nucleated patch, KO (HC (CA1)); [15]: IB, KO (CB)/ Acute slices, KO (CB (PCs)
Shisa7 (CKAMP59)	- [16]	- [16]	$\downarrow$ [16]	N/A	N/A	[16]: IB (HC)/Acute slices, KO (HC (CA1))
Shisa8 (CKAMP39)	N/A	N/A	N/A	N/A	N/A	N/A

$\uparrow$ depicts increase, $\downarrow$ decrease, $\leftrightarrow$ variable effects/conflicting results, – no effect. *Effect depends on type of GluA subunit. N/A, not available; CN, cultured neurons; IB, immunoblotting; KO, knockout; KD, knockdown; CB, cerebellum; HC, hippocampus; DG, dentate gyrus; dLGN: dorsal lateral geniculate nucleus; PCs, Purkinje cells. The table represents our best approximation of data in the literature.

Figure 1

Schematic showing a broad range of AMPAR auxiliary proteins at varying stages of their life cycle, highlighting their association with newly synthesized AMPARs in the endoplasmic reticulum (ER) and with AMPARs at the postsynaptic density (PSD). AMPARs and auxiliary subunits are assembled in the ER and Golgi cell soma and subsequently, upon ER exit (1, ER exit) and Golgi exit and via microtubular vesicular trafficking (2, vesicular transport), delivered to the plasma membrane (3, exocytosis). Via lateral diffusion, surface AMPARs reach the postsynapse. Via endocytosis (4, endocytosis) AMPARs can be retrieved from the plasma membrane (recycling). Studies have shown that several auxiliary subunits affect AMPAR trafficking (Shisa9 [17]; TARP $\gamma 2$ [21,22,25,28,29]; $\gamma 8$ [81]; CNIH-2 [39]; GSG1L [46]; CPT1c [78]; SynDIG4 [55]), with some of the auxiliary proteins acting at specific stages, such as ER exit (TARP $\gamma 2$ [21]; $\gamma 8$ [81]) and lateral diffusion (TARP $\gamma 2$ [30,35]; Noelin1 [59]). Proteins that have been specifically shown to associate with AMPARs within the ER and postsynaptic zone are shown in the figure (supporting literature is given in parentheses). However, the precise auxiliary subunit composition, stoichiometry, as well as the interaction of auxiliary subunits with each other, remains to be determined. Of note, with the exception of SynDIG4 and Noelin1, auxiliary subunits present in the postsynaptic zone are most likely assembled in the ER.

Shisa9 protein interacts with PDZ domain proteins, such as PSD-95, thereby providing indirect binding of AMPARs to the PSD [61,62]. Interference with these Shisa9-PDZ interactions profoundly affects synaptic AMPAR function [62].

Shisa9 modulates AMPAR gating properties different from TARP (Tables 1 and 2). Initial studies in both *Xenopus* oocytes and CA1 pyramidal neurons have shown that overexpression of Shisa9 affects AMPAR channel kinetics, such as deactivation and desensitization rates, whereas *Shisa9* deletion remained without

effect in CA1 neurons (Table 2) [10]. In DG granule cells, in which Shisa9 is clearly expressed [17], *Shisa9* deletion resulted in faster deactivation, slower desensitization, and faster recovery from desensitization of AMPAR-mediated currents [17]. Depending on the endogenous expression of Shisa9, its effect on AMPAR surface expression has yielded diverse results (Tables 1 and 2). Although Shisa9 expression in CA1 neurons did not show profound effects in terms of AMPAR surface expression [10,63], overexpression in DG granule cells increased surface expression [17], whereas heterologous expression in HEK cells reduced surface

expression [13]. Co-expression of both Shisa9 and TARP $\gamma 8$ was proven essential for efficient AMPAR surface expression in the DG (Table 2) [17], with additive effects observed upon a double deletion. Deletion of TARP $\gamma 8$ decreased paired-pulse ratio (PPR) at medial and lateral perforant path (MPP, LPP) synapses of DG granule cells, whereas deletion of *Shisa9* increased short-term plasticity [10,17]. On the other hand, deletion of TARP $\gamma 8$ fully abolished LTP at MPP synapses of DG granule cells, but deletion of *Shisa9* had no effect [17].

Shisa6

Shisa6 (CKAMP52) was recently identified as an additional constituent of native AMPAR complexes and is closely related to the AMPAR auxiliary protein Shisa9 [8,10,14]. Shisa6 shares the majority of its structural features with Shisa9, as both are type-I single-pass transmembrane proteins that contain an extracellular cysteine-rich domain, a proline-rich intracellular region, and a C-terminal type II PDZ-ligand motif [14,64]. Shisa6 is highly expressed in the cerebellum and all hippocampal subregions [14,65]. Unlike Shisa9, Shisa6 is highly expressed in pyramidal neurons and has moderate expression in two classes of SST⁺ interneurons in the hippocampus CA1 [60]. Shisa6 interacts with AMPAR subunits GluA1, GluA2, and GluA3 in both HEK293 cells and hippocampal tissue, and does so without an evident subunit preference [14]. In addition, Shisa6 interacts with PDZ-domain-containing scaffold proteins, such as PSD-95, through its C-terminal EVTV domain [14]. Shisa6 co-localizes with protein PSD-95 at excitatory postsynaptic sites and is enriched within purified PSD fractions. Overexpression of Shisa6 within hippocampal neurons was shown to constrain lateral surface mobility of AMPARs and confine them to PSD by association with PDZ proteins. Expression of a truncated version of Shisa6, lacking the C-terminal EVTV motif, did not alter AMPAR mobility [14].

Shisa6 modulates AMPAR-mediated currents directly upon co-expression with homomeric GluA1 and heteromeric GluA1/2 receptors within HEK293 cells. The presence of Shisa6 decreased the AMPAR deactivation rate and slowed recovery from desensitization, similar to what was reported for Shisa9 (Table 1) [10]. However, in contrast to the effect of Shisa9, Shisa6 was found to decrease the rate of desensitization and to greatly enhance the steady-state current upon prolonged glutamate application [14]. In CA1 pyramidal cells, deletion of *Shisa6* leads to a decrease in both rise and decay time of AMPAR-mediated miniature excitatory postsynaptic currents (mEPSCs), without affecting the amplitude or frequency [14]. Moreover, the absence of *Shisa6* profoundly impacted short-term plasticity in a frequency-dependent manner [14], resulting in enhanced synaptic depression during high-frequency stimulation (50 Hz).

Combined with the enhanced sustained current observed in HEK293 cells, the presence of Shisa6 is suggested to facilitate synaptic transmission by protecting postsynaptic AMPARs against full desensitization.

Just recently, an essential role for Shisa6 in the facilitation of cerebellar procedural memory formation has been described [15]. In the study of Peter *et al.*, it was shown that Shisa6 is prominently expressed by cerebellar Purkinje cells (PCs). Shisa6 proteins colocalize and associate with GluA2-containing AMPARs in cerebellar PCs and modulate AMPAR gating properties, such as AMPAR deactivation [15]. The absence of *Shisa6* resulted in a reduction of mEPSC amplitude and rise time and an increased deactivation of AMPAR-mediated currents in PCs, as well as impaired facilitation of functional parallel fiber (PF) to PC excitation. Whereas AMPAR expression, measured in the synaptic membrane fraction, does not seem to be affected by deletion of *Shisa6* in the hippocampus [14], cerebellar levels of AMPAR subunits are significantly reduced upon *Shisa6* deletion [15], in agreement with the effects observed on mEPSC amplitude in both regions. Furthermore, Shisa6 is required for PF-PC synapse LTP, whereas deletion of the AMPAR-interacting protein does not affect LTD of those synapses [15].

Shisa7

Shisa7 (CKAMP59), recently identified as an AMPAR-interacting protein [13,16], shares all major structural characteristics present in Shisa6 and Shisa9, including the N-terminal Cysteine-rich region and the C-terminal PDZ type II binding motif [16]. Shisa7 is highly expressed in the hippocampus, striatum, amygdala, cortex, and olfactory bulb [13,16]. *In situ* hybridization analyses from the Allen Brain Atlas shows high expression of *Shisa7* in both CA regions and the DG of the hippocampus, similar to Shisa6 [65]. Shisa7 is expressed in pyramidal neurons in the hippocampus CA1, with low interneuron expression [60]. Shisa7 specifically associates with the AMPAR in native hippocampal protein complexes and directly binds to homomeric GluA1, GluA2, and GluA3 receptors upon HEK293 co-expression, with no evident preference for a specific subunit [10,13,14,16]. PSD-95 is the most prominent PDZ domain-containing protein that interacts with Shisa7 in the hippocampus, with a crucial role for the C-terminal EVTV motif in this interaction [16]. In comparison to Shisa6, Shisa7 AMPAR-mediated AMPAR mobility has not been investigated.

In 2015, Farrow *et al.* reported that co-expression of Shisa7 in HEK293 cells does not affect most AMPAR-mediated current kinetics, except for the current amplitude mediated by GluA2-containing AMPARs [13]. More recently, it was established that Shisa7 can alter several AMPAR desensitization kinetics

(desensitization time, steady-state current, and recovery from desensitization), whereas channel kinetics such as AMPAR deactivation and rise time are indeed not affected [16]. Moreover, AMPAR expression levels in the synaptic membrane fraction in *Shisa7* KO mice do not differ from expression levels in *Shisa7* WT mice, resembling findings of *Shisa6* studies (Table 2) [14,16]. In contrast to both *Shisa9* and *Shisa6*, *Shisa7* does not affect short-term plasticity in the hippocampus [10,14,16]. These findings suggest that also *Shisa7* modulates AMPAR gating properties in a unique way, compared to other members of the Shisa protein family (Tables 1 and 2). Furthermore, *Shisa7* seems to have a crucial role in both the initiation as well as the maintenance of hippocampal LTP [16]. *Shisa7* KO mice showed a slower initiation of LTP as well as a decrease in the maintenance phase, compared to WT animals.

Interestingly, in 2019, *Shisa7* was identified as a GABA_AR-interacting protein [66]. Han et al. showed that *Shisa7* localized at GABAergic inhibitory synapse and that it is a critical protein for inhibitory transmission. Using cocultures of hippocampal neurons with HEK293 cells, co-expression of *Shisa7* and GABA_ARs resulted in a reduction of both the decay time constant of spontaneous inhibitory postsynaptic currents (sIPSCs) as well as the time constant of deactivation, indicating modulation of the kinetics of GABAergic transmission and GABA_ARs by *Shisa7*. Moreover, deletion of *Shisa7* increased the decay time constant of mIPSCs in CA1 pyramidal cells and decreased the expression of GABA_AR subunits at the inhibitory PSD [66]. In opposition to previous studies [13,16], Han et al. have reported that *Shisa7* does not colocalize with glutamatergic synaptic markers in primary cultures [66].

Shisa8

The least studied AMPAR-interacting member of the Shisa protein family is *Shisa8* (CKAMP39). The highest expression of *Shisa8* was observed in the cerebellum and olfactory bulb, with almost no expression in other brain regions [13]. In the study of Farrow et al., co-expression of *Shisa8* and AMPAR subunits GluA1 and GluA2 in *Xenopus* oocytes showed that it co-immunoprecipitated with both subunits, and modulated AMPAR gating properties [13]. In HEK293 cells, *Shisa8* significantly affected AMPAR-mediated current kinetics, such as current amplitude, as well as AMPAR desensitization and recovery from desensitization. However, some differences were observed for deactivation- and desensitization kinetics between GluA1-mediated currents and GluA2-mediated currents, when modulated by *Shisa8*. The effects of *Shisa8* on AMPAR gating properties in neurons remains unknown since this effect has not been studied to date. With lacking data on the endogenous neuronal function, *Shisa8* can currently not be regarded as a true AMPAR auxiliary protein.

Perspective: pharmacological targeting of interfaces between AMPARs and auxiliary proteins

Auxiliary proteins interacting with the AMPAR play a critical role in the CNS by modulating receptor synthesis, trafficking, surface expression, and gating properties. With current knowledge, a broad range of AMPAR auxiliary proteins seems to modulate the receptor's function at different cellular locations, from the newly synthesized tetramers at the early stage of biogenesis to mature AMPARs in the cell membrane. Throughout the years, various 'families' of proteins have been identified and today the approximate full complement of AMPAR auxiliary proteins is known. With this parts list identified, the journey into the discovery of specific functions of individual proteins or combinations thereof has only begun. After the discovery of the Shisa protein family, the precise examination of their role in allosteric modulation of AMPAR function and membrane diffusion has significantly contributed to the detailed understanding of mechanisms acting on AMPAR-mediated fast synaptic transmission and plasticity. This holds true for many of the identified AMPAR-associated proteins.

Interestingly, the expression levels of various auxiliary subunits have been identified by determining molecular profiles of AMPAR auxiliary subunits across distinct brain regions and recently even down to single cells. Distinct cellular expression of auxiliary proteins may lead to functionally unique combinatorial actions on the AMPAR [17]. With respect to this, the composition and stoichiometry of auxiliary subunits also vary greatly within AMPAR complexes [4,67]. The stoichiometry of TARPs is best understood since recent cryo-electron microscopy (cryo-EM) studies have provided structural information on the AMPAR-TARP complex [68–70]. In addition, structural analyses have revealed the molecular mechanism underlying the modulation of AMPAR function by Stargazin/TARP $\gamma 2$ [71,72]. However, structural analyses of the AMPAR interaction with other auxiliary subunits, such as Shisa proteins, or even with multiple different auxiliary proteins, are absent.

Importantly, both functional and structural information of the combinatorial action of auxiliary proteins on the AMPAR is opening new avenues for the rational design of compounds acting at the molecular interfaces of the auxiliary proteins and AMPAR [73]. This may provide unprecedented possibilities for modulating AMPAR receptor function. In conclusion, examining the subcellular points of interaction between auxiliary subunits and AMPARs, the cell-type-specific auxiliary subunit expression, the precise auxiliary subunit composition, and stoichiometry, as well as the interaction of Shisa proteins with each other and/or with other auxiliary subunits together at the AMPAR, might all contribute to potential new pharmacological interventions.

Author contributions

August B Smit: Conceptualization, Writing and editing, Supervision. **Mazyar Abdollahi Nejat:** Writing – original draft, Writing. **Sabine Spijker:** Visualization, editing. **Remco Klaassen:** Investigation, editing.

Conflict of interest statement

Nothing declared.

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
- ** of outstanding interest

1. Isaac JTR, Ashby M, McBain CJ: **The role of the GluR2 subunit in AMPA receptor function and synaptic plasticity.** *Neuron* 2007, **54**:859–871.
2. Shepherd JD, Huganir RL: **The cell biology of synaptic plasticity: AMPA receptor trafficking.** *Annu Rev Cell Dev Biol* 2007, **23**:613–643.
3. Jackson AC, Nicoll RA: **The expanding social network of ionotropic glutamate receptors: TARPs and other transmembrane auxiliary subunits.** *Neuron* 2011, **70**:178–199.
4. Greger IH, Watson JF, Cull-Candy SG: **Structural and functional architecture of AMPA-type glutamate receptors and their auxiliary proteins.** *Neuron* 2017, **94**:713–730.
5. Haering SC, Tapken D, Pahl S, Hollmann M: **Auxiliary subunits: shepherding AMPA receptors to the plasma membrane.** *Membranes* 2014, **4**:469–490.
6. Noebels JL, Qiao X, Bronson RT, Spencer C, Davisson MT: **Stargazer: a new neurological mutant on chromosome 15 in the mouse with prolonged cortical seizures.** *Epilepsy Res* 1990, **7**:129–135.
7. Letts VA, Felix R, Biddlecome GH, Arikath J, Mahaffey CL, Valenzuela A, Bartlett FS, Mori Y, Campbell KP, Frankel WN: **The mouse stargazer gene encodes a neuronal Ca²⁺-channel γ subunit.** *Nat Genet* 1998, **19**:340–347.
8. Schwenk J, Harmel N, Brechet A, Zolles G, Berkefeld H, Müller CS, Bildl W, Baehrens D, Hüber B, Kulik A, *et al.*: **High-resolution proteomics unravel architecture and molecular diversity of native AMPA receptor complexes.** *Neuron* 2012, **74**:621–633.
9. Chen N, Pandya NJ, Koopmans F, Castelo-Székely V, Van Der Schors RC, Smit AB, Li KW: **Interaction proteomics reveals brain region-specific AMPA receptor complexes.** *J Proteome Res* 2014, **13**:5695–5706.
10. Von Engelhardt J, Mack V, Sprengel R, Kavenstock N, Li KW, Stern-Bach Y, Smit AB, Seeburg PH, Monyer H: **CKAMP44: a brain-specific protein attenuating short-term synaptic plasticity in the dentate gyrus.** *Science (80-)* 2010, **327**:1518–1522.
11. Nakagawa T: **The biochemistry, ultrastructure, and subunit assembly mechanism of AMPA receptors.** *Mol Neurobiol* 2010, **42**:161–184.
12. Tomita S: **Regulation of ionotropic glutamate receptors by their auxiliary subunits.** *Physiology* 2010, **25**:41–49.
13. Farrow P, Khodosevich K, Sapir Y, Schulmann A, Aslam M, Stern-Bach Y, Monyer H, von Engelhardt J: **Auxiliary subunits of the CKAMP family differentially modulate AMPA receptor properties.** *Elife* 2015, **4**.
14. Klaassen RV, Stroeder J, Coussen F, Hafner AS, Petersen JD, Renancio C, Schmitz LJM, Normand E, Lodder JC, Rotaru DC, *et al.*: **Shisa6 traps AMPA receptors at postsynaptic sites and prevents their desensitization during synaptic activity.** *Nat Commun* 2016, **7**:10682.
15. Peter S, Urbanus BHA, Klaassen RV, Wu B, Boele HJ, Azizi S, Slotman JA, Houtsmuller AB, Schonewille M, Hoebeek FE, *et al.*: **AMPA auxiliary protein SHISA6 facilitates Purkinje cell synaptic excitability and procedural memory formation.** *Cell Rep* 2020, **31**:107515.
- ** This study shows that the Shisa6 protein is prominently expressed in Purkinje cells and plays an essential role in the facilitation of cerebellar procedural memory formation. Moreover, the authors find that Shisa6 modulates AMPAR kinetics and synaptic function of Purkinje cells in mice.
16. Schmitz LJM, Klaassen RV, Ruiperez-Alonso M, Zamri AE, Stroeder J, Rao-Ruiz P, Lodder JC, van der Loo RJ, Mansvelder HD, Smit AB, *et al.*: **The AMPA receptor-associated protein Shisa7 regulates hippocampal synaptic function and contextual memory.** *Elife* 2017, **6**.
17. Khodosevich K, Jacobi E, Farrow P, Schulmann A, Rusu A, Zhang L, Sprengel R, Monyer H, von Engelhardt J: **Coexpressed auxiliary subunits exhibit distinct modulatory profiles on AMPA receptor function.** *Neuron* 2014, **83**:601–615.
18. Zheng Y, Mellem JE, Brockie PJ, Madsen DM, Maricq AV: **SOL-1 is a CUB-domain protein required for GLR-1 glutamate receptor function in *C. elegans*.** *Nature* 2004, **427**:451–457.
19. Chen L, Chetkovich DM, Petralia RS, Sweeney NT, Kawasaki Y, Wenthold RJ, Brecht DS, Nicoll RA: **Stargazin regulates synaptic targeting of AMPA receptors by two distinct mechanisms.** *Nature* 2000, **408**:936–943.
20. Bedoukian MA, Weeks AM, Partin KM: **Different domains of the AMPA receptor direct stargazin-mediated trafficking and stargazin-mediated modulation of kinetics.** *J Biol Chem* 2006, **281**:23908–23921.
21. Tomita S, Chen L, Kawasaki Y, Petralia RS, Wenthold RJ, Nicoll RA, Brecht DS: **Functional studies and distribution define a family of transmembrane AMPA receptor regulatory proteins.** *J Cell Biol* 2003, **161**:805–816.
22. Vandenberghe W, Nicoll RA, Brecht DS: **Stargazin is an AMPA receptor auxiliary subunit.** *Proc Natl Acad Sci U S A* 2005, **102**:485–490.
23. Shanks NF, Maruo T, Farina AN, Ellisman MH, Nakagawa T: **Contribution of the global subunit structure and stargazin on the maturation of AMPA receptors.** *J Neurosci* 2010, **30**:2728–2740.
24. Kato AS, Zhou W, Milstein AD, Knierman MD, Siuda ER, Dotzlaw JE, Yu H, Hale JE, Nisenbaum ES, Nicoll RA, *et al.*: **New transmembrane AMPA receptor regulatory protein isoform, γ -7, differentially regulates AMPA receptors.** *J Neurosci* 2007, **27**:4969–4977.
25. Kato AS, Siuda ER, Nisenbaum ES, Brecht DS: **AMPA receptor subunit-specific regulation by a distinct family of type II TARPs.** *Neuron* 2008, **59**:986–996.
26. Soto D, Coombs ID, Renzi M, Zonouzi M, Farrant M, Cull-Candy SG: **Selective regulation of long-form calcium-permeable AMPA receptors by an atypical TARP, γ -5.** *Nat Neurosci* 2009, **12**:277–285.
27. Milstein AD, Nicoll RA: **Regulation of AMPA receptor gating and pharmacology by TARP auxiliary subunits.** *Trends Pharmacol Sci* 2008, **29**:333–339.
28. Tomita S, Adesnik H, Sekiguchi M, Zhang W, Wada K, Howe JR, Nicoll RA, Brecht DS: **Stargazin modulates AMPA receptor gating and trafficking by distinct domains.** *Nature* 2005, **435**:1052–1058.
29. Milstein AD, Zhou W, Karimzadegan S, Brecht DS, Nicoll RA: **TARP subtypes differentially and dose-dependently control synaptic AMPA receptor gating.** *Neuron* 2007, **55**:905–918.
30. Bats C, Groc L, Choquet D: **The interaction between stargazin and PSD-95 regulates AMPA receptor surface trafficking.** *Neuron* 2007, **53**:719–734.
31. Zeng M, Díaz-Alonso J, Ye F, Chen X, Xu J, Ji Z, Nicoll RA,

- Zhang M: **Phase separation-mediated TARP/MAGUK complex condensation and AMPA receptor synaptic transmission.** *Neuron* 2019, **104**:529–543. e6.
32. Tomita S, Stein V, Stocker TJ, Nicoll RA, Brecht DS: **Bidirectional synaptic plasticity regulated by phosphorylation of stargazin-like TARPs.** *Neuron* 2005, **45**:269–277.
 33. Opazo P, Labrecque S, Tigaret CM, Frouin A, Wiseman PW, De Koninck P, Choquet D: **CaMKII triggers the diffusional trapping of surface AMPARs through phosphorylation of stargazin.** *Neuron* 2010, **67**:239–252.
 34. Hafner AS, Penn AC, Grillo-Bosch D, Retaillieu N, Poujol C, Philippat A, Coussen F, Sainlos M, Opazo P, Choquet D: **Lengthening of the stargazin cytoplasmic tail increases synaptic transmission by promoting interaction to deeper domains of PSD-95.** *Neuron* 2015, **86**:475–489.
 35. Sumioka A, Yan D, Tomita S: **TARP phosphorylation regulates synaptic AMPA receptors through lipid bilayers.** *Neuron* 2010, **66**:755–767.
 36. Park J, Chávez AE, Mineur YS, Morimoto-Tomita M, Lutz S, Kim KS, Picciotto MR, Castillo PE, Tomita S: **CaMKII phosphorylation of TARP γ -8 is a mediator of LTP and learning and memory.** *Neuron* 2016, **92**:75–83.
 37. Schwenk J, Harmel N, Zolles G, Bildl W, Kulik A, Heimrich B, Chisaka O, Jonas P, Schulte U, Fakler B, *et al.*: **Functional proteomics identify cornichon proteins as auxiliary subunits of AMPA receptors.** *Science (80-)* 2009, **323**:1313–1319.
 38. Bökel C, Dass S, Wilsch-Bräuninger M, Roth S: **Drosophila Cornichon acts as cargo receptor for ER export of the TGF α -like growth factor Gurken.** *Development* 2006, **133**:459–470.
 39. Herring BE, Shi Y, Suh YH, Zheng CY, Blankenship SM, Roche KW, Nicoll RA: **Cornichon proteins determine the subunit composition of synaptic AMPA receptors.** *Neuron* 2013, **77**:1083–1096.
 40. Nakagawa T: **Structures of the AMPA receptor in complex with its auxiliary subunit cornichon.** *Science (80-)* 2019, **366**:1259–1263.
 41. Gill MB, Kato AS, Roberts MF, Yu H, Wang H, Tomita S, Brecht DS: **Cornichon-2 modulates AMPA receptor-transmembrane AMPA receptor regulatory protein assembly to dictate gating and pharmacology.** *J Neurosci* 2011, **31**:6928–6938.
 42. Gill MB, Kato AS, Wang H, Brecht DS: **AMPA receptor modulation by cornichon-2 dictated by transmembrane AMPA receptor regulatory protein isoform.** *Eur J Neurosci* 2012, **35**:182–194.
 43. Brown PMGE, McGuire H, Bowie D: **Stargazin and cornichon-3 relieve polyamine block of AMPA receptors by enhancing blocker permeation.** *J Gen Physiol* 2018, **150**:67–82.
 44. Kato AS, Gill MB, Ho MT, Yu H, Tu Y, Siuda ER, Wang H, Qian YW, Nisenbaum ES, Tomita S, *et al.*: **Hippocampal AMPA receptor gating controlled by both TARP and cornichon proteins.** *Neuron* 2010, **68**:1082–1096.
 45. Shanks NF, Savas JN, Maruo T, Cais O, Hirao A, Oe S, Ghosh A, Noda Y, Greger IH, Yates JR, *et al.*: **Differences in AMPA and kainate receptor interactomes facilitate identification of AMPA receptor auxiliary subunit GSG1L.** *Cell Rep* 2012, **1**:590–598.
 46. Gu X, Mao X, Lussier MP, Hutchison MA, Zhou L, Hamra FK, Roche KW, Lu W: **GSG1L suppresses AMPA receptor-mediated synaptic transmission and uniquely modulates AMPA receptor kinetics in hippocampal neurons.** *Nat Commun* 2016, **7**:1–18.
 47. Twomey EC, Yelshanskaya MV, Grassucci RA, Frank J, Sobolevsky AI: **Structural bases of desensitization in AMPA receptor-auxiliary subunit complexes.** *Neuron* 2017, **94**:569–580. e5.
 48. Mao X, Gu X, Lu W: **GSG1L regulates the strength of AMPA receptor-mediated synaptic transmission but not AMPA receptor kinetics in hippocampal dentate granule neurons.** *J Neurophysiol* 2017, **117**:28–35.
 49. Keifer J, Tiwari NK, Buse L, Zheng Z: **Subunit-specific synaptic delivery of AMPA receptors by auxiliary chaperone proteins TARP γ 8 and GSG1L in classical conditioning.** *Neurosci Lett* 2017, **645**:53–59.
 50. Kalashnikova E, Lorca RA, Kaur I, Barisone GA, Li B, Ishimaru T, Trimmer JS, Mohapatra DP, Díaz E: **SynDIG1: an activity-regulated, AMPA- receptor-interacting transmembrane protein that regulates excitatory synapse development.** *Neuron* 2010, **65**:80–93.
 51. Lovero KL, Blankenship SM, Shi Y, Nicoll RA: **SynDIG1 promotes excitatory synaptogenesis independent of AMPA receptor trafficking and biophysical regulation.** *PLoS One* 2013, **8**, e66171.
 52. Jordan BA, Fernholz BD, Boussac M, Xu C, Grigorean G, Ziff EB, Neubert TA: **Identification and verification of novel rodent postsynaptic density proteins.** *Mol Cell Proteomics* 2004, **3**:857–871.
 53. Kirk LM, Ti SW, Bishop HI, Orozco-Llamas M, Pham M, Trimmer JS, Díaz E: **Distribution of the SynDIG4/proline-rich transmembrane protein 1 in rat brain.** *J Comp Neurol* 2016, **524**:2266–2280.
 54. Matt L, Kirk LM, Chenuaux G, Specia DJ, Puhger KR, Pride MC, Qneibi M, Haham T, Plambeck KE, Stern-Bach Y, *et al.*: **SynDIG4/Prtr1 is required for excitatory synapse development and plasticity underlying cognitive function.** *Cell Rep* 2018, **22**:2246–2253.
- In this study, SynDIG4 (Prtr1) has been shown to modify AMPAR gating properties in a subunit-dependent manner. The authors show that SynDIG4 colocalizes with GluA1 at non-synaptic sites. They suggest that SynDIG4 maintains a pool of extrasynaptic AMPARs necessary for synapse development.
55. Troyano-Rodriguez E, Mann S, Ullah R, Ahmad M: **PRRT1 regulates basal and plasticity-induced AMPA receptor trafficking.** *Mol Cell Neurosci* 2019, **98**:155–163.
- Using biochemistry and electrophysiology, this study demonstrated that Prtr1 (SynDIG4) controls the levels of AMPARs at the cell surface and is an important regulator of AMPAR stabilization.
56. Brechet A, Buchert R, Schwenk J, Boudkazi S, Zolles G, Siquier-Pernet K, Schaber I, Bildl W, Saadi A, Bole-Feyssot C, *et al.*: **AMPA-receptor specific biogenesis complexes control synaptic transmission and intellectual ability.** *Nat Commun* 2017, **8**:1–14.
 57. Han W, Wang H, Li J, Zhang S, Lu W: **Ferric chelate reductase 1 like protein (FRS1L) associates with dynein vesicles and regulates glutamatergic synaptic transmission.** *Front Mol Neurosci* 2017, **10**:402.
 58. Schwenk J, Boudkazi S, Kocylowski MK, Brechet A, Zolles G, Bus T, Costa K, Kollwe A, Jordan J, Bank J, *et al.*: **An ER assembly line of AMPA-receptors controls excitatory neurotransmission and its plasticity.** *Neuron* 2019, **104**:680–692. e9.
- This study provides insight into the biogenesis of AMPARs in native ER membranes. They establish that FRS1L is a potent regulator of synaptic signaling and synaptic plasticity.
59. Pandya NJ, Seeger C, Babai N, Gonzalez-Lozano MA, Mack V, Lodder JC, Gouwenberg Y, Mansvelder HD, Danielson UH, Li KW, *et al.*: **Noelin1 affects lateral mobility of synaptic AMPA receptors.** *Cell Rep* 2018, **24**:1218–1230.
- This study shows that the secreted protein Noelin1 (Olfactomedin-1) interacts with a high affinity to AMPARs. The authors show that Noelin1 reduces synaptic AMPAR lateral mobility, limiting synaptic plasticity. This is the first evidence for a secreted auxiliary subunit in the mobility and plasticity of glutamate receptors.
60. Zeisel A, Moz-Manchado AB, Codeluppi S, Lönnerberg P, Manno G La, Jureus A, Marques S, Munguba H, He L, Betsholtz C, *et al.*: **Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq.** *Science (80-)* 2015, **347**:1138–1142.
 61. Kunde SA, Rademacher N, Zieger H, Shoichet SA: **Protein kinase C regulates AMPA receptor auxiliary protein Shisa9/**

- CKAMP44 through interactions with neuronal scaffold PICK1.** *FEBS Open Bio* 2017, **7**:1234–1245.
62. Karataeva AR, Klaassen RV, Ströder J, Ruiperez-Alonso M, Hjorth JJJ, van Nierop P, Spijker S, Mansvellder HD, Smit AB: **C-terminal interactors of the AMPA receptor auxiliary subunit Shisa9.** *PLoS One* 2014, **9**, e87360.
 63. Guzman SJ, Jonas P: **Beyond TARPs: the growing list of auxiliary AMPAR subunits.** *Neuron* 2010, **66**:8–10.
 64. Pei J, Grishin NV: **Unexpected diversity in Shisa-like proteins suggests the importance of their roles as transmembrane adaptors.** *Cell Signal* 2012, **24**:758–769.
 65. Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, Boe AF, Boguski MS, Brockway KS, Byrnes EJ, *et al.*: **Genome-wide atlas of gene expression in the adult mouse brain.** *Nature* 2007, **445**:168–176.
 66. Han W, Li J, Pelkey KA, Pandey S, Chen X, Wang YX, Wu K, Ge L, Li T, Castellano D, *et al.*: **Shisa7 is a GABA_A receptor auxiliary subunit controlling benzodiazepine actions.** *Science (80-)* 2019, **366**:246–250.
- In this study, Shisa7 has been identified as a GABA_AR-interacting protein. They show that Shisa7 localizes at GABAergic inhibitory synapses and that it is a critical protein for inhibitory transmission. In opposition to previous studies, the authors have reported that Shisa7 does not colocalize with glutamatergic synaptic markers in primary cultures.
67. Miguez-Cabello F, Sánchez-Fernández N, Yefimenko N, Gasull X, Gratacòs-Batlle E, Soto Del Cerro D: **AMPA/TARP stoichiometry differentially modulates channel properties.** *Elife* 2020, **9**:1.
- The authors of this study have found that AMPARs are modulated by TARP $\gamma 2$ in a stoichiometry-dependent manner. They show the effects of stoichiometry on AMPAR biophysical properties and they show important differences in this stoichiometric modulation between calcium-permeable and calcium-impermeable AMPARs.
68. Twomey EC, Yelshanskaya MV, Grassucci RA, Frank J, Sobolevsky AI: **Elucidation of AMPA receptor-stargazin complexes by cryo-electron microscopy.** *Science (80-)* 2016, **353**: 83–86.
 69. Herguedas B, Watson JF, Ho H, Cais O, García-Nafria J, Greger IH: **Architecture of the heteromeric GluA1/2 AMPA receptor in complex with the auxiliary subunit TARP $\gamma 8$.** *Science (80-)* 2019:364.
- This study presents structural (cryo-EM) and functional studies that elucidate the organization of GluA1/A2 in complex with two TARP $\gamma 8$ proteins, which is the principal AMPAR complex at hippocampal synapses.
70. Dohrke J-N, Watson JF, Birchall K, Greger IH: **Characterising the binding and function of TARP $\gamma 8$ -selective AMPA receptor modulators.** *J Biol Chem* 2020, <https://doi.org/10.1074/jbc.ra120.014135>.
 71. Zhao Y, Chen S, Yoshioka C, Bacongus I, Gouaux E: **Architecture of fully occupied GluA2 AMPA receptor-TARP complex elucidated by cryo-EM.** *Nature* 2016, **536**:108–111.
 72. Ben-Yaacov A, Gillor M, Haham T, Parsai A, Qneibi M, Stern-Bach Y: **Molecular mechanism of AMPA receptor modulation by TARP/stargazin.** 2017, <https://doi.org/10.1016/j.neuron.2017.01.032>.
 73. Kato AS, Witkin JM: **Auxiliary subunits of AMPA receptors: the discovery of a forebrain-selective antagonist, LY3130481/CERC-611.** *Biochem Pharmacol* 2018, **147**: 191–200.
- This review discusses a novel approach to drug discovery based on the identification and characterization of auxiliary proteins.
74. Shi Y, Suh YH, Milstein AD, Iozaki K, Schmid SM, Roche KW, Nicoll RA: **Functional comparison of the effects of TARPs and cornichons on AMPA receptor trafficking and gating.** *Proc Natl Acad Sci U S A* 2010, **107**:16315–16319.
 75. Priel A, Kollek A, Ayalon G, Gillor M, Osten P, Stern-Bach Y: **Stargazin reduces desensitization and slows deactivation of the AMPA-type glutamate receptors.** *J Neurosci* 2005, **25**: 2682–2686.
 76. Coombs ID, MacLean DM, Jayaraman V, Farrant M, Cull-Candy SG: **Dual effects of TARP $\gamma 2$ on glutamate efficacy can account for AMPA receptor autoinactivation.** *Cell Rep* 2017, **20**:1123–1135.
 77. Cho CH, St-Gelais F, Zhang W, Tomita S, Howe JR: **Two families of TARP isoforms that have distinct effects on the kinetic properties of AMPA receptors and synaptic currents.** *Neuron* 2007, **55**:890–904.
 78. Gratacòs-Batlle E, Yefimenko N, Cascos-García H, Soto D: **AMPA interacting protein CPT1C enhances surface expression of GLuA1-containing receptors.** *Front Cell Neurosci* 2015, **8**.
 79. Menuz K, O'Brien JL, Karmizadegan S, Bredt DS, Nicoll RA: **TARP redundancy is critical for maintaining AMPA receptor function.** *J Neurosci* 2008, **28**:8740–8746.
 80. Turetsky D, Garringer E, Patneau DK: **Stargazin modulates native AMPA receptor functional properties by two distinct mechanisms.** *J Neurosci* 2005, **25**:7438–7448.
 81. Rouach N, Byrd K, Petralia RS, Elias GM, Adesnik H, Tomita S, Karimzadegan S, Kealey C, Bredt DS, Nicoll RA: **TARP $\gamma 8$ controls hippocampal AMPA receptor number, distribution and synaptic plasticity.** *Nat Neurosci* 2005, **8**: 1525–1533.
 82. Fadó R, Soto D, Miñano-Molina AJ, Pozo M, Carrasco P, Yefimenko N, Rodríguez-Álvarez J, Casals N: **Novel regulation of the synthesis of α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic Acid (ampa) receptor subunit glua1 by carnitine palmitoyltransferase 1C (CPT1C) in the Hippocampus.** *J Biol Chem* 2015, **290**:25548–25560.
 83. Chen X, Aslam M, Gollisch T, Allen K, Von Engelhardt J: **CKAMP44 modulates integration of visual inputs in the lateral geniculate nucleus.** *Nat Commun* 2018, **9**:1–13.
- This study clearly shows that CKAMP44 (Shisa9) is an essential auxiliary subunit of the AMPAR in relay neurons of the dLGN, affecting several AMPAR gating properties. They provide evidence that CKAMP44 is responsible for short-term depression in retinogeniculate synapses.