VU Research Portal

Comment on: First observation of the rotational constants for the homonuclear ion He₂⁺

van der Zande, W.J.; Ubachs, W.M.G.

published in

Physical Review Letters
2000

DOI (link to publisher)

[10.1103/PhysRevLett.84.3212](https://doi.org/10.1103/PhysRevLett.84.3212)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

van der Zande, W. J., & Ubachs, W. M. G. (2000). Comment on: First observation of the rotational constants for the homonuclear ion He₂⁺. *Physical Review Letters*, 84, 3212-3212.
<https://doi.org/10.1103/PhysRevLett.84.3212>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Comment on "First Measurement of the Rotational Constants for the Homonuclear Molecular Ion He_2^+ "

In a recent Letter [1], Coman *et al.* describe a technique to derive spectroscopic information on the structure of the parent ion, using information on slow atoms emerging from a dissociative recombination process. This form of translational energy spectroscopy was used to experimentally determine the rotational constant in the electronic ground state of the He_2^+ ion. Neutral atoms are formed in dissociative recombination of the ion with low energy electrons in a plasma discharge. As a consequence of accidental near degeneracies between molecular ion energies and the internal energy of the neutral He atom in the $(1s3s)$ state, the fragment atoms are very slow. Their kinetic energy can therefore be determined with high accuracy.

The analysis of the experimental data seems to be flawed in a fundamental way. In Table I of Ref. [1] experimental entries for energy levels are displayed showing a linear progression as a function of $J(J+1)$, which is indicative of a rotational progression. In a diatomic species, consisting of two indistinguishable spin-0 particles, the symmetrization postulate of quantum mechanics forbids the existence of quantum states with a certain parity. It is easily proven (see, e.g., Ref. [2]) that for a $^2\Sigma_u^+$ state only levels of negative parity can exist, where the parity operator is the inversion in a space-fixed frame I_{SF} or the reflection in a plane σ_v in a body-fixed frame of the molecule. Hence quantum levels with an even number of rotational angular momentum (here defined with N) do not exist for the $X^2\Sigma_u^+$ electronic ground state of the He_2^+ ion. The fact that in a state of doublet character a spin-rotation interaction $\vec{N} \cdot \vec{S}$ splits the level into two nearly degenerate components ($J = N + 1/2$ and $J = N - 1/2$) does not alter the argument. Here $\vec{S}$ is the spin of the unpaired electron.

No violations of this symmetry postulate have been detected so far. Predictions on possible small violations of this postulate have led to experimental tests searching for very weak occurrences of forbidden lines in the spectrum of $^{16}\text{O}_2$, also consisting of two indistinguishable spin-0 particles. These studies have produced negative results at the 10^{-6} – 10^{-7} sensitivity level [3–5].

The effect of nuclear spin on the populations of rotational levels was observed in an experiment on the photodissociation of H_2^+ , also in combination with high-resolution translational spectroscopy [6]. The characteristic 3:1 ortho-para ratio was clearly obtained. Lewis *et al.* [7] analyzed the spectrum in excitation to the $f^1\Sigma_u^+$ state of $^{16}\text{O}_2$, in which half of the rotational levels were shown to be missing due to the restriction of the symmetrization postulate. Except for the singlet spin structure the symmetry of that state is the same as that of the ground state of the $^4\text{He}_2^+$ ion. In conclusion we comment that the results of the experiment on He_2^+ would have to be reinterpreted without invoking the existence of quantum states of even rotational angular momentum.

W.J. van der Zande

FOM-Institute for Atomic and Molecular Physics
Kruislaan 407
1098 SJ Amsterdam, The Netherlands

W. Ubachs

Department of Physics and Astronomy
Vrije Universiteit
De Boelelaan 1081
1081 HV Amsterdam, The Netherlands

Received 21 October 1999

PACS numbers: 33.15.Mt

- [1] L. Coman, M. Guna, L. Simons, and K. A. Hardy, *Phys. Rev. Lett.* **83**, 2715 (1999).
- [2] P.R. Bunker, *Molecular Symmetry and Spectroscopy* (Academic Press, New York, 1979).
- [3] M. de Angelis, G. Gagliardi, L. Gianfrani, and G. M. Tino, *Phys. Rev. Lett.* **76**, 2840 (1996).
- [4] R.C. Hilborn and C.L. Yuca, *Phys. Rev. Lett.* **76**, 2844 (1996).
- [5] H. Naus, A. de Lange, and W. Ubachs, *Phys. Rev. A* **56**, 4755 (1997).
- [6] W. Koot, W.J. van der Zande, and D.P. de Bruijn, *Chem. Phys.* **115**, 297 (1987).
- [7] B. R. Lewis, J. P. England, R. J. Winkel, Jr., S. S. Banerjee, P. M. Dooley, S. T. Gibson, and K. G. H. Baldwin, *Phys. Rev. A* **52**, 2717 (1995).