VU Research Portal

EUNETHYDIS - Searching for valid aetiological candidates of Attention-Deficit Hyperactivity Disorder of Hyperkinetic Disorder

Sergeant, J.A.

published in

European Child and Adolescent Psychiatry
2004

DOI (link to publisher)

[10.1007/s00787-004-1005-7](https://doi.org/10.1007/s00787-004-1005-7)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Sergeant, J. A. (2004). EUNETHYDIS - Searching for valid aetiological candidates of Attention-Deficit Hyperactivity Disorder of Hyperkinetic Disorder. *European Child and Adolescent Psychiatry*, 13(S1), 43-49. <https://doi.org/10.1007/s00787-004-1005-7>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Joseph Sergeant

EUNETHYDIS – Searching for valid aetiological candidates of Attention-Deficit Hyperactivity Disorder or Hyperkinetic Disorder

■ **Abstract** *Background* To step up research in ADHD, exchange of ideas, working together on key theoretical models and cooperative studies are necessary. *Objective* To

report about a European approach with strong links to the rest of the world. *Method* European Network on Hyperkinetic Disorders (Eunethydis) studies of Attention-Deficit Hyperactivity Disorder (ADHD) or Hyperkinetic Disorder (HKD) is briefly reviewed in the context of the international effort to discover the aetiology of the disorder. *Results* There are promising neurobiological, neurophysiological and neuropsychological candi-

dates to explain the nature of ADHD/HKD. *Conclusion* Eunethydis has shown to be a fruitful platform for ADHD research and has good resources for its further development.

■ **Key words** ADHD – HKD – executive functioning – inhibition – working memory – arousal – cognitive-energetic – delay aversion – frontal lobe – cerebellum – review

Prof. Dr. J. Sergeant (✉)
Free University
Clinical Neuropsychology
De Boelelaan 1105
1081 HV Amsterdam, The Netherlands
Tel.: +31-20/444-8757
Fax: +31-20/444-8832
E-Mail: j.a.sergeant@psy.vu.nl

Eunethydis

Eunethydis (European Network on Hyperkinetic Disorders) is a group of child psychiatrists, paediatricians, psychologists, neurologists and neuroscientists who gather annually to discuss research in Attention-Deficit Hyperactivity Disorder (ADHD) or Hyperkinetic Disorder (HKD) and work on key theoretical models of ADHD/HKD [65]. Eunethydis explicitly attempts to forge theoretical links between clinical and fundamental science. This paper briefly sketches the main neurobiological models used by Eunethydis. The paper has a cognitive neuroscience character and will, therefore, not deal with clinical science issues of taxonomy [27], epidemiology [20] longitudinal research [96] or treatment studies in Europe [17].

■ Neurobiology of ADHD

One of the main contributions of Eunethydis to the field has been to encourage the development of neurobiological models and mechanisms of HKD/ADHD. The pri-

mary model from this work proposes that ADHD/HKD is due to hypofunctioning of dopamine (DA), which produces altered reinforcement processes and deficient extinction [65, 66]. A key mechanism involved in reinforcement process is the nucleus accumbens [56]. An alternative “hyper”–“hypo” model of DA has also been proposed [52]. Hypofunctioning of DA is thought to give rise to delay aversion, impulsiveness, deficient response suppression and increased behavioural variability. The dopaminergic hypothesis of ADHD/HKD is crucial to both animal and molecular genetic studies of ADHD and explaining methylphenidate’s therapeutic effect.

One candidate mechanism is how perinatal anoxia and dopamine relate. Brake, Sullivan, & Gratton [9] found that, while the left prefrontal cortex (PFC) DA response was unaffected by perinatal anoxia, the right PFC response was dramatically suppressed. Moreover, dopamine transporter (DAT) levels were elevated selectively in the right PFC.

The Naples group [22] identified an epigenetic mechanism [41] suggesting that tyrosine hydroxylase and dopamine transporter gene expression are significantly and transiently reduced in one rat ADHD model (SHR) midbrain during the first month of postnatal develop-

ment. High-affinity DA uptake activity was significantly reduced in synaptosomes from the striatum of 1-month-old SHR compared to controls. Down-regulation of DA neurotransmission occurs in the midbrain of this ADHD animal model in a developmentally regulated temporal window during postnatal development. These findings suggest that epigenetic factors acting during critical periods of post-natal development may interact with genetic determinants in ADHD.

The dopamine dysfunction hypothesis has led to the search for candidate dopamine gene(s) encoding the dopamine transporter. Currently, several Eunethydis centres are involved in the international molecular ADHD genetic study (Image) led by Dr S. Faraone. Previously, Faraone, Doyle, Mick & Biederman [19] performed a meta-analysis of case-control and family-based studies. Faraone et al. found a small association between ADHD and the dopamine receptor gene D_4 (DRD_4). Mill et al. [46] genotyped five polymorphisms across D_4 and noted that a haplotype of markers may confer susceptibility to ADHD, confirming another meta-analysis of the involvement of dopamine system genes in ADHD [44]. Holmes et al. [30] reported that DRD_4 was significantly associated with ADHD children comorbid for Conduct Disorder (CD). While this search for candidate genes for ADHD is given considerable interest, it is unclear in how far the candidates are specific for ADHD/HKD or are also associated with a region implicated in autism [84].

■ Behavioural genetics and cognition

The dopamine-dysfunction hypothesis in ADHD/HKD has inspired research into the relation between cognitive performance, particularly response suppression, and molecular as well as behavioural genetics [42]. Swanson et al. [93] examined the relation between the 7-repeat allele of the dopamine receptor D_4 gene and cognitive neuropsychological performance. No relation was found between the 7-repeat and inhibitory deficits in ADHD, possibly due to the small sample size. However, Kuntsi & Stevenson [37] found in a twin study that variability of speed on the Stop Signal task [53] was associated with genetic effects on extreme hyperactivity ratings, and a significant proportion was shared with the genetic effects on the variability in speed of processing. This latter variable emerged most consistently and was the best discriminator between the hyperactive and control groups in the non-genetic analyses [38]. Using a familial design Crosbie and Schachar [16] also found deficient response inhibition to be a possible marker of familial ADHD. In a sib-proband study, Slaats-Willemse et al. [82] found that non-affected siblings of ADHD probands had response inhibition deficits, but that their performance on the inhibition

measures did not significantly differ from their siblings with ADHD.

The hunt for ADHD genes and their relation to cognitive neuropsychological characteristics of ADHD/HKD is now in full swing in various European centres and in the coming ten years significant relationships may be expected to be found.

■ Neurophysiological mechanisms and neuroanatomy

Neurophysiological research in ADHD/HKD originally sought to determine the arousal/activation deficit in ADHD [68], stimulus seeking behaviour [1] and, recently, whether disinhibition of the nervous system can explain ADHD/HKD [51]. A variety of techniques have been used to reach these goals. Power spectrum analysis indicated less right lateralized asymmetry in ADHD boys than in ADHD girls [6]. Both event-related potential (ERP) paradigms and whole head analyses of the electroencephalogram (EEG) with low-resolution electromagnetic tomography (LORETA) have been used in several studies of ADHD/HKD children [10]. The ERP research indicated that ADHD have reduced motor preparation [26], and reduced P300 to both cued and non-cued targets [31, 40, 54] suggesting under activation of the posterior attention system in ADHD/HKD children [4].

Additional evidence has been found that children with HKD have reduced P300 amplitudes to cues and distracters suggesting a deficiency in attentional orienting/inhibition [4]. Global field power of the cue-CNV microstate related to anticipation and preparation was reduced in HKD but not in children comorbid for HKD and CD (HKD-CD). Reduction in global power is consistent with a model that HKD children (without ODD/CD) suffer from an energetical state dysregulation [75] rather than only a response suppression deficit [5]. Subsequent analysis demonstrated that HKD children but not children with HKD-CD were impaired in processing a signal requiring motor preparation [4]. ERP paradigms differentiate ADHD/HKD from ADHD/HKD comorbid for CD or Tic disorder [61, 103] consistent with MRI studies of ADHD and Tourette syndrome [32]. ERP research has successfully shown the therapeutic effects of methylphenidate in ADHD as effecting central and motor output processes [31] and differentiates methylphenidate from desipramine and l-dopa [55].

Another window into inhibition in ADHD has been to use transcranial magnetic stimulation (TMS). Moll et al. [47] found reduced intracortical inhibition in ADHD children but not in the cortical silent period, suggesting a specific inhibition, motor dysfunction in ADHD. This finding is consistent with structural MRI findings of smaller basal ganglia volumes in ADHD children [2, 11, 12, 21, 97] and smaller cerebellar volume in ADHD [7].

Developmental transitions in brain structure and function may be crucial in ADHD. For example, the caudate nucleus differences between ADHD and control children disappear in mid adolescence [14]. Amplitude reductions of brain electric visual evoked potentials and high ratio of slow frequencies in standard EEG have been found to normalize from eight to thirteen years, suggesting that it is only later that the sensory pathways of hyperkinetic children presumably reach the same functional level as that of healthy children [60]. It remains to be determined whether reduced intracortical inhibition in ADHD follows a similar developmental path and cannot be considered a biological marker of ADHD [13]. Further, differences in imaging technique are producing different results between centres. In contrast to the reduction in volume reported in earlier studies, more recent work is showing that an *enlarged* superior frontal lobe is correlated with higher hyperactivity scores [91]. The significance of these results remains to be determined.

Finally, sleep research in ADHD has gained the insight that a forced ultradian cycling appears characteristic for the sleep of children with ADHD, which may be related to alterations of brain monoamines and cortical inhibitory control [33].

■ Neuropsychological mechanisms

Recent reviews have summarized the two main goals of neuropsychological research in ADHD as: (1) determination of the neural networks in the disorder and (2) establish the specificity of the neuropsychological dysfunction [77]. The five candidate neuropsychological mechanisms of ADHD are a prefrontal inhibition dysfunction [5], working memory/selective attention dysfunction [13, 94], a timing/motor deficit [63, 64] and a deficiency of state regulation [75]. These candidates are briefly discussed below.

There has been a systematic evaluation of the inhibition hypothesis from preschool [90] to school-aged children [69]. More than 20 studies of the Stop Signal Task [43] have been published indicating that ADHD children have a longer, variable, inhibitory function than controls but this finding is not specific to ADHD but is common to ODD and CD [53]. Furthermore, both children with a fast-guess reading style [72] and children with Higher Function Autism (HFA) have also been demonstrated to have poorer Stop Signal Task performance than controls [23] and HFA to have this more than children with Tourette's syndrome [101]. Rubia et al. [63] found that the right prefrontal cortex and left caudate nucleus showed less activation in ADHD patients than controls. Taken together these results support an inhibition deficit hypothesis in ADHD/HKD but this is not specific to ADHD or to externalizing disorders.

Scheres et al. [70] demonstrated that the disinhibition effect did not generalize to other forms of inhibitory control, suggesting that a generalized loss of inhibition in ADHD is not an adequate account of ADHD [5]. This result is consistent with the TMS study that methylphenidate enhances intracortical facilitation but not intracortical inhibition [48]. Solanto et al. [86] demonstrated that in addition to inhibition, delay aversion could explain independent variance [89]. This finding has led to a reformulated model of ADHD in which both delay aversion and inhibition are conceived of as dual pathways in ADHD [87, 88].

Two key candidates in ADHD research are, first, the relation between attention and Working Memory and, second, the relation between inhibition and response conflict. Nigg [51] proposed that inhibition deficits in HFA might be explained by WM, whereas in ADHD inhibition deficits can be better explained by a motor dysregulation [5]. Geurts et al. [23] found that this intriguing dichotomy did not hold up in an executive function study of both children with HFA and ADHD. A recent meta-analysis of response conflict, as measured by the Stroop, indicates that Stroop response conflict is an unattractive measure in ADHD [49]. It remains to be determined whether specific working memory candidates can be discovered which can separate ADHD from HFA children.

One explanation of ADHD has been that performance deficits are due to poor selective attention [18]. Early research defined selective attention as a limitation in the rate of controlled processing located in short-term memory [71, 80]. This definition led to a series of studies using both a memory and visual load paradigm [74]. The conclusion from that research programme was that ADHD children did not suffer from a selective attention deficit (see for review [73]). Hence, a selective attention deficit does not appear to be implicated in ADHD.

The role of executive functions [57] and the frontal lobes [81] has been a candidate mechanism. However, MRI reports [7] and variability in timing [62, 63] have refocused research on the role of the cerebellum in ADHD [13, 85]. Response selection difficulties [45] as well as poor balancing have been noted in ADHD [58] suggesting possible cerebellar deficiency in ADHD. These findings have revived interest in minor neurological dysfunction [35, 99, 100] and differentiation of ADHD children from children with a neurological disorder [34]. In the near future, there may be an attempt to differentiate ADHD from HFA children [23] or children with clumsiness [25] via timing and cerebellar neuropsychological tasks. Alternatively, pragmatic dysfunction in both HFA and to a lesser degree in ADHD [24] suggests that this particular aspect of executive functioning may be relevant in some subgroups [95], although one study found no significant interactions be-

tween hyperactivity and Specific Language Impairment on any neuropsychological measure [102].

The cognitive energetic dysregulation hypothesis argues that ADHD performance deficits are due to non-optimal energetic allocation, which leads to dysfunctioning at either a specific energetic pool, at the motor stage of processing or at a higher level of executive management [8, 73, 75]. This model can be traced to earlier work in which ADHD children were shown to be less physiologically aroused and faster habituators than normal controls [67]. Herpertz et al. [29] found recently that children with CD and ADHD + CD showed decreased electrodermal responses and accelerated habituation compared to children with ADHD alone and controls, thus suggesting possible energetic differences in externalizing disorders.

Differences in energetic allocation have been reported in response conflict tasks [59], which substantially effect responses inhibition in ADHD [83]. Sonuga-Barke [87] found in two studies that AD/HD children exhibited a quadratic performance function as predicted by the cognitive-energetic model of AD/HD. Steger et al. [92] found that in long intervals in a motor task revealed increased errors and onset variability. Using the Flanker task, Crone, Jennings and van der Molen [15] reported that ADHD children responded less accurately under the threat of punishment. Phasic heart rate did not differ between groups, but immediate reward feedback induced greater heart rate responses in control than in ADHD children. Inappropriate energetic allocation has also been observed in vigilance studies using EEG measures [3, 28]. Further gamma band responses showed a right lateral preference in ADHD children to both attended and non-attended stimuli, suggesting poor energetic allocation [104]. An interesting link be-

tween molecular genetic studies and the dopamine transporter (DAT) has been laid by a review of SPECT studies in adult ADHD patients: Patients with ADHD and with a history of nicotine abuse both displayed lower values of DAT density in SPECT than non-smokers with ADHD. DAT seem to be elevated in non-smoking ADHD patients suffering from the purely inattentive subtype of ADHD as well as in those with the combined or purely hyperactive/impulsive subtype [36]. Thapar et al. [98] reported that, although genetic influences accounted for most of the variance in offspring ADHD, maternal smoking during pregnancy accounted for a significant environmentally mediated association. This suggests that ADHD patients may, in some cases, be employing nicotine as a method for self-therapy to compensate for their non-optimal energetic allocation and be related to brain activation patterns [50].

While ADHD is often suggested to be a good example of dysexecutive functioning [5], comparison of ADHD children with HFA suggests that HFA children have a much more profound and extensive deficit in executive functioning than ADHD children [23, 76, 79], although this may depend upon the age at which the child is tested [6, 90]. More work is required to determine norms for preschool severity in ADHD than currently available. It is clear that a much wider neural network is involved in ADHD than one specific to the prefrontal cortex and involves some features of WM, timing and energetic allocation [13, 77]. Differentiation of ADHD from correlated disorders such as dyscalculia [78], developmental coordination disorder [39], HFA, Tourette and Fragile-X syndrome is a matter of urgent study. The specific candidates have been described above and will be tested for their validity in future work associated with Eunethydis.

References

1. Antrop I, Roeyers H, van Oost P, Buysee A (2000) Stimulation seeking and hyperactivity in children with ADHD. *J Child Psychol Psychiatry* 41:225–231
2. Aylward EH, Reiss AL, Reader MJ, Singer HS, Brown JE, Denckla MB (1996) Basal ganglia volumes in children with attention-deficit hyperactivity disorder. *J Child Neurol* 11:112–125
3. Banaschewski T, Brandeis D, Heinrich H, Albrecht B, Brunner E, Rothenberger A (2003) Association of ADHD and conduct disorder – brain electrical evidence for the existence of a distinct subtype. *J Child Psychol Psychiatry* 43:1–21
4. Banaschewski T, Brandeis D, Heinrich H, Albrecht B, Brunner E, Rothenberger A (2004) Questioning inhibitory control as the specific deficit of ADHD – evidence from brain electrical activity. *J Neural Transm* 111:841–864 Epub 2003 Oct 24
5. Barkley RA (1997) Behavioural inhibition, sustained attention, and executive functions: constructing a unifying theory of ADHD. *Psychol Bull* 121:65–94
6. Baving L, Laucht M, Schmidt MH (1999) Atypical frontal brain activation in ADHD: Preschool and elementary school boys and girls. *Journal of the American Acad Child Adolesc Psychiatry* 38:1363–1371
7. Berquin PC, Giedd JN, Jacobsen LK, Hamburger SD, Krain AL, Rapoport JL, Castellanos FX (1998) Cerebellum in attention-deficit hyperactivity disorder: a morphometric MRI study. *Neurology* 50:1087–1093
8. Borger N, van der Meere JJ (2000) Visual behaviour of ADHD children during an attention test: an almost forgotten variable. *Attention-Deficit Hyperactivity Disorder. J Child Psychol Psychiatry* 41: 525–532
9. Brake WG, Sullivan RM, Gratton A (2000) Perinatal distress leads to lateralized medial prefrontal cortical dopamine hypofunction in adult rats. *J Neurosci* 20:5538–5543
10. Brandeis D, Banaschewski T, Baving L, Georgiewa P, Blanz P, Schmidt MH, Warnke A, Steinhausen HC, Rothenberger A, Scheuerpflug P (2002) Multicenter P300 brain mapping of impaired attention to cues in hyperkinetic children. *Journal of the American Acad Child Adolesc Psychiatry* 41:990–998
11. Casey BJ, Castellanos FX, Giedd JN, Marsh WL, Hamburger SD, Schubert AB, Vauss YC, Vaituzis CK, Dickstein DP, Sarfatti SE, Rapoport JL (1997) Implication of right frontostriatal circuitry in response inhibition and attention deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 36: 374–383

12. Castellanos FX, Giedd JN, Eckburg P, Marsh WL, Vaituzis C, Kaysen D, Hamburger S, Hamburger S, Rapoport JL (1994) Quantitative morphology of the caudate nucleus in attention deficit hyperactivity disorder. *Am J Psychiatry* 151:1791–1796
13. Castellanos FX, Tannock R (2002) Neuroscience of Attention Deficit/Hyperactivity Disorder. *Nat Neurosci Rev* 3: 617–628
14. Castellanos FX, Lee PP, Sharp W, Jeffries NO, Greenstein DK, Clasen LS, Blumenthal JD, James RS, Ebens CL, Walter JM, Zijdenbos A, Giedd JN, Rapoport JL (2002) Developmental trajectories of brain volume abnormalities in children and adolescents with attention-deficit/hyperactivity disorder. *J Am Medical Assoc* 288:1740–1748
15. Crone EA, Jennings R, van der Molen M (2003) Sensitivity to interference and response contingencies in attention-deficit/hyperactivity disorder. *J Child Psychol Psychiatry* 44:214–226
16. Crosbie J, Schachar R (2001) Deficient inhibition as a marker for familial ADHD. *Am J Psychiatry* 158:1884–1890
17. Döpfner M, Gerber WD, Banaschewski T, Breuer D, Freisleder FJ, Gerber-von Müller G, Günter M, Hässler F, Ose C, Rothenberger A, Schmeck K, Sinzig J, Stadler C, Uebel H, Lehmkuhl G (2004) Comparative efficacy of once-a-day extended-release methylphenidate, two-times-daily immediate release methylphenidate, and placebo in a laboratory school setting. *Eur Child Adolesc Psychiatry* 13(Suppl 1):I80–I92
18. Douglas VI (1999) Cognitive control processes in attention-deficit/hyperactivity disorder. In: Quay HC, Hogan A (eds) *Handbook of Disruptive Behavior Disorders*. New York, Plenum Press, pp 105–138
19. Faraone SV, Doyle AE, Mick E, Biederman J (2001) Meta-analysis of the association between the 7-repeat allele of the dopamine D(4) receptor gene and attention deficit hyperactivity disorder. *Am J Psychiatry* 158:1052–1057
20. Faraone SV, Sergeant J, Gillberg C, Biederman J (2003) The worldwide prevalence of ADHD: is it an American condition? *World Psychiatry* 2:104–113
21. Filipek PA, Semrud-Clikeman M, Steingard RJ, Kennedy DN, Biederman J (1997) Volumetric MRI analysis comparing attention-deficit hyperactivity disorder and normal controls. *J Child Neurol* 48:589–601
22. Fresiello A, Grammatikopoulos G, Pignatelli M, Sadleir AG (2002) Environmental factors during postnatal period modify activity and non-selective attention in the Naples High-Excitability rat. *Behav Brain Res* 130:111–115
23. Geurts HM, Verté S, Oosterlaan J, Roeyers R, Sergeant JA (2003) How specific are executive functioning deficits in attention deficit hyperactivity disorder and autism? *J Child Psychol Psychiatry* 43:1–19
24. Geurts HM, Verté S, Oosterlaan J, Roeyers H, Hartman CA, Mulder EJ, van Berckelaer-Onnes IA, Sergeant JA (2004) Can the Children's Communication Checklist Differentiate between Children with Autism, Children with ADHD, and Normal Controls? *J Child Psychol Psychiatry* (in press)
25. Gillberg C, Kadesjo B (2003) Why bother about clumsiness? The implications of having developmental coordination disorder (DCD). *Neural Plasticity* 10:59–68
26. Grünewald-Züebler E, Grünewald G, Rasche A, Netz J (1978) Contingent negative variation and alpha attenuation responses in children with different abilities to concentrate. *Electroencephalogr Clin Neurophysiol* 44:37–47
27. Hartman CA, Hox J, Mellenbergh GJ, Boyle MH, Offord DR, Racine Y, McNamee J, Gadow KD, Sprafkin J, Kelly KL, Nolan EE, Tannock R, Schachar R, Schut H, Postma I, Drost R, Sergeant JA (2001) When taxonomy meets data. *J Child Psychol Psychiatry* 42:817–836
28. Hennighausen K, Schulte-Körne G, Warnke A, Remschmidt H (2000) Contingent negative variation (CNV) in children with hyperkinetic syndrome – an experimental study using the Continuous Performance Test (CPT). *Zeitschrift für Kinder und Jugendpsychiatrie Psychotherapie* 28:239–246
29. Herpertz SC, Mueller B, Wenning B, Qunaibi M, Lichterfeld C, Herpertz-Dahlmann B (2003) Autonomic responses in boys with externalizing disorders. *Journal of Neural Transmission* 110:1181–1195
30. Holmes J, Payton A, Barrett J, Harrington R, McGuffin P, Owen M, Ollier W, Worthington J, Gill M, Kirley A, Hawi Z, Fitzgerald M, Asherson P, Curran S, Mill J, Gould A, Taylor E, Kent L, Craddock N, Thapar A (2002) Association of DRD4 in children with ADHD and comorbid conduct problems. *Am J Med Gen (Neuropsychiatric Genetics)* 114: 150–153
31. Jonkman LM, Kemner C, Verbaten MN, Van Engeland H, Kenemans JL, Camfferman G, Buitelaar JK, Koelega HS (1999) Perceptual and response interference in children with attention-deficit hyperactivity disorder, and the effects of methylphenidate. *Psychophysiology* 36:419–429
32. Kates WR, Frederikse M, Mostofsky SH, Folley BS, Cooper K, Mazur-Hopkins P, Kofman O, Singer HS, Denckla MB, Pearlson GD, Kaufmann WE (2002) MRI parcellation of the frontal lobe in boys with attention deficit hyperactivity disorder or Tourette syndrome. *Psychiatry Research Neuroimaging* 116: 63–81
33. Kirov R, Kinkelbur J, Heipke S, Kostanecka-Endress T, Westhoff M, Cohrs S, Rütger E, Hajak G, Banaschewski T, Rothenberger A (2004) Is there a specific polysomnographic sleep pattern in children with ADHD. *J Sleep Res* 13: 87–93
34. Konrad K, Gauggel S, Manz A, Schöll M (2000) Inhibitory control in children with traumatic brain injury (TBI) and children with attention deficit/hyperactivity disorder (ADHD). *Brain Injury* 14:859–875
35. Kramer F, Pollnow H (1932) Über eine hyperkinetische Erkrankung im Kindesalter. *Monatsschrift Psychiatrie und Neurologie* 82:1–40
36. Krause KH, Dreselb SH, Krause J, la Fougere C, Ackenheil M (2003) The dopamine transporter and neuroimaging in attention deficit hyperactivity disorder. *Neurosci Biobehav Rev* 27: 605–613
37. Kuntsi J, Stevenson J (2001) Psychological mechanisms in hyperactivity: II The role of genetic factors. *J Child Psychol Psychiatry* 42:211–219
38. Kuntsi J, Oosterlaan J, Stevenson J (2001) Psychological mechanisms in hyperactivity: I Response inhibition deficit, working memory impairment, delay aversion, or something else? *J Child Psychol Psychiatry* 42:199–210
39. Landgren M, Kjellman B, Gillberg C (1998) Attention deficit disorder with developmental coordination disorders. *Arch Dis Childhood* 79:207–212
40. Leeuwen TH van, Steinhausen HC, Overtom CCE, Pasqual-Marqui RD, Rothenberger A, Sergeant JA, Brandeis D (1998) The continuous performance test revisited with neuroelectric mapping: impaired orienting in children with attention deficits. *Behav Brain Res* 94:97–110
41. Leo D, Sorrentino E, Volpicella F, Eymanb M, Greco D, Viggiano D, di Porzio U, Perrone-Capano C (2003) Altered midbrain dopaminergic neurotransmission during development in an animal model of ADHD. *Neurosci Biobehav Rev* 27:661–669
42. Levy F, Hay D (2001) (eds) *Attention, Genes and ADHD*. Hove, Brunner-Routledge
43. Logan GD, Cowan WB (1984) On the ability to inhibit thought and action: A theory of an act of control. *Psychol Rev* 91:295–327

44. Maher BS, Marazita ML, Ferrell RE, Vanyukov MM (2002) Dopamine system genes and attention deficit hyperactivity disorder: a meta-analysis. *Psychiatr Gen* 12:207–215
45. Meere JJ van der, Baal M van, Sergeant JA (1989) The additive factor method: A differential diagnostic tool in hyperactivity and learning disablement. *J Abnorm Child Psychol* 17:409–422
46. Mill J, Fisher N, Curran S, Richards S, Taylor E, Asherson P (2003) Polymorphisms in the dopamine D₄ receptor gene and attention-deficit hyperactivity disorder. *Neuro Report* 14: 1463–1466
47. Moll GH, Heinrich H, Trott G-E, Wirth S, Rothenberger A (2000) Deficient intracortical inhibition in drug-naïve children with attention-deficit hyperactivity disorder is enhanced by methylphenidate. *Neuroscience Letters* 284:121–125
48. Moll GH, Heinrich H, Rothenberger A (2003) Methylphenidate and intracortical excitability: opposite effects in healthy subjects and attention-deficit hyperactivity disorder. *Acta Psychiatr Scand* 107:69
49. Mourik van R, Oosterlaan J, Sergeant JA (2004) The Stroop Revisited: A Meta-Analysis of Interference Control in AD/HD (submitted)
50. Newhouse PA, Potter A, Singh A (2004) Effects of nicotine stimulation on cognitive performance. *Curr Opin Pharmacol* 4:1–11
51. Nigg JT (2000) On inhibition/disinhibition in developmental psychopathology: views from cognitive and personality psychology and a working inhibition taxonomy. *Psychol Bull* 126: 220–246
52. Oades R (2002) Dopamine may be 'hyper' with respect to noradrenalin metabolism, but 'hypo' with respect to serotonin metabolism in children with attention-deficit hyperactivity disorder. *Behav Brain Res* 130:97–102
53. Oosterlaan J, Logan GD, Sergeant JA (1998) Response inhibition in AD/HD, CD, comorbid AD/HD+CD, anxious and control children: a meta-analysis of studies with the stop task. *J Child Psychol Psychiatry* 39:411–426
54. Overtom CC, Verbaten MN, Kemner C, Kenemans JL, van Engeland H, Buitelaar JK, Camfferman G, Koelega HS (1998) Associations between event related potentials and measures of attention and inhibition in the continuous performance task in children with ADHD and normal controls. *J Am Acad Child Adolesc Psychiatry* 37:977–985
55. Overtom CCE, Verbaten MN, Kemner C, Kenemans JL, van Engeland H, Buitelaar JK, van der Molen MW, van der Gugten J, Westenberg H, Maes RAA, Koelega HS (2003) Effects of methylphenidate, desipramine, and l-dopa on attention and inhibition in children with Attention Deficit Hyperactivity Disorder. *Behavioural Brain Research* 145:7–15
56. Papa M, Berger DE, Sagvolden T, Sergeant JA, Sadile AG (1998) A quantitative cytochrome oxidase mapping study, cross-regional and neurobehavioural correlations in the anterior forebrain of an animal model of attention deficit hyperactivity disorder. *Behav Brain Res* 94:197–212
57. Pennington BF, Ozonoff S (1996) Executive functions and developmental psychopathology. *J Child Psychol Psychiatry* 37:51–87
58. Raberger T, Wimmer H (2003) On the automaticity/cerebellar deficit hypothesis of dyslexia: balancing and continuous rapid naming in dyslexic and ADHD children. *Neuropsychologia* 41: 1493–1497
59. Ridderinkhof KR, Scheres A, Oosterlaan J, Sergeant JA (2004) Distribution-analytical techniques in the study of ADHD: delta plot analyses reveal deficits in response suppression that are eliminated by methylphenidate treatment. *J Abnorm Psychol* (in press)
60. Rothenberger A (1995) Electrical brain activity in children with hyperkinetic syndrome: evidence of frontal cortical dysfunction. In: Sergeant J (ed) *Eunethydis – European Approaches to Hyperkinetic Disorder*. Trümpi, Zürich, pp 255–270
61. Rothenberger A, Banaschewski T, Heinrich H, Moll GH, Schmidt MH, van't Klooster B (2000) Comorbidity in ADHD-children: effects of coexisting conduct disorder or tic disorder on event-related potentials in an auditory selective-attention task. *Eur Arch Psychiatry Clin Neurosci* 250:101–110
62. Rubia K, Overmeyer S, Taylor E, Brammer M, Williams S, Simmons A, Bullmore ET (1999) Hypofrontality in attention deficit hyperactivity disorder during higher-order motor control: A study with functional MRI. *Am J Psychiatry* 156:891–896
63. Rubia K, Taylor A, Taylor E, Sergeant JA (1999) Synchronization, anticipation, and consistency in motor timing of children with dimensionally defined attention deficit hyperactivity behaviour. *Perceptual and Motor Skills* 89: 1237–1258
64. Rubia K, Noorlos J, Smith A, Gunning B, Sergeant J (2003) Motor timing deficits in community and clinical boys with hyperactive behavior: The effect of methylphenidate on motor timing. *J Abnorm Child Psychol* 31:301–313
65. Sagvolden T, Sergeant JA (1998) Attention-deficit hyperactivity disorder: from brain dysfunctions to behaviour. *Behav Brain Res* 94:1–10
66. Sagvolden T, Johansen EB, Aase H, Russell VA (2004) A dynamic developmental theory of attention-deficit/hyperactivity disorder. *Behavioral and Brain Sciences* (in press)
67. Satterfield JH, Dawson ME (1971) Electrodermal correlates of hyperactivity in children. *Psychophysiology* 8:191–197
68. Satterfield JH, Cantwell DP, Satterfield BT (1974) Pathophysiology of the hyperactive child syndrome. *Arch Gen Psychiatry* 31:839–844
69. Scheres A, Oosterlaan J, Sergeant JA (2001) Response execution and inhibition in children with AD/HD and other disruptive disorders: The role of behavioral activation. *J Child Psychol Psychiatry* 42:347–357
70. Scheres A, Oosterlaan J, Swanson J, Morein-Zamir S, Meiran N, Schut H, Vlasveld L, Sergeant JA (2003) The effect of methylphenidate on three forms of response inhibition in boys with AD/HD. *J Abnorm Child Psychol* 31: 107–122
71. Schneider W, Shiffrin RM (1977) Controlled and automatic human information processing: I Detection, search and attention. *Psychol Rev* 84:1–66
72. Schoot van der M, Licht R, Horsley TM, Sergeant JA (2000) Inhibitory deficits in reading disability depend on subtype: Guessers but not spellers. *Child Neuropsychol* 6:297–312
73. Sergeant JA, Meere JJ van der (1990) Additive factor methodology applied to psychopathology with special reference to hyperactivity. *Acta Psychol* 74: 277–295
74. Sergeant JA, Scholten CA (1983) A stages-of-information processing approach to hyperactivity. *J Child Psychol Psychiatry* 22:49–60
75. Sergeant JA, Oosterlaan J, Meere van der JJ (1999) Information processing and energetic factors in Attention-Deficit/Hyperactivity Disorder. In: Quay HC; Hogan A (eds) *Handbook of Disruptive Behavior Disorders*. New York, Plenum Press, pp 75–104
76. Sergeant JA, Geurts H, Oosterlaan J (2002) How specific is a deficit of executive functioning for Attention-Deficit/Hyperactivity Disorder? *Behav Brain Res* 130:3–28
77. Sergeant JA, Geurts H, Huijbregts S, Scheres A, Oosterlaan J (2003) The top and the bottom of ADHD: a neuropsychological perspective. *Neurosci Biobehav Rev* 27:583–592
78. Shalev RS (2002) Developmental dyscalculia. In: Segalowitz SJ, Rapin I (eds) *Handbook of Neuropsychology*, 2nd ed, Vol. 8, Part II. Amsterdam, Elsevier, pp 118–143

79. Shallice T, Marzocchi GM, Coser S, Del Savio M, Meuter RF, Rumiati R (2002) Executive function profile of children with attention deficit hyperactivity disorder. *Devel Neuropsychol* 21:43–71
80. Shiffrin RM, Schneider W (1977) Controlled and automatic information processing: II. Perceptual learning, automatic attending, and a general theory. *Psychol Rev* 84:127–190
81. Shue KL, Douglas VI (1992) Attention deficit hyperactivity disorder and the frontal lobe syndrome. *Brain and Cognition* 20:104–124
82. Slaats-Willemse D, Swaab-Barneveld H, De Sonneville L, van der Meulen E, Buitelaar J (2003) Deficient response inhibition as a cognitive endophenotype of ADHD. *J Am Acad Child Adolesc Psychiatry* 42:1242–1248
83. Slusarek M, Velling S, Bunk D, Eggers C (2001) Motivational effects on inhibitory control in children with ADHD. *J Am Acad Child Adolesc Psychiatry* 40:355–363
84. Smalley SL, Kustanovich V, Minassian SL, Stone JL, Ogdie MN, James J, McGough JJ, McCracken JT, MacPhie L, Francks C, Simon E, Fisher SE, Cantor RM, Monaco AP, Nelson SF (2002) Genetic linkage of attention-deficit/hyperactivity disorder on chromosome 16p13, in a region implicated in autism. *Am J Human Gen* 71:959–963
85. Smith Taylor, Lidzba, Rubia (2003) A right hemispheric frontocerebellar network for time discrimination of several hundreds of milliseconds. *NeuroImage* 20:344–350
86. Solanto MV, Abikoff H, Sonuga-Barke E, Schachar R, Logan GD, Wigal T, Hechtman L, Hinshaw S, Turkel E (2001) The ecological validity of delay aversion and response inhibition as measures of impulsivity in AD/HD: a supplement to the NIMH multimodal treatment study of AD/HD. *J Abnorm Child Psychol* 29: 215–228
87. Sonuga-Barke EJS (2002) Psychological heterogeneity in AD/HD – a dual pathway model of behaviour and cognition. *Behav Brain Res* 130:29–36
88. Sonuga-Barke EJS (2002) Interval length and time-use by children with AD/HD: a comparison of four models. *J Abnorm Child Psychol* 30:257–264
89. Sonuga-Barke EJS, Taylor E, Sembi S, Smith J (1992) Hyperactivity and delay aversion-I. The effect of delay choice. *J Child Psychol Psychiatry* 33:387–398
90. Sonuga-Barke EJS, Dalen L, Daley D, Remington B (2002) Are planning, working memory, and inhibition associated with individual differences in preschool ADHD symptoms? *Devel Neuropsychol* 21:255–272
91. Sowell ER, Thompson PM, Welcome SE, Henkenius AL, Toga AW, Peterson BS (2003) Cortical abnormalities in children and adolescents with attention-deficit hyperactivity disorder. *Lancet* 362:1699–1707
92. Steger J, Imhof K, Coutts E, Gundelfinger R, Steinhausen HC, Brandeis D (2001) Attentional and neuromotor deficits in ADHD. *Devel Med Child Neurol* 43:172–179
93. Swanson J, Oosterlaan J, Murias M, Schuck S, Flodman P, Spence MA, Waddell M, Ding Y, Smith M, Mann M, Carlson C, Kennedy JL, Sergeant JA, Leung P, Zhang YP, Sadeh A, Chen C, Whalen CK, Babb KA, Moyzis R, Posner MI (2000) Attention deficit/hyperactivity disorder children with a 7-repeat allele of the dopamine receptor D₄ gene have extreme behavior but normal performance on critical neuropsychological tests of attention. *Proc Natl Acad Sci USA* 97:4754–4759
94. Tannock R (1998) Attention deficit hyperactivity disorder. *J Child Psychol Psychiatry* 39:65–99
95. Tannock R, Martinussen R, Frijters J (2000) Naming speed performance and stimulant effects indicate effortful, semantic processing deficits in attention-deficits/hyperactivity disorder. *J Abnorm Child Psychol* 28:237–252
96. Taylor E, Chadwick O, Heptinstall E, Danckaerts M (1996) Hyperactivity and conduct problems as risk factors for adolescent development. *J Am Acad Child Adolesc Psychiatry* 35: 1213–1226
97. Teicher MH, Anderson CM, Polcari A, Glod CA, Maas LC, Renshaw PF (2000) Functional deficits in basal ganglia of children with attention-deficit/hyperactivity disorder shown with functional magnetic resonance imaging relaxometry. *Nat Med* 6:470–473
98. Thapar A, Fowler T, Rice F, Scourfield J, van den Bree M, Thomas H, Harold G, Hay D (2003) Maternal smoking during pregnancy and attention deficit hyperactivity disorder symptoms in offspring. *Am J Psychiatry* 160:1985–1989
99. Touwen BCL (1979) *The Neurological Examination of the Child with Minor Nervous Dysfunction* 2nd edition. London, William Heinemann Medical Books, p 105
100. Tupper DE (1987) (ed) *Soft Neurological Signs*. Orlando, Grune and Stratton
101. Verté S, Geurts HM, Roeyers H, Oosterlaan J, Sergeant JA (2004) Executive Functioning in Children with Autism and Tourette Syndrome. *J Autism Res* (submitted)
102. Williams D, Stott CM, Goodyer IM, Sahakian BJ (2000) Specific language impairment with or without hyperactivity: neuropsychological evidence for frontostriatal dysfunction. *Devel Med Child Neurol* 42:368–375
103. Yordanova J, Dumais-Huber C, Rothenberger A, Woerner W (1997) Frontocortical activity in children with comorbidity of tic disorder and attention-deficit hyperactivity disorder. *Biol Psychiatry* 41:585–594
104. Yordanova J, Banaschewski T, Kolev V, Woerner W, Rothenberger A (2001) Abnormal early stages of task stimulus processing in children with attention-deficit hyperactivity disorder – evidence from event-related gamma oscillations. *Clin Neurophysiol* 112: 1096–1108

Copyright of European Child & Adolescent Psychiatry is the property of Springer - Verlag New York, Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.