VU Research Portal

Online Intervention for Prevention of Major Depression-Reply.

Ebert, D.D.; Buntrock, C.; Cuijpers, Pim

published in

JAMA

2016

DOI (link to publisher)

[10.1001/jama.2016.9586](https://doi.org/10.1001/jama.2016.9586)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Ebert, D. D., Buntrock, C., & Cuijpers, P. (2016). Online Intervention for Prevention of Major Depression-Reply. *JAMA*, 316(8), 881-882. <https://doi.org/10.1001/jama.2016.9586>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/306434602>

Online Intervention for Prevention of Major Depression–Reply

Article in JAMA The Journal of the American Medical Association · August 2016

DOI: 10.1001/jama.2016.9586

CITATIONS

0

READS

152

3 authors:

[David Daniel Ebert](#)

Friedrich-Alexander-University of Erlangen-Nür...

93 PUBLICATIONS 794 CITATIONS

[SEE PROFILE](#)

[Claudia Buntrock](#)

Friedrich-Alexander-University of Erlangen-Nür...

10 PUBLICATIONS 114 CITATIONS

[SEE PROFILE](#)

[Pim Cuijpers](#)

VU University Amsterdam

732 PUBLICATIONS 23,656 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:

NESDA : The Netherlands Study of Depression and Anxiety [View project](#)

Prevention Of Depression in Back Pain Patients - The PROD-BP pragmatic randomised controlled trial.
[View project](#)

All content following this page was uploaded by [David Daniel Ebert](#) on 03 September 2016.

The user has requested enhancement of the downloaded file.

Sciences Alliance, Dartmouth College, Healthcare Financial Management Association, New York City Health and Hospitals Corporation, Robert W. Baird & Co, Healthcare Billing and Management Association, Cadence Health, Pompeu Fabra University, Aon Hewitt, American Health Lawyers Association, Parenteral Drug Association, UBS, Aetna, Toshiba, Ernst and Young, Yale University, Bank of America, and New York University. No other disclosures were reported.

Online Intervention for Prevention of Major Depression

To the Editor The article by Ms Buntrock and colleagues¹ presented data on primary prevention of major depression in patients who did not exhibit prior depressive episodes or secondary prevention in patients who had prior depressive episodes using an online intervention. Only 6 one-hour sessions provided a substantial and sustained effect for 1 year. If a major depressive episode is already present, usually 12 to 20 individual sessions of cognitive behavioral therapy are needed to achieve response or remission,² although fewer may also be effective.³ However, some issues should be clarified.

First, could the authors explain the abrupt decrease in number of participants without a major depressive episode found in both treatment groups, but especially in the control group, after 26 weeks (Figure 2 in the article)?

Second, an online trainer was used in the intervention group. Although the trainer did not perform psychotherapy, it would be important to know if the trainer intervened if more severe depressive symptoms, such as suicidal thoughts, were mentioned by the participants. Could this have influenced the outcome of the trial? How many hours were online trainers engaged in the guidance of patients?

Third, inpatient treatment rates and sickness leaves attributable to depression have substantially increased over the past decade^{4,5} in Germany. Therefore, an intervention like the one described in the study could not only provide benefit for people with incipient symptoms and their health insurance arrangements, but also for psychiatrists, because they could focus on more severely ill patients who have already developed the full clinical syndrome but remain on long waiting lists for care. Furthermore, web-based treatment provides an opportunity for patients living in more remote areas to receive intervention.

Ion Angheliescu, MD

Author Affiliation: Department of Psychiatry, Clinic Dr Fontheim, Liebenburg, Germany.

Corresponding Author: Ion Angheliescu, MD, Department of Psychiatry, Clinic Dr Fontheim, Lindenstrasse 15, 38704 Liebenburg, Germany (ion.angheliescu@charite.de).

Conflict of Interest Disclosures: The author has completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

1. Buntrock C, Ebert DD, Lehr D, et al. Effect of a web-based guided self-help intervention for prevention of major depression in adults with subthreshold depression: a randomized clinical trial. *JAMA*. 2016;315(17):1854-1863.

2. Gabbard GO, ed. *Gabbard's Treatments of Psychiatric Disorders*. 5th ed. Arlington, VA: American Psychiatric Association; 2014.

3. Nieuwsma JA, Trivedi RB, McDuffie J, Kronish I, Benjamin D, Williams JW. Brief psychotherapy for depression: a systematic review and meta-analysis. *Int J Psychiatry Med*. 2012;43(2):129-151.

4. Gaebel W, Kowitz S, Zielasek J. The DGPPN research project on mental healthcare utilization in Germany: inpatient and outpatient treatment of

persons with depression by different disciplines. *Eur Arch Psychiatry Clin Neurosci*. 2012;262(suppl 2):S51-S55.

5. Faktenblatt BKK. Gesundheitsatlas 2015—Blickpunkt Psyche [in German]. http://www.bkk-dachverband.de/fileadmin/publikationen/gesundheitsatlas/Faktenblatt_BKK_Gesundheitsatlas_2015.pdf. Accessed June 17, 2016.

In Reply We agree with Dr Angheliescu that it is remarkable that a major depressive episode can be prevented in people with subthreshold depression with a brief 6-session intervention. However, although this study was the first one showing that it is possible to prevent a major depressive episode using a web-based guided self-help intervention, evidence is accumulating that this can be achieved with brief face-to-face interventions.¹

Providing low-threshold interventions for the treatment of subthreshold depression and the prevention of major depressive disorder may thus not only reduce substantial disease burden for the individual but also reduce societal costs. However, most clinical guidelines do not recommend psychological treatments for subthreshold depression. Given that effect sizes can be expected to be smaller than for the treatment of major depressive disorder, cost-effectiveness studies are needed to determine whether such an approach is good value for the money.

Angheliescu points out that there was a decrease in the number of participants without a major depressive episode in the survival analyses at 26 weeks. Participants who could not be reached at the last assessment point by diagnostic assessors after several attempts were censored at week 26 (mid-point assessment). Thus, we do not know whether the decrease reflects a loss in efficacy, because we do not know whether participants who were not reached experienced a major depressive episode or stayed healthy.

Another question referred to intensity and type of guidance provided by the online trainer, such as whether they intervened when suicidal thoughts were mentioned, and whether this might have influenced the results. The purpose of the guidance was not to deliver individual psychotherapy but to support participants in adhering to the treatment modules, which is typical for guided self-help interventions.² The total time a trainer spent per participant was 2 to 3 hours.³ The online trainer provided semistandardized feedback on each of the completed modules using a standardized, preformulated text that was individually adapted based on what the participant wrote in the exercises.⁴ Moreover, the online trainer regularly checked whether participants completed the intervention sessions on time and, if not, reminded them to do so. Trainers were advised not to deliver individualized psychotherapeutic strategies (such as techniques on how to handle suicidal thoughts), and this was assessed through fidelity checks by a supervisor. Hence, although we assume that the use of online trainers increased adherence and therefore the effectiveness of the intervention,⁵ it is unlikely that effects were mediated by individually delivered psychological techniques.

Angheliescu also points out other potential benefits of web-based preventive approaches. Based on the results of this study and other emerging empirical evidence on preventive interventions for depression, implementing and disseminating such

low-threshold interventions on a large scale may eventually contribute to a reduction in the prevalence and burden of major depressive disorder.

David Daniel Ebert, PhD
Claudia Buntrock, MSc
Pim Cuijpers, PhD

Author Affiliations: Department of Clinical Psychology and Psychotherapy, Friedrich-Alexander-University Erlangen-Nuremberg, Erlangen, Germany (Ebert); Division of Online Health Training, Leuphana University, Lueneburg, Germany (Buntrock); Department of Clinical, Neuro, and Developmental Psychology, Vrije Universiteit, Amsterdam, the Netherlands (Cuijpers).

Corresponding Author: David Daniel Ebert, PhD, Department of Clinical Psychology and Psychotherapy, Friedrich-Alexander University Erlangen-Nuremberg, 90453 Erlangen, Germany (david.ebert@fau.de).

Conflict of Interest Disclosures: The authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Dr Ebert reported holding shares of the Institute for Online Health Training, which licenses the intervention studied from Leuphana University. No other disclosures were reported.

1. van Zoonen K, Buntrock C, Ebert DD, et al. Preventing the onset of major depressive disorder: a meta-analytic review of psychological interventions. *Int J Epidemiol*. 2014;43(2):318-329.
2. Cuijpers P, Donker T, van Straten A, Li J, Andersson G. Is guided self-help as effective as face-to-face psychotherapy for depression and anxiety disorders? a systematic review and meta-analysis of comparative outcome studies. *Psychol Med*. 2010;40(12):1943-1957.
3. Buntrock C, Ebert D, Lehr D, et al. Effectiveness of a web-based cognitive behavioural intervention for subthreshold depression: pragmatic randomised controlled trial. *Psychother Psychosom*. 2015;84(6):348-358.
4. Buntrock C, Ebert DD, Lehr D, et al. Evaluating the efficacy and cost-effectiveness of web-based indicated prevention of major depression: design of a randomised controlled trial. *BMC Psychiatry*. 2014;14:25.
5. Baumeister H, Reichler L, Munzinger M, Lin J. The impact of guidance on internet-based mental health interventions—a systematic review. *Internet Interventions*. 2014;1(4):205-215.

Association of Infection in Early Life and Risk of Developing Type 1 Diabetes

To the Editor A research letter by Dr Beyerlein and colleagues focused on infections in early life and the risk of developing type 1 diabetes (T1D) in a large pediatric cohort.¹ We would like to address some relevant points about this study.

The authors did not assess some early determinants, such as gestational age, feeding practice (ie, breastfeeding), and perinatal use of antibiotics. These factors influence both gut development and function,² the risk of infections, and the consequent risk of T1D.

Also, choosing juvenile idiopathic arthritis (JIA) as a control autoimmune disease is questionable. Some proportion of children diagnosed with JIA in fact have a nonautoimmune systemic form, Still disease.³

Celiac disease, rather than JIA, would have been a better control condition because celiac disease shares with T1D a strong genetic background (ie, *HLA-DQ2*), as well as a common association with increased intestinal permeability.⁴

The gut microbiome is important in developmental programming and balanced helper T cell subtype 1 and subtype 2 (T_H1 and T_H2) immune homeostasis,⁵ which is disrupted during autoimmune conditions such as T1D. Modulating the bacterial interface is crucial during the first 6 months of life. Thus

potential influences on the microbiota should be considered in evaluating the risk to develop T1D.

Hakim Rahmoune, MD
Nada Boutrid, MD
Belkacem Bioud, MD

Author Affiliations: Pediatrics Department, Setif University Hospital, Setif, Algeria.

Corresponding Author: Hakim Rahmoune, MD, Pediatrics Department, Setif University Hospital, Setif-1 University, ALN Avenue, 19000 Setif, Algeria (rahmounehakim@gmail.com).

Conflict of Interest Disclosures: The authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

1. Beyerlein A, Donnachie E, Jergens S, Ziegler AG. Infections in early life and development of type 1 diabetes. *JAMA*. 2016;315(17):1899-1901.
2. Houghteling PD, Walker WA. Why is initial bacterial colonization of the intestine important to infants' and children's health? *J Pediatr Gastroenterol Nutr*. 2015;60(3):294-307.
3. Thierry S, Fautrel B, Lemelle I, Guillemin F. Prevalence and incidence of juvenile idiopathic arthritis: a systematic review. *Joint Bone Spine*. 2014;81(2):112-117.
4. Visser J, Rozing J, Sapone A, Lammers K, Fasano A. Tight junctions, intestinal permeability, and autoimmunity: celiac disease and type 1 diabetes paradigms. *Ann N Y Acad Sci*. 2009;1165(1):195-205.
5. Walker WA, Iyengar RS. Breast milk, microbiota, and intestinal immune homeostasis. *Pediatr Res*. 2015;77(1-2):220-228.

To the Editor The study by Dr Beyerlein and colleagues¹ is subject to an important confounder that challenges the validity of the association between infections and T1D. The prior administration of antibiotics has been strongly linked with the development of T1D both in empirical animal studies and in epidemiological studies.^{2,3} The gut microbiota is thought to be critical to the development of the immune system. Eradication or depletion of the gut fauna adversely affects this process, increasing the risk of autoimmunity possibly by inducing anergy in pathways involved in self-tolerance.^{2,3} Empirical animal models suggest that antibiotics in childhood, rather than infections, increase the risk of T1D.^{2,3} The antibiotic model also would explain the precipitous rise in incidence of T1D over the past decades, at a mean annual rate of 3% as reported by the World Health Organization.⁴ The misuse of antibiotics is an increasing global problem, with antibiotics administered inappropriately in viral infections.⁵ A dose-response correlation is observed, whereby the greater the number of courses of antibiotics the greater the risk of T1D. Empirical and epidemiological studies both point to antimicrobials against gram-positive organisms as those most strongly associated with T1D.² A definitive interpretation of the study by Beyerlein and colleagues is difficult without adjusting for this confounder.

Ossie Ferdinand Uzoigwe, BSc

Author Affiliation: University of Sheffield, Sheffield, United Kingdom.

Corresponding Author: Ossie Ferdinand Uzoigwe, BSc, University of Sheffield, Western Bank, Sheffield, South Yorkshire S10 2TN, United Kingdom (ofuzoigwe@yahoo.co.uk).

Conflict of Interest Disclosures: The author has completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

1. Beyerlein A, Donnachie E, Jergens S, Ziegler AG. Infections in early life and development of type 1 diabetes. *JAMA*. 2016;315(17):1899-1901.