VU Research Portal

Constraint-induced movement therapy

van der Lee, J.H.; Beckerman, H.; Lankhorst, G.J.; Bouter, L.M.

published in

Archives of Physical Medicine and Rehabilitation
1999

DOI (link to publisher)

[10.1016/S0003-9993\(99\)90339-8](https://doi.org/10.1016/S0003-9993(99)90339-8)
[10.1016/S0003-9993\(99\)90339-8](https://doi.org/10.1016/S0003-9993(99)90339-8)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

van der Lee, J. H., Beckerman, H., Lankhorst, G. J., & Bouter, L. M. (1999). Constraint-induced movement therapy. *Archives of Physical Medicine and Rehabilitation*, 80(12), 1606-1606. [https://doi.org/10.1016/S0003-9993\(99\)90339-8](https://doi.org/10.1016/S0003-9993(99)90339-8), [https://doi.org/10.1016/S0003-9993\(99\)90339-8](https://doi.org/10.1016/S0003-9993(99)90339-8)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

DEPARTMENTS

*Letters to the Editor***Constraint-Induced Movement Therapy**

Kunkel and colleagues¹ recently published an article on constraint-induced (CI) movement therapy for motor recovery in chronic stroke patients. They presented the results of an uncontrolled series of 5 stroke patients undergoing CI therapy. The authors also presented a review of five studies, including their own, concerning the effectiveness of CI therapy based on a calculation of effect sizes. We have several concerns with this paper.

In our opinion, research in physical medicine and rehabilitation has progressed to a point at which patient series no longer make any substantial contribution to the accumulation of knowledge. Controlled studies, preferably randomized, are needed to determine the effectiveness of treatment modalities. Uncontrolled studies can be misleading and will almost always report grossly overestimated treatment effects.² The importance of a control group, when evaluating CI therapy, is emphasized by the interim conclusions of a controlled study carried out by Wittenberg and colleagues,³ who found that patients who received standard therapy also showed improvement, possibly due to expectation bias. The results of a randomized clinical trial conducted at our hospital, which will be published shortly, confirm this.⁴

In their literature review, Kunkel and colleagues claim to have identified two controlled studies. However, the publication on one of these studies mentions a control group of 15 patients, but provides no data concerning improvement in either the experimental or the control group.⁵ Therefore, we were puzzled by the fact that Kunkel presented data on the experimental group of 20 patients in this study, but no data on the control group.

Finally, we strongly disagree with the interpretation of effect sizes based on trends over time in uncontrolled series as important evidence in favor of CI movement therapy. In fact, when using within-group time trends to calculate effect sizes, all the methodologic advantages of having a control group are ignored. We conclude that Kunkel failed to demonstrate that CI therapy is efficacious for chronic stroke patients.

**Johanna H. van der Lee, MD
Heleen Beckerman, PhD, PT
Gustaaf J. Lankhorst, MD, PhD
University Hospital Vrije Universiteit
Lex M. Bouter, PhD
Institute for Research in Extramural Medicine
Vrije Universiteit
Amsterdam, The Netherlands**

References

1. Kunkel A, Kopp B, Muller G, Villringer K, Villringer A, Taub E, et al. Constraint induced movement therapy for motor recovery in chronic stroke patients. *Arch Phys Med Rehabil* 1999;80:624-8.
2. Sackett DL, Haynes RB, Guyatt GH, Tugwell P. *Clinical epidemiology. A basic science for clinical medicine*. Boston/Toronto/London: Little, Brown and Company; 1995.
3. Wittenberg G, Chen R, Ishii K, Croarkin E, Eckloff S, Gerber N, et al. Effect of constraint-induced movement therapy on motor function and cortical physiology in chronic stroke. Presented at the 2nd World Congress on Neurological Rehabilitation, Toronto, April 1999.

4. Van der Lee JH, Wagenaar RC, Lankhorst GJ, Vogelaar TW, Devillé WL, Bowler LM. Forced use of the upper extremity in chronic stroke patients: results from a single-blind randomized clinical trial. *Stroke* (Forthcoming).
5. Taub E, Crago JE, Uswatte G. Constraint-induced movement therapy: a new approach to treatment in physical rehabilitation. *Rehabil Psychol* 1998;43:152-70.

The authors reply

Van der Lee and colleagues are correct in emphasizing that randomized controlled studies are preferable to uncontrolled studies; however, replications of clinical series still have a place in rehabilitation research. As noted in our report, we sought to perform an independent replication of the effects of constraint-induced movement therapy, which had only been fully tested in coauthor Taub's laboratory.^{1,2} A similar replication without a control group appeared recently in *Stroke*.³ Since CI treatment is a new and promising therapy, we believe that these independent replications are still viable, as was also suggested by Ottenbacher.⁴ The paper by Wittenbacher and associates⁵ is a preliminary report with still-unpublished data and no conclusions can be drawn from it. The meta-analysis was limited to pre-post effect sizes, since only two of the studies we reported contained a control group. Control group effect sizes could thus not have been pooled for the demonstration of the overall effect we wanted to report. Finally, we want to point out that we tested the effects of CI therapy in a sample of highly chronic stroke patients where spontaneous recovery has never been demonstrated and we want to emphasize that the effects reported here were substantial and long-lasting.

We believe that these independent replications in two laboratories have established the efficacy of CI therapy for the treatment of chronic stroke without claiming the specificity of the observed effect. We agree with van der Lee that now would be the time to compare these effects to those of other treatment approaches to evaluate to what extent the effects we observed are actually related to specific behavioral training rather than reflecting nonspecific therapy effects.

**Herta Flor, PhD
Bruno Kopp, PhD
Annett Kunkel
Department of Psychology,
Humboldt-University
D-10117 Berlin, Germany**

References

1. Taub E, Miller NE, Novack TA, Cook EW III, Fleming WC, Nepomuceno CS, et al. Technique to improve chronic motor deficit after stroke. *Arch Phys Med Rehabil* 1993;74:347-54.
2. Taub E, Wolf SL. Constraint induced movement techniques to facilitate upper extremity use in stroke patients. *Top Stroke Rehabil* 1997;3:38-61.
3. Miltner WH, Bauder H, Sommer M, Dettmers C, Taub E. Effects of constraint-induced movement therapy on patients with chronic motor deficits after stroke: a replication. *Stroke* 1999;30:586-92.
4. Ottenbacher KJ. Why rehabilitation research does not work (as well as we think it should). *Arch Phys Med Rehabil* 1995;76:123-9.
5. Wittenberg G, Chen R, Ishii K, Croarkin E, Eckloff S, Gerber NL, et al. Effect of constraint-induced movement therapy on motor function and cortical physiology in chronic stroke. Poster presented at