VU Research Portal

The efficacy of the back school for patients with non-specific low back pain

Keijsers, Jolanda F.E.M.; Meertens, Ree M.; Kok, Gerjo J.; Bouter, Lex M.

published in

Physiotherapy Theory and Practice
1992

DOI (link to publisher)

[10.3109/09593989209108085](https://doi.org/10.3109/09593989209108085)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Keijsers, J. F. E. M., Meertens, R. M., Kok, G. J., & Bouter, L. M. (1992). The efficacy of the back school for patients with non-specific low back pain: An overview. *Physiotherapy Theory and Practice*, 8(2), 85-88.
<https://doi.org/10.3109/09593989209108085>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

The efficacy of the back school for patients with non-specific low back pain: An overview

Jolanda F. E. M. Keijsers, Lex M. Bouter, Ree M. Meertens and Gerjo J. Kok

This overview represents the experimental intervention used in the study by Keijsers et al reported on pp. 79-83, this issue.

INTRODUCTION

Low back pain is a frequent complaint. It is likely that up to 80% of all people will experience back pain to some extent during their active lives (Nachemson, 1976). In most cases, the complaints are self-limiting, with recovery within 2 months in 90% of cases (Bergquist-Ullman and Larsson, 1977; Frymoyer, 1988). However, recurrences are frequent (40-60%; Haanen, 1984). The complaints are in most cases non-specific, which means that no underlying pathology has been established. In the Netherlands, 80% of all consultations with general practitioners (GPs) involve such non-specific complaints (Hoekstra, 1985). It is commonly recognised that, in addition to physical factors, psychosocial factors are also involved in both the aetiology and prognosis of low back pain (Meilman, 1984; Nachemson, 1979; Turk and Flor, 1984). These factors are of equal importance in terms of treatment and they necessitate a multi-

dimensional approach to non-specific low back pain.

One of the possible treatments for low back pain is attendance at a so-called back school, which is of Swedish origin (Zachrisson Forssell, 1981). Back schools offer an education and skills programme in a group setting aimed primarily at pain management: information is given about ways of dealing with pain, so that the patient is able to control the pain problem better (Linton and Kamwendo, 1987). Ultimately, this should lead to a decrease in work absenteeism and medical utilisation. The efficacy of back schools is controversial (Fisk, Dimonte and McKay Courington, 1983; Keijsers, Bouter, Steenbakkers and Meertens, 1989a; Linton and Kamwendo, 1987; Terpstra and Bouter, 1988).

In the existing literature, six studies can be found in which patients were randomly assigned to an experimental and a control group (randomised clinical trial). Patients in the control group received no treatment or an alternative treatment. The reason for excluding all other studies into the efficacy of back schools is that the results of such studies can hardly be interpreted and are therefore not useful. In the six available randomised clinical trials, the efficacy of non-clinical back schools for patients with non-specific low back pain was assessed. In two of the six studies, the back school turned out to be ineffective (Berwick, Budman and Feldstein, 1989;

J. F. E. M. Keijsers, Dutch Centre of Health Education and Health Promotion, PO Box 5104, 3502 JC Utrecht, The Netherlands

R. M. Meertens, G. J. Kok, Department of Health Education, **L. M. Bouter**, Department of Epidemiology and Health Care Research, University of Limburg, PO Box 616, 6200 MD Maastricht, The Netherlands
(Reprint requests to JFEMK)

Accepted for publication July 1991

Lankhorst et al, 1983). In one study, the efficacy of the back school is doubtful (Keijsers et al, 1989b). In the remaining three studies, the authors concluded that the back school is indeed effective (Bergquist-Ullman and Larsson, 1977; Hurri, 1989; Klaber Moffett, Chase, Portek and Ennis, 1986). Several problems concerning the validity of these 'positive' studies make such conclusions at least doubtful. The duration of the follow-up, for example, was relatively short (16 weeks) in the study by Klaber Moffett et al (1986). Furthermore, the often large number of effect parameters used in the studies, increases substantially the chances of finding at least one statistically significant effect.

Since many doubts remained concerning the efficacy of back schools, we conducted another randomised clinical trial (Keijsers et al, 1990b). The results of this randomised clinical trial will be presented below.

At the Department of Health Education of the University of Limburg, the so-called Maastricht back school was developed, in cooperation with the University Hospital. The Maastricht back school consists of seven sessions, each lasting 2.5 h, plus a refresher session after 6 months. A course instructor is present at every session. In addition, various guest lecturers are invited to give information and training. At the end of each session, a written summary is handed out. The Maastricht back school is considered to be a combination of all those elements which we felt ought to be included in a back school information and training programme (Keijsers, Steenbakkers, Gerards and Meertens, 1990a). In order to determine the efficacy of the Maastricht back school in primary health care, a randomised clinical trial was conducted. The experimental group that followed the sessions of the back school was compared to a waiting list control group, who were promised entry into the back school programme at the end of the study. All of the participants could continue using traditional care for their complaints.

SELECTION CRITERIA

The inclusion criteria for the study were (1)

complaints to be non-specific, (2) duration of complaints to be between 2 months and 3 years and (3) patient to be able to perform physical and relaxation exercises. Prospective participants were judged on all these criteria by the GPs recruiting the study population.

EFFECT PARAMETERS

The most important measures of effect were (1) pain management, (2) pain, (3) medical utilisation (subdivided into consultations with health care providers, number of types of treatment and medication) and (4) absenteeism from work, expressed in numbers of days. Other effect parameters were functional disability, knowledge of the subjects discussed in the course, general well-being and satisfaction about the Maastricht back school. Data on all of the effect parameters were collected at baseline and 2 and 8 months after randomisation. As much as possible, internationally acknowledged effect parameters, validated for the Dutch language, were chosen.

RESULTS

Univariate and multivariate analyses of the repeated measures data were performed on every effect parameter, in order to determine whether there were differences across time between the experimental and control groups, i.e. whether the experimental group showed more progress. Furthermore, univariate and multivariate analyses of covariance of the repeated measures data were performed on most of the effect parameters. This was done in order to determine the efficacy of the Maastricht back school after adjustment for differences between the experimental and the control groups at baseline.

In Table 1, the means of the experimental and control groups are given for the most important effect parameters (pain management, pain, medical utilisation and absenteeism from work) and for each episode of data collection. Data on medical utilisation, as well as on absenteeism from work, were collected only at baseline and 8 months after randomisation. None of the dif-

Table 1
Means of the experimental and control groups for the most important effect parameters and for each data collection episode*

	Experimental group (n=37)			Control group (n=40)		
	Baseline	2 months	8 months	Baseline	2 months	8 months
Pain management (range 1–10)	5.2	4.1	4.2	4.6	4.4	3.8
Pain (range 1–10)	6.5	5.4	5.4	5.8	5.3	4.1
Medical utilisation						
consultations (range 0–∞)	23.3		9.0	16.1		6.6
treatments (range 0–22)	3.3		1.8	2.6		1.0
medication (range 0–7)	1.2		0.8	0.6		0.3
Absenteeism from work (range 0–∞)	27.0		19.0	21.3		3.4

*For each effect parameter, a higher figure corresponds to a less desirable situation.

ferences between the experimental and control groups turned out to be statistically significant.

Summarising the results, it can be concluded that:

1. Both the experimental and the waiting list control groups showed progress with time. However, the experimental group did not show significantly more progress than the control group.
2. Although the patients were randomly assigned to the treatment groups, there were substantial differences between the groups at baseline.
3. After adjustment for these differences, there was still no difference in effect between the group receiving the Maastricht back school treatment and the waiting list control group.

CONCLUSION

In addition to the apparent inefficacy of the Maastricht back school, some problems concerning the validity of our study need also to be discussed. For example, the differences at baseline may be due to the data collected after randomisation rather than before. Filling out a questionnaire when knowing whether or not one can enter a back school programme might bias one's answers. Moreover, the number of patients in the study was relatively small ($n=77$). As a result, only major differences between the experi-

mental and control groups could reach statistical significance. Nevertheless, it can be concluded that it is unlikely that the Maastricht back school is an effective treatment for patients with non-specific low back pain. The limited evidence available from other randomised clinical trials indicates at most borderline effects for other back schools. Very large trials with perfect methodology might yet show some beneficial effects. The question is, however, whether such trials deserve high priority.

References

- Bergquist-Ullman M, Larsson U 1977 Acute low back pain in industry. *Acta Orthopaedica Scandinavica* 170: 1–117 (suppl)
- Berwick DM, Budman S, Feldstein M 1989 No clinical effect of back schools in an HMO: A randomized prospective trial. *Spine* 3: 338–344
- Fisk JR, Dimonte P, McKay Courington S 1983 Back schools, past, present and future. *Clinical Orthopaedics and Related Research* 179: 18–23
- Frymoyer JW 1988 Back pain and sciatica. *New England Journal of Medicine* 318: 291–300
- Haanen HCM 1984 Een epidemiologisch onderzoek naar lage rugpijn. Dissertation, Erasmus Universiteit Rotterdam
- Hoekstra GR 1985 De rugpijnpatiënt in de huisartspraktijk. In: Kingma MJ, Dokter HJ (eds) *Rugpijn*, pp 40–66. Bohn, Scheltema and Holkema, Utrecht
- Hurri H 1989 The Swedish back school in chronic low back pain: Part I. Benefits. *Scandinavian Journal of Rehabilitation Medicine* 21: 33–40
- Keijsers JFEM, Bouter LM, Steenbakkers WHL, Meertens RM 1989a Methodologische kwaliteit en onderlinge vergelijkbaarheid van onderzoek naar de effectiviteit van rugscholen. *Nederlands Tijdschrift voor Fysiotherapie* 5: 112–116

- Keijsers JFEM, Groenman NH, Gerards FM, van Oudheusden E, Steenbakkens WHL 1989b A back school in the Netherlands: Evaluating the results. *Patient Education and Counseling* 14: 31-44
- Keijsers JFEM, Steenbakkens WHL, Gerards FM, Meertens RM (1990a) The efficacy of the back school: An analysis of the literature. *Arthritis Care and Research* 3: 210-215
- Keijsers JFEM, Steenbakkens WHL, Meertens RM, Bouter LM, Kok GJ (1990b) The efficacy of the back school: A randomized trial. *Arthritis Care and Research* 3: 204-209
- Klaber Moffett JA, Chase SM, Portek I, Ennis JR 1986 A controlled, prospective study to evaluate the effectiveness of a back school in the relief of chronic low back pain. *Spine* 11: 120-122
- Lankhorst GJ, Van de Stadt RJ, Vogelaar TW, Van der Korst LK, Prevo AJH 1983 The effect of the Swedish back school in chronic idiopathic low back pain. *Scandinavian Journal of Rehabilitation Medicine* 15: 141-145
- Linton SJ, Kamwendo K 1987 Low back schools: A critical review. *Physical Therapy* 67: 1375-1383
- Meilman PW 1984 The management of the chronic pain patient: Clinical considerations. *Journal of Orthopaedic Sport Physical Therapy* 5: 305-317
- Nachemson AL 1976 The lumbar spine: An orthopaedic challenge. *Spine* 1: 59-71
- Nachemson A 1979 A critical look at the treatment for low back pain. *Scandinavian Journal of Rehabilitation Medicine* 11: 143-147
- Terpstra SJ, Bouter LM 1988 Het effect van de Rugschool, overzicht van de literatuur. *Nederlands Tijdschrift voor Fysiotherapie* 3: 60-63
- Turk DC, Flor H 1984 Etiological theories and treatment for chronic back pain. II. Psychological models and interventions. *Pain* 19: 209-233
- Zachrisson Forssell M 1981 The Swedish back school. *Physiotherapy* 66: 112-114