VU Research Portal

Prevention of depressive disorders: Towards a further reduction of the disease burden of mental disorders

Cuijpers, P.

published in

Early Intervention in Psychiatry
2011

DOI (link to publisher)

[10.1111/j.1751-7893.2011.00282.x](https://doi.org/10.1111/j.1751-7893.2011.00282.x)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Cuijpers, P. (2011). Prevention of depressive disorders: Towards a further reduction of the disease burden of mental disorders. *Early Intervention in Psychiatry*, 5(3), 179-180. <https://doi.org/10.1111/j.1751-7893.2011.00282.x>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Editorial

Prevention of depressive disorders: towards a further reduction of the disease burden of mental disorders

From a public health perspective, depression is one of the most important mental disorders. It is highly prevalent,^{1,2} has a high incidence³ and is associated with a substantial loss of quality of life for patients and their relatives,^{4,5} increased mortality rates,⁶ high levels of service use and enormous economic costs.^{7–9} Major depression is currently ranked fourth worldwide in disease burden, and it is expected to rank first in disease burden in high-income countries by the year 2030.¹⁰ Although current treatments are effective in treating depressive disorders, it is estimated that treatment alone can reduce the disease burden of depression by only about 35%, but only in optimal conditions.¹¹ Prevention of the incidence of new cases of depression has been suggested as a new strategy next to treatment, which may help to further lower the disease burden.^{12,13}

In the past 15 years, considerable progress has been made in examining the possibilities to prevent the onset of depressive disorders. In a recent meta-analysis, we were able to include 19 randomized trials examining interventions aimed at preventing the onset of depressive disorders in participants with no disorder at baseline,¹⁴ and currently almost 30 trials have been conducted.¹⁵ These studies show that preventive interventions are capable of preventing about one-quarter of the new incident cases.

These studies vary considerably from each other, with some studies being aimed at adolescents, whereas others focus on adults and some on older adults. The setting in which the studies are conducted also differs considerably, ranging from studies aimed at post-partum depression to studies aimed at patients with general medical disorders and primary care patients. Some studies are aimed at universal prevention, which is aimed at the general population or parts of the general population, regardless of whether they are at elevated risk of developing a disorder (e.g. school programmes or mass media campaigns). Other studies focus on selective prevention, which is aimed at high-risk groups, who have not yet developed a mental disorder. Still other studies focus on indicated prevention

(aimed at individuals who have some symptoms of a mental disorder but do not meet diagnostic criteria). The interventions used in these studies also vary considerably and include cognitive behavioural interventions, interventions based on interpersonal psychotherapy and psychoeducational interventions. Some studies explicitly focus on the first ever depressive disorder, whereas others allow lifetime depressive disorders.

So, the field is still very heterogeneous in terms of target groups, setting and type of intervention. In our meta-analysis, we did not find high levels of statistical heterogeneity, which may indicate that our finding of about 25% reduction of incidence is robust across target groups, settings and interventions. It also indicates, however, that the field is still very young, and definite conclusions about the effects of these interventions cannot yet be drawn.

Prevention of mental disorders in general is still at an early stage of development. Although a relatively large number of studies have examined the possibilities of preventing depressive disorders, we still have very little knowledge about effective interventions and how to apply them. Although several studies have examined the possibilities to prevent anxiety disorders,^{16–18} eating disorders¹⁹ and psychotic disorders,²⁰ this is nothing compared with the thousands of studies that have examined treatment of existing disorders. Although treatment studies have examined the effects, acceptability and feasibility of many different types of treatment in many different settings, the possibilities of preventing the onset of mental disorders have only just started to be explored in a few dozen studies.

Research on the possibilities of preventing mental disorders has been developed within subfields, largely independent of other subfields and with little exchange. Prevention of depressive disorders, anxiety disorders, psychotic disorders, eating disorders, conduct disorders in children and all fields has been developed from research on treatments of these disorders and epidemiological research examining each of these disorders. This has also led to differences among subfields. For example, in the

field of prevention of depression, many hope that universal prevention through school programmes will prove to be possible and feasible,²¹ whereas in the field of preventing eating disorders, universal prevention is almost impossible,¹³ and universal prevention of psychotic disorders is not considered to be possible either at this moment by most researchers. In the field of prevention of psychotic disorders, indicated prevention and early interventions are the best options, with few possibilities for selective prevention, whereas selective prevention of depression is probably effective in well-chosen target groups.¹⁴ Prevention and early intervention of psychotic disorders are aimed at preventing or delaying the first psychotic episode in young people, whereas prevention of depression is also aimed at older adults and recurrent episodes.

Despite these differences, it is time that these subfields start working together and learn from each other's experiences. In the prodromal and early phases of mental disorders, it may not yet be completely clear which disorder will develop, and generic preventive strategies using transdiagnostic approaches may be able to prevent different types of disorders. Clinical staging provides a good framework for the further development of these preventive strategies.²²

There is no doubt that prevention of mental disorders is one of the major challenges for the next decades. Prevention of depression has been shown to be effective and feasible in demonstration projects. Collaboration with other fields and disseminating this knowledge to practice will be the next steps in further reducing the disease burden of mental disorders.

Pim Cuijpers

*Department of Clinical Psychology and
EMGO Institute for Health and Care Research,
VU University and VU University Medical Center,
Amsterdam, The Netherlands*

REFERENCES

1. Kessler RC, McGonagle KA, Zhao S *et al.* Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry* 1994; **51**: 8–19.
2. Alonso J, Angermeyer MC, Bernert S *et al.* European Study of the Epidemiology of Mental Disorders (ESEMeD) Project: prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) Project. *Acta Psychiatr Scand Suppl* 2004; **420**: 21–7.
3. Waraich P, Goldner EM, Somers JM, Hsu L. Prevalence and incidence studies of mood disorders: a systematic review of the literature. *Can J Psychiatry* 2004; **49**: 124–38.
4. Ustun TB, Ayuso-Mateos JL, Chatterji S, Mathers C, Murray CIL. Global burden of depressive disorders in the year 2000. *Br J Psychiatry* 2004; **184**: 386–92.
5. Saarni SI, Suvisaari J, Sintonen H *et al.* Impact of psychiatric disorders on health-related quality of life: general population survey. *Br J Psychiatry* 2007; **190**: 326–32.
6. Cuijpers P, Smit F. Excess mortality in depression: a meta-analysis of community studies. *J Affect Disord* 2002; **72**: 227–36.
7. Berto P, D'Ilario D, Ruffo P, Di Virgilio R, Rizzo F. Depression: cost-of-illness studies in the international literature, a review. *J Ment Health Policy Econ* 2000; **3**: 3–10.
8. Greenberg PE, Birnbaum HG. The economic burden of depression in the US: societal and patient perspectives. *Exp Opin Pharmacother* 2005; **6**: 369–76.
9. Smit F, Cuijpers P, Oostenbrink J, Batelaan N, de Graaf R, Beekman A. Excess costs of common mental disorders: population based cohort study. *J Ment Health Policy Econ* 2006; **9**: 193–200.
10. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006; **3**: e442.
11. Andrews G, Issakidis C, Sanderson K, Corry J, Lapsley H. Utilising survey data to inform public policy: comparison of the cost-effectiveness of treatment of ten mental disorders. *Br J Psychiatry* 2004; **184**: 526–33.
12. Smit F, Ederveen A, Cuijpers P, Deeg D, Beekman A. Opportunities for cost-effective prevention of late-life depression: an epidemiological approach. *Arch Gen Psychiatry* 2006; **63**: 290–6.
13. Cuijpers P. Examining the effects of prevention programs on the incidence of new cases of mental disorders: the lack of statistical power. *Am J Psychiatry* 2003; **160**: 1385–91.
14. Cuijpers P, van Straten A, Smit F, Mihalopoulos C, Beekman A. Preventing the onset of depressive disorders: a meta-analytic review of psychological interventions. *Am J Psychiatry* 2008; **165**: 1272–80.
15. Muñoz RF, Cuijpers P, Smit F, Barrera AZ, Leykin Y. Prevention of major depression. *Annu Rev Clin Psychol* 2010; **6**: 181–212.
16. Gardenschwartz CA, Craske MG. Prevention of panic disorder. *Behav Ther* 2001; **32**: 725–37.
17. Higgins DM, Hecker JE. A randomized trial of brief cognitive-behavioral therapy for prevention of generalized anxiety disorder. *J Clin Psychiatry* 2008; **69**: 1337.
18. Schmidt NB, Eggleston AM, Woolaway-Bickel K, Fitzpatrick KK, Vasey MW, Richey JA. Anxiety Sensitivity Amelioration Training (ASAT): a longitudinal primary prevention program targeting cognitive vulnerability. *J Anxiety Disord* 2007; **21**: 302–19.
19. Stice E, Shaw H, Marti CN. A meta-analytic review of eating disorder prevention programs: encouraging findings. *Annu Rev Clin Psychol* 2007; **3**: 207–31.
20. Harvey PO, Lepage M, Malla A. Benefits of enriched intervention compared with standard care for patients with recent-onset psychosis: a metaanalytic approach. *Can J Psychiatry* 2007; **52**: 464–72.
21. Spence S, Sheffield J, Donovan C. Preventing adolescent depression: an evaluation of the problem solving for life program. *J Consult Clin Psychol* 2003; **71**: 3–13.
22. McGorry PD, Nelson B, Goldstone S, Yung AR. Clinical staging: a heuristic and practical strategy for new research and better health and social outcomes for psychotic and related mood disorders. *Can J Psychiatry* 2010; **55**: 486–97.