VU Research Portal

Establishing Guidelines for Executing and Reporting Internet Intervention Research

Proudfoot, J.; Klein, B.; Barak, A.; Carlbring, P.; Cuijpers, P.; Lange, A.; Ritterband, L.; Andersson, G.

published in

Cognitive Behaviour Therapy
2011

DOI (link to publisher)

[10.1080/16506073.2011.573807](https://doi.org/10.1080/16506073.2011.573807)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Proudfoot, J., Klein, B., Barak, A., Carlbring, P., Cuijpers, P., Lange, A., Ritterband, L., & Andersson, G. (2011). Establishing Guidelines for Executing and Reporting Internet Intervention Research. *Cognitive Behaviour Therapy*, 40, 82-97. <https://doi.org/10.1080/16506073.2011.573807>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Cognitive Behaviour Therapy

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/sbeh20>

Establishing Guidelines for Executing and Reporting Internet Intervention Research

Judith Proudfoot ^a, Britt Klein ^b, Azy Barak ^c, Per Carlbring ^d, Pim Cuijpers ^e, Alfred Lange ^f, Lee Ritterband ^g & Gerhard Andersson ^h

^a Black Dog Institute and School of Psychiatry, University of New South Wales, Sydney, New South Wales, Australia

^b National eTherapy Centre and the Brain and Psychological Sciences Research Centre (BPsyC) eTherapy Unit, Faculty of Life and Social Sciences, Swinburne University, Melbourne, Victoria, Australia

^c Department of Counseling and Human Development, University of Haifa, Haifa, Israel

^d Department of Psychology, Umeå University, Umeå, Sweden

^e Department of Clinical Psychology, Faculty of Psychology and Education, VU University, Amsterdam, The Netherlands

^f Department of Clinical Psychology, University of Amsterdam, Amsterdam, The Netherlands

^g Behavioral Health and Technology, Department of Psychiatry and Neurobehavioral Sciences, University of Virginia, Charlottesville, Virginia, USA

^h Department of Behavioural Sciences and Learning, Linköping University, Linköping, Sweden

Available online: 24 Jun 2011

To cite this article: Judith Proudfoot, Britt Klein, Azy Barak, Per Carlbring, Pim Cuijpers, Alfred Lange, Lee Ritterband & Gerhard Andersson (2011): Establishing Guidelines for Executing and Reporting Internet Intervention Research, *Cognitive Behaviour Therapy*, 40:2, 82-97

To link to this article: <http://dx.doi.org/10.1080/16506073.2011.573807>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan, sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Establishing Guidelines for Executing and Reporting Internet Intervention Research

Judith Proudfoot¹, Britt Klein², Azy Barak³, Per Carlbring⁴, Pim Cuijpers⁵, Alfred Lange⁶, Lee Ritterband⁷ and Gerhard Andersson⁸

¹*Black Dog Institute and School of Psychiatry, University of New South Wales, Sydney, New South Wales, Australia;* ²*National eTherapy Centre and the Brain and Psychological Sciences Research Centre (BPsyC) eTherapy Unit, Faculty of Life and Social Sciences, Swinburne University, Melbourne, Victoria, Australia;* ³*Department of Counseling and Human Development, University of Haifa, Haifa, Israel;* ⁴*Department of Psychology, Umeå University, Umeå, Sweden;* ⁵*Department of Clinical Psychology, Faculty of Psychology and Education, VU University, Amsterdam, The Netherlands;* ⁶*Department of Clinical Psychology, University of Amsterdam, Amsterdam, The Netherlands;* ⁷*Behavioral Health and Technology, Department of Psychiatry and Neurobehavioral Sciences, University of Virginia, Charlottesville, Virginia, USA;* ⁸*Department of Behavioural Sciences and Learning, Linköping University, Linköping, Sweden*

Abstract. The field of Internet interventions is growing rapidly. New programs are continually being developed to facilitate health and mental health promotion, disease and emotional distress prevention, risk factor management, treatment, and relapse prevention. However, a clear definition of Internet interventions, guidelines for research, and evidence of effectiveness have been slower to follow. This article focuses on the quality standardization of research on Internet-delivered psychological and behavioural interventions. Although the science underpinning Internet interventions is just starting to be established, across research studies there are often conceptual and methodological difficulties. The authors argue that this situation is due to the lack of universally accepted operational guidelines and evaluation methods. Following a critical appraisal of existing codes of conduct and guidelines for Internet-assisted psychological and health interventions, the authors developed a framework of guidelines for Internet intervention research utilizing aspects of facet theory (Guttman & Greenbaum, 1998). The framework of facets, elements, and guidelines of best practice in reporting Internet intervention research was then sent to several leading researchers in the field for their comment and input, so that a consensus framework could be agreed on. The authors outline 12 key facets to be considered when evaluating and reporting Internet intervention studies. Each facet consists of a range of recommended elements, designed as the minimum features for reporting Internet intervention studies. The authors propose that this framework be utilized when designing and reporting Internet intervention research, so results across studies can be replicated, extended, compared, and contrasted with greater ease and clarity. *Key words:* Internet interventions; research; evaluation; guidelines; reporting criteria.

Received 13 September, 2010; Accepted 17 March, 2011

Correspondence address: Judith Proudfoot, PhD, Black Dog Institute and School of Psychiatry, University of New South Wales, Hospital Road, Randwick, New South Wales 2031, Australia. Tel: + 612 9382 3767. Fax: + 612 9382 8207. E-mail: j.proudfoot@unsw.edu.au

The health and mental health fields have lagged behind other fields in the use of interactive communication technologies, but e-health in

general and Internet-assisted psychological and medical services in particular have been growing rapidly in recent years. The Internet

offers a variety of multimedia interactivity and connectivity formats, with tailoring to specific needs and interests of individual users. Online applications are ideal for use within the helping professions because they allow for the dissemination of standardized yet personalised treatments; they can be used 24 hours per day, 7 days a week, without affecting efficiency; they avoid the need for waiting lists; they allow privacy and consistency of care; and symptom monitoring and outcome measures can be built in relatively easily (Andersson, 2009; Andrews, Cuijpers, Craske, McEvoy, & Titov, 2010; Cuipers et al., 2009; National Institute of Clinical Excellence, 2004). Consequently, new Internet-based interventions for psychological and behavioural health problems are appearing at an increasing rate. These interventions include interactive therapy programs that provide highly specialized treatment and feedback tailored to the characteristics of the user, delivered with and without guidance from a human therapist; "expert systems" in which assessment and therapeutic techniques based on decision rules and behaviour change strategies are integrated; and education programs, digitized information, and online therapeutic communication services such as online counseling, online discussion and support groups, and "ask an expert" websites (Barak, Klein, & Proudfoot, 2009).

For users, the advantages of online psychological interventions include the anonymity and invisibility that some programs afford (with associated reduction in attributed stigma); the accessibility to treatments for those restricted by geographical, transport, personal disability, or financial barriers; the availability of treatment or support at any time; consumer empowerment regarding managing one's own health care; and the flexibility of use in terms of self-determined pace and opportunity to review the material as often as desired. These advantages have created growing interest in the use of online interventions (Andersson, 2009; Andrews et al., 2010; Mohr et al., 2010). Until recently, the evidence base for the usability, clinical efficacy, and cost-effectiveness of online interventions was limited; hence, comparison of web-based behavioural and mental health programs was difficult to pursue. However, more recently, there has been an increase in the number and nature of methodologically sound studies examining such interventions. Most of these studies have

demonstrated positive outcomes for a variety of psychological and behavioural health issues, such as depressive symptoms/depression (Andersson et al., 2005; Christensen, Griffiths, & Jorm, 2004; Clarke et al., 2005; Robertson, Smith, Castle, & Tannenbaum, 2006; Ruwaard et al., 2009; Spek et al., 2007; Titov et al., 2010; Vernmark et al., 2010), panic disorder (Carlbring, Westling, Ljungstrand, Ekselius & Andersson, 2001; Klein & Richards, 2001; Klein, Austin, et al., 2009; Klein, Richards, & Austin, 2006; Pier et al., 2008; Ruwaard, Broeksteeg, Schrieken, Emmelkamp, & Lange, 2010; Schneider, Mataix-Cols, Marks, & Bachofen, 2003), social anxiety disorder (Andersson et al., 2006; Carlbring, Gunnarsdóttir, et al., 2007; Titov, Andrews, Choi, Schwenke, & Mahoney, 2008; Titov, Andrews, & Schwenke, 2008), posttraumatic stress symptoms/disorder (Klein, Mitchell, et al., 2009; Klein et al., 2010; Lange & Ruwaard, 2010; Lange et al., 2003), anxiety prevention (Bennett, Reynolds, Christensen, & Griffiths, 2010; Christensen, Griffiths, et al., 2010; Kenardy, McCafferty, & Rosa, 2003), work-related stress (Ruwaard, Lange, Bouwman, Broeksteeg, & Schrieken, 2007), eating disorders/weight problems (Ljótsson, Mitsell, Lundin, Carlbring, & Ghaderi, 2007; Tate, Jackvony, & Wing, 2006; Winzelberg et al., 2000), encopresis (Ritterband et al., 2003), smoking (Swartz, Noell, Schroeder, & Ary, 2006), insomnia (Ritterband, Thorndike, Gonder-Frederick, et al., 2009; Ström, Pettersson, & Andersson, 2004), physical health conditions (Kuhl, Sears, & Conti, 2006; Ljótsson et al., 2011; Stinson, Wilson, Gill, Yamada, & Holt, 2009), well-being (Mitchell, Stanimirovic, Klein, & Vella-Brodick, 2009), and resilience (Abbott, Klein, Hamilton, & Rosenthal, 2009) to name a few. A number of reviews have also been carried out, with results demonstrating feasibility and efficacy (e.g. Andersson & Cuijpers, 2009; Andersson, Ljótsson, & Weise, 2011; Cavanagh & Shapiro, 2004; Griffiths, Farrer, & Christensen, 2010; Macea, Gajos, Calil, & Fregni, 2010; Myung, McDonnell, Kazinets, Seo, & Moskowitz, 2009; Neve, Morgan, Jones, & Collins, 2010; Reger & Gahm, 2009; Rooke, Thorsteinsson, Karpin, Copeland, & Allsop, 2010; Spek et al., 2007; Tait & Christensen, 2010; van't Hof, Cuijpers, & Stein, 2009; White et al., 2010), as well as a comprehensive meta-analysis across 92 published studies (Barak, Hen, Boniel-Nissim,

& Shapira, 2008), which also showed highly positive results.

Despite the recent growth in research, much heterogeneity remains between the studies, limiting the comparisons able to be drawn and thus confounding interpretation. In addition, the literature is still in development in terms of the science underpinning the programs (Barak et al., 2009; Ritterband & Tate, 2009), and across research studies there are methodological difficulties such as a lack of common terminology and definitions, selection biases, inappropriate research designs, study attrition, and questionable conclusions drawn from the findings (Danaher & Seeley, 2009). We argue that these difficulties are, in part, due to a lack of a set of universally accepted research guidelines because of the comparative newness of the field. Existing quality standards for Internet-delivered psychological and health interventions, as reviewed below, are insufficient for guiding research. Thus, we outline a framework of guidelines specifically tailored to the components, science, and quality of Internet intervention research. We aim at the construction of a comprehensive framework of guidelines so that they are relevant for a wide scope of online interventions, across communication modality, population, nature of distress, type of intervention, and the other important delivery characteristics.

Existing quality standards

Although consumers have increasingly sought Internet-based psychological and health-related information and care services over the last decade, the quality of such information is variable and largely unregulated and has frequently been questioned (e.g. Bremner, Quinn, Quinn, & Veledar, 2006; Childs, 2005; Risk & Dzenowagis, 2001). In response to concerns about incomplete, misleading, inaccurate, or dangerous online mental and physical health information and practices, two major approaches have arisen in an attempt to guide developers/researchers of health websites and to assist consumers in identifying information resources of good quality.

The first approach has involved the development of principles or codes of conduct to guide creators of health information to

produce good-quality websites (e.g. American Medical Association, n.d.; European Union eHealth, 2010; Rippen & Risk, 2000). Certain organizations, such as the Health on the Net Foundation (1997) have augmented their codes of conduct by certifying websites that comply with their codes. Additionally, websites could apply for third-party accreditation such as TRUSTe (n.d.) for a fee or, alternatively, seek inclusion in a "quality filtered" database or portal, for example, Organising Medical Networked Information (OMNI, n.d.), evaluated by a third party.

A second, parallel approach relating to the delivery of online therapeutic services has focused on the development of ethical standards of conduct (but not research per se) by various professional associations (e.g. American Counseling Association, 2005; Australian Psychological Society, 2011) or specialist societies (e.g. International Society for Mental Health Online, 2000). Here, members are encouraged to follow standards set by their particular professional society when incorporating web-based technology into their work, encompassing such issues as informed consent, confidentiality, research using the Internet, record keeping, and managing professional boundaries when using the web. In addition, several articles have been published discussing best practices and ethical and legal considerations for e-therapy (Abbott, Klein, & Ciechomki, 2008; Dever Fitzgerald, Hunter, Hadjistavropoulos, & Koocher, 2010; Whitehead & Proudfoot, 2010), as well as Beacon (n.d.; www.beacon.anu.edu.au) which is an online portal providing quality ratings of health websites (Christensen, Murray, et al., 2010).

We believe that the existing codes of conduct and standards of web content quality are important scientific resources for guiding intervention practice, but are insufficient as professional standards to guide Internet intervention research activities. Similarly, reports of efficacy, which are common in other research guidelines, are necessary but not sufficient when reporting Internet intervention research. To this end, we developed a set of best-practice guidelines, specifically for Internet interventions, to assist researchers who work in this field.

Method

To establish the guidelines, we utilized aspects of facet theory (Guttman & Greenbaum, 1998), a systematic approach to conceptualizing and defining constructs to assist with theory construction and research design in the behavioural and social sciences. Facet theory provides a structure for defining the “universe of attributes” (in this case, within Internet intervention research) as well as the facets for which we argue that guidelines are needed. First, we outlined a “mapping sentence,” a statement concerning the domain of Internet intervention research. The mapping sentence serves to define a priori exactly what is being studied: the population, the content variables, and the range of possible responses (Guttman & Greenbaum, 1998). Second, using an iterative approach based on empirical findings and research experience, we proposed a number of core “facets” (or necessary components) of reporting Internet intervention research. Third, we developed “elements” for each facet. Elements refer to the range of possible values (or response categories) that a facet contains. On the basis of this conceptualization, we derived guidelines of best practice for each facet, designed for executing and reporting Internet intervention research. The resultant multidimensional model—consisting of facets, elements, and guidelines—was then sent to several leading researchers in the field of Internet interventions for their critical comments and suggestions. The outcome of this process, the consensus framework of guidelines for Internet interventions, is outlined next and summarised in Figure 1.

Framework of guidelines for Internet intervention research

First, the mapping sentence is presented. It is followed by 12 facets, each with their constituent elements and a recommended guideline.

Mapping sentence: the domain of Internet intervention research

An Internet-based intervention is a self-guided or human-assisted program for health promotion, disease and emotional distress prevention, risk factor management, treatment, or relapse prevention that is executed by

means of a prescriptive online intervention and/or communication operated through the Internet and used by consumers and patients seeking health- and mental health-related assistance. The program itself attempts to create positive change and/or enhance knowledge, awareness, and understanding via the provision of sound health-related material and use of web-based components, with or without support from health professionals or others (Barak et al., 2009). The scope of Internet interventions is broad and includes structured websites, chat, and Internet-enabled mobile phone applications. Internet intervention research consists of any formative or summative evaluation of an Internet intervention.

Facet 1: focus and target population

A range of different programs exists under the umbrella term “Internet interventions,” including programs for specific mental health and behavioural health problems as well as those for general mental health and well-being. The target population may be defined in terms of demographic factors (e.g. age, gender, ethnic group, educational level), problem area or symptoms (e.g. insomnia, depression, anxiety, weight loss, diabetes), psychological indices (e.g. treatment history, learning style, self-efficacy, readiness for change, motivation, locus of control), therapeutic factors (e.g. time since diagnosis), and technologically related considerations (e.g. previous experience with computers).

Guideline. The primary focus of the Internet intervention is clearly displayed along with, if relevant, a diagnostic label, cutoff score on a reliable and valid assessment, structured diagnostic interview, or case formulation to identify the problem area or symptoms. A clear statement is also provided about the target group for the intervention, with age limits and other eligibility criteria.

Facet 2: authorship details

Authorship information, such as the identity of the program developers, their affiliation, copyrights and intellectual property ownership, whether there are advertising or commercial sponsors, the country of origin, last update, and the existence of an editorial board or governance structure, provides relevant indicators of the reliability and quality of an Internet intervention.

Guideline. The names, credentials, and affiliations of the program's developers are cited as well as the ownership and affiliation of the program. A link to further information is provided, such as whether and by whom the intervention is sponsored, date on which the program was developed or updated, and its country of origin.

Facet 3: model of change

Process variables or mediators, both specific and nonspecific, are most likely responsible for therapeutic change, although these variables have been continuously studied and are not fully clear (Walker et al., 2006). Specific process variables relate to the theoretical basis of the Internet intervention being evaluated, whereas nonspecific variables include common factors such as empathy for participants' distress, communication of hope for improvement, maintaining participants' motivation, and checking their understanding of and satisfaction with the program (Proudfoot et al., 2003). To date, the theories of therapeutic change relating to Internet interventions have been primarily derived from face-to-face interventions and include approaches such as cognitive behavioural theory (Beck, 1976; Ellis, 1994; Meichenbaum, 1985), behaviour therapy (Lazarus, 1997; Watson & Tharp, 2002), health beliefs model (Glanz, Rimer, & Lewis, 2002), social cognitive/learning theory (Bandura, 1997), trans-theoretical model of behaviour change (Prochaska & DiClemente, 1986), solution-focused theory of change (Egan, 1998), narrative psychotherapies (White & Epston, 1990), and positive psychology (Seligman & Csikszentmihalyi, 2000). Recently, however, Ritterband, Thorndike, Cox and colleagues (2009) have proposed a holistic model of behaviour change and symptom improvement pertaining specifically to Internet interventions. Involving nine components, the model consists of (1) the user, influenced by (2) environmental factors that affect (3) website use and adherence, which are influenced by (4) support and (5) website characteristics. This leads to (6) behavior change and (7) symptom improvement through various (8) mechanisms of change, which are sustained via (9) treatment maintenance.

Guideline. The theory of change underpinning the intervention is clearly articulated when

discussing outcomes. Because Internet intervention research is still in its infancy, we believe it is premature at this stage to require researchers to demonstrate that the process of change was achieved by the intervention's hypothesized mechanism of action; however, we see the field evolving to this point in the future.

Facet 4: type and dose of intervention

Several types of Internet intervention exist, and the number continues to grow as technology develops and new electronic approaches are applied to psychological and behavioural problems. Types of Internet interventions include but are not limited to the following:

1. Targeted prevention, early intervention, treatment, and self-management programs for specific conditions that include registration and screening/assessment functions, are tailored to individual circumstances and are supported by a database that enables users (and, in some circumstances, clinicians) to track user progress and receive feedback. They may be offered in conjunction with professional or other assistance (see later discussion of Facet 6).
2. Prevention, early intervention, treatment, and self-management programs that do not require registration and/or screening and do not include tailoring to specific circumstances or tracking of progress. These are usually offered on a large scale to whole populations or at-risk sections of the public without professional or other support.
3. e-Counselling via various modalities (e.g. e-mail, text chat, audio chat, video chat).
4. Psycho-education for specific conditions that may or may not require screening.
5. Self-screening and self-assessment websites.
6. Moderated forums.

This facet also includes information about the duration between sessions. Some programs, for example, prescribe weekly sessions and others daily sessions, while still others do not stipulate a number of sessions but rather allow unstructured access over a period of time. Some are more variable, for example,

starting with high session frequency and moving to a less frequent schedule after a critical point.

Guideline. The type of the intervention is clearly stated, including its dose (the prescribed and actual number and frequency of sessions or modules undertaken by participants and/or the period of time over which the intervention is accessed), whether it is tailored to individual circumstances, and whether it allows users to track their progress and receive feedback. When assessments are used, it is stated whether they are validated for online administration.

Facet 5: ethical issues

Ethical issues associated with the use of Internet interventions include the provision of relevant information to allow individuals to make an informed decision about whether they want to use the intervention and take part in the study, obtaining acceptance regarding the website's "terms and conditions" or informed consent from those who do want to participate in the research, assuring their confidentiality, maintaining the security of the site or program, outlining users' rights and responsibilities, ensuring that appropriate risk assessment and follow-up procedures are an integral part of the research procedure, and establishing appropriate use for different samples (e.g. issues pertaining to the use by minors) (Whitehead & Proudfoot, 2010). Some potential participants (e.g. those with suicidal plans or histories of abuse) insist on full anonymity if they are to take part in the research, which can pose ethical dilemmas for researchers. Researchers have managed these situations by having participants consent to the terms and conditions without being required to register or provide any identifiable information. More generally, the use of unique user names and passwords, encryption methods, firewalls and back-up procedures, secured sites and transfer of confidential information, and verification of the participant's and therapist's identities (where applicable) and the therapist's credentials are important legal and safety elements.

Guideline. Researchers report the process of using the program, its potential risks and benefits, safeguards against the risks, procedures to follow in an emergency, and sources of help, especially when offering

open-access anonymous online services where registration is not required. In the majority of cases where participants are required to register, they should use a unique user name and password to protect their privacy. Where the site is open access and anonymous, participants should consent to the terms and conditions of the service. The site, software, and data transmission are secure, and data are encrypted. A statement is included about how data are used and stored and when they will be destroyed. Users' rights and responsibilities are outlined as well as the program's liability. If e-mail or another procedure of synchronous or asynchronous communication is used that is not built into the system, participants are informed that it is not entirely secure. Researchers report their clinical trial registration and ethics approval number.

Facet 6: professional support

Professional support refers to the assistance provided by health professionals as an adjunct to the Internet intervention. Not all Internet interventions involve professional support, but when it is provided, elements include the following:

1. Health professional providing the support (e.g. clinical psychologist, psychiatrist, general practitioner, nurse, counselor, or specifically trained helper).
2. Timing and frequency of the support, including the way in which the communication is initiated (e.g. as needed, scheduled on a regular basis, only in response to patients' questions) and how the professional support is timed.
3. Type of assistance provided: assistance with intervention techniques, feedback about homework tasks, tracking progress, crisis management detection and assistance procedures, providing reminders/prompts, redirecting and reinforcing efforts, moderating bulletin board postings or forums, technical support.
4. Delivery modality of support: e-mail, forum, webcam, audio chat, telephone, SMS (texting), face-to-face support, closed message system requiring log-in by the therapist and the client.

Guideline. A statement is provided as to whether or not the Internet intervention is

delivered with professional support. For programs in which it is included, the following further details are specified: type and qualifications of health professional(s) providing the support, type of assistance offered, timing and frequency of the support, how it is initiated, and the medium by which the assistance is delivered.

Facet 7: other support

This facet refers to assistance provided by people other than health professionals. Similar to Facet 6, not all Internet interventions include other support, but when it is provided, four elements are involved:

1. Type of person providing support: for example, research assistant, technician, teacher, mentor, parent, participants' interaction with each other, "informed supporters" (expert patients who are successfully managing their condition).
2. Level of support: none, individually provided, group setting.
3. Type of support: providing information, offering practical advice and strategies, referring to additional help resources, sharing ideas, providing emotional help, following a therapeutic manual or template to assist participant's use of the intervention, technical support.
4. Medium of delivery of support: e-mail, forum, webcam chat, audio chat, telephone, face-to-face support, SMS (texting), chat room.

Guideline. Researchers report whether or not the Internet intervention includes support from someone other than a health professional. The type of person assisting and, where relevant, their qualifications are specified, together with the precise nature of their role, the type and availability of assistance offered, and whether it is anonymous as well as the medium of delivery. The supervisory arrangements should also be reported.

Facet 8: program interactivity

Program interactivity and feedback pertains to the dynamic computer-generated activity offered to users of an intervention and is a key feature of Internet interventions. Levels of interactivity are intentionally varied for the purposes of enhancing engagement, motivation, and adherence; increasing active rather

than passive behaviour; tailoring content to individual users; increasing involvement in decision making; improving learning; increasing user control; and/or enhancing the impact of the intervention (Hawkins et al., 2010; Walther, Pingree, Hawkins, & Buller, 2005). Elements may include questionnaires and quizzes, interactive exercises, action planning, journaling, inputting self-monitoring data, tailored feedback, tasks to complete offline between sessions, chat or chat room discussions, questions and answers, and virtual games.

Guideline. The degree of user interactivity offered by the program is described by researchers, and examples are provided of component interactive features, together with their purpose and the amount of time a typical participant spends on the program (both its online and offline components).

Facet 9: multimedia channel of delivery

Interventions vary according to the number and type of multimedia channels used to communicate information, provide support, build skills, and provide additional assistance. Text, graphics, animations, images, 3-D virtual reality environments, audio, and video are currently used. For example, an intervention may include text and graphics to communicate information, audio instructions for progressive relaxation, video presenting case studies, as well as e-mail support from a therapist and/or the use of automated e-mail or SMS reminders. Graphics and animations may be interactive. Different channels may be chosen to align with intended users' characteristics, degree of distress, e-health literacy, treatment readiness, learning style, motivation, and self-efficacy as well as practical considerations. They are also used to enhance user engagement in the intervention.

Guideline. The communication delivery channels used in the intervention are fully described and explained.

Facet 10: degree of synchronicity

This facet refers to the timing and delay of the communication, feedback, or support from clinicians, from other users, or from the program itself. The facet pertains to all communication channels included in the program, because degree of communication synchronicity might be a principal research (Paulus & Phipps, 2008) or intervention

(Cress, Kimmerle, & Hesse, 2009; Pullen & Snow, 2007) value. The facet includes the following elements:

1. Synchronous: communication or support is immediate (e.g. real-time chat).
2. Asynchronous: communication is delayed, such as with e-mail and forums. The delays may be short (e.g. a daily e-mail) or longer (e.g. therapist's e-mail response every 3 days).
3. Random, periodic, or as required: either predetermined or in response to the user.

Guideline. The synchronicity of support and feedback are clearly reported within the program as well as the duration between program sessions, where relevant.

Facet 11: audience reach

This facet relates to the accessibility of an Internet intervention. Some online programs, particularly those with no professional or other (human) assistance, offer open access via the web or free and open-to-all online communication and thus have extremely broad reach (Christensen et al., 2004; Clarke et al., 2005; Farvolden, Denisoff, Selby, Bagby, & Rudy, 2005; Proudfoot et al., 2007) but potentially with limited treatment adherence. Others, such as those coupled with mandatory a priori clinical assessments or therapist assistance, have a more limited and narrow reach (Carlbring, Westling, Ljungstrand, Ekselius, & Andersson, 2001; Klein et al., 2006, 2010; Klein, Austin, et al., 2009; Klein, Mitchell, et al., 2009; Ström, Pettersson, & Andersson, 2000; Thorndike et al., 2008) but usually lead to greater adherence to treatment. A third type of intervention limits availability to users affiliated with a certain group or institution, such as military personnel, a certain clinic or hospital patients, or university students (Bergström et al., 2010; Hazzard, Celano, Collins, & Markov, 2002; Winzelberg et al., 2000).

Guideline. Details of the participant's mode of access to the intervention are clearly defined, as are the reasons for exclusion (if applicable). Additional information regarding other potential sources of support is provided in case participants drop out.

Facet 12: program evaluation

In considering guidelines related to efficacy, effectiveness, efficiency, and safety, those

proposed for empirically supported psychological interventions (Chambless & Hollon, 1998; Chambless & Ollendick, 2001) and those established by the Society for Prevention Research (Flay et al., 2005) serve as a basic foundation. However, the processes of evaluation in the field of e-health are necessarily different and require their own research methodology.

While randomized controlled trials are prioritized in outcome research, other types of evaluation are also indicated in Internet intervention research. For example, formative evaluation is recommended to explore ways of meeting cultural, socioeconomic, and technological literary challenges of underserved groups (Ahern, 2007; Ahern, Kreslake, & Phalen, 2006) or those lacking the motivation or resources to access the Internet (Campbell-Grossman, Hudson, Keating-Lefler, & Heusinkvelt, 2009; Finfgeld-Connett & Madsen, 2008). Standardization of measures is desirable to aid comparisons across studies, and online tools should be validated against their paper-and-pencil or face-to-face counterparts (e.g. Carlbring, Brunt, et al., 2007; Herrero & Meneses, 2006; Lygidakis, Cambiaso, Cuzzo, & Bella, 2010; Thorndike et al., 2009, in press; Vallejo, Mañanes, Comeche, & Díaz, 2008). The use of online tests and questionnaires is feasible but should be cautiously handled in clinical-related assessments (Barak, 2010; Buchanan, 2003, 2007). In addition to symptom and/or behaviour change, users' functioning and quality of life can be measured as well as health service usage. Process measures, where applicable, such as usage, traffic, and utilization as well as dropout, are also important because they can shed light on delivery mechanisms, users' behaviour, differential attrition, and outcomes (Ahern et al., 2006; Christensen, Griffiths, & Farrer, 2009). Internet interventions are especially suited to monitoring effectiveness in clinical practice, such as assessing the client-therapist alliance, tracking processes of change, and assessing posttreatment and long-term functioning. Measures of user acceptance should go beyond the collection of satisfaction ratings from intervention "completers" to include acceptability surveys of nonusers and potential users, qualitative studies of how users interact with a system (Beattie, Shaw, Kaur, & Kessler, 2009), measures of

take-up rates, and reasons for disengagement. Feasibility and cost-effectiveness are yet other important factors, especially in relation to traditional interventions, and should be estimated or assessed (Abroms, Gill, Windsor, & Simon-Morton, 2009; Hedman et al., 2010; Klein et al., 2010; Schoenberg et al., 2008; Tate, Finkelstein, Khavjou, & Gustafson, 2009; Titov, Andrews, Johnston, Schwenke, & Choi, 2009; Warmerdam, Smit, van Straten, Riper, & Cuijpers, 2010). New approaches to data analysis (e.g. preference vs. randomized designs, statistical simulations, permutation analyses, tracking analyses) may be necessary in Internet intervention research as the field develops and evolves (Danaher & Seeley, 2009; National Cancer Institute & Robert Wood Johnson Foundation, 2001). Mixed-method research that utilizes qualitative methods to help understand the quantitative results are also sometimes necessary (Glasgow, 2009). Some Internet intervention research resembles the evaluation of "complex interventions" (Campbell et al., 2000). In these cases, recognition of additional complexities, such as participant preferences, selection biases, differential dropout rates between intervention and control groups, as well as the use of appropriate quantitative and qualitative approaches, is recommended.

Guidelines.

1. **Efficacy:** Guidelines here are similar to those in CONSORT-R for randomized controlled trials. Researchers define the size and characteristics of their sample on the basis of a power analysis (including the proportions of participants taking medication and changing their dosage during the trial), outline the method of group allocation conducted (including how it was concealed), define the comparison groups, cite the psychometric properties of the measures used, outline the data collection procedures and the schedule of measurements implemented (including long-term follow up), and describe the type of data analyses undertaken, including whether they are completer or intention-to-treat analyses. Importantly, they provide details of the sample attrition at each study time point and the methods of analysis employed to handle dropouts if

appropriate. Details of adherence to the intervention are also provided, including participants' reasons for nonadherence where relevant. Findings of the study and the conclusions drawn are reported.

2. **Effectiveness:** Information about the intervention's effectiveness in real-world conditions is reported, including the size and characteristics of the sample, the measures applied, and the time schedule over which the assessments were taken. Routine outcome measurement is reported as well as participants' adherence to the intervention. An outline is provided of the generalizability of the program's effects and user suitability characteristics.
3. **Readiness for mass dissemination:** Cost-effectiveness data are reported (as per the efficacy guidelines above), along with information about the capacity for the program to be scaled up for widespread release, including inherent safety provisions.

Conclusion

The field of Internet interventions is growing. Evidence can be seen in the increasing penetration of such interventions into therapist training programs and into the roles of new behavioural and mental health professionals (such as low-intensity practitioners; Bennett-Levy et al., 2010), by the expanding number of intervention websites, and by the fast-growing number of people in need who seek and receive professional help online. Simultaneously, research in this area is increasing rapidly too. This growth can be facilitated by considering clear, broadly accepted, substantiated guidelines for the execution and reporting of scientific research, which will also promote efficient and improved communication among researchers in the field.

The guidelines were derived using an iterative model of discussion leading to consensus. It is important to note that any set of guidelines is not simply and automatically adopted by its relevant users unless there are broadly accepted regulations assigned to it. This means that the set of guidelines proposed here must be discussed by the professional community before acceptance and broad implementation can be expected. Consequent practical steps may include

adoption by relevant bodies and professional organizations as well as by journals and other publication outlets. Needless to say, governing committees and national agencies will need to be involved in actual implementation. This has been exemplified and stressed by developers of equivalent guidelines for Internet-based experimenting (Reips, 2002), Internet-based research (Spyridakis, Wei, Barrick, Uddihy, & Maust, 2005), and Internet-delivered testing (International Test Commission, 2006). With the information presented here, researchers have guidelines to follow when executing and reporting their research as well as an infrastructure for further discussion and development in the field of Internet intervention research.

Acknowledgements

The authors thank and remember Jen McLaren for her assistance in editing the paper. JP is grateful to the National Health and Medical Research Council (Program Grant 510135) for salary support. No competing financial interests exist.

References

- Abbott, J., Klein, B., & Ciechomski, L. (2008). Best practices in online therapy. *Journal of Technology in Human Services*, 26, 360–375.
- Abbott, J., Klein, B., Hamilton, C., & Rosenthal, A. (2009). The impact of online resilience training for sales managers on wellbeing and work performance. *Electronic Journal of Applied Psychology*, 5, 89–95.
- Abroms, L. J., Gill, J., Windsor, R., & Simon-Morton, B. (2009). A process evaluation of e-mail counseling for smoking cessation in college students: Feasibility, acceptability and cost. *Journal of Smoking Cessation*, 4, 26–33.
- Ahern, D. (2007). Challenges and opportunities of ehealth research. *American Journal of Preventive Medicine*, 32, s75–s85.
- Ahern, D., Kreslake, J., & Phalen, J. (2006). What is ehealth (6): Perspectives on the evolution of ehealth research. *Journal of Medical Internet Research*, 8(1), p. e4.
- American Counseling Association (2005). *ACA code of ethics*. Retrieved March 10, 2011, from <http://www.counseling.org/Files/FD.ashx?guid=ab7c1272-71c4-46cf-848c-f98489937dda>.
- American Medical Association (n.d). *AMA code of medical ethics*. Retrieved May 1, 2011, from <http://www.ama-assn.org/ama/pub/physician-resources/medical-ethics/code-medical-ethics.page>
- Andersson, G. (2009). Using the Internet to provide cognitive behaviour therapy. *Behaviour Research and Therapy*, 47, 175–180.
- Andersson, G., Bergström, J., Hollädare, F., Carlbring, P., Kaldö, V., & Ekselius, L. (2005). Internet-based self-help for depression: A randomised controlled trial. *British Journal of Psychiatry*, 187, 456–461.
- Andersson, G., Carlbring, P., Holmström, A., Sparthán, E., Furmark, T., Nilsson-Ihrfelt, E., ... Ekselius, L. (2006). Internet-based self-help with therapist feedback and in-vivo group exposure for social phobia: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 74, 677–686.
- Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized psychological treatments for adult depression: A meta-analysis. *Cognitive Behaviour Therapy*, 38, 196–205.
- Andersson, G., Ljotsson, B., & Weise, C. (2011). Internet-delivered treatment to promote health. *Current Opinion in Psychiatry*, 24, 168–172.
- Andrews, G., Cuijpers, P., Craske, M. G., McEvoy, P., & Titov, N. (2010). Computer therapy for the anxiety and depressive disorders is effective, acceptable and practical health care: A meta-analysis. *PloS ONE*, 5(10), p. e13196.
- Australian Psychological Society (2011). *Guidelines for providing psychological services and products using the Internet and telecommunications technologies*. Retrieved March 8, 2011 from <http://www.psychology.org.au/Assets/Files/EG-Internet.pdf>.
- Bandura, A. (1997). *Self-efficacy: The exercise of control*. New York: WH Freeman.
- Barak, A. (2010). Internet-supported psychological testing and assessment. In R. Kraus, G. Stricker, & C. Speyer (Eds.), *Online counseling: A handbook for mental health professionals* (2nd ed., pp. 225–255). San Diego, CA: Elsevier.
- Barak, A., Hen, L., Boniel-Nissim, M., & Shapira, N. (2008). A comprehensive review and a meta-analysis of the effectiveness of Internet-based psychotherapeutic interventions. *Journal of Technology in Human Services*, 26, 109–160.
- Barak, A., Klein, B., & Proudfoot, J. (2009). Defining Internet-supported therapeutic interventions. *Annals of Behavioral Medicine*, 38, 4–17.
- Beacon (n.d). *Find self-help for mental and physical health*. Retrieved May 20, 2010, from <http://www.webcitation.org/5prLbw654>.
- Beattie, A., Shaw, A., Kaur, S., & Kessler, D. (2009). Primary-care patients' expectations and experiences of online cognitive behavioural therapy for depression: A qualitative study. *Health Expectations*, 12, 45–59.
- Beck, A. (1976). *Cognitive therapy and emotional disorders*. New York: Meridian.
- Bennett, K., Reynolds, J., Christensen, H., & Griffiths, K. M. (2010). e-Hub: An online self-help mental health service in the community. *Medical Journal of Australia*, 192(11), S48–S52.
- Bennett-Levy, J., Richards, D., Farrand, P., Christensen, H., Griffiths, K. M., Kavanagh,

- D., ..., & Williams, C. (Eds.). (2010). *Oxford guide to low intensity CBT interventions*. New York: Oxford University Press.
- Bergström, J., Andersson, G., Ljótsson, B., Rück, C., Andréewitch, S., Karlsson, A., & Lindefors, N. (2010). Internet- versus group-administered cognitive behaviour therapy for panic disorder in a psychiatric setting: A randomised trial. *BMC Psychiatry*, 10, p. 54.
- Bremner, J. D., Quinn, J., Quinn, W., & Veledar, E. (2006). Surfing the Net for medical information about psychological trauma: An empirical study of the quality and accuracy of trauma-related websites. *Medical Informatics and the Internet in Medicine*, 31, 227–236.
- Buchanan, T. (2003). Internet-based questionnaire assessment: Appropriate use in clinical contexts. *Cognitive Behaviour Therapy*, 32, 100–109.
- Buchanan, T. (2007). Personality testing on the Internet: What we know, and what we do not. In A. Joinson, K. McKenna, T. Postmes, & U. Reips (Eds.), *The Oxford handbook of Internet psychology* (pp. 447–460). Oxford, UK: Oxford University Press.
- Campbell, M., Fitzpatrick, R., Haines, A., Kinmouth, A. L., Sandercock, P., Spiegelhalter, D., & Trier, P. (2000). Framework for design and evaluation of complex interventions to improve health. *British Medical Journal*, 321, 694–696.
- Campbell-Grossman, C. K., Hudson, D. B., Keating-Lefler, R., & Heusinkvelt, S. (2009). New mothers network: The provision of social support to single, low-income, African American mothers via e-mail messages. *Journal of Family Nursing*, 15, 220–236.
- Carlbring, P., Brunt, S., Bohman, S., Austin, D., Richards, J., Öst, L., & Andersson, G. (2007). Internet vs. paper and pencil administration of questionnaires commonly used in panic/agoraphobia research. *Computers in Human Behavior*, 23, 1421–1434.
- Carlbring, P., Gunnarsdóttir, M., Hedensjö, L., Andersson, G., Ekselius, L., & Furmark, T. (2007). Treatment of social phobia: Randomised trial of Internet-delivered cognitive-behavioural therapy with telephone support. *British Journal of Psychiatry*, 190, 123–128.
- Carlbring, P., Westling, B., Ljungstrand, P., Ekselius, L., & Andersson, G. (2001). Treatment of panic disorder via the Internet: A randomized trial of a self-help program. *Behavior Therapy*, 32, 751–764.
- Cavanagh, K., & Shapiro, D. A. (2004). Computer treatment for common mental health problems. *Journal of Clinical Psychology*, 60, 239–251.
- Chambless, D., & Hollon, S. (1998). Defining empirically supported therapies. *Journal of Consulting and Clinical Psychology*, 66, 7–18.
- Chambless, D., & Ollendick, T. (2001). Empirically supported psychological interventions: Controversies and evidence. *Annual Review of Psychology*, 52, 685–716.
- Childs, S. (2005). Judging the quality of Internet-based health information. *Performance Measurement and Metrics*, 6, 80–96.
- Christensen, H., Griffiths, K. M., & Farrer, L. (2009). Adherence in Internet interventions for anxiety and depression: Systematic review. *Journal of Medical Internet Research*, 11(2), p. e13.
- Christensen, H., Griffiths, K. M., & Jorm, A. F. (2004). Delivering interventions for depression by using the Internet: Randomised controlled trial. *British Medical Journal*, 328, 265–269.
- Christensen, H., Griffiths, K. M., Mackinnon, A., Kalia, K., Batterham, P. J., Kenardy, J., ... Bennett, K. (2010). Protocol for a randomised controlled trial investigating the effectiveness of an online e health application for the prevention of generalised anxiety disorder. *BMC Psychiatry*, 10, p. 25.
- Christensen, H., Murray, K., Cleave, A., Bennett, K., Bennett, A., & Griffiths, K. (2010). Beacon: A web portal to high-quality mental health websites for use by health professional and the public. *Medical Journal of Australia*, 192, 40–44.
- Clarke, G., Eubanks, D., Reid, E., Kelleher, C., O'Connor, E., DeBar, L. L., ... Gullion, C. (2005). Overcoming depression on the Internet (ODIN) (2): A randomized trial of a self-help depression skills program with reminders. *Journal of Medical Internet Research*, 7(2), p. e16.
- Cress, U., Kimmerle, J., & Hesse, F. W. (2009). Impact of temporal extension, synchronicity, and group size on computer-supported information exchange. *Computers in Human Behavior*, 25, 731–737.
- Cuijpers, P., Marks, I. M., van Straten, A., Cavanagh, K., Gega, L., & Andersson, G. (2009). Computer-aided psychotherapy for anxiety disorders: A meta-analytic review. *Cognitive Behaviour Therapy*, 38, 66–82.
- Danaher, B. G., & Seeley, J. R. (2009). Methodological issues in research on web-based behavioral interventions. *Annals of Behavioral Medicine*, 38, 28–39.
- Dever Fitzgerald, T., Hunter, P. V., Hadjistavropoulos, T., & Koocher, G. P. (2010). Ethical and legal considerations for Internet-based psychotherapy. *Cognitive Behaviour Therapy*, 39, 173–187.
- Egan, G. (1998). *The skilled helper: A problem-management approach to helping*. Pacific Grove, CA: Brooks/Cole.
- Ellis, A. (1994). *Reason and emotion in psychotherapy revised*. New York: Kensington.
- European Union eHealth (2010). *Quality criteria*. Retrieved May 1, 2011, from http://ec.europa.eu/health-eu/care_for_me/quality_assurance/index-en.htm.
- Farvolden, P., Denisoff, E., Selby, P., Bagby, R., & Rudy, L. (2005). Usage and longitudinal effectiveness of a web-based self-help cognitive behavioral therapy program for panic disorder. *Journal of Medical Internet Research*, 7(1), p. e7.
- Fingfeld-Connett, D., & Madsen, R. (2008). Web-based treatment of alcohol problems among rural women: Results of a randomized pilot

- investigation. *Journal of Psychosocial Nursing*, 46, 46–53.
- Flay, B. R., Biglan, A., Boruch, R. F., Castro, F. G., Gottfredson, D., Kellam, S., ... Ji, P. (2005). Standards of evidence: Criteria for efficacy, effectiveness and dissemination. *Prevention Science*, 6, 151–175.
- Glanz, K., Rimer, B., & Lewis, F. (2002). *Health behavior and health education: Theory, research and practice*. San Francisco, CA: Wiley.
- Glasgow, R. (2009). Enhancing the scientific foundation of Internet intervention research. *Annals of Behavioral Medicine*, 38, 46–47.
- Griffiths, K. M., Farrer, L., & Christensen, H. (2010). The efficacy of Internet interventions for depression and anxiety disorders: A review of randomised controlled trials. *Medical Journal of Australia*, 192(11), S4–S11.
- Guttman, R., & Greenbaum, C. (1998). Facet theory: Its development and current status. *European Psychologist*, 3, 13–36.
- Hawkins, R. P., Han, J. -Y., Pingree, S., Shaw, B. R., Baker, T. B., & Roberts, L. J. (2010). Interactivity and presence of three health interventions. *Computers in Human Behavior*, 26, 1081–1088.
- Hazzard, A., Celano, M., Collins, M., & Markov, Y. (2002). Effects of STARBRIGHT world on knowledge, social support, and coping in hospitalized children with sickle cell disease and asthma. *Children's Health Care*, 31, 69–86.
- Health on the Net Foundation (1997). *HON code of conduct (HONcode) for medical and health websites*. Retrieved May 20, 2010, from <http://www.webcitation.org/5prKsG5jU>.
- Hedman, E., Ljótsson, B., Andersson, E., Rück, C., Andersson, G., & Lindefors, N. (2010). Effectiveness and cost offset analysis of group CBT for hypochondriasis delivered in a psychiatric setting: An open trial. *Cognitive Behaviour Therapy*, 39, 239–250.
- Herrero, J., & Meneses, J. (2006). Short web-based versions of the perceived stress (PSS) and Center for Epidemiological Studies-Depression (CESD) Scales: A comparison to pencil and paper responses among Internet users. *Computers in Human Behavior*, 22, 830–846.
- International Society for Mental Health Online (2000). *Suggested principles for the online provision of mental health services*. Retrieved May 20, 2010, from <http://www.webcitation.org/5prLQPG9x>.
- International Test Commission (2006). International guidelines of computer-based and Internet-delivered testing. *International Journal of Testing*, 6, 181–188.
- Kenardy, J., McCafferty, K., & Rosa, V. (2003). Internet-delivered indicated prevention for anxiety disorders: A randomized controlled trial. *Behavioural and Cognitive Psychotherapy*, 31, 279–289.
- Klein, B., Austin, D., Pier, C., Kiropoulos, L., Shandley, K., Mitchell, J., ... Ciechowski, L. (2009). Internet-based treatment for panic disorder: Does frequency of therapist contact make a difference? *Cognitive Behavior Therapy*, 38, 100–113.
- Klein, B., Mitchell, J., Abbott, J., Shandley, K., Austin, D., Gilson, K., ... Redman, T. (2010). A therapist-assisted cognitive behaviour therapy Internet intervention for posttraumatic stress disorder: Pre-, post- and 3 month follow-up results from an open trial. *Journal of Anxiety Disorders*, 24, 635–644.
- Klein, B., Mitchell, J., Gilson, K., Shandley, K., Austin, D., Kiropoulos, L., ... Cannard, G. (2009). A therapist-assisted Internet-based CBT intervention for post-traumatic stress disorder: Preliminary results. *Cognitive Behavioural Therapy*, 38, 121–131.
- Klein, B., & Richards, J. C. (2001). A brief Internet-based treatment for panic disorder. *Behavioural and Cognitive Psychotherapy*, 29, 113–117.
- Klein, B., Richards, J. C., & Austin, D. (2006). Efficacy of Internet therapy for panic disorder. *Journal of Behavior Therapy and Experimental Psychiatry*, 37, 231–238.
- Kuhl, E. A., Sears, S. F., & Conti, J. B. (2006). Internet-based behavioral change and psychosocial care for patients with cardiovascular disease: A review of cardiac disease-specific applications. *Heart and Lung*, 35, 374–382.
- Lange, A., Rietdijk, D., Hudcovicova, M., Van De Ven, J., Schrieken, B., & Emmelkamp, P. (2003). Interapy: A controlled randomized trial of the standardized treatment of posttraumatic stress through the Internet. *Journal of Consulting and Clinical Psychology*, 71, 901–909.
- Lange, A., & Ruwaard, J. (2010). Ethical dilemmas in online research & treatment of sexually abused adolescents. *Journal of Medical Internet Research*, 12(5), p. e58.
- Lazarus, A. (1997). *Brief but comprehensive psychotherapy: The multimodal way*. New York: Springer.
- Ljótsson, B., Hedman, E., Lindfors, P., Hursti, T., Lindefors, N., Andersson, G., & Rück, C. (2011). Long-term follow-up of Internet-delivered exposure and mindfulness based treatment for irritable bowel syndrome. *Behaviour Research and Therapy*, 49, 58–61.
- Ljótsson, B., Mitsell, K., Lundin, C., Carlbring, P., & Ghaderi, A. (2007). Remote treatment of bulimia nervosa and binge eating disorder: A randomized trial of Internet-based cognitive behavioral therapy. *Behavior Research and Therapy*, 45, 649–661.
- Lygidakis, C., Cambiaso, S., Cuzzo, F., & Bella, C. D. (2010). A web-based versus paper questionnaire on alcohol and tobacco in adolescents. *Telemedicine and E-Health*, 16, 925–930.
- Macea, D. D., Gajos, K., Calil, Y. A. D., & Fregni, F. (2010). The efficacy of web-based cognitive behavioral interventions for chronic pain: A systematic review and meta-analysis. *Journal of Pain*, 11, 917–929.
- Meichenbaum, D. (1985). *Stress inoculation training*. New York: Pergamon Press.

- Mitchell, J., Stanimirovic, R., Klein, B., & Vella-Brodick, D. (2009). A randomised controlled trial of a self-guided Internet intervention promoting well-being. *Computers in Human Behavior*, 25, 749–760.
- Mohr, D. C., Siddique, J., Ho, J., Duffecy, J., Jin, L., & Fokuo, J. K. (2010). Interest in behavioral and psychological treatments delivered face-to-face, by telephone, and by Internet. *Annals of Behavioral Medicine*, 40, 89–98.
- Myung, S. -K., McDonnell, D. D., Kazinets, G., Seo, H. G., & Moskowitz, J. M. (2009). Effects of web- and computer-based smoking cessation programs: Meta-analysis of randomized controlled trials. *Archives of Internal Medicine*, 169, 929–937.
- National Cancer Institute and Robert Wood Johnson Foundation (2001). *Online behavior change and disease management research: A research dialog 2001*. Retrieved May 20, 2010, from <http://www.webcitation.org/5prOXCaou>.
- National Institute of Clinical Excellence (2004). *Depression: Management of depression in primary and secondary care*. London: Author.
- Neve, M., Morgan, P. J., Jones, P. R., & Collins, C. E. (2010). Effectiveness of web-based interventions in achieving weight loss and weight loss maintenance in overweight and obese adults: A systematic review with meta-analysis. *Obesity Reviews*, 11, 306–321.
- OMNI. (n.d). *Organising Medical Networked Information now known as the 'Medicine gateway of Intute: Health and Life Sciences'*. Retrieved May 20, 2010, from <http://www.webcitation.org/5prLEwEoo>.
- Paulus, T. M., & Phipps, G. (2008). Approaches to case analyses in synchronous and asynchronous environments. *Journal of Computer-Mediated Communication*, 13, 459–484.
- Pier, C., Austin, D. W., Klein, B., Mitchell, J., Schattner, P., Ciechomski, L., ... Wade, V. (2008). A controlled trial of Internet-based cognitive-behavioural therapy for panic disorder with face-to-face support from a general practitioner or email support from a psychologist. *Mental Health in Family Medicine*, 5, 29–39.
- Prochaska, J., & DiClemente, C. (1986). Toward a comprehensive model of change. In W. Miller & N. Heather (Eds.), *Treating addictive behaviors: Processes of change* (pp. 3–27). New York: Plenum Press.
- Proudfoot, J., Parker, G., Hyett, M., Manicavasagar, V., Smith, M., Grdovic, S., & Greenfield, L. (2007). Next generation of self-management education: Web-based bipolar disorder program. *Australian and New Zealand Journal of Psychiatry*, 41, 903–909.
- Proudfoot, J., Swain, S., Widmer, S., Watkins, E., Goldberg, D., Marks, I., ... Gray, J. A. (2003). The development and beta-test of a computer therapy program for anxiety and depression: Hurdles and lessons. *Computers in Human Behavior*, 19, 277–289.
- Pullen, J. M., & Snow, C. (2007). Integrating synchronous and asynchronous Internet distributed education for maximum effectiveness. *Education and Information Technologies*, 12, 137–148.
- Reger, M. A., & Gahm, G. A. (2009). A meta-analysis of the effects of Internet- and computer-based cognitive-behavioral treatments for anxiety. *Journal of Clinical Psychology*, 65, 53–75.
- Reips, U. (2002). Standards for Internet-based experimenting. *Experimental Psychology*, 49, 243–256.
- Rippen, H., & Risk, A. (2000). e-Health code of ethics (May 24). *Journal of Medical Internet Research*, 2(2), p. e9.
- Risk, A., & Dzenowagis, J. (2001). Review of Internet health information quality initiatives. *Journal of Medical Internet Research*, 3(4), p. e28.
- Ritterband, L. M., Cox, D., Walker, L., Kovatchev, B., McKnight, L., & Patel, K. (2003). An Internet intervention as adjunctive therapy for pediatric encopresis. *Journal of Consulting and Clinical Psychology*, 71, 910–917.
- Ritterband, L. M., & Tate, D. (2009). The science of Internet interventions. *Annals of Behavioral Medicine*, 38, 1–3.
- Ritterband, L. M., Thorndike, F. P., Cox, D. J., Kovatchev, B. P., & Gonder-Frederick, L. A. (2009). A behavior change model for Internet interventions. *Annals of Behavioral Medicine*, 38, 18–27.
- Ritterband, L. M., Thorndike, F., Gonder-Frederick, L., Magee, J., Bailey, E., Saylor, D., & Morin, C. M. (2009). Efficacy of an Internet-based behavioral intervention for adults with insomnia. *Archives of General Psychiatry*, 66, 692–698.
- Robertson, L., Smith, M., Castle, D., & Tannenbaum, D. (2006). Using the Internet to enhance the treatment of depression. *Australasian Psychiatry*, 14, 413–417.
- Rooke, S., Thorsteinsson, E., Karpin, A., Copeland, J., & Allsop, D. (2010). Computer-delivered interventions for alcohol and tobacco use: A meta-analysis. *Addiction*, 135, 1381–1390.
- Ruwaard, J., Broeksteeg, J., Schrieken, B., Emmelkamp, P., & Lange, A. (2010). Web-based therapist-assisted cognitive behavioural treatment of panic symptoms: A randomised controlled trial with a three-year follow-up. *Journal of Anxiety Disorders*, 24, 387–396.
- Ruwaard, J., Lange, A., Bouwman, M., Broeksteeg, J., & Schrieken, B. (2007). E-mailed standardized cognitive behavioural treatment of work-related stress: A randomized controlled trial. *Cognitive Behaviour Therapy*, 36, 179–192.
- Ruwaard, J., Schrieken, B., Schrijver, M., Broeksteeg, J., Dekker, J., Vermeulen, H., & Lange, A. (2009). Standardized web-based CBT of mild to moderate depression: A randomized controlled trial with a long-term follow-up. *Cognitive Behaviour Therapy*, 38, 206–222.

- Schneider, A., Mataix-Cols, D., Marks, I., & Bachofen, M. (2003). Internet-guided self-help with or without exposure therapy for phobic and panic disorders. *Psychotherapy and Psychosomatics*, 74, 154–164.
- Schoenberg, M. R., Ruwe, W. D., Dawson, K., McDonald, N. B., Houston, B., & Forducey, P. G. (2008). Comparison of functional outcomes and treatment cost between a computer-based cognitive rehabilitation teletherapy program and a face-to-face rehabilitation program. *Professional Psychology: Research and Practice*, 39, 169–175.
- Seligman, M. E. P., & Csikszentmihalyi, M. (2000). Positive psychology: An introduction. *American Psychologist*, 55, 5–14.
- Spek, V., Cuijpers, P., Nyklicek, I., Riper, H., Keyzer, J., & Pop, V. (2007). Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: A meta-analysis. *Psychological Medicine*, 37, 319–328.
- Spyridakis, J., Wei, C., Barrick, J., Uddihy, E., & Maust, B. (2005). Internet-based research: Providing a foundation for web-design guidelines. *IEEE Transactions on Professional Communication*, 48, 242–260.
- Stinson, J., Wilson, R., Gill, N., Yamada, J., & Holt, J. (2009). A systematic review of Internet-based self-management interventions for youth with health conditions. *Journal of Pediatric Psychology*, 34, 495–510.
- Ström, L., Pettersson, R., & Andersson, G. (2000). Controlled trial of self-help treatment of recurrent headache conducted via the Internet. *Journal of Consulting and Clinical Psychology*, 68, 722–727.
- Ström, L., Pettersson, R., & Andersson, G. (2004). Internet-based treatment for insomnia: A controlled evaluation. *Journal of Consulting and Clinical Psychology*, 72, 113–120.
- Swartz, L., Noell, J., Schroeder, S., & Ary, D. (2006). A randomised control study of a fully automated Internet-based smoking cessation program. *Tobacco Control*, 15, 7–12.
- Tait, R., & Christensen, H. (2010). Internet-based interventions for young people with problematic substance use: A systematic review. *Medical Journal of Australia*, 192, 15–21.
- Tate, D. F., Finkelstein, E. A., Khavjou, O., & Gustafson, A. (2009). Cost effectiveness of Internet interventions: Review and recommendations. *Annals of Behavioral Medicine*, 38, 40–45.
- Tate, D. F., Jackvony, E., & Wing, R. A. (2006). Randomized trial comparing human e-mail counseling, computer-automated tailored counseling, and no counseling in an Internet weight loss program. *Archives of Internal Medicine*, 166, 1620–1625.
- Thorndike, F. P., Carlbring, P., Smyth, F. L., Magee, J., Gonder-Frederick, L., Öst, L. -G., & Ritterband, L. M. (2009). Web-based measurement: Effect of completing single or multiple items per webpage. *Computers in Human Behaviour*, 25, 393–401.
- Thorndike, F. P., Ritterband, L. M., Saylor, D. K., Magee, J. C., Gonder-Frederick, L. A., & Morin, C. M. (in press). Validation of the Insomnia Severity Index as a web-based measure. *Behavioral Sleep Medicine*.
- Thorndike, F. P., Saylor, D. K., Bailey, E. T., Gonder-Frederick, L., Morin, C. M., & Ritterband, L. M. (2008). Development and perceived utility and impact of an Internet intervention for insomnia. *E-Journal of Applied Psychology*, 4(2), 32–42.
- Titov, N., Andrews, G., Choi, I., Schwenke, G., & Mahoney, A. (2008). Shyness 3: Randomised controlled trial of guided versus unguided Internet-based CBT for social phobia. *Australian and New Zealand Journal of Psychiatry*, 42, 1030–1040.
- Titov, N., Andrews, G., Davies, M., McIntyre, K., Robinson, E., & Solley, K. (2010). Internet treatment for depression: A randomized controlled trial comparing clinician vs. technician assistance. *PLoS ONE*, 5(6), p. e10939.
- Titov, N., Andrews, G., Johnston, L., Schwenke, G., & Choi, I. (2009). Shyness programme: Longer term benefits, cost-effectiveness, and acceptability. *Australian and New Zealand Journal of Psychiatry*, 43, 36–44.
- Titov, N., Andrews, G., & Schwenke, G. (2008). Shyness 2: Treating social phobia online: Replication and extension. *Australian and New Zealand Journal of Psychiatry*, 42, 595–605.
- TRUSTe (n.d). *TRUSTe seals*. Retrieved May 1, 2011, from <http://www.truste.org>
- Vallejo, M. A., Mañanes, G., Comeche, M. I., & Díaz, M. I. (2008). Comparison between administration via Internet and paper-and-pencil administration of two clinical instruments: SCL-90-R and GHQ-28. *Journal of Behavior Therapy and Experimental Psychiatry*, 39, 201–208.
- Van't Hof, E., Cuijpers, P., & Stein, D. J. (2009). Self-help and Internet-guided interventions in depression and anxiety disorders: A systematic review of meta-analyses. *CNS Spectrums*, 14, 34–40.
- Vernmark, K., Lenndin, J., Bjärehed, J., Carlsson, M., Karlsson, J., Öberg, J., ... Andersson, G. (2010). Internet administered guided self-help versus individualized e-mail therapy: A randomized trial of two versions of CBT for major depression. *Behaviour Research and Therapy*, 48, 368–376.
- Walker, M., Toneatto, T., Potenza, M. N., Petry, N., Ladouceur, R., Hodgins, D. C., ... Blaszczynski, A. (2006). A framework for reporting outcomes in problem gambling treatment research: The Banff, Alberta consensus. *Addiction*, 101, 504–511.
- Walther, J., Pingree, S., Hawkins, R., & Buller, D. (2005). Attributes of interactive online health information systems. *Journal of Medical Internet Research*, 7(3), p. e33.
- Warmerdam, L., Smit, F., van Straten, A., Riper, H., & Cuijpers, P. (2010). Cost-utility and cost-effectiveness of Internet-based treatment for

- adults with depressive symptoms: Randomized trial. *Journal of Medical Internet Research*, 12(5), p. e53.
- Watson, D., & Tharp, R. (2002). *Self directed behavior: Self modification for personal adjustment* (8th ed.). Pacific Grove, CA: Brooks/Cole.
- White, A., Kavanagh, D., Stallman, H., Klein, B., Kay-Lambkin, F., Proudfoot, J., ... Hines, E. (2010). Online alcohol interventions: A systematic review. *Journal of Medical Internet Research*, 12, p. e62.
- White, M., & Epston, D. (1990). *Narrative means to therapeutic ends*. New York, NY: Norton.
- Whitehead, L., & Proudfoot, J. (2010). Standards and operating guidelines for Internet interventions. In J. Bennett-Levy, D. Richards, P. Farrand, P. H. Christensen, K. Griffiths, D. J. Kavanagh, ... C. Williams (Eds.), *Oxford guide to low intensity CBT interventions* (pp. 247–252). Oxford, UK: Oxford University Press.
- Winzelberg, A., Eppstein, D., Eldredge, K., Wilfley, D., Dasmahapatra, R., Dev, P., & Taylor, C. B. (2000). Effectiveness of an Internet-based program for reducing risk factors for eating disorders. *Journal of Consulting and Clinical Psychology*, 68, 346–350.