

# VU Research Portal

## Host immune response in children with Down syndrome related to respiratory tract infection

Broers, C.J.M.

2016

### **document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

### **citation for published version (APA)**

Broers, C. J. M. (2016). *Host immune response in children with Down syndrome related to respiratory tract infection*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# Chapter 5

## **Frequency of lower respiratory tract infections in relation to adaptive immunity in children with Down syndrome compared to their healthy siblings**

Chantal J.M. Broers  
Reinoud J.B.J. Gemke  
Michel E. Weijerman  
Dirk-J. Kuik  
Ingrid M.W. van Hoogstraten  
A. Marceline van Furth

## Abstract

**Aim:** Children with Down syndrome (DS) experience respiratory tract infections (RTIs) more frequently than healthy children. We investigated whether this is related to different immunological characteristics associated with DS.

**Methods:** The study group consisted of 22 children with DS and 22 of their healthy, age-range matched siblings. Data were collected from children who had experienced infections and hospitalisation due to lower RTIs. Immunoglobulin and IgG subclass levels in blood, as well as lymphocyte and T cell (subset) counts, were determined.

**Results:** The children with DS had a significantly higher frequency of lower RTIs and related hospitalisation than their siblings. We also found significantly reduced IgG2 levels as well as significantly lower counts of total lymphocytes, CD4<sup>+</sup> T lymphocytes, CD4<sup>+</sup> invariant natural killer (iNKT) cells, and regulatory T cells in the DS group.

**Conclusion:** In children with DS, reduced levels of IgG2, total lymphocytes, T lymphocytes, iNKT cells, and regulatory T cells might contribute to their higher susceptibility to lower RTIs.

**Key notes:** The study shows that children with Down syndrome experience more lower RTIs and related hospitalisation than their healthy siblings. They show significantly diminished subpopulations of T cells, including invariant natural killer T cells and regulatory T cells. The diminished subpopulations of T cells might contribute to their higher susceptibility.

## Introduction

The prevalence of Down syndrome (DS), or trisomy 21, in the Netherlands is 16 per 10,000 live births.<sup>1</sup> In addition to mental retardation, children with DS have a higher incidence of physical disorders, such as congenital heart disease (CHD), gastrointestinal malformations, hypothyroidism, and leukaemia, and are also more likely to develop autoimmune disease than children without this condition. Children with DS are also prone to infections, especially respiratory tract infections (RTIs).<sup>2,3</sup> This higher susceptibility can be due to anatomical,<sup>4,5</sup> neurological<sup>6</sup> or immunological abnormalities as concluded in many reviews.<sup>7,8</sup>

To date, most immunological studies of children with DS have reported defects in both the innate and adaptive immune systems, such as a deficiency of mannan-binding lectin,<sup>9</sup> a high number of proinflammatory CD14<sup>dim</sup>CD16<sup>+</sup> monocytes,<sup>10</sup> changes in T and B lymphocyte counts,<sup>8,11,12</sup> an intrinsic defect of T and B lymphocytes,<sup>8,13,14</sup> and deficiencies in the IgG2 and IgG4 subclasses.<sup>8,14,15</sup> However, in one study, DS children were found to have normal immunoglobulin levels and normal lymphocyte counts (total lymphocytes, CD4<sup>+</sup> T cells, CD8<sup>+</sup> T cells, and natural killer cells).<sup>16</sup> A possible reason for these conflicting results might be the fact that different types of control group were used in the studies. Different studies have used historical controls, healthy controls, children with mental retardation who do not have DS or no controls at all; thus, the control groups had different environmental and genetic backgrounds.<sup>11,15,16</sup>

The aim of the present study was to investigate the role of adaptive immunity in relation to lower RTIs in children with DS. To minimize genetic, environmental, and age-related differences, we used their healthy, age-ranged matched siblings as a control group. Given that 50% of children with DS have CHD<sup>17</sup> and CHD may also lead to an increase in RTIs,<sup>18</sup> we also analysed the presence of CHD in these children in relation to RTIs.

## Patients and methods

### Patient evaluation

Twenty-two children with DS and 22 of their healthy siblings were recruited from the DS outpatient clinic at the VU University Medical Center in Amsterdam, the Netherlands. Children with DS who met the following criteria were included: diagnosis of DS proven by chromosome analysis, age >3 months, and no infection present on the day the blood sample

was taken. Siblings of the children in the DS group were included if they met the following criteria: age >3 months, and no infection present on the day the blood sample was taken. Immunoglobulin levels and lymphocyte counts vary with age; thus, we chose age categories on the basis of the literature.<sup>19</sup> A DS-sibling pair was included in the study only when both the child with DS and the matched sibling were in the same age category.

To rule out the presence of an infection on the day of venipuncture, parents were interviewed about their child's recent health and a physical examination of the child was performed. The appointment was postponed if the interview, physical examination or laboratory results (including leukocyte count, leukocyte differentiation, and C-reactive protein) revealed an infection.

One of the investigators (C.J.M. Broers, paediatrician) conducted a structured interview with the parents about the total number of infections that their children had had since birth, and any related hospitalisation. To confirm these data, we asked the parents for written consent to obtain a copy of their children's medical data. From the medical records, the frequency and type of infection were retrieved. Lower RTIs were defined as: subglottic laryngitis, pneumonia, bronchitis, and bronchiolitis. The medical records of all the participants were checked for the frequency of infections, presence of CHD, any history of heart surgery and remaining heart defects, and in children with DS, the results of chromosome analysis.

## **Immunological evaluation**

Immunoglobulin and IgG subclass analysis was performed using immunonephelometry (IMMAGE 800 Immunochemistry System; Beckman Coulter, Fullerton, CA, USA) in a routine diagnostic clinical chemistry laboratory.

Peripheral blood lymphocytes were phenotyped by monoclonal antibody (mAb) staining of whole blood, and flow cytometric analysis after lysing red cells with Lysing Solution (BD Biosciences, San Jose, CA, USA) and fixing with 1% paraformaldehyde. Flow cytometric analysis was performed using a four-colour FACSCalibur flow cytometer (BD Biosciences).

Absolute numbers of T lymphocytes were determined by adding fixed volumes of FlowCount fluorospheres (Beckman Coulter) to the pretreated cell sample, just before flow cytometric evaluation. Lymphocyte gates in the scatter diagram were determined in each blood sample by CD14/CD45 mAb staining (BD Biosciences). T helper (Th) cells and cytotoxic T cells were characterized using the markers CD3, CD4, and CD8. CD25 on T cells was also evaluated (all mAb of BD Biosciences).

T regulatory cells (Tregs) were identified on the basis of coexpression of CD3 [fluorescein isothiocyanate (FITC)-labelled mAb], CD4 [phycoerythrin (PE)-labelled mAb], CD45 [peridinin chlorophyll protein complex (PerCP)-labelled mAb], and high expression of CD25 [allophycocyanin (APC)-labelled mAb] (all mAb of BD-Biosciences).

V $\alpha$ 24<sup>+</sup> V $\beta$ 11<sup>+</sup> invariant natural killer T (iNKT) cells were determined by flow cytometry using a FITC-labelled monoclonal antibody (mAb) against human V $\alpha$ 24, a PE-labelled mAb against human V $\beta$ 11 (Immunotech, Marseille, France), a PerCP-Cy5.5-labelled mAb against CD3, and an APC-labelled mAb against CD4 (BD Biosciences). V $\alpha$ 24<sup>+</sup> V $\beta$ 11<sup>+</sup> iNKT cells were evaluated as a fraction of CD3<sup>+</sup> lymphocytes in the blood. A minimum of 100,000 viable lymphocytes were acquired from each patient for the determination of iNKT cells.

### Data analysis

Fisher's exact test (SPSS version 15.0) was used to analyse the response categories in the questionnaires. The paired *t* test (SPSS version 15.0) was used to analyse the data for lymphocytes and immunoglobulins. A *p*-value <0.05 was considered significant.

### Ethical approval

The study was approved by the Ethics Committee of the VU University Medical Center in Amsterdam.

### Results

All 22 children with DS had trisomy 21 (no translocation or mosaic trisomy 21). The number of DS-sibling pairs in each age category was as follows: 5 DS-5 siblings in age category 2–5 years; 7 DS-7 siblings in age category 5–10 years; 6 DS-6 siblings in age category 10–16 years; and 4 DS-4 siblings in age category >16 years.

Children with DS had significantly more lower RTIs in comparison to their siblings, and they also were significantly more hospitalised because of infection (Table 5.1). Concerning RTIs, the level of agreement between the data obtained from the parental interview and from the medical records was 95%.

As seen in Figure 5.1 levels of IgM, IgG2, and IgG4 were significantly lower in the DS group than in the sibling group, whereas levels of IgG1 and IgG3 were significantly higher in the

**Table 5.1 The number of children with at least one infection and the number of children with hospitalization because of infection. (DS vs sibling group).**

	Down syndrome (n=22)	Sibling (n=22)	p-value
Patient characteristics			
Sex (% male)	15 /22 (68%)*	8 /22 (36%)	0.07
Age (in years)	10.0 ± 1.3**	10.4 ± 1.2	0.82
History of infections			
Lower RTI	13 /22 (59%)*	2 /22 (9%)	0.001
Urinary tract infection	2 /22 (9%)	2 /22 (9%)	1.00
Gastroenteritis	9 /22 (41%)	10 /22 (45%)	1.00
Hospitalization <sup>1</sup>	11 /22 (50%)	2 /22 (9%)	0.007
Hospitalization <sup>2</sup>	7 /22 (32%)	0 /22 (0%)	0.009

\* number (%); \*\* mean (SEM); <sup>1</sup> because of infection; <sup>2</sup> because of lower RTI.


DS group. There was no significant difference in total IgG and IgA levels between the two groups. In the sibling group, all immunoglobulin levels were within the normal range.

As seen in Figure 5.2 the numbers of total lymphocytes, CD4<sup>+</sup> T lymphocytes, CD4<sup>+</sup> iNKT cells, and CD4<sup>+</sup> CD25<sup>high</sup> T cells (Tregs) were all significantly lower in the DS group than in the sibling group. In the sibling group, all counts for total lymphocytes, T lymphocytes, CD4<sup>+</sup> T lymphocytes, and CD8<sup>+</sup> T lymphocytes were within the normal range. No normal values were available for CD4<sup>+</sup> iNKT cells or Tregs. The CD4/CD8 T cell ratio was significantly lower in the DS group than in the sibling group (data not shown).

In the DS group, 10 out of 22 children had CHD and cardiac surgery for CHD was performed in 5 patients. At the time of our study 4 of the 10 patients with CHD had a remaining minor heart defect, either post-surgery or because of an incomplete spontaneous closure. No significant difference was found between the number of hospitalizations due to lower RTIs in the 10 DS children with CHD as compared with the 12 DS children without CHD. In addition, no significant difference was found between the 4 patients with a remaining heart defect and 6 patients without a remaining heart defect in terms of the number of hospitalizations related to lower RTIs (data not shown). None of the 22 siblings had CHD.


## Discussion

The results of the study demonstrated that the children with DS had a significantly higher frequency of lower RTIs and related hospitalization than their siblings. Immunological defects



**Figure 5.1** Immunoglobulin levels (mean) of DS children and siblings, including p-values. n.s. = not significant.





**Figure 5.2** T cell numbers (mean) of Down syndrome children and siblings, including p-values. n.s. = not significant.

associated with DS are reported frequently as a cause of RTI. To date, the interpretation of immunological studies in children with DS have been hampered by the fact that control groups have been lacking or have consisted of historical controls.<sup>11,16</sup> One strength of the present study is that we included healthy, age-range matched siblings as controls, thereby circumventing environmental, genetic, and age-related differences between the patient and control groups as much as possible.

For the most part, the immunoglobulin levels that were found in the children with DS in the present study confirmed previously reported results,<sup>15</sup> namely increased levels of IgG1 and IgG3 and reduced levels of IgM, IgG2, and IgG4. In general, reduced IgG2 levels can cause an increase in RTIs.<sup>20</sup>

The results of the results study clearly demonstrate that reduced numbers of lymphocytes and T cells are associated with DS, especially CD4<sup>+</sup> T cells, CD4<sup>+</sup> iNKT cells, and Tregs. iNKT cells constitute a subset of T lymphocytes that have a hybrid phenotype of NK cells and conventional T cells and can be both of thymic and extrathymic (e.g. bone marrow or blood) origin.<sup>21</sup> The majority of iNKT cells express CD4, but most of the remaining cells express neither CD4 nor CD8, although humans contain a small subset of CD8<sup>+</sup> iNKT cells.<sup>22,23</sup> The T cell receptors of iNKT cells recognise glycolipid antigens that are presented through CD1d molecules on antigen-presenting cells.<sup>21</sup> iNKT cells play an important role in the defence to bacteria and viruses and after activation CD4<sup>+</sup> iNKT cells can produce both Th1-type and Th-2 type cytokines.<sup>22,23</sup>

In the present study, we found that children with DS had more lower RTIs than their healthy siblings. It is well known that respiratory syncytial and influenza A viruses are important causes of severe viral disease in DS.<sup>2,24</sup> Enhanced production of pro-inflammatory cytokines, which leads to excessive inflammation, and a more severe clinical course may be an explanation for the increased severity of influenza A infection in DS children.<sup>25,26</sup> Also iNKT cells play an important role in the defence against influenza A virus by enhancing the maturation of dendritic cells, and the antigen-specific antibody response by B cells, and regulating CD8<sup>+</sup> T cell functions through the production of interferon- $\gamma$ .<sup>21</sup> Lower counts of iNKT cells in children with DS, as we and others found<sup>10</sup> might also contribute to the severity of influenza A infection.

Tregs are a naturally occurring subpopulation of CD4<sup>+</sup> CD25<sup>high</sup> T cells, which are produced by the thymus. Forkhead box protein 3 is a transcription factor that is expressed in CD4<sup>+</sup> Tregs and is required for Treg development and function. Tregs suppress host immune responses against self or nonself antigens, and thus play a crucial role in the prevention of autoimmune disease and in the modulation of immune responses to bacterial and viral pathogens. During infection, Tregs suppress the inflammatory response via the production of anti-inflammatory cytokines, such as transforming growth factor  $\beta$  and interleukin-10.<sup>27</sup> During infection with swine-origin H1N1 influenza virus in children, low percentages of Tregs are present in patients with complications, which suggests dysregulation of the host

immune response.<sup>28</sup> The low number of Tregs that we found in children with DS could also lead to a hyperinflammatory response and therefore increased severity of influenza A infection.

Another important cause of RTI in children with DS is the presence of CHD in up to 33%.<sup>3,29</sup> Children with DS and CHD were more often hospitalized for RTI and needed more often ventilatory support than children with DS but no CHD.<sup>3,30</sup> In our study the presence of CHD in children with DS did not enhance the frequency of lower RTI, but we have to take into account that the subgroups (CHD vs. non-CHD) are small and thus the subgroupanalysis might be underpowered.

The limitations of our study are that we only had access to a small, however useful, group of children with DS and their siblings and that our study design was retrospective as far as the history of infections was concerned.

In conclusion, children with DS have reduced levels of IgG2 and reduced numbers of lymphocytes, CD4<sup>+</sup> T cells, CD4<sup>+</sup> iNKT cells, and Tregs, as compared with their healthy siblings. This could contribute to an increased frequency of lower RTI and related hospitalization.

## **Acknowledgements**

We would like to thank Dr. M.M. van Weissenbruch of the VU University Medical Center, the Netherlands, for her critical revision of the manuscript.

## References

1. Weijerman ME, van Furth AM, Vonk Noordegraaf A, van Wouwe JP, Broers CJM, Gemke RJB. Prevalence, neonatal characteristics, and first-year mortality of Down syndrome: a national study. *J Pediatr* 2008;152:15-9.
2. Bloemers BLP, van Furth AM, Weijerman ME, Gemke RJB, Broers CJM, van den Ende K, et al. Down syndrome: a novel risk factor for respiratory syncytial virus bronchiolitis--a prospective birth-cohort study. *Pediatrics* 2007;120:e1076-e81.
3. Hilton JM, Fitzgerald DA, Cooper DM. Respiratory morbidity of hospitalized children with Trisomy 21. *J Paediatr Child Health* 1999;35:383-6.
4. Shott SR. Down syndrome: common otolaryngologic manifestations. *Am J Med Genet Part C Semin Med Genet* 2006;142C:131-40.
5. McDowell KM, Craven DI. Pulmonary complications of Down syndrome during childhood. *J Pediatr* 2011;158:319-25.
6. Frazier JB, Friedman B. Swallow function in children with Down syndrome: a retrospective study. *Dev Med Child Neurol* 1996;38:695-703.
7. Bloemers BLP, Broers CJM, Bont L, Weijerman ME, Gemke RJB, van Furth AM. Increased risk of respiratory tract infections in children with Down syndrome: the consequence of an altered immune system. *Microbes Infect* 2010;12:799-808.
8. Kusters MAA, Verstegen RHJ, Gemen EFA, de Vries E. Intrinsic defect of the immune system in children with Down syndrome: a review. *Clin Exp Immunol* 2009;156:189-93.
9. Nisihara RM, Utiyama SRR, Oliveira NP, Messias-Reason JJ. Mannan-binding lectin deficiency increases the risk of recurrent infections in children with Down's syndrome. *Hum Immunol* 2010;71:63-6.
10. Bloemers BLP, van Bleek GM, Kimpen JLL, Bont L. Distinct abnormalities in the innate immune system of children with Down syndrome. *J Pediatr* 2010;156:804-9.
11. de Hingh YCM, van der Vossen PW, Gemen EFA, Mulder AB, Hop WCJ, Brus F, et al. Intrinsic abnormalities of lymphocyte counts in children with down syndrome. *J Pediatr* 2005;147:744-7.
12. Bloemers BLP, Bont L, de Weger RA, Otto SA, Borghans JA, Tesselaar K. Decreased thymic output accounts for decreased naive T cell numbers in children with Down syndrome. *J Immunol* 2011;186: 4500-7.
13. Kusters MAA, Gemen EFA, Verstegen RHJ, Wever PC, de Vries E. Both normal memory counts and decreased naive cells favour intrinsic defect over early senescence of Down syndrome T lymphocytes. *Pediatr Res* 2010;67:557-62.
14. Verstegen RHJ, Kusters MAA, Gemen EFA, de Vries E. Down syndrome B-lymphocyte subpopulations, intrinsic defect or decreased T-lymphocyte help. *Pediatr Res* 2010;67:563-9.

15. Anneren G, Magnusson CG, Lilja G, Nordvall SL. Abnormal serum IgG subclass pattern in children with Down's syndrome. *Arch Dis Child* 1992;67:628-31.
16. Cocchi G, Mastrola M, Capelli M, Bastelli A, Vitali F, Corvaglia L. Immunological patterns in young children with Down syndrome: is there a temporal trend? *Acta Paediatr* 2007;96:1479-82.
17. Weijerman ME, de Winter JP. Clinical practice. The care of children with Down syndrome. *Eur J Paediatr* 2010;169:1445-52.
18. Kristensen K, Stensballe LG, Bjerre J, Roth D, Fisker N, Kongstad T, et al. Risk factors for respiratory syncytial virus hospitalisation in children with heart disease. *Arch Dis Child* 2009;94:785-9.
19. Comans-Bitter WM, de Groot R, van den Beemd R, Neijens HJ, Hop WCJ, Groeneveld K, et al. Immunophenotyping of blood lymphocytes in childhood. Reference values for lymphocyte subpopulations. *J Paediatr* 1997;130:388-93.
20. Ozkan H, Atlihan F, Genel F, Targan S, Gunvar T. IgA and/or IgG subclass deficiency in children with recurrent respiratory infections and its relationship with chronic pulmonary damage. *J Invest Allergol Clin Immunol* 2005;15:69-74.
21. Kulkarni RR, Haeryfar SM, Sharif S. The invariant NKT cell subset in anti-viral defenses: a dark horse in anti-influenza immunity? *J Leukoc Biol* 2010;88:635-43.
22. La Cava A, Van Kaer L, Fu-Dong-Shi. CD4+CD25+ Tregs and NKT cells: regulators regulating regulators. *Trends Immunol* 2006;27:322-7.
23. Tupin E, Kinjo Y, Kronenberg M. The unique role of natural killer T cells in the response to microorganisms. *Nat Rev Microbiol* 2007;5:405-17.
24. Pérez-Padilla R, Fernández R, García-Sancho C, Franco-Marina F, Aburto O, López-Gatell H, et al. Pandemic (H1N1) 2009 virus and Down syndrome patients. *Emerg Infect Dis* 2010;16:1312-4.
25. Broers CJM, Gemke RJJ, Weijerman ME, van der Sluijs KF, van Furth AM. Increased pro-inflammatory cytokine production in Down syndrome children upon stimulation with live influenza A virus. *J Clin Immunol* 2012;32:323-29.
26. Rouse BT, Sehrawat S. Immunity and immunopathology to viruses; what decides the outcome? *Nat Rev Immunol* 2010;10:514-26.
27. Vignali DAA, Collison LW, Workman CJ. How regulatory T cells work. *Nat Rev Immunol* 2008;8:523-32.
28. Frisullo G, Iorio R, Plantone D, Nociti V, Patanela AK, Marti A, et al. Involvement of type I immune responses in swine-origin H1N1 influenza virus infection. *Hum Immunol* 2011;72:632-5.
29. Ribeiro LMA, Jacob CMA, Pastorino AC, Kim CAE, Fomin ABF, Castro APBM. [Evaluation of factors associated with recurrent and/or severe infections in patients with Down's syndrome]. *J Paediatr (Rio J)* 2003;79:141-8.

30. Medrano C, Garcia-Guereta L, Grueso J, Insa B, Ballesteros F, Casaldaliga J, et al. Respiratory viral infection in congenital cardiac disease. Hospitalizations in young children in Spain during 2004 and 2005: the CIVIC Epidemiologic Study. *Cardiol Young* 2007;17:360-71.

