VU Research Portal

A review of biomechanical studies on stoop and squat lifting

van Dieen, J.H.; Hoozemans, M.J.M.; Toussaint, H.M.

published in

IEA 2000/ HFES 2000 Congress
2000

[Link to publication in VU Research Portal](#)

citation for published version (APA)

van Dieen, J. H., Hoozemans, M. J. M., & Toussaint, H. M. (2000). A review of biomechanical studies on stoop and squat lifting. In *IEA 2000/ HFES 2000 Congress* (pp. 643-646). HFES.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

A REVIEW OF BIOMECHANICAL STUDIES ON STOOP AND SQUAT LIFTING

Jaap H. van Dieën¹, Marco J.M. Hoozemans², Huub M. Toussaint¹

¹ Amsterdam Spine Unit, Institute for Fundamental and Clinical Human Movement Sciences, Faculty of Human Movement Sciences, 'Vrije Universiteit', Amsterdam, The Netherlands

² Coronel Institute for Occupational and Environmental Health, Academic Medical Center / University of Amsterdam, Amsterdam, The Netherlands

To assess the rationale of advocating the squat as opposed to the stoop technique, biomechanical studies comparing the two were reviewed. With the exception of some specific lifting tasks, net moments and compression forces were estimated to be equal or higher in squat lifting. Shear force and spinal bending moments appeared lower in squat lifting. Net moments and compression forces probably can cause injury, whereas the other load components remain below injury threshold. In conclusion, the literature does not provide support for advocating squat lifting.

INTRODUCTION

Recently, Volinn (1999) has argued that studies designed after randomised clinical trials are needed to prove the effectiveness of ergonomics interventions with respect to low back pain (LBP) prevention. Indeed an 'evidence-based' ergonomics practice seems overdue. However, given the high costs involved with this type of studies ample consideration of the risk factors addressed and preventive strategy implemented is necessary. Several recent epidemiological review studies conclude that lifting is the best documented risk factor for LBP (Burdorf & Sorock, 1997; Ferguson & Marras, 1997; Frank et al., 1996; Kuiper et al., in press). In line with this, it would seem appropriate to consider preventive strategies involving measures aimed at reducing back load associated with lifting tasks.

In practice, administrative controls such as training and instruction in particular with respect to lifting technique are widely used (Chavalinitikul, Nopteepkangwan, & Kanjanopas, 1995; Lagerstrom, Josephson, Pingel, Tjernstrom, & Hagberg, 1998; Nevalapuranen, 1996; Nygard, Merisalo, Arola, Manka, & Huhtala, 1998; Schenk, Doran, & Stachura, 1996; St-Vincent, Tellier, & Lortie, 1989; Videman et al., 1989). The most commonly advised lifting technique is the so-called squat technique or leg lift, in which the back remains as erect as possible and in which the knees are flexed (Garg & Moore, 1992). However, biomechanical studies have shown conflicting results on the effectiveness of this technique. The aim of the present review, therefore, was to evaluate the evidence that the lifting technique is an important determinant of the mechanical load during lifting. This review was limited to studies comparing symmetric stoop and squat lifting, as these are well defined and frequently studied techniques in manual materials handling.

METHODS

This review was based on a literature search in the several electronic databases. The references were screened on the basis of titles and abstracts. The literature retrieved was

supplemented with references from reviews with a somewhat broader scope (Hsiang, Brogmus, & Courtney, 1997; Kroemer, 1992; Yu, Roht, Wise, Kilian, & Weir, 1984) and studies cited in the previously retrieved papers.

It is unknown what structures are responsible for LBP, and it seems likely that different structures may be involved in different cases. Therefore, all mechanical loads likely to induce injury to structures in the low back will be considered, this includes loads on spinal structures (e.g. ligaments, intervertebral disc, vertebrae) and musculotendinous structures (e.g. muscle, musculotendinous junction, tendinous insertion). Muscular damage is most likely to occur when high forces are sustained or produced repeatedly. The mechanical load on the osteoligamentous spine during symmetric lifting consist of three components: compression, shear and bending, each of which according to in vitro studies has the potential to cause injury (Adams, Green, & Dolan, 1994a; Adams & Hutton, 1982; Brinckmann, Biggeman, & Hilweg, 1989; Cyron, Hutton, & Troup, 1976; Lamy, Bazergui, Kraus, & Farfan, 1975; Perey, 1957; Yingling & McGill, 1999). Unfortunately, none of the four variables of interest can be measured directly. Therefore, we considered indicators of these parameters, based on model calculations, including net moments, estimated muscle forces or muscle moments, estimated compression and shear forces, and predicted bending moments resisted by the osteoligamentous spine, or tensile forces in individual ligaments.

RESULTS

In total 25 studies comparing stoop and squat lifting with respect to the mechanical load on the back were included in this review (Anderson & Chaffin, 1986; Buseck, Schipplein, Andersson, & Andriacchi, 1988; Bush-Joseph, Schipplein, Andersson, & Andriacchi, 1988; Chaffin & Page, 1994; Dieën, Creemers, Draisma, Toussaint, & Kingma, 1994; Dolan, Earley, & Adams, 1994a; Dolan, Mannion, & Adams, 1994b; Ekholm, Arborelius, & Nemeth, 1982; Garg & Herrin, 1979; Hagen & Harmsringdahl, 1994; Kjellberg,

Lindbeck, & Hagberg, 1988; Leskinen, 1985; Leskinen, Stalhammar, Kuorinka, & Troup, 1983; Lindbeck & Arborelius, 1991; Looze, Dolan, Kingma, & Baten, 1998; Looze, Kingma, Thunnissen, Vanwijk, & Toussaint, 1994; Mittal & Malik, 1991; Park & Chaffin, 1974; Potvin, McGill, & Norman, 1991a; Potvin, Norman, Eckenrath, McGill, & Bennet, 1992; Toussaint, Baar, Langen, Looze, & Dieën, 1992; Toussaint, Commissaris, & Beek, 1997; Troup, Leskinen, Stålhammar, & Kuorinka, 1983; Wax, Flenghi, & Meyer, 1987). In several studies the comparison of the techniques was confounded with the horizontal distance (Ekholm et al., 1982; Garg & Herrin, 1979; Mittal & Malik, 1991; Wax et al., 1987), the mass lifted (Mittal & Malik, 1991), or the velocity of lifting (Buseck et al., 1988; Dieën et al., 1994).

The studies using a static linked segment model (LSM) to estimate the net moment yielded varying results. The majority of studies predicted a substantial reduction (10 to 34%) in back load when using the squat technique in at least one of the experimental conditions. However, two of these studies (Ekholm et al., 1982; Garg & Herrin, 1979), showed that when horizontal distance is constant the effect of technique disappears. The thirteen studies in which the net moment or extensor moment was estimated using dynamic analysis techniques found back loads in the two techniques to be either significantly higher (4 to 18%) in squat lifting or not significantly different. In eleven of these studies, the loads were lifted from a position in front of the feet. In three studies the horizontal position of the load was not described, nor could it be derived from any of the figures. Two studies did not report whether the differences found were significant. One of these did report a substantially lower (13%) net moment in squat lifting, but actually compared to a free technique, which was performed at a substantially higher velocity.

Overall, the studies providing compression force estimates yielded results in line with those of the moment estimates described above (Anderson & Chaffin, 1986; Chaffin & Page, 1994; Leskinen, 1985; Leskinen et al., 1983; Potvin et al., 1991a; Potvin et al., 1992; Troup et al., 1983). Most of these studies used fairly crude anatomical models of the lumbar spinal musculature or were based on predicted or statically analysed kinematics. The two studies (Potvin et al., 1991a; Potvin et al., 1992) using a detailed anatomical model and dynamically calculated net moments indicated higher compression in squat lifting.

Shear forces reported in one study only (Potvin et al., 1991a), were higher in stoop lifting than in squat lifting, as were bending moments (and ligament stresses) (Anderson & Chaffin, 1986; Dolan et al., 1994a; Dolan et al., 1994b).

DISCUSSION

Positive effects of squat lifting with respect to estimated moments and compression forces were found only when squat lifting allowed for lifting from a position between the feet. In practice, lifting from a position between the feet is often not possible and these results thus appear to be valid

for a limited range of tasks. Actually in a study by Dieën et al. (1994), subjects lifting a barbell preferred a larger horizontal distance when using a squat technique as compared to a stoop technique. In addition, all studies showing this benefit of squat lifting used a static LSM. Hence the validity of the positive findings on squat lifting in these cases may be questioned. A striking finding in this respect is the fact that the positive effect of squat lifting in one study disappeared, when reanalysing the same data using a dynamic LSM (Lindbeck & Arborelius, 1991).

In lifting tasks where the load is not lifted from a position between the feet, the net moment and compression tended to be lower using the stoop technique. In contrast, in all studies reporting shear and bending moments, these were higher in stoop lifting. Consequently, the parameters of back load these indicators stand for need to be weighted with respect to each other. When a parameter increases with a change in lifting technique, but remains well below injury threshold, this increase can be considered of little importance. Thus the injury potential of the parameters of back load could be used to obtain such a weighting.

The net moments in lifting are within the range of the maximum isometric moments of healthy male subjects (Dieën, 1996; Dieën, Böke, Oosterhuis, & Toussaint, 1996; Dieën & Heijblom, 1996; Dieën, Oude Vrielink, Housheer, Lötters, & Toussaint, 1993) and probably above the strength of inactive males or older populations (Chaffin & Andersson, 1991; Dieën & Heijblom, 1996; McNeill, Warwick, & Andersson, 1980). The incidence of back injuries appears to increase when lifting moments exceed the isometric strength (Chaffin & Park, 1973; Herrin, Jaraiedi, & Anderson, 1986).

Compression forces of 3 to 5 kN, as occur during lifting, are high enough to cause failure in females over 20 and in males over 40 years old (Jäger & Luttmann, 1997) and probably in younger males under repetitive loading (Brinckmann, Biggeman, & Hilweg, 1988; Hansson, Keller, & Spengler, 1987). In the context of the comparison of stoop and squat lifting, it is important to note that compression strength is not significantly affected by flexion of the motion segment (Adams, McNally, Chinn, & Dolan, 1994b). It has been hypothesised that a major proportion of all LBP cases is attributable to excessive compression during tasks such as lifting (Dieën, Weinans, & Toussaint, 1999), which is supported by recent epidemiological research (Granata & Marras, 1999). Shear forces reported in one study only, were below strength values (Cyron et al., 1976; Lamy et al., 1975), suggesting that shear forces during lifting do not pose a serious injury risk (Cyron et al., 1976). However, it should be kept in mind that in repetitive loading lower shear forces may cause damage (Cyron & Hutton, 1981) and shear forces estimates during lifting are strongly dependent on the functional and anatomical assumptions in the model used (Dieën & Looze, 1999; Potvin, Norman, & McGill, 1991b). The bending moments carried by the osteoligamentous spine during lifting generally remain well below the injury threshold (Adams & Hutton, 1986; Dolan et al., 1994a). In conclusion, the conclusion based on net moments and compression forces are definitely relevant,

whereas the relevance of shear remains to be shown and bending moments appear to be of little importance.

In conclusion, the present review shows that there is no substantial biomechanical evidence to support training and instruction in which the squat technique is advocated. Evidence obtained with other approaches such as psychophysics and exercise physiology, generally appears to support this conclusion (Duplessis et al., 1998; Garg & Herrin, 1979; Hagen, Hallen, & Harms-Ringdahl, 1993; Kumar, 1984; Welbergen, Kemper, Knibbe, Toussaint, & Clijssen, 1991). It is, therefore, suggested that further study of controls for preventing low back pain should be focussed on other aspects of lifting.

REFERENCES

- Adams, M. A., Green, T. P., & Dolan, P. (1994a). The strength in anterior bending of lumbar intervertebral discs. *Spine*, 19(19), 2197-2203.
- Adams, M. A., & Hutton, W. C. (1982). Prolapsed intervertebral disc: a hyperflexion injury. *Spine*, 7, 184-191.
- Adams, M. A., & Hutton, W. C. (1986). Has the lumbar spine a margin of safety in forward bending? *Clinical Biomechanics*, 1, 3-6.
- Adams, M. A., McNally, D. S., Chinn, H., & Dolan, P. (1994b). Posture and compressive strength of the lumbar spine. *Clinical Biomechanics*, 9, 5-14.
- Anderson, C. K., & Chaffin, D. B. (1986). A biomechanical evaluation of five lifting techniques. *Applied Ergonomics*, 17, 2-8.
- Brinckmann, P., Biggemann, M., & Hilweg, D. (1988). Fatigue fracture of human lumbar vertebrae. *Clinical Biomechanics, Supplement 1*, 1-27.
- Brinckmann, P., Biggemann, M., & Hilweg, D. (1989). Prediction of the compressive strength of human lumbar vertebrae. *Clinical Biomechanics, Supplement 12*, 1-27.
- Burdorf, A., & Sorock, G. (1997). Positive and negative evidence of risk factors for back disorders. *Scandinavian Journal of Work, Environment and Health*, 23, 243-256.
- Buseck, M., Schipplein, O., Andersson, G. B. J., & Andriacchi, T. P. (1988). Influence of dynamic factors and external loads on the moment at the lumbar spine in lifting. *Spine*, 13, 918-921.
- Bush-Joseph, C., Schipplein, O., Andersson, G. B. J., & Andriacchi, T. P. (1988). Influence of dynamic factors on the lumbar spine moment in lifting. *Ergonomics*, 31(2), 211-216.
- Chaffin, D. B., & Andersson, G. B. J. (1991). *Occupational Biomechanics*. (2nd ed.). New York: Wiley.
- Chaffin, D. B., & Page, G. B. (1994). Postural effects on biomechanical and psychophysical weight-lifting limits. *Ergonomics*, 37(4), 663-676.
- Chaffin, D. B., & Park, K. S. (1973). A longitudinal study of low back pain as associated with occupational weight lifting factors. *American Industrial Hygiene Association Journal*, 34, 513-525.
- Chavalititkul, C., Nopteepkangwan, N., & Kanjanopas, F. (1995). Improvement of lifting heavy objects work. *Journal of Human Ergology*, 24, 55-58.
- Cyron, B. M., & Hutton, W. C. (1981). The fatigue strength of the lumbar neural arch in spondylosis. *Journal of Bone and Joint Surgery - British Volume*, 60B, 234-238.
- Cyron, B. M., Hutton, W. C., & Troup, J. D. G. (1976). Spondylytic fractures. *Journal of Bone and Joint Surgery - British Volume*, 58B, 462-466.
- Dieën, J. H. v. (1996). Asymmetry of erector spinae muscle-activity in twisted postures and consistency of muscle activation patterns across subjects. *Spine*, 21(22), 2651-2661.
- Dieën, J. H. v., Böke, B., Oosterhuis, W., & Toussaint, H. M. (1996). The influence of torque and velocity on erector spinae muscle fatigue and its relationship to changes of electromyogram spectrum density. *European Journal of Applied Physiology*, 72, 310-315.
- Dieën, J. H. v., Creemers, M., Draisma, I., Toussaint, H. M., & Kingma, I. (1994). Repetitive lifting and spinal shrinkage, effects of age and lifting technique. *Clinical Biomechanics*, 9, 367-374.
- Dieën, J. H. v., & Heijblom, P. (1996). Reproducibility of isometric trunk extension torque, trunk extensor endurance, and related electromyographic parameters in the context of their clinical applicability. *Journal of Orthopaedic Research*, 14, 139-143.
- Dieën, J. H. v., & Looze, M. P. d. (1999). Sensitivity of single-equivalent trunk extensor muscle models to anatomical and functional assumptions. *Journal of Biomechanics*, 32(2), 195-198.
- Dieën, J. H. v., Oude Vrielink, H. H. E., Housheer, F. A. F., Lötters, F. J. B., & Toussaint, H. M. (1993). Trunk extensor endurance and its relationship to electromyogram parameters. *European Journal of Applied Physiology*, 66, 388-396.
- Dieën, J. H. v., Weinans, H., & Toussaint, H. M. (1999). Fractures of the lumbar vertebral endplate in the etiology of low back pain. A hypothesis on the causative role of spinal compression in a-specific low back pain. *Medical Hypotheses*, 53(3), 246-252.
- Dolan, P., Earley, M., & Adams, M. A. (1994a). Bending and compressive stresses acting on the lumbar spine during lifting activities. *Journal of Biomechanics*, 27(10), 1237.
- Dolan, P., Mannion, A. F., & Adams, M. A. (1994b). Passive tissues help the back muscles to generate extensor moments during lifting. *Journal of Biomechanics*, 27(8), 1077-1085.
- Duplessis, D. H., Greenway, E. H., Keene, K. L., Lee, I. E., Clayton, R. L., Metzler, T., & Underwood, F. B. (1998). Effect of Semi-Rigid Lumbosacral Orthosis Use On Oxygen Consumption During Repetitive Stoop and Squat Lifting. *Ergonomics*, 41(6), 790-797.
- Ekholm, J., Arborelius, U. P., & Nemeth, G. (1982). The load on the lumbo-sacral joint and trunk muscle activity during lifting. *Ergonomics*, 25(2), 145-161.
- Ferguson, S. A., & Marras, W. S. (1997). A literature review of low back disorder surveillance measures and risk factors. *Clinical Biomechanics*, 12(4), 211-226.
- Frank, J. W., Kerr, M. S., Brooker, A. S., DeMaio, S. E., Maetzel, A., Shannon, H. S., Sullivan, T. J., Norman, R. W., & Wells, R. P. (1996). Disability resulting from low back pain. Part I: What do we know about primary prevention? A review of the scientific evidence on prevention before disability begins. *Spine*, 21(24), 2908-2917.
- Garg, A., & Herrin, G. D. (1979). Stoop or squat, a biomechanical and metabolic evaluation. *A. I. E. Transactions*, 11, 293-302.
- Garg, A., & Moore, J. S. (1992). Prevention strategies and the low back in industry. *Occupational Medicine*, 7(4), 629-640.
- Granata, K. P., & Marras, W. S. (1999). Relation between spinal load factors and the high-risk probability of occupational low-back disorders. *Ergonomics*, 42(9), 1187-1199.
- Hagen, K. B., Hallen, J., & Harms-Ringdahl, K. (1993). Physiological and subjective responses to maximal repetitive lifting employing stoop and squat technique. *European Journal of Applied Physiology*, 67, 291-297.
- Hagen, K. B., & Harmsringdahl, K. (1994). Ratings of perceived thigh and back exertion in forest workers during repetitive lifting using squat and stoop techniques. *Spine*, 19(22), 2511-2517.
- Hansson, T. H., Keller, T. S., & Spengler, D. M. (1987). Mechanical behaviour of the human lumbar spine II. Fatigue strength during dynamic compressive loading. *Journal of Orthopaedic Research*, 5, 479-487.
- Herrin, G. D., Jaraiedi, M., & Anderson, C. K. (1986). Prediction of overexertion injuries using biomechanical and psychophysical

- models. *American Industrial Hygiene Association Journal*, 47, 322-330.
- Hsiang, S. M., Brogmus, G. E., & Courtney, T. K. (1997). Low-back-pain (lbp) and lifting technique - a review. *International Journal of Industrial Ergonomics*, 19(1), 59-74.
- Jäger, M., & Lüttmann, A. (1997). *Assessment of low-back load during manual materials handling*. Paper presented at the 13th Triennial Congress of the International Ergonomics Association, Tampere.
- Kjellberg, K., Lindbeck, L., & Hagberg, M. (1988). Method and performance: two elements of work technique. *Ergonomics*, 41(6), 798-816.
- Kroemer, K. H. E. (1992). Personnel training for safer manual handling. *Ergonomics*, 35(9), 1119-1134.
- Kuiper, J., Burdorf, A., Verbeek, J. H. A. M., Frings-Dresen, M. H. W., Beek, A. J. v. d., & Viikari-Juntura, E. R. A. (in press). Epidemiologic evidence on manual materials handling as a risk factor for back disorders: A systematic review. *International Journal of Industrial Ergonomics*.
- Kumar, S. (1984). The physiological cost of three different methods of lifting in sagittal and lateral planes. *Ergonomics*, 27(4), 425-433.
- Lagerstrom, M., Josephson, M., Pingel, B., Tjernstrom, G., & Hagberg, M. (1998). Evaluation of the implementation of an education and training programme for nursing personnel at a hospital in Sweden. *International Journal of Industrial Ergonomics*, 21(1), 79-90.
- Lamy, C., Bazergui, A., Kraus, H., & Farfan, H. F. (1975). The strength of the neural arch and the etiology of spondylolysis. *Orthopaedic Clinics of North America*, 6, 215-231.
- Leskinen, T. P. J. (1985). Comparison of static and dynamic biomechanical models. *Ergonomics*, 28, 285-291.
- Leskinen, T. P. J., Stalhammar, H. R., Kuorinka, I. A. A., & Troup, J. G. D. (1983). A dynamic analysis of spinal compression with different lifting techniques. *Ergonomics*, 26(6), 595-604.
- Lindbeck, L., & Arborelius, U. P. (1991). Inertial effects from single body segments in dynamic analysis of lifting. *Ergonomics*, 34, 421-433.
- Looze, M. P. d., Dolan, P., Kingma, I., & Baten, C. T. M. (1998). Does an asymmetric straddle-legged lifting movement reduce the low-back load? *Human Movement Science*, 17, 243-259.
- Looze, M. P. d., Kingma, I., Thunnissen, W., Vanwijk, M. J., & Toussaint, H. M. (1994). The evaluation of a practical biomechanical model estimating lumbar moments in occupational activities. *Ergonomics*, 37(9), 1495-1502.
- McNeill, T., Warwick, D., & Andersson, G. (1980). Trunk strengths in attempted flexion, extension, and lateral bending in healthy subjects and patients with low-back disorders. *Spine*, 5, 529-538.
- Mittal, M., & Malik, S. L. (1991). Biomechanical evaluation of lift postures in adult Koli female workers. *Ergonomics*, 34(1), 103-108.
- Nevalapuranen, N. (1996). Effects of occupationally-oriented rehabilitation on farmers work techniques, musculoskeletal symptoms, and work ability. *Journal of Occupational Rehabilitation*, 6(3), 191-200.
- Nygard, C. H., Merisalo, T., Arola, H., Manka, M. L., & Huhtala, H. (1998). Effects of work changes and training in lifting technique on physical strain - a pilot study among female workers of different ages. *International Journal of Industrial Ergonomics*, 21(1), 91-98.
- Park, K. S., & Chaffin, D. B. (1974). A biomechanical evaluation of two methods of manual load lifting. *A. I. E. Transactions*, 6, 105-113.
- Perey, O. (1957). Fracture of the vertebral endplate in the human spine. *Acta Orthopædica Scandinavica, Suppl.* 25.
- Potvin, J. R., McGill, S. M., & Norman, R. W. (1991a). Trunk muscle and lumbar ligament contributions to dynamic lifts with varying degrees of trunk flexion. *Spine*, 16, 1099-1107.
- Potvin, J. R., Norman, R. W., Eckenrath, M. E., McGill, S. M., & Bennet, G. M. (1992). Regression models for the prediction of dynamic L4/L5 compression forces during lifting. *Ergonomics*, 35, 187-201.
- Potvin, J. R., Norman, R. W., & McGill, S. M. (1991b). Reduction in anterior shear forces on the L4/L5 disc by the lumbar musculature. *Clinical Biomechanics*, 6, 88-96.
- Schenk, R. J., Doran, R. L., & Stachura, J. J. (1996). Learning effects of a back education program. *Spine*, 21(19), 2183-2188.
- St-Vincent, M., Tellier, C., & Lortie, M. (1989). Training in handling: an evaluative study. *Ergonomics*, 32(2), 191-210.
- Toussaint, H. M., Baar, C. E. v., Langen, P. P. v., Looze, M. P. d., & Dieën, J. H. v. (1992). Coordination of the leg muscles in back-lift and leglift. *Journal of Biomechanics*, 25, 1279-1289.
- Toussaint, H. M., Commissaris, D. A. C. M., & Beek, P. J. (1997). Anticipatory postural adjustments in the back and leg lift. *Medicine and Science in Sports and Exercise*, 29(9), 1216-1224.
- Troup, J. D. G., Leskinen, T. P. J., Stålhammar, H. R., & Kuorinka, I. A. A. (1983). A comparison of intraabdominal pressure increases, hip torque, and lumbar intervertebral compression in different lifting techniques. *Human Factors*, 25(5), 517-525.
- Videman, T., Rauhala, H., Asp, S., Lindstrom, K., Cedercrutz, G., Kamppi, M., Tola, S., & Troup, J. D. G. (1989). Patient-handling skill, back injuries, and back pain: An intervention study in nursing. *Spine*, 14, 148-156.
- Volinn, E. (1999). Do workplace interventions prevent low-back disorders? If so, why?: a methodologic commentary. *Ergonomics*, 42(1), 258-272.
- Wax, C., Flenghi, D., & Meyer, J. (1987). Comparison of two load lifting techniques. Biomechanical analysis and physiological costs. *Travail Humain*, 50(4), 334-345.
- Welbergen, E., Kemper, H. C. G., Knibbe, J. J., Toussaint, H. M., & Clijssen, L. (1991). Efficiency and effectiveness of stoop and squat lifting at different techniques. *Ergonomics*, 34, 613-624.
- Yingling, V. R., & McGill, S. M. (1999). Anterior shear of spinal motion segments - Kinematics, kinetics, and resultant injuries observed in a porcine model. *Spine*, 24(18), 1882-1889.
- Yu, T. S., Roht, L. H., Wise, R. A., Kilian, D. J., & Weir, F. W. (1984). Low-back pain in industry. An old problem revisited. *Journal of Occupational Medicine*, 26(7), 517-524.