VU Research Portal

Analysis of motor control in patients with low back pain

Van Dieën, Jaap H.; Peter Reeves, N.; Kawchuk, Greg; Van Dillen, Linda R.; Hodges, Paul W.

published in

Journal of Orthopaedic and Sports Physical Therapy
2019

DOI (link to publisher)

[10.2519/jospt.2019.7916](https://doi.org/10.2519/jospt.2019.7916)

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Van Dieën, J. H., Peter Reeves, N., Kawchuk, G., Van Dillen, L. R., & Hodges, P. W. (2019). Analysis of motor control in patients with low back pain: A key to personalized care? *Journal of Orthopaedic and Sports Physical Therapy*, 49(6), 380-388. <https://doi.org/10.2519/jospt.2019.7916>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

1 **Analysis of motor control in low-back pain patients, a key to personalized care?**

2 Jaap H. van Dieën, PhD

3 Department of Human Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam Movement
4 Sciences, Amsterdam, The Netherlands

5
6 N. Peter Reeves, PhD

7 Michigan State University Center for Orthopedic Research, Michigan State University, Lansing,
8 Michigan, USA

9 Department of Osteopathic Surgical Specialties, Michigan State University, East Lansing, Michigan,
10 USA

11
12 Greg Kawchuk, PhD

13 Faculty of Rehabilitation Medicine, Department of Physical Therapy, University of Alberta, Edmonton,
14 Canada

15
16 Linda van Dillen, PT, PhD

17 Program in Physical Therapy and Department of Orthopaedic Surgery, Washington University in St.
18 Louis School of Medicine, St. Louis, MO, USA

19
20 Paul W. Hodges, PT, PhD

21 The University of Queensland, Centre for Clinical Research Excellence in Spinal Pain, Injury and
22 Health, School of Health & Rehabilitation Sciences, Brisbane Queensland, 4072, Australia

23
24
25 **Correspondence**

26 prof.dr. Jaap H. van Dieën

27 Department of Human Movement Sciences

28 VU University Amsterdam

29 van der Boechorststraat 9

30 NL-1081 BT Amsterdam

31 Netherlands

32 t: (31) 20 5988501, f: (31) 20 5988529, e: j.van.dieen@vu.nl

33
34 **conflict of interest:** The authors have no conflict of interest to report.

35
36 This paper reports a literature review, not supported by specific funding. Ethical approval does not
37 apply.

38

39

40 **Synopsis**

41 Motor control exercise has been shown to be effective in the management of low-back
42 pain (LBP), but effect sizes are modest, possibly due to the fact that studies have used
43 a one-size-fits-all approach, whereas literature suggests that patients may differ in
44 presence or type of motor control issues. In this commentary, we address the question
45 whether consideration of such variation in motor control issues might contribute to
46 more personalized motor control exercise for patients with LBP. Such an approach is
47 plausible, since motor control changes may play a role in persistence of pain through
48 effects on tissue loading that may cause nociceptive afference in particular in case of
49 peripheral sensitization. Subgrouping systems used in clinical practice which comprise
50 motor control aspects allow reliable classification that is in part aligned with findings
51 in studies on motor control in patients with LBP. Motor control issues may have
52 heuristic value for treatment allocation, as the different presentations observed suggest
53 different targets for motor control exercise, but this remains to be proven. Finally,
54 clinical assessment of patients with LBP should take into account more aspects than
55 motor control alone, including pain mechanisms, musculoskeletal health and
56 psychosocial factors, and may need to be embedded in a stratification approach based
57 on prognosis to avoid undue diagnostic procedures.

58

59 **Keywords:** back pain, postural control, exercise, diagnostics, subgrouping

60 In the treatment of low back pain (LBP), exercise that targets motor control is
61 commonly used and with some success.^{10, 49, 75} Motor control can be defined as the way
62 in which the nervous system controls posture and movement to perform a given motor
63 task and includes consideration of all the associated motor, sensory and integrative
64 processes. Here we use the term “motor control exercise” (MCE) to refer to exercise
65 that aims to change the manner in which a person controls their body (including
66 posture/alignment, movement, muscle activation) to modify loading of the spine and
67 adjacent structures.

68

69 The effectiveness of MCE has been the subject of several systematic reviews that have
70 undertaken different comparisons.^{10,49,75} A consistent outcome is that MCE is better than
71 minimal intervention in reducing pain in the short-, intermediate- and long-term, and in
72 reducing disability at long-term follow-up.⁴⁹ The pooled effect size was ~14% for pain
73 and ~11% for disability when compared to minimal intervention.⁴⁹ Effects were better
74 than for many other interventions, although they were still modest and only better than
75 other exercise interventions in the short-term.⁴⁹ Recent systematic reviews provide
76 contrasting evidence for comparison of effects of MCE and general exercise on
77 disability: one reported better outcomes for MCE;¹⁰ the other concluded there is low to
78 high quality evidence that MCE is not clinically more effective than other exercises.⁷⁵

79 Of note, most large clinical trials with modest effects investigated application of MCE
80 in a standardised manner to a heterogeneous group of patients with non-specific LBP.

81 This contrasts the prevailing clinical view that treatment effects may be larger if
82 treatments are targeted to the right patients, at the right time, and in a tailored,
83 individualized manner. This has been the topic of considerable research and clinical
84 attention.

It has been suggested that specific patient characteristics may predict who will or will not benefit from MCE,⁴⁸ or guide how it should be tailored to the individual patient. As reviewed by van Dieën et al.,⁹⁴ laboratory studies of motor control in individuals with LBP and healthy subjects demonstrate high variability between studies,^{e.g. 52, 95} and between individuals with LBP within studies.^{e.g. 16, 72} This concurs with the proposal that tailored rehabilitation programs are likely to be required to address the specific changes in motor control that are unique for the individual.

This commentary aims to address the overall question whether features of motor control could form an important element of a subgrouping scheme. Individualisation of MCE could involve identification of subgroups of patients with similar motor control issues or similar response to treatment, or individualising treatment to match each individual patient's presenting characteristics. A further aim is to highlight the research and development that is needed to address the major issues of subgrouping, particularly related to motor control, for application in clinical practice.

2 Subgrouping of patients with LBP

Based on diversity in presentation among individuals with LBP, it has been argued that no single treatment is likely to be effective for all patients and various authors have emphasized the need to administer more personalized treatment.^{6, 7, 27, 97} Subgrouping of patients is generally considered to be a step towards personalization, and LBP is seen as a disorder for which subgrouping may be particularly useful in view of the large and heterogeneous patient population, the large variation in treatment outcomes, and the variety of available treatment options with varying costs and risks. Among clinicians it is generally believed that LBP includes many different conditions.²⁷ Consensus on the best way to subgroup patients or to personalize treatment is, however,

110 lacking^{38, 97} and there is no strong evidence yet for effectiveness of subgroup-based
111 treatment.^{5, 24, 33, 45, 54}

112 Towards resolution of the issues addressed above, Foster et al.²⁶ proposed a set
113 of requirements for subgrouping in LBP. First, the subgrouping system should be
114 plausible; in other words, it should be compatible with current knowledge about
115 pathology of and risk factors for LBP. Second, subgrouping should be reliable; for
116 instance, repeated testing or testing by different clinicians should assign the same
117 patients to the same subgroups. Third, methods need to be simple enough to allow
118 application in clinical practice. The simplicity of a method must be balanced with
119 acceptability to patients and clinicians, and cost-effectiveness. Very sophisticated
120 diagnostic instruments can be useful if the outcomes allow more effective treatment at
121 a lower overall cost. Fourth, for clinical utility a subgrouping system should yield
122 mutually exclusive subgroups, meaning all cases, at one point in time, should fit into
123 only one subgroup and this subgroup membership should guide a unique treatment
124 choice. In the following sections, we review motor control subgrouping based on the
125 criteria proposed by Foster et al..²⁶

126

127 **3 Is subgrouping based on motor control plausible?**

128 For subgrouping based on motor control to be plausible, issues with motor
129 control would have to be relevant for the development or continuation of LBP and
130 relevant variation in motor control presentation would have to exist in the population
131 of individuals with LBP.

132 With respect to the first question, the nature of loads on the spine and adjacent
133 structures depends on the quality of motor control, in combination with anatomical
134 factors (e.g. muscle moment arms) and motor tasks that are performed. However,

135 whether loading of these structures is relevant with respect to development of LBP has
136 been heavily debated.^{3, 4, 42, 43, 53, 66, 83, 93} Recent systematic reviews and meta-analyses,
137 however, provide consistent evidence for a prospective association between some
138 activities and tasks that induce high mechanical loads on the back and LBP.^{11, 14, 30} In
139 addition, variables that quantify (cumulative) mechanical load on lumbar tissues, such
140 as lumbar moments and compression forces, are associated with LBP incidence or
141 prevalence.^{12, 13, 40, 51, 61} Another line of evidence for the plausibility of a causal relation
142 between mechanical loading and LBP stems from biomechanical studies in animal
143 models and on human cadaveric material. Such studies indicate that loads on spinal
144 tissues that occur in daily life can cause injury^{8, 81} and, even without injury, ongoing
145 mechanical stimulation of tissues can potentially activate nociceptors and initiate an
146 inflammatory response.⁴⁷ Although, it is difficult to confirm the presence of micro-
147 trauma let alone non-injurious noxious stimulation of tissues in the back in individuals
148 with LBP, a range of literature supports the plausibility of a causal relation between
149 mechanical loading and the development of LBP.⁹⁶ Finally, several mechanisms can
150 play a role in transition to chronic LBP, specifically non-healing of injured tissues,
151 ongoing nociceptive input, central sensitization and neuropathic pain development.
152 Mechanical loading of tissues would be relevant in relation to the first two of these. It
153 may both hamper and stimulate tissue healing, likely dependent on intensity and
154 frequency of loading and time after injury,^{23, 46, 82} and also in the absence of frank injury
155 it can promote ongoing nociceptive input, especially in the presence of peripheral
156 sensitization.^{19, 59, 103}

157 With respect to the question whether there is relevant variation in motor control
158 presentation among individuals with LBP, a recent review of the literature concluded
159 that the group with LBP may show overlap with or be at either extreme of the

160 distribution in motor control found in healthy participants.⁹⁴ The groups deviating from
161 normal motor control can be divided based on the mechanical consequences of the
162 changes in motor control. One pattern of change involves increased activation of trunk
163 muscles and may provide tight control over lumbar movements, but at the cost of higher
164 loads on muscles and on the spine.⁹¹ The opposite pattern, involves lower muscle
165 activation and might avoid high muscle forces and compressive loading, but with the
166 cost of a loose control over movement and a potential result of higher tensile strains of
167 tissues. In the following we will refer to these two ends of a spectrum as “tight” and
168 “loose” control. Clearly tight and loose control would have different mechanical
169 consequences that could both be relevant for development and continuation of LBP, but
170 they also suggest different targets for MCE.

171

172 **4 Is subgrouping based on motor control practically applicable and reliable?**

173 Studies on motor control in LBP, summarized in van Dieën et al.,⁹⁴ have used a
174 broad range of laboratory-based measurement techniques to characterize motor control.
175 In principle, these techniques could provide a basis for the development of clinical tests
176 to assess motor control to inform clinicians regarding subgrouping. However, generally
177 speaking application of these techniques involves substantial costs and requires specific
178 expertise that is not readily available. Therefore, the following considers the extent to
179 which subgrouping systems already applied in clinical practice take motor control
180 aspects into account and to what extent this results in reliable classification.

181 Several systems for subgrouping or profiling that are in common use clinically
182 incorporate motor control aspects in the assessment of patients with LBP. Those that
183 have been studied most extensively are, the “Treatment Based Classification” (TBC),
184 the “Multi-Dimensional Clinical” framework (MDC) (formerly named the “O’Sullivan

Classification”), and the “Movement System Impairment” classification (MSI). If these assessments capture the differences in motor control that have been identified in laboratory-based motor control measures, this would indicate that assessment of motor control issues based on clinically applicable tools can yield reliable outcomes.

4.1 Treatment Based Classification

The TBC system, originally proposed by Delitto et al.,¹⁸ and updated by Fritz et al.²⁸ and Alrwaily et al.¹ proposes four LBP subgroups, each named for the treatment to which the patient is most likely to respond; (1) manipulation, (2) stabilization, (3) specific exercise, and (4) traction. The inter-rater reliability of examiners (physical therapists who are familiar with the classification system) to classify patients is clinically acceptable.⁹⁷

With respect to the current understanding of motor control changes in LBP,⁹⁴ the criterion of hypomobility of the lumbar spine, as one of the criteria for allocation to the TBC *manipulation* subgroup, could be considered to align with a group of patients with LBP who present with tight motor control. Importantly, other criteria for subgroup allocation (e.g. time since symptom onset, age) cannot be considered specific to this motor control phenotype. Furthermore, it would seem plausible that the TBC *stabilization* subgroup could involve individuals who use loose motor control,⁹⁴ as this group are described to require restriction of excessive segmental motion. Consistent with this proposal, studies report that individuals classified into this group more often have excessive segmental rotations or translation on flexion/extension radiography than others,²⁹ more aberrant segmental lumbar movement on flexion/extension radiography,⁸⁴ poorer ability to contract the transversus abdominis muscle in isolation from other abdominal muscles,⁸⁵ and lower multifidus activation,³² which could all be considered to align the loose motor control phenotype.

210 4.2 *Multi-Dimensional Clinical framework*

211 The MDC framework has evolved from a subgrouping approach⁶² to a
212 multidimensional clinical profiling approach.⁶⁵ Within the MDC, motor responses are
213 described in three broad contexts: adaptive/protective motor responses to an acute tissue
214 injury and or underlying pathological process (i.e. “movement impairment”), motor
215 responses secondary to dominant central pain mechanisms, or maladaptive/provocative
216 motor responses that may contribute to the pain (i.e. “motor control impairment”).
217 These presentations may be associated with directional patterns of pain provocation
218 (flexion, extension, rotation, side bending) or multiple directions (multidirectional).⁶⁹
219 Reliability testing among trained physical therapists has shown good to excellent inter-
220 rater reliability in classification of patients.^{17, 99}

221 There is strong potential alignment between the MDC characterisation of motor
222 responses and the tight and loose motor control phenotypes of LBP. The movement
223 impairment presentation aligns well with motor control changes interpreted as tight
224 motor control. The MDC movement impairment is characterized by abnormally high
225 levels of muscle guarding and co-contraction of trunk muscles.⁶² Whether the
226 subdivision on the basis of the movement direction avoided by the individual aligns
227 with detailed assessment of motor control has not been tested.⁶⁹ The motor control
228 impairment presentation, which is described as demonstrating “an impairment or deficit
229 in the control of the symptomatic spinal segment in the primary direction of pain”, can
230 be hypothesized to overlap with the loose control end of the spectrum of motor control
231 changes. This applies in particular to the flexion presentation, who tend to adopt flexed
232 trunk postures, which provoke pain. These individuals gradually increase trunk flexion
233 over time when cycling,⁹ or when seated,^{16, 64} less accurately resume a “neutral” trunk
234 posture (perhaps caused by proprioceptive impairment^{60, 63}), may have lumbar

235 hypermobility in forward bending,⁴¹ and lower lumbar muscle activity in sitting.¹⁵ The
236 “passive extension” sub-group, who tend to hinge into extension with low trunk muscle
237 activity,⁶² may also align with a loose control group, while the “active extension”
238 subgroup, who tend to adopt extended trunk postures characterized by high muscle
239 activity,^{15, 16} appear more aligned to a tight control phenotype.

240 **4.3 Movement System Impairment classification**

241 The MSI classification system, developed and described by Sahrman,⁷³ has the
242 underlying assumption that people with LBP tend to move one or more lumbar joints
243 more readily than adjacent joints/segments (e.g. thoracic or hip joints). This is thought
244 to result from habitual movement patterns during daily activity, eventually leading to
245 excessive loading of tissues associated with the specific joint. Five LBP subgroups are
246 proposed, named for the specific direction(s) of lumbar movement considered to
247 contribute to the patient’s symptoms: flexion, extension, rotation, rotation with flexion,
248 and rotation with extension. Trained physical therapists can attain fair to excellent
249 reliability in MSI classification.⁹⁷

250 The MSI system describes motor impairments in LBP as a failure to constrain
251 movement of some lumbar joints in a specific direction. This concurs with the notion
252 of loose control, and the MSI system differentiates separate subgroups based on
253 movement direction in which the impairment is most apparent and linked to pain
254 provocation. Whether the direction inferred from MSI classification parallels direction-
255 specific differences in trunk mechanics or muscle activity requires clarification. Also,
256 it is unclear how a tight control subgroup might relate to the MSI classification.

257 **4.4 Do clinical tools allow reliable classification of motor control?**

258 Current subgrouping methods were not specifically developed to classify
259 patients based on motor control issues. Nevertheless, the fact that these methods

reliably arrive at subgroups that likely show partial overlap with those that might be found using the laboratory-based biomechanical and electromyography measurements used in motor control studies is promising. Objective measurements may add to consistency, validity and reliability of subgrouping and might have as additional benefit that they would permit consideration as a measure of treatment effects, if found responsive. In several of the classification systems, motor control is assessed in a direction specific manner. The relation between directional specificity of the clinical presentation and underlying changes in motor control and their effects require further study.

5. Is subgrouping based on motor control clinically useful?

Subgrouping based on motor control can be considered of clinical value if it has heuristic value, meaning, if assignment of a patient to a specific subgroup implies a specific treatment and if such targeted care is more effective than a one-size-fits all approach. Review of biomechanical, electromyography and modelling studies reveals a spectrum of changes in motor control in LBP with extremes of tight control and loose control.⁹⁴ Motor control changes at both ends of this spectrum have the potential to lead to suboptimal mechanical loading of the spine, but in different ways. This implies that modification of motor control has potential benefit with opposite treatment targets for the subgroups at either end. Loose control implies that enhancement of muscle activity is required, whereas tight control implies an emphasis on reduction of muscle activity.³⁶ It should be kept in mind that these interpretations are based on the assumption that these motor control patterns are maladaptive and clinical benefit will be derived from “correction” of the strategy. For each of the motor control measures that have been used in research, there is a subgroup of individuals with LBP who show ‘normal’ motor

control,⁹⁴ which suggests that this subgroup would *not* benefit from MCE. There is some evidence to support this hypothesis. Two clinical trials have shown less clinical improvement for individuals without evidence of a motor control deficit (poor control of transversus abdominis) at baseline.^{25, 87} On the other hand, baseline findings on trunk muscle control were not correlated to clinical improvements in two other studies.^{50, 102}

The question whether subgrouping based on motor control is useful can only be answered after appropriate clinical trials have been performed. To date there is mixed evidence whether interventions that target treatment based on motor control subgrouping achieve better outcomes than non-targeted treatments for LBP. Two RCTs with a focus on matching exercise to movement subgroups showed no benefit over general exercise in the long-term primary outcomes of pain and disability in chronic LBP.^{2, 45, 74} In contrast, several recent RCTs demonstrated superior long term outcomes with individualized MCE in people with chronic LBP, based on an integrated subgrouping approach, one included assistance of a wearable biofeedback device³⁹ and another used an individualized approach to targeting relevant cognitive, motor control and lifestyle factors in people with chronic LBP.⁹⁸ A missing link is whether the clinical effects in these trials were related to a change in motor control. The possibility that other factors mediated the positive outcomes remains to be excluded. Given the preceding discussions it can be concluded that an affirmative answer is plausible and hence subgrouping based on motor control would merit further research.

305

306 **6. Are subgroups based on motor control mutually exclusive?**

Mutual exclusivity of subgroups implies that an individual can only be allocated to a single subgroup and would only be expected to respond to the ascribed course of management. With the exception of the MDC, existing clinical approaches, described

310 above, force assessors to allocate patients to a single subgroup, making it difficult to
311 evaluate whether subgroups are mutually exclusive. Some differences in subgroup
312 allocation between testers (inter-tester variability) implies that overlap may exist.

313 The tight and loose control subgroups that are apparent in biomechanical and
314 electromyography studies would appear to be mutually exclusive, but with some
315 considerations. First, how the groups are separated is not yet clear. Literature indicates
316 that a group with “normal” control sits between those with tight and loose control. The
317 measures that would be considered to differentiate the groups and the cut-off scores
318 have not been established. Second, some patients may even present with elements of
319 both subgroups: an overall tight presentation may be combined with elements of low
320 stiffness in specific directions or of specific joints. For instance, increased activity of
321 some muscles with pain, causing an overall increase in trunk stiffness, may coincide
322 with reduced activity in other muscles.³⁵ While the overall change in muscle activity
323 would allow tight control over thorax movements, it might coincide with a reduced
324 control over segmental movements in a specific direction in view of the inhibition of
325 some muscles. Third, motor control patterns are somewhat context dependent. It cannot
326 be excluded that an individual may show ‘loose’ control in one situation, and show tight
327 control in another situation; for example, a more threatening task may elicit a
328 compensatory strategy with high levels of muscle activity regardless of strategy adopted
329 in a less threatening situation.⁹²

330 Subgrouping of patients with LBP purely on the basis of motor control assumes
331 that motor control and tissue loading is relevant for the underlying persistence of pain
332 in all patients, yet not all pain is the same. As highlighted earlier, pain can be broadly
333 considered to primarily involve nociceptive, neuropathic or central sensitization
334 mechanisms. In the presence of a primary nociceptive mechanism, loading of tissue is

335 likely to be relevant. The motor control adaptation may be adaptive and potentially
336 helpful or maladaptive and relevant for persistence. When the mechanism is
337 neuropathic, loading may be relevant with respect to loading of neural tissue.

338 In the presence of primarily central sensitisation pain, pain may persist despite
339 absence of ongoing nociceptive input from the tissue and treatment targeted to
340 optimisation of tissue loading through motor control training is unlikely to address the
341 underlying mechanism, but could aid recovery through exposure to healthy movement.
342 Consideration of pain mechanisms in a motor control subgrouping approach could take
343 two main paths. First, the approach may involve a hierarchical process where the first
344 step is to identify the primary pain mechanism. If a nociceptive (and perhaps
345 neuropathic) mechanism is identified, then the patient would be characterized
346 according to motor control presentation. If central pain mechanisms are identified then
347 an alternative course of management is planned to address the pain mechanism (pain
348 coping training, pain education, fear-deconditioning, etc), without primary
349 consideration of motor control. Second, the approach could also involve a parallel
350 process whereby all patients are assessed on the basis of pain mechanism and motor
351 control and a treatment package is developed that includes components of intervention
352 targeted to both domains, based on the presenting features. This latter model assumes
353 that pain mechanism and motor control phenotypes are not mutually exclusive and
354 some central sensitisation may be present in those with nociceptive/neuropathic pain
355 (which is highly probable) and some nociceptive input may contribute to maintenance
356 of pain state. In each case assessment of the dominant pain mechanism requires
357 attention. Several instruments have been proposed.^{67, 68, 76-80} These assessments require
358 further validation and development towards a clinical tool.

359 To be comprehensive, in addition to pain mechanism, the diagnostic system
 360 requires evaluation of patients across multiple biological, psychological and social
 361 dimensions. These would include features relevant to motor control such as patterns of
 362 pain provocation and relief,^{20-22, 62, 73} muscle atrophy and weakness,^{55, 56} proprioceptive
 363 impairment,^{63, 86} as well as differentiation of psychological features including pain
 364 beliefs and fear of pain or re-injury,^{57, 100} depression, catastrophising, self-efficacy, and
 365 social issues.⁷⁰ An important consideration is that domains are not independent. For
 366 instance, measures of motor control may reflect psychological factors such as fear of
 367 pain.^{31, 44, 58, 71, 88-90} Overlap of domains, particularly some of the sensory and motor
 368 domains may reflect redundancy and may allow simplification of diagnostic schemes.
 369 Furthermore, in many cases characterization of patients occurs along a continuous
 370 scale, not necessarily yielding exclusive subgroups.^{c.f. 67} In the parallel model, rather
 371 than fitting explicit subgroups, it may be more ideal to profile patients across these
 372 dimensions rather than fitting into explicit subgroups, allowing outcomes to be
 373 monitored with respect to each of the dimensions, in line with the MDC approach.⁶⁷
 374 Comprehensive profiling of patients or subgrouping may also benefit from
 375 being embedded in a system with stratification based on prognosis.^{c.f. 1} Prognostic
 376 stratification tools such as StartBack³⁴ are based on the belief that many LBP cases
 377 recover within several weeks irrespective of treatment,^{37, 101} and that more
 378 comprehensive management should be reserved for those with greater likelihood of
 379 poor outcome. These tools attempt to predict which patients belong to this group, to
 380 avoid unnecessary diagnostic procedures and over-treatment in the “low-risk” group.
 381 The StartBack tool specifically identifies greater psychological prognostic barriers for
 382 recovery in the “high-risk” group and recommends psychologically informed treatment.
 383 In the “moderate-risk” group, comprehensive treatment is recommended and our model

384 of patient characterisation across multiple domains including motor control (with or
385 without allocation to subgroups) is likely to be most relevant in this group.

386

387 7. *Potential role for objective tests of motor control in patient assessment*

388 Although clinical assessments can be used to reliably allocate patients to
389 subgroups, there may be additional benefit for interpretation of underlying mechanisms
390 and objectively and sensitively tracking recovery by objective measurements. Further
391 research is needed to verify that individuals can consistently be classified into motor
392 control-based categories based on a minimal battery of objective tests.

393 Motor control of the trunk comprises modulation of intrinsic stiffness through
394 tonic muscle activity, anticipatory control, and feedback control.⁹⁴ To characterize trunk
395 control in LBP it may be necessary to evaluate these different aspects with dedicated
396 tests. Given the emphasis on directional preferences or directional impairments in
397 current classification systems, objective testing should probably be multi-directional.
398 The potential existence of positive (adaptive) and negative (maladaptive) subcategories
399 of both tight and loose control requires further consideration. An additional
400 consideration is that adapted motor control may be context dependent; for example,
401 individuals with LBP may show more pronounced changes when they perceive the task
402 that they perform as threatening in terms of pain provocation or re-injury. These
403 considerations would suggest that a comprehensive set of tests and test conditions is
404 necessary to characterize motor control in LBP. This might cast some doubt on the
405 practical applicability of subgrouping based on objective measures of motor control.
406 As an alternative approach, assessment of trunk control in daily life could be considered
407 as an efficient way to obtain a large amount of ecologically valid information with
408 limited effort, although substantial work would be required to develop and test such an

409 analysis. Comprehensive testing may be shown to yield redundant information. If motor
410 control impairments in LBP can be sufficiently characterized based on a limited number
411 of tests, this would greatly simplify clinical implementation.

412

413

414 **8. Conclusions**

415 Targeting of treatment for the management of LBP based on motor control
416 presentation may be helpful. Although clinical trials provide evidence for some aspects
417 of the approach and motor control literature provides support for the plausibility, there
418 are major gaps remaining in the literature. Large RCTs are required to compare the
419 benefit of interventions that are matched to motor control presentation against
420 treatments that are not matched. Further insight might be gained from the establishment
421 of a minimal battery of objective tests that aid in the identification of the specific motor
422 control phenotypes. Approaches to allocate patients to subgroups to guide treatment or
423 alternatively to evaluate patients across a range of domains and measures should be
424 compared for their effectiveness. Both imply personalisation of care to the individual
425 patient, and both methods have positive and negative features.

426

427 **Acknowledgements**

428 The authors would like to thank dr. Peter O' Sullivan and dr. Julie Fritz for reviewing
429 a previous version of this paper.

430

431 References

- 432 1. Alrwaily M, Timko M, Schneider M, et al. Treatment-Based Classification
433 System for Low Back Pain: Revision and Update. *Phys Ther.*
434 2015;96:1057-1066.
- 435 2. Azevedo DC, Ferreira PH, Santos HO, Oliveira DR, de Souza JVL, Costa LOP.
436 Movement System Impairment-Based Classification Treatment Versus
437 General Exercises for Chronic Low Back Pain: Randomized Controlled
438 Trial. *Phys Ther.* 2018;98:28-39.
- 439 3. Bakker EW, Verhagen AP, van Trijffel E, Lucas C, Koes BW. Spinal
440 mechanical load as a risk factor for low back pain: a systematic review of
441 prospective cohort studies. *Spine.* 2009;34:E281-293.
- 442 4. Balague F, Mannion AF, Pellise F, Cedraschi C. Non-specific low back pain.
443 *Lancet.* 2012;379:482-491.
- 444 5. Beneciuk JM, Bishop MD, George SZ. Clinical prediction rules for physical
445 therapy interventions: a systematic review. *Phys Ther.* 2009;89:114-124.
- 446 6. Borkan JM, Cherkin DC. An agenda for primary care research on low back
447 pain. *Spine.* 1996;21:2880-2884.
- 448 7. Bouter LM, Pennick V, Bombardier C, Editorial Board of the Back Review
449 G. Cochrane back review group. *Spine.* 2003;28:1215-1218.
- 450 8. Brinckmann P, Biggemann M, Hilweg D. Fatigue fracture of human lumbar
451 vertebrae. *Clin Biomech.* 1988;3:s1-s28.
- 452 9. Burnett AF, Cornelius MW, Dankaerts W, O'Sullivan P B. Spinal kinematics
453 and trunk muscle activity in cyclists: a comparison between healthy
454 controls and non-specific chronic low back pain subjects-a pilot
455 investigation. *Man Ther.* 2004;9:211-219.
- 456 10. Bystrom MG, Rasmussen-Barr E, Grooten WJ. Motor control exercises
457 reduces pain and disability in chronic and recurrent low back pain: a
458 meta-analysis. *Spine.* 2013;38:E350-358.
- 459 11. Coenen P, Gouttebauge V, van der Burght ASAM, et al. The effect of lifting
460 during work on low-back pain – a health impact assessment based on a
461 meta-analysis. *Occup Environ Med.* 2014;71:871-877.
- 462 12. Coenen P, Kingma I, Boot CR, Bongers PM, van Dieen JH. Cumulative
463 mechanical low-back load at work is a determinant of low-back pain.
464 *Occup Environ Med.* 2014;71:332-337.
- 465 13. Coenen P, Kingma I, Boot CR, Twisk JW, Bongers PM, van Dieen JH.
466 Cumulative low back load at work as a risk factor of low back pain: a
467 prospective cohort study. *J Occup Rehabil.* 2013;23:11-18.
- 468 14. da Costa BR, Vieira ER. Risk factors for work-related musculoskeletal
469 disorders: A systematic review of recent longitudinal studies. *Am J Ind*
470 *Med.* 2010;53:285-323.
- 471 15. Dankaerts W, O'Sullivan P, Burnett A, Straker L. Altered patterns of
472 superficial trunk muscle activation during sitting in nonspecific chronic
473 low back pain patients: importance of subclassification. *Spine.*
474 2006;31:2017-2023.
- 475 16. Dankaerts W, O'Sullivan P, Burnett A, Straker L. Differences in sitting
476 postures are associated with nonspecific chronic low back pain disorders
477 when patients are subclassified. *Spine.* 2006;31:698-704.

17. Dankaerts W, O'Sullivan PB, Straker LM, Burnett AF, Skouen JS. The inter-examiner reliability of a classification method for non-specific chronic low back pain patients with motor control impairment. *Man Ther.* 2006;11:28-39.
18. Delitto A, Erhard RE, Bowling RW. A treatment-based classification approach to low back syndrome: identifying and staging patients for conservative treatment. *Phys Ther.* 1995;75:470-485; discussion 485-479.
19. Diers M, Koeppe C, Diesch E, et al. Central processing of acute muscle pain in chronic low back pain patients: An EEG mapping study. *J Clin Neurophysiol.* 2007;24:76-83.
20. Donelson R, Aprill C, Medcalf R, Grant W. A prospective study of centralization of lumbar and referred pain. A predictor of symptomatic discs and anular competence. *Spine.* 1997;22:1115-1122.
21. Donelson R, Grant W, Kamps C, Medcalf R. Pain response to sagittal end-range spinal motion. A prospective, randomized, multicentered trial. *Spine.* 1991;16:S206-212.
22. Donelson R, Silva G, Murphy K. Centralization phenomenon. Its usefulness in evaluating and treating referred pain. *Spine.* 1990;15:211-213.
23. Durgam S, Stewart M. Cellular and Molecular Factors Influencing Tendon Repair. *Tissue Eng Part B Rev.* 2017;23:307-317.
24. Fairbank J, Gwilym SE, France JC, et al. The role of classification of chronic low back pain. *Spine.* 2011;36:S19-42.
25. Ferreira PH, Ferreira ML, Maher CG, Refshauge K, Herbert RD, Hodges PW. Changes in recruitment of transversus abdominis correlate with disability in people with chronic low back pain. *Br J Sports Med.* 2010;44:1166-1172.
26. Foster NE, Hill JC, Hay EM. Subgrouping patients with low back pain in primary care: are we getting any better at it? *Man Ther.* 2011;16:3-8.
27. Foster NE, Hill JC, O'Sullivan P, Hancock M. Stratified models of care. *Best Pract Res Clin Rheumatol.* 2013;27:649-661.
28. Fritz JM, Cleland JA, Childs JD. Subgrouping patients with low back pain: evolution of a classification approach to physical therapy. *J Orthop Sports Phys Ther.* 2007;37:290-302.
29. Fritz JM, Piva SR, Childs JD. Accuracy of the clinical examination to predict radiographic instability of the lumbar spine. *Eur Spine J.* 2005;14:743-750.
30. Griffith LE, Shannon HS, Wells RP, et al. Individual participant data meta-analysis of mechanical workplace risk factors and low back pain. *Am J Public Health.* 2012;102:309-318.
31. Hebert JJ, Koppenhaver SL, Magel JS, Fritz JM. The relationship of transversus abdominis and lumbar multifidus activation and prognostic factors for clinical success with a stabilization exercise program: a cross-sectional study. *Arch Phys Med Rehabil.* 2010;91:78-85.
32. Hebert JJ, Koppenhaver SL, Magel JS, Fritz JM. The relationship of transversus abdominis and lumbar multifidus activation and prognostic factors for clinical success with a stabilization exercise program: a cross-sectional study. *Arch Phys Med Rehabil.* 2010;91:78-85.
33. Henry SM, Van Dillen LR, Ouellette-Morton RH, et al. Outcomes are not different for patient-matched versus nonmatched treatment in subjects

- 526 with chronic recurrent low back pain: a randomized clinical trial. *Spine J.*
527 2014;14:2799-2810.
- 528 34. Hill JC, Whitehurst DGT, Lewis M, et al. Comparison of stratified primary
529 care management for low back pain with current best practice (STarT
530 Back): a randomised controlled trial. *Lancet.* 2011;378:1560-1571.
- 531 35. Hodges PW, Coppieters MW, MacDonald D, Cholewicki J. New insight into
532 motor adaptation to pain revealed by a combination of modelling and
533 empirical approaches. *Eur J Pain.* 2013;17:1138-1146.
- 534 36. Hodges PW, van Dillen LR, McGill SM, Brumagne S, Hides JA, Moseley GL.
535 Integrated clinical approach to motor control interventions in low back
536 and pelvic pain. In: Hodges PW, Cholewicki J, van Dieen JH, eds. *Spinal*
537 *Control: The Rehabilitation of Back Pain.* Edinburgh: Elsevier; 2013:243-
538 310.
- 539 37. Hoogen JMMvd, Koes BW, Devillé W, Eijk JTMv, Bouter LM. The prognosis
540 of low back pain in general practice. *Spine.* 1997;22:1515-1521.
- 541 38. Karayannis NV, Jull GA, Hodges PW. Movement-based subgrouping in low
542 back pain: synergy and divergence in approaches. *Physiother.* 2015;
- 543 39. Kent P, Laird R, Haines T. The effect of changing movement and posture
544 using motion-sensor biofeedback, versus guidelines-based care, on the
545 clinical outcomes of people with sub-acute or chronic low back pain-a
546 multicentre, cluster-randomised, placebo-controlled, pilot trial. *BMC*
547 *Musculoskelet Disord.* 2015;16:131.
- 548 40. Kerr MS, Frank JW, Shannon HS, et al. Biomechanical and psychosocial
549 risk factors for low back pain at work. *Am J Public Health.* 2001;91:1069-
550 1075.
- 551 41. Kim MH, Yoo WG. Comparison of the lumbar flexion angle and
552 repositioning error during lumbar flexion-extension in young computer
553 workers in Korea with differing back pain. *Work.* 2015;51:223-228.
- 554 42. Kuijer PP, Takala EP, Burdorf A, et al. Low back pain: doesn't work matter
555 at all? *Occup Med.* 2012;62:152-153.
- 556 43. Kwon DK, Roffey DM, Bishop P, Dagenais S, Wai EK. Systematic review:
557 occupational physical activity and low back pain. *Occ Med.* 2011;61:541-
558 548.
- 559 44. Lamoth CJ, Daffertshofer A, Meijer OG, Lorimer Moseley G, Wuisman PI,
560 Beek PJ. Effects of experimentally induced pain and fear of pain on trunk
561 coordination and back muscle activity during walking. *Clin Biomech.*
562 2004;19:551-563.
- 563 45. Lomond KV, Henry SM, Hitt JR, DeSarno MJ, Bunn JY. Altered postural
564 responses persist following physical therapy of general versus specific
565 trunk exercises in people with low back pain. *Man Ther.* 2014;19:425-432.
- 566 46. Lu HH, Thomopoulos S. Functional attachment of soft tissues to bone:
567 development, healing, and tissue engineering. *Annu Rev Biomed Eng.*
568 2013;15:201-226.
- 569 47. Lynn B. Neurogenic inflammation caused by cutaneous polymodal
570 receptors. *Prog Brain Res.* 1996;113:361-368.
- 571 48. Macedo LG, Maher CG, Hancock MJ, et al. Predicting response to motor
572 control exercises and graded activity for patients with low back pain:
573 preplanned secondary analysis of a randomized controlled trial. *Phys*
574 *Ther.* 2014;94:1543-1554.

- 575 49. Macedo LG, Maher CG, Latimer J, McAuley JH. Motor control exercise for
576 persistent, nonspecific low back pain: a systematic review. *Phys Ther.*
577 2009;89:9-25.
- 578 50. Mannion AF, Caporaso F, Pulkovski N, Sprott H. Spine stabilisation
579 exercises in the treatment of chronic low back pain: a good clinical
580 outcome is not associated with improved abdominal muscle function. *Eur*
581 *Spine J.* 2012;21:1301-1310.
- 582 51. Marras WS, Lavender SA, Ferguson SA, Splittstoesser RE, Yang G.
583 Quantitative dynamic measures of physical exposure predict low back
584 functional impairment. *Spine.* 2010;35:914-923.
- 585 52. Mazaheri M, Coenen P, Parnianpour M, Kiers H, van Dieen JH. Low back
586 pain and postural sway during quiet standing with and without sensory
587 manipulation: A systematic review. *Gait Posture.* 2013;37:12-22.
- 588 53. McGill SM. Re: Causal assessment of occupational lifting and low back
589 pain: results of a systematic review by Wai et al. *Spine J.* 2011;11:365-366.
- 590 54. Mistry D, Patel S, Hee SW, Stallard N, Underwood M. Evaluating the
591 quality of subgroup analyses in randomized controlled trials of therapist-
592 delivered interventions for nonspecific low back pain: a systematic
593 review. *Spine.* 2014;39:618-629.
- 594 55. Mooney V, Andersson GB. Trunk strength testing in patient evaluation and
595 treatment. *Spine.* 1994;19:2483-2485.
- 596 56. Mooney V, Gulick J, Perlman M, et al. Relationships between myoelectric
597 activity, strength, and MRI of lumbar extensor muscles in back pain
598 patients and normal subjects. *J. Spinal Dis.* 1997;10:348-356.
- 599 57. Moseley GL. Evidence for a direct relationship between cognitive and
600 physical change during an education intervention in people with chronic
601 low back pain. *Eur J Pain.* 2004;8:39-45.
- 602 58. Moseley GL, Hodges PW. Reduced variability of postural strategy prevents
603 normalization of motor changes induced by back pain: a risk factor for
604 chronic trouble? *Behav Neurosci.* 2006;120:474-476.
- 605 59. Moseley GL, Vlaeyen JW. Beyond nociception: the imprecision hypothesis
606 of chronic pain. *Pain.* 2015;156:35-38.
- 607 60. Noh KH, Oh JS, Yoo WG. Comparison of lumbar repositioning error
608 according to different lumbar angles in a flexion pattern (FP) subgroup of
609 patients with non-specific chronic low back pain. *J Phys Ther Sci.*
610 2015;27:293-294.
- 611 61. Norman R, Wells R, Neumann P, Frank J, Shannon H, Kerr M. A
612 comparison of peak vs cumulative physical work exposure risk factors for
613 the reporting of low back pain in the automotive industry. *Clin Biomech.*
614 1998;13:561-573.
- 615 62. O'Sullivan P. Diagnosis and classification of chronic low back pain
616 disorders: maladaptive movement and motor control impairments as
617 underlying mechanism. *Man Ther.* 2005;10:242-255.
- 618 63. O'Sullivan PB, Burnett A, Floyd AN, et al. Lumbar repositioning deficit in a
619 specific low back pain population. *Spine.* 2003;28:1074-1079.
- 620 64. O'Sullivan PB, Mitchell T, Bulich P, Waller R, Holte J. The relationship
621 between posture and back muscle endurance in industrial workers with
622 flexion-related low back pain. *Man Ther.* 2006;11:264-271.

- 623 65. O'Sullivan P, Dankaerts W, O'Sullivan K, Fersum KP. Multidimensional
624 approach for targeted management of low back pain. In: Jull G, Moore A,
625 Falla D, Lewis J, McCarthy C, Sterling M, eds. *Grieve's Modern*
626 *Musculoskeletal Physiotherapy*. Elsevier; 2015:465-469.
- 627 66. Olson O. Re: Bakker EW, Verhagen AP, van Trijffel E, et al. Spinal
628 mechanical load as a risk factor for low back pain: a systematic review of
629 prospective cohort studies. *Spine*. 2009;35:E576.
- 630 67. Rabey M, Beales D, Slater H, O'Sullivan P. Multidimensional pain profiles
631 in four cases of chronic non-specific axial low back pain: An examination
632 of the limitations of contemporary classification systems. *Man Ther*.
633 2015;20:138-147.
- 634 68. Rabey M, Slater H, O'Sullivan P, Beales D, Smith A. Somatosensory
635 nociceptive characteristics differentiate subgroups in people with chronic
636 low back pain: a cluster analysis. *Pain*. 2015;156:1874-1884.
- 637 69. Rabey M, Smith A, Slater S, Beales D, O'Sullivan P. Pain Provocation
638 Following Sagittal Plane Repeated Movements in People With Chronic
639 Low Back Pain: Associations with Pain Sensitivity and Psychological
640 Profiles. *Scand J Pain*. 2017;16:22-28.
- 641 70. Ramond-Roquin A, Bouton C, Begue C, Petit A, Roquelaure Y, Huez JF.
642 Psychosocial Risk Factors, Interventions, and Comorbidity in Patients
643 with Non-Specific Low Back Pain in Primary Care: Need for
644 Comprehensive and Patient-Centered Care. *Front Med*. 2015;2:73.
- 645 71. Ramprasad M, Shenoy DS, Sandhu JS, Sankara N. The influence of
646 kinesiophobia on trunk muscle voluntary responses with pre-
647 programmed reactions during perturbation in patients with chronic low
648 back pain. *J Bodywork Mov Ther*. 2011;15:485-495.
- 649 72. Reeves NP, Cholewicki J, Silfies SP. Muscle activation imbalance and low-
650 back injury in varsity athletes. *J Electromyogr Kinesiol*. 2006;16:264-272.
- 651 73. Sahrmann SA. *Diagnosis and treatment of movement impairment*
652 *syndromes*. St Louis, MO, USA: Mosby; 2002.
- 653 74. Saner J, Kool J, Sieben JM, Luomajoki H, Bastiaenen CH, de Bie RA. A
654 tailored exercise program versus general exercise for a subgroup of
655 patients with low back pain and movement control impairment: A
656 randomised controlled trial with one-year follow-up. *Man Ther*.
657 2015;20:672-679.
- 658 75. Saragiotto BT, Maher CG, Yamato TP, et al. Motor control exercise for
659 chronic non-specific low-back pain. *Cochrane Database Syst Rev*.
660 2016;1:CD012004.
- 661 76. Scholz J, Mannion RJ, Hord DE, et al. A Novel Tool for the Assessment of
662 Pain: Validation in Low Back Pain. *PLoS Med*. 2009;6:e1000047.
- 663 77. Smart KM, Blake C, Staines A, Doody C. The Discriminative validity of
664 "nociceptive," "peripheral neuropathic," and "central sensitization" as
665 mechanisms-based classifications of musculoskeletal pain. *Clin J Pain*.
666 2011;27:655-663.
- 667 78. Smart KM, Blake C, Staines A, Thacker M, Doody C. Mechanisms-based
668 classifications of musculoskeletal pain: part 1 of 3: symptoms and signs of
669 central sensitisation in patients with low back (+/- leg) pain. *Man Ther*.
670 2012;17:336-344.

79. Smart KM, Blake C, Staines A, Thacker M, Doody C. Mechanisms-based classifications of musculoskeletal pain: part 2 of 3: symptoms and signs of peripheral neuropathic pain in patients with low back (+/- leg) pain. *Man Ther.* 2012;17:345-351.
80. Smart KM, Blake C, Staines A, Thacker M, Doody C. Mechanisms-based classifications of musculoskeletal pain: part 3 of 3: symptoms and signs of nociceptive pain in patients with low back (+/- leg) pain. *Man Ther.* 2012;17:352-357.
81. Solomonow M, Baratta RV, Zhou B-H, Burger E, Zieske A, Gedalia A. Muscular dysfunction elicited by creep of lumbar viscoelastic tissues. *J Electromyogr Kinesiol.* 2003;13:381-396.
82. Steele J, Bruce-Low S, Smith D, Osborne N, Thorkeldsen A. Can specific loading through exercise impart healing or regeneration of the intervertebral disc? *Spine J.* 2015;15:2117-2121.
83. Takala EP. Lack of "statistically significant" association does not exclude causality. *Spine J.* 2010;10:944.
84. Teyhen DS, Flynn TW, Childs JD, Abraham LD. Arthrokinematics in a subgroup of patients likely to benefit from a lumbar stabilization exercise program. *Phys Ther.* 2007;87:313-325.
85. Teyhen DS, Miltenberger CE, Deiters HM, et al. The use of ultrasound imaging of the abdominal drawing-in maneuver in subjects with low back pain. *J Orthop Sports Phys Ther.* 2005;35:346-355.
86. Tong MH, Mousavi SJ, Kiers H, Ferreira P, Refshauge K, van Dieën JH. Is there a relationship between lumbar proprioception and low back pain? A systematic review with meta-analysis. *Arch Phys Med Rehabil.* 2017;98:120-136.
87. Unsgaard-Tondel M, Lund Nilsen TI, Magnussen J, Vasseljen O. Is activation of transversus abdominis and obliquus internus abdominis associated with long-term changes in chronic low back pain? A prospective study with 1-year follow-up. *Br J Sports Med.* 2012;46:729-734.
88. Unsgaard-Tondel M, Nilsen TIL, Magnussen J, Vasseljen O. Are fear avoidance beliefs associated with abdominal muscle activation outcome for patients with low back pain? *Physiother Res Int.* 2013;18:131-139.
89. van der Hulst M, Vollenbroek-Hutten MM, Rietman JS, Schaaake L, Groothuis-Oudshoorn KG, Hermens HJ. Back Muscle Activation Patterns in Chronic Low Back Pain During Walking: A "Guarding" Hypothesis. *Clin J Pain.* 2010;26:30-37.
90. van der Hulst M, Vollenbroek-Hutten MM, Schreurs KM, Rietman JS, Hermens HJ. Relationships between coping strategies and lumbar muscle activity in subjects with chronic low back pain. *Eur J Pain.* 2010;14:640-647.
91. van Dieen JH, Cholewicki J, Radebold A. Trunk muscle recruitment patterns in patients with low back pain enhance the stability of the lumbar spine. *Spine.* 2003;28:834-841.
92. van Dieen JH, Kingma I, van der Bug P. Evidence for a role of antagonistic cocontraction in controlling trunk stiffness during lifting. *J Biomech.* 2003;36:1829-1836.

- 719 93. van Dieën JH, Kuijer PP, Burdorf A, Marras WS, Adams MA. Non-specific
720 low back pain. *Lancet*. 2012;379:1874.
- 721 94. van Dieën JH, Reeves NP, Kawchuk G, van Dillen LR, Hodges PW. Motor
722 control changes in low-back pain: Divergence in presentations and
723 mechanisms. *J Orthop Sports Phys Ther*. submitted;
- 724 95. van Dieën JH, Selen LP, Cholewicki J. Trunk muscle activation in low-back
725 pain patients, an analysis of the literature. *J Electromyogr Kinesiol*.
726 2003;13:333-351.
- 727 96. van Dieën JH, van der Beek AJ. Work-Related Low-Back Pain:
728 Biomechanical Factors and Primary Prevention. In: S. K, eds. *Ergonomics
729 for Rehabilitation Professionals*. Boca Raton, FL: CRC Press; 2009:359-395.
- 730 97. van Dillen LR, van Tulder M. Targeting interventions to patients:
731 development and evaluation. In: Hodges PW, Cholewicki J, Dieën JHv, eds.
732 *Spinal control: the rehabilitation of back pain. State of the art and science*.
733 Edinburgh: Churchill Livingstone; 2013:195-205.
- 734 98. Vibe Fersum K, O'Sullivan P, Skouen JS, Smith A, Kvale A. Efficacy of
735 classification-based cognitive functional therapy in patients with non-
736 specific chronic low back pain: a randomized controlled trial. *Eur J Pain*.
737 2013;17:916-928.
- 738 99. Vibe Fersum KV, O'Sullivan P, Kvale A, Skouen J. Inter-examiner reliability
739 of a classification system for patients with non-specific low back pain.
740 *Man Ther*. 2009;14:555-561.
- 741 100. Vlaeyen JWS, Kole-Snijders AMJ, Roteveel AM, Ruesink R, Heuts PHTG.
742 The role of fear of movement/(re)injury in pain disability. *J Occup Rehabil*.
743 1995;5:235-252.
- 744 101. Von Korff M, Saunders K. The course of back pain in primary-care. *Spine*.
745 1996;21:2833-2837.
- 746 102. Wong AY, Parent EC, Funabashi M, Stanton TR, Kawchuk GN. Do various
747 baseline characteristics of transversus abdominis and lumbar multifidus
748 predict clinical outcomes in nonspecific low back pain? A systematic
749 review. *Pain*. 2013;154:2589-2602.
- 750 103. Woolf CJ. Central sensitization: implications for the diagnosis and
751 treatment of pain. *Pain*. 2011;152:S2-S15.
- 752