VU Research Portal

Validity of the Flemish working alliance inventory in a Dutch physiotherapy setting in patients with shoulder pain

Karel, Yasmaline; Thoomes-de Graaf, Marloes; Scholten-Peeters, Gwendolijne; Ferreira, Paulo H; Rizopoulos, Dimitris; Koes, Bart W; Verhagen, Arianne P

published in

Physiotherapy, Theory and Practice
2018

DOI (link to publisher)

[10.1080/09593985.2017.1400141](https://doi.org/10.1080/09593985.2017.1400141)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Karel, Y., Thoomes-de Graaf, M., Scholten-Peeters, G., Ferreira, P. H., Rizopoulos, D., Koes, B. W., & Verhagen, A. P. (2018). Validity of the Flemish working alliance inventory in a Dutch physiotherapy setting in patients with shoulder pain. *Physiotherapy, Theory and Practice*, 34(5), 384-392.
<https://doi.org/10.1080/09593985.2017.1400141>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Physiotherapy Theory and Practice

An International Journal of Physical Therapy

ISSN: 0959-3985 (Print) 1532-5040 (Online) Journal homepage: <https://www.tandfonline.com/loi/iptp20>

Validity of the Flemish working alliance inventory in a Dutch physiotherapy setting in patients with shoulder pain

Yasmaine Karel, M. Thoomes-de Graaf, Gwendolijne Scholten-Peeters, Paulo Ferreira, Dimitris Rizopoulos, Bart W. Koes & Arianne P. Verhagen

To cite this article: Yasmaine Karel, M. Thoomes-de Graaf, Gwendolijne Scholten-Peeters, Paulo Ferreira, Dimitris Rizopoulos, Bart W. Koes & Arianne P. Verhagen (2018) Validity of the Flemish working alliance inventory in a Dutch physiotherapy setting in patients with shoulder pain, *Physiotherapy Theory and Practice*, 34:5, 384-392, DOI: [10.1080/09593985.2017.1400141](https://doi.org/10.1080/09593985.2017.1400141)

To link to this article: <https://doi.org/10.1080/09593985.2017.1400141>

Published online: 09 Nov 2017.

Submit your article to this journal [↗](#)

Article views: 129

View Crossmark data [↗](#)

Citing articles: 1 View citing articles [↗](#)

DESCRIPTIVE REPORT

Validity of the Flemish working alliance inventory in a Dutch physiotherapy setting in patients with shoulder pain

Yasmine Karel MSc PT^{a,b}, M. Thoomes-de Graaf MSc PT^{a,b}, Gwendolijne Scholten-Peeters PhD^{a,c}, Paulo Ferreira PhD^d, Dimitris Rizopoulos PhD^e, Bart W. Koes Professor^b, and Arianne P. Verhagen PhD^{a,b}

^aResearch Group Diagnostics, Avans University of Applied Sciences, Breda, The Netherlands; ^bDepartment of General Practice, Erasmus Medical Center, Rotterdam, The Netherlands; ^cFaculty of Behavioural and Movement Sciences, MOVE research Institute Amsterdam, VU University of Amsterdam, Amsterdam, The Netherlands; ^dFaculty of Health Sciences, University of Sydney, Sydney, Australia; ^eDepartment of Biostatistics, Erasmus Medical Center, Rotterdam, The Netherlands

ABSTRACT

Background: Working alliance is the interaction between the patient and therapist. It is a crucial part of the physiotherapeutic process. One instrument to measure working alliance is available in Dutch/Flemish language and validated in psychotherapy setting. **Objective:** This study aims to validate the Working Alliance Inventory Short-Form in a Dutch physiotherapy setting. **Design:** A prospective cohort study in primary-care physiotherapy. **Method:** To validate the Dutch/Flemish version of the working alliance inventory short-form (WAV-12) a RASCH analysis was used. **Results:** Sixty-six physiotherapists enrolled in total 389 patients with an average age of 50 years and a mean duration of shoulder pain of 33 weeks. A total of 274 patients filled in one or more items of the WAV-12. The WAV-12 shows good discriminative abilities and all items contributed to a one-dimensional construct. Due to the selective nature of the missing items, we believed rewording was necessary to make it more suitable to the physiotherapy setting. We performed a Delphi study and revised the WAV-12 into the PAS (Physio Alliance Scale). The validity of the revised version is unknown and is therefore not sufficiently strong to be implemented as a measurement tool. **Limitations:** The response rate for three items especially was low and we found ceiling effects in ten items. **Conclusion:** Although the measurement instrument shows good internal consistency and reliability, we made adjustments to the WAV-12 for Dutch physiotherapy setting.

ARTICLE HISTORY

Received 29 June 2016
Revised 16 November 2016
Accepted 22 February 2017

KEYWORDS

Working alliance;
therapeutic alliance; rasch
analysis; WAV-12;
physiotherapy; physician-
patient relations

Introduction

In physiotherapy practice patients usually follow a treatment regimen provided in coherence with the physiotherapist. This interaction between patient and therapist is referred to as a working alliance (WA). WA is first described in psychotherapy as the extent to which a client and therapist work collaboratively, purposefully, and connect emotionally. WA is defined as a combination of three factors; agreement about the goals of treatment, the tasks of treatment, and the bond between client and therapist (Bordin, 1979).

For a treatment to be effective, one important factor is that the patient complies with the regimen, after which health outcomes are more likely to improve (Bennett, Fuertes, Keitel, Phillips, 2011). Therefore, it is essential for the therapist to provide a proper transfer of information about the goals and tasks of treatment for the patient in order to carry out the treatment regimen (Crow et al., 1999; Sluijs, Kok, Van Der Zee,

1993). Besides agreement about treatment goals and tasks, co-operation and compliance are achieved by means of bonding and trust between the therapist and the patient. Patients consult a physiotherapist because they seek help and they are in that case vulnerable. Help must therefore be offered and accepted based on trust. How this relationship will develop during the treatment period can have a significant impact on treatment outcome.

Several reviews have found that WA is a strong predictor of improvement in psychotherapy and psychology practices (Horvath and Symonds, 1991; Martin, Garske, Davis, 2000). Later research has established the importance of a good alliance also in other medical settings, such as in patients with ulcer disease, hypertension, and diabetes (Kaplan, Greenfield, Ware, 1989; Lee and Lin, 2009). One review included 14 studies examining the patient–therapist relationship in physical rehabilitation setting (Hall et al., 2010). In nine studies, a registered physiotherapist delivered the

interventions. Results of the individual studies indicated that WA has a consistent positive correlation with treatment outcomes of pain, disability, physical/mental health, and patient satisfaction (Hall et al., 2010). A recent observational study of therapeutic alliance in patients with chronic low back pain confirmed these findings and found WA to be a consistent predictor of function, pain, and disability measures (Ferreira et al., 2013). WA might be more important in some therapies especially in those where treatment adherence represents an important component for treatment effect (Siev, Huppert, Chambless, 2009).

The Working Alliance Inventory (WAI) is one of the most commonly used and validated questionnaires to measure the working alliance (Hall et al., 2010). It has been originally developed as a 36-item questionnaire based on Bordin's model measuring three domains; goal, task, and bond (Horvath and Greenberg, 1989; Horvath and Lubrowsky, 1993). The WAI exists of one questionnaire for the client (WAI (C)) and one for the therapist (WAI (T)). Evidence suggests that the clients WA rating at the beginning of treatment is superior over the therapist-rated version in predicting outcome (Horvath and Symonds, 1991).

The WAI was translated to Flemish, which is closely related to Dutch, named the "werk alliantie vragenlijst" (WAV). The 12 most indicative items were selected using confirmatory factor analysis to form the WAV-12 short form (Tracey and Kokotovic, 1989). The WAV-12 has been used and validated in patients receiving psychotherapy in Belgium (Stinckens, Ulburghs, Claes, 2009). This study found a good internal consistency for the three-factor model according to Bordin (1979): (1) task scale—correlation coefficient $\alpha = 0.85$; (2) bond scale—correlation coefficient $\alpha = 0.82$; and (3) goal scale correlation coefficient $\alpha = 0.83$. Correlations between the task and goal scales were good (correlation coefficient $r = 0.80$) but correlations between the other scales were both lower (Cronbach's $\alpha = 0.49$). The WAV-12 used a five-point likert scale instead of a seven-point likert scale in the original WAV-36. Therefore, it is difficult to compare results from this validation study with other data. Literature does describe slightly higher correlation coefficients for the English and French short versions (Corbière, Bisson, Lauzon, Ricard, 2006; Tracey and Kokotovic, 1989). A review has shown that translated versions of a measurement instrument for the neck do not guarantee similar measurement properties compared with the original instrument (Schellingerhout et al., 2011). Cross-cultural validation in the Dutch population and physiotherapy setting is an important step to evaluate whether the underlying construct still holds for the WAV-12. Therefore, this study aims to investigate

whether the WAV-12 is a valid measurement instrument in terms of the construct and discriminative abilities for a population of patients with shoulder pain in physiotherapy care.

Methods

Study design

The study population consisted of patients with shoulder pain that participated in a prospective cohort study in patients consulting a physiotherapist for shoulder pain (Karel et al., 2013). Recruitment period was from November 2011 through December 2012. The Research Committee of the Erasmus Medical Centre in Rotterdam approved the project (MEC-2011-414). After signing an informed consent, patients were included and followed up for 6 months.

Participants

A total of 125 physiotherapists were invited to enroll patients. Patients consulting a physiotherapist were included if they suffered from shoulder pain, were aged ≥ 18 years and had adequate understanding of the Dutch language. Patients were excluded if they had serious pathologies (infection, cancer or fracture), surgery of the shoulder in the previous 12 months, or had received diagnostic imaging techniques such as musculoskeletal ultrasound, magnetic resonance imaging or X-ray of the shoulder in the 3 months prior to start of the study. Patients included in the cohort study were followed for 6 months and received usual physiotherapy care. Questionnaires were sent by email at 6, 12, and 26 weeks and two reminders were sent after 2 and 4 days whenever the patient had not responded to the questionnaire.

Working alliance

Working alliance (WA) was measured 6 weeks after baseline for both the patient and physiotherapist, because earlier assessment would not clearly reflect the WA. We used the Flemish version of the WAI (WAV-12). It contains 12 items scored on a five-point scale ranging from 1 ("never") to 5 ("always") and scoring is done for the total score and each subscale (goal, task and bond). The total score ranges from 12 (low WA) to 60 (high WA), and subscales range from 4 to 20. Where the patient had to fill in the name of the therapist we replaced the empty space with the words: "my therapist."

Statistical analysis

Descriptive data for demographic and symptom severity are presented as percentages for nominal variables (i.e. gender, level of education, cause of injury, first episode, and reasons for stopping treatment) and as means for continuous variables (i.e. age and symptom duration). T-tests were used to test for differences in demographics between participants scoring all WAV-12 items and those who did not. Cronbach's alpha was used to assess the internal consistency of the WAV-12 and we assessed the correlation between patient and therapist scores using Pearson's correlation coefficient. Coefficients equal or more than 0.7 were regarded as acceptable. R and SPSS v20.0 were used to conduct the analysis.

Validation

Performance of the items in the WAV-12 questionnaire was assessed with a partial credit Rasch model (Masters, 1982). The response patterns from the set of available items in the questionnaire were tested against what is expected by the model that works according to a probabilistic form of Guttman scaling (Guttman, 1950). This scale assumes a deterministic pattern with a hierarchical ordering of items (low and high level of item scale). When a higher level of the item is affirmed, there must be a high probability that lower items will also be affirmed. The analysis gives the probability that a person will affirm an item of the difference between the person's level of working alliance and the level of working alliance expressed by the item.

The Rasch model was used to test: (1) internal validity of the construct; (2) whether specific items exhibit different properties in different subgroups in the population (differential item functioning); and (3) whether item redundancy can be considered (Tennant and Conaghan, 2007). Analysis was carried out using the ltm package in the statistical programming language R (Rizopoulos, 2006).

First a one partial credit model with the discrimination parameter fixed at one was tested to check whether it fits the data. If this model did not fit the data an extended partial credit model with a common discrimination parameter not constrained at one or separate discrimination parameters for each parameter was considered. Uni-dimensionality could further be examined to investigate if the test variance is attributable to the principal factor or construct, estimated with Cronbach's alpha. Due to the fact that some patient responses were missing, multiple imputations were utilized to calculate Cronbach's alpha.

Differential item functioning was examined based on a likelihood ratio X^2 test implemented in the Lordif package in R. Expected scores for each item should

remain the same whether, an older or younger person (<50, which was the mean age) and a man or women scores the same item.

Rasch analysis can be useful and psychometrically sound in modifying measurement instruments (Veloza, Lai, Mallinson, Hauselman, 2001). Different criteria could be considered for item redundancy: High Item Characteristic Curve (ICC), low ICC or items having similar calibrations.

Results

Study population

Sixty-six physiotherapists enrolled in total 389 patients. Physiotherapists were 72% male and had a mean working experience of 15 years. Of the 389 patients, 43% were male, average age was 50 years with a mean duration of shoulder pain of 33 weeks (Table 1). At baseline, only 4% of the patients did not fill out the baseline questionnaire. At 6 weeks, 30% of the responses were lost to follow up.

Working alliance

Seventy-eight patients (22%) filled in all the WAV-12 questions, enabling us to calculate a total score. The mean WAV score was 45 on a total range of 24 to 60, which is slightly above 50% of the maximum score. Most patients did not answer one or more questions of the WAV-12. The population that had responded to all WAV-12 questions did not significantly differ at baseline with the patients that did not (Table 1). Even though not statistically significant, the difference for duration of complaints appeared to be large. Selective responses can therefore not be excluded. The questions with the most missing values are questions 1, 3, 7, and 9 (Figure 1). Question 3, 7, and 9 are part of the "bond" subscale and question 1 is part of the "goal" subscale. The working alliance score of therapists was 52 and for patients 45. WAV-12 scores between patient and therapist had a poor correlation ($r = 0.30$).

Validity of WAV-12

Of all patients, 274 had at least filled in one or more items of the WAV-12. Three models were fitted to the data. The first model (RASCH) assumes the discrimination parameter is equal for all items and fixed at one. The second model (1PL) assumes the discrimination parameter is equal for all items but is estimated from the data and the third model (gpcm) assumes the discrimination parameter is free to vary across items.

Table 1. Baseline characteristics.

Characteristics of cohort	Total n = 389	Participants filling in all items of WAV-12; n = 87	Participants, missing 1 or more items of WAV-12; n = 302
Male (%)	170 (43)	41 (49)	129 (44)
Age (SD)	50 (13)	50 (14)	50 (13)
Duration of complaint in weeks (SD)	33 (82)	27 (58)	34 (88)
Comorbidity (%)			
No	128 (35)	25 (29)	103 (34)
Yes	236 (65)	62 (71)	199 (66)
Medication use (%)	183 (47)	40 (49)	144 (50)
Highest education (%)			
Primary school	40 (10)	12 (15)	28 (10)
High school	199 (51)	44 (54)	155 (54)
University or applied sciences	127 (33)	25 (31)	102 (36)
Paid job (%)	261 (67)	53 (65)	208 (72)
Profession (%)			
Physically intensive job	65 (17)	13 (25)	52 (25)
Static repetitive job	88 (23)	14 (27)	74 (35)
Job with awkward positions/ postures	11 (3)	3 (6)	8 (4)
Other	99 (25)	22 (42)	77 (36)
NRS median (IQR)	6.0 (3.0)	6.0 (2.0)	6.0 (3.0)
SDQ (SD)	62 (23)	63 (24)	62 (23)
EQ-5D (SD)	0.83 (0.08)	0.82 (0.07)	0.83 (0.09)

NRS Numeric Rating Scale, SDQ Shoulder Disability Index, EQ-5D EuroQol 5 Dimensions, SD standard deviation

Likelihood ratio tests between these models showed that the third model provided the best fit to the data ($p \leq 0.001$).

Item properties

All but two items (item 1 and 2), showed ceiling effects, meaning that most of the patients scored a good working alliance. Appendix 1 displays the item characteristic curves for the 12 items from the WAV-12. Items 5, 6, and 8 have a high slope and are endorsed at higher levels of working alliance. Items 1, 2, and 4 have a low slope (discrimination) and are endorsed at lower levels of WA. Considerable variation exists between item discrimination indicating the WAV-12 questionnaire includes items measuring the whole construct and items discriminating at lower and higher levels of working alliance (Table 2). The item information curve showed the amount of information given by the questionnaire is highest between an ability of -2 and 0 , implying that the item set is most useful in discriminating among individuals at the lower end of the working alliance trait.

Unidimensionality

Five imputed datasets were created. Cronbach's alpha's were calculated for the 12 items in each dataset and led to a pooled Cronbach's alpha coefficient of 0.89 , indicating that items correlate highly and measure the same explanatory concept.

Differential item functioning

The X^2 tested three models. Model 1 is a standard model where the ability for each person remains the same. Model 2 tests whether levels of ability differ among groups and model 3 adds an interaction term for the level of ability and the group in order to test whether discrimination parameters differ among groups.

Figure 1. Relative response rate per item of WAV-12.
I item 1–12 of the WAV 12

Table 2. Discrimination values of WAV-12 items.

Item	Discrimination	Standard error	Z value
1	0.496	0.103	4.793
2	0.443	0.088	5.066
3	1.286	0.225	5.716
4	0.761	0.118	6.424
5	2.212	0.457	4.842
6	2.067	0.338	6.114
7	1.377	0.234	5.895
8	2.266	0.369	6.139
9	1.151	0.208	5.537
10	1.068	0.158	6.742
11	1.414	0.224	6.319
12	1.107	0.167	6.613

Age was dichotomized in younger patients (under the mean age of 50) and older patients (50 and over). The X^2 tested flagged item one for differential item functioning where all models were statistically significant. No differential item functioning was found between men and women. Slightly higher factor scores (mean difference = 0.0385) for the WA in patients being treated by a physiotherapist with less than 13 years of experience but was not statistically significant ($p = 0.73$).

Rasch analysis for the WAV-12 questionnaire indicates that items have good discriminative abilities for the lower end of the construct. High correlations coefficients indicate items measure one construct and other factors like age and experience of the physiotherapist did not influence item scoring. Validity for the items in the questionnaire appears to be sound but due to the difference in the percentage of missing data among the items and observed ceiling effects we advise linguistic (Dutch) and contextual (physiotherapeutic setting) adjustments.

Modification of the WAV-12

We believed rewording was necessary due to the selective number of missing responses in some items of the questionnaire and because the researchers had received comments from several patients and physiotherapists about items 3, 7, and 9 of the WAV-12. Therefore, we decided to make adjustments in the questionnaire and did a Delphi study. A two round survey was employed to ask the panels opinion on the adjustments in the WAV-12. The panel consisted of 11 members (six clinical/research experts and five patients). Panel members were sent a questionnaire via email and these were sent separately to ensure panel members were unaware each of other's identity. For each item, the panel member had to give his/her opinion about the adjustments with a five-point Likert scale. If the score was below three (neutral, disagree, totally disagree) the panel member were asked to give their reasoning and/or a

suggestion for adjustment. If consensus for one item was <80% after the first round it was included in the second round containing the suggestions of all panel members (anonymous). Full consensus (100% response rate) was reached after the second round and the adjusted questionnaire can be found in Appendix 2.

Discussion

Main findings

Just a small proportion of patients filled in the complete WAV-12 compared with other questionnaires at 6-weeks follow-up. A large number of participants only completed a limited number of items. This might indicate that the measurement instrument is not appropriate either in terms of language, setting, or participants had other specific reasons not to complete the questionnaire. The principal investigator also received comments from several patients and therapists, involved in the study, about items 3, 7, and 9 in the WAV-12 questionnaire. The construct theory of the WAV appeared to be sound but ceiling-effects were found in ten items. Rewording was necessary for the WAV-12.

Comparison with the literature

Items correlated highly and measured the same explanatory concept which is found by several other translated versions of the WAI (Corbière, Bisson, Lauzon, Ricard, 2006; Horvath and Greenberg, 1989; Tracey and Kokotovic, 1989). A French validation study found a very high correlation between the three subscales indicating that we cannot significantly distinct these subscales (Corbière, Bisson, Lauzon, Ricard, 2006).

The poor correlation between patient and therapist WA score is consistent with other studies indicating that the two perspectives are not associated, which is confirmed by other studies as well (Bachelor, 2013; Huddy et al., 2012). To ensure unbiased results, the patient and the physiotherapist completed the rating forms independently of each other. Nevertheless, contact between the therapist and patient could not have been avoided, resulting in the possibility of deliberation between them.

WA was measured at 6 weeks when alliance might already have evolved into a stable situation; whereas, the first clinical experience between patient and therapist could determine more valid WA scores (Horvath, 2001). The literature is still inconsistent about what the optimal timing would be for measuring WA and some studies report that early WA predicted recovery after controlling for symptom change (Anker, Owen,

Duncan, Sparks, 2010; Barber et al., 2000; De Bolle, Johnson, De Fruyt, 2010; Klein et al., 2003), while others have found a reduction of the predictive value of WA (Barber et al., 1999; DeRubeis and Feeley, 1990; Puschner, Wolf, Kraft, 2008). In this study WA was measured at six weeks as the first questionnaire was filled in before the first treatment. Nevertheless, we believe multiple measurements during the treatment period might yield more insight into the concept of WA.

Although WA is a valid construct within psychological interventions and research, whether it predicts recovery in a patient population in physiotherapy setting remains unknown. Psychological interventions are usually based on behavioral therapy that physiotherapists mostly use in chronic patients. The patient population in this study has a new episode of shoulder pain where WA might be less relevant for the therapeutic process.

Strengths and limitations

This is the first study to perform a validation analysis on the Flemish version of the working alliance inventory in a physiotherapy setting. The measurement tool was able to discriminate between patients that experience a good or poor alliance. In ten items, we observed ceiling effects, which might have been due to the fact that patients give socially desirable answers or that the items do not properly assess the total construct. There appeared to be a pattern in missing responses, where four items showed more missing than others, indicating that these might need adjustment. The questionnaire was developed in Belgium and applied in a Dutch setting which might not be appropriate given some linguistic characteristic differences of the Belgian Dutch (Flemish) and the Dutch language in the Netherlands. Due to the high number of missing responses in specific items (item 1, 3, and 9) and low discriminative values (item 1 and 2), we made changes in terms of adjustments in language and specific to the context of physiotherapy.

Implications for future research

The new questionnaire from our Delphi study has not been tested and therefore future research should test the psychometric properties of this questionnaire and evaluate the possible predictive value of the WA throughout the whole process of treatment in patients with musculoskeletal complaints. Whether measuring WA at the beginning or later in therapy is more predictive remains unknown. Studying a relationship between WA and recovery is

complex because other factors, like self-adherence, compliance, might influence the relationship and therefore a mediation analysis might find more valid results.

Conclusions

The WAV-12 measurement tool is not suitable for implementation in clinical or research practice yet. However, WA is a concept that needs attention within the field of physiotherapy and therefore we made adjustments to the questionnaire. Previous research has shown a positive correlation between working alliance and recovery in physiotherapy setting. Since shoulder pain can become a chronic condition in more than 50% of patients, interventions from physiotherapy need to be effective and a good WA can possibly contribute to optimal treatment effects.

Acknowledgments

We thank the Delphi panel and Davy Paap for contributing to the adjustments of the questionnaire.

Declaration of interest

The authors report no declarations of interest.

Funding

This study is financed by the SIA-RAAK grant. The ministry of education has made this funding available for the innovation and promotion of research. This study is partly funded by a program grant of the Dutch Arthritis Foundation.

References

- Anker M, Owen J, Duncan B, Sparks JA 2010 The alliance in couple therapy: Partner influence, early change, and alliance patterns in a naturalistic sample. *Journal of Consulting and Clinical Psychology* 78: 635–645.
- Bachelor A 2013 Clients' and therapists' views of the therapeutic alliance: Similarities, differences and relationship to therapy outcome. *Clinical Psychology and Psychotherapy* 20: 118–135.
- Barber J, Connolly M, Crits-Christoph P, Gladis L, Siqueland L 2000 Alliance predicts patient's outcome beyond in-treatment change in symptoms. *Journal of Consulting and Clinical Psychology* 68: 1027–1032.
- Barber J, Luborsky L, Crits-Christoph P, Thase M, Weiss R, Frank A, Onken L, Gallop R 1999 Therapeutic alliance as a predictor of outcome in treatment of cocaine dependence. *Psychotherapy Research* 9: 54–73.
- Bennett J, Fuertes J, Keitel M, Phillips R 2011 The role of patient attachment and working alliance on patient adherence, satisfaction and health related quality of life in lupus treatment. *Patient Education and Counseling* 85: 53–59.

- Bordin E 1979 The generalizability of the psychoanalytic concepts of the working alliance. *Psychotherapy: Theory, Research and Practice* 16: 252–260.
- Corbière M, Bisson J, Lauzon S, Ricard N 2006 Factorial validation of a French short-form of the Working Alliance Inventory. *International Journal of Methods in Psychiatric Research* 15: 36–45.
- Crow R, Gage H, Hampson S, Hart J, Kimber A, Thomas H 1999 The role of expectancies in the placebo effect and their use in the delivery of health care: A systematic review. *Health Technology Assessment* 3: 1–96.
- De Bolle M, Johnson J, De Fruyt F 2010 Patient and clinician perceptions of therapeutic alliance as predictors of improvement in depression. *Psychotherapy and Psychosomatics* 79: 378–385.
- DeRubeis R, Feeley M 1990 Determinants of change in cognitive therapy for depression. *Cognitive Therapy and Research* 14: 469–482.
- Ferreira P, Ferreira M, Maher C, Refshauge K, Latimer J, Adams R 2013 The therapeutic alliance between clinicians and patients predicts outcome in chronic low back pain. *Physical Therapy* 93: 470–478.
- Guttman L 1950 *The Basis for Scalogram Analysis*, 4th edn. New York, Wiley.
- Hall A, Ferreira P, Maher C, Latimer J, Ferreira M 2010 The influence of the therapist-patient relationship on treatment outcome in physical rehabilitation: A systematic review. *Physical Therapy* 90: 1099–1110.
- Horvath A 2001 The alliance. *Psychotherapy* 38: 365–372.
- Horvath A, Greenberg L 1989 Development and validation of the working alliance inventory. *Journal of Counseling Psychology* 36: 223–233.
- Horvath A, Lubrowsky L 1993 The role of the therapeutic alliance in psychotherapy. *Journal of Consulting and Clinical Psychology* 61: 561–573.
- Horvath A, Symonds D 1991 Relation between working alliance and outcome in psychotherapy: A meta-analysis. *Journal of Counseling Psychology* 38: 139–149.
- Huddy V, Reeder C, Kontis D, Wykes T, Stahl D 2012 The effect of working alliance on adherence and outcome in cognitive remediation therapy. *Journal of Nervous and Mental Disease* 200: 614–619.
- Kaplan S, Greenfield S, Ware J 1989 Assessing effects of physician-patient interactions on the outcomes of chronic disease. *Medical Care* 27: S110–127.
- Karel YH, Scholten-Peeters WG, Thoomes-De Graaf M, Duijn E, Ottenheijm R, Van Den Borne MP, Koes BW, Verhagen AP, Dinant GJ, Tetteroo E, et al 2013 Current management and prognostic factors in physiotherapy practice for patients with shoulder pain: Design of a prospective cohort study. *BMC Musculoskeletal Disorders* 14: 62.
- Klein D, Schwartz J, Santiago N, Vivian D, Vocisano C, Castonguay L, Arnow B, Blalock JA, Manber R, Markowitz JC, et al 2003 Therapeutic alliance in depression treatment: Controlling for prior change and patient characteristics. *Journal of Consulting and Clinical Psychology* 71: 997–1006.
- Lee Y, Lin J 2009 The effects of trust in physician on self-efficacy, adherence and diabetes outcomes. *Social Science and Medicine* 68: 1060–1068.
- Martin D, Garske J, Davis K 2000 Relation of the therapeutic alliance with outcome and other variables: A meta-analytic review. *Journal of Consulting and Clinical Psychology* 68: 438–450.
- Masters G 1982 A Rasch model for partial credit scoring. *Psychometrika* 47: 149–174.
- Puschner B, Wolf M, Kraft S 2008 Helping alliance and outcome in psychotherapy: what predicts what in routine outpatient treatment? *Psychotherapy Research* 18: 167–178.
- Rizopoulos D 2006 ltm: An R package for latent variable modeling and item response theory analyses. *Journal of Statistical Software* 17(5): 1–25.
- Schellingerhout J, Heymans M, Verhagen A, De Vet H, Koes B, Terwee C 2011 Measurement properties of translated versions of neck-specific questionnaires: A systematic review. *BMC Medical Research Methodology* 11: 87.
- Siev J, Huppert J, Chambless D 2009 The dodo bird, treatment technique, and disseminating empirically supported treatments. *Behavior Therapist* 32: 69–75.
- Sluijs E, Kok G, Van Der Zee J 1993 Correlates of exercise compliance in physical therapy. *Physical Therapy* 73: 771–782.
- Stinckens N, Ulburghs A, Claes L 2009 De werkaliantie als sleutelement van de WAV-12, de Nederlandstalige verkorte versie van de working alliance inventory. *Tijdschrift Voor Klinische Psychologie* 39: 44–60.
- Tennant A, Conaghan P 2007 The Rasch measurement model in rheumatology: What is it and when use it? When should it be applied and what should one look for in a rasch paper? *Arthritis Care and Research* 57: 1358–1362.
- Tracey T, Kokotovic A 1989 Factor structure of the working alliance inventory. *Psychological Assessment* 1: 207–210.
- Veloze C, Lai J, Mallinson T, Hauselman E 2001 Maintaining instrument quality while reducing items: Application of Rasch analysis to a self-report of visual function. *Journal of Outcome Measurement* 4: 667–680.

Appendix 1

Item characteristic curves for the items in the WAV-12 questionnaire. Probability of working alliance score on the

total construct for each response category of the item in different colors (1–5 likert scale).

Item response category characteristic curve item 1

Item response category characteristic curve item 2

Item response category characteristic curve item 3

Item response category characteristic curve item 4

Item response category characteristic curve item 5

Item response category characteristic curve item 6

Item response category characteristic curve item 7

Item response category characteristic curve item 8

Item response category characteristic curve item 9

Item response category characteristic curve item 10

Item response category characteristic curve item 11

Item response category characteristic curve item 12

Appendix 2

Physio Alliance Scale (PAS)

Naam Datum:

Instructies:

Hieronder en op de volgende pagina worden een aantal omschrijvingen gegeven over de wijze waarop patiënten kunnen denken of voelen omtrent de relatie met de fysiotherapeut. Onder elke uitspraak bevinden zich vijf mogelijkheden om te antwoorden:

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

Indien de uitspraak aangeeft hoe u zich altijd voelt (of hoe u altijd denkt), omcirkelt u de antwoordmogelijkheid ALTIJD. Als ze nooit op u van toepassing is, omcirkelt u de antwoordmogelijkheid ZELDEN OF NOOIT. Gebruik de alternatieven tussenin om de variaties tussen deze extremen te beschrijven.

Geef een antwoord op alle uitspraken.

- (1) Een resultaat van de therapie is dat ik weet hoe ik mijn klacht kan beïnvloeden.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (2) De therapie geeft mij een nieuwe kijk op mijn klacht.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (3) Ik geloof dat mijn fysiotherapeut(e) mij een prettig persoon vindt.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (4) De fysiotherapeut(e) betreft mij bij het bepalen van de doelstellingen voor de therapie.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (5) Mijn fysiotherapeut(e) en ik respecteren elkaar.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (6) Mijn fysiotherapeut(e) en ik werken naar de doelstellingen toe waar we het beide over eens zijn.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (7) Ik heb het gevoel dat mijn fysiotherapeut(e) mij waardeert.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (8) Mijn fysiotherapeut(e) en ik zijn het erover eens wat voor mij belangrijk is om aan te werken.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (9) Ik heb het gevoel dat mijn fysiotherapeut(e) het beste met mij voor heeft, zelfs wanneer ik dingen doe waar hij/zij het niet mee eens is.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (10) Ik heb het gevoel dat de dingen die ik tijdens de therapie doe, mij zullen helpen om mijn doelen voor de therapie te bereiken.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (11) Mijn fysiotherapeut(e) en ik hebben dezelfde opvattingen over de veranderingen die goed zouden zijn voor mij.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

- (12) Ik geloof dat de manier waarop we aan mijn klacht werken, de juiste is.

ZELDEN OF NOOIT/SOMS/VAAK/ZEER VAAK/ALTIJD

Kijk of u alle antwoorden heeft ingevuld.
Hartelijk bedankt voor uw medewerking!