

VU Research Portal

Traumatic spinal fractures the fall and rise

Smits, A.J.

2020

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Smits, A. J. (2020). *Traumatic spinal fractures the fall and rise*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

PART 2

OPERATIVE TREATMENT OF SPINAL FRACTURES

CHAPTER

3

COMBINED ANTEROPOSTERIOR FIXATION USING A TITANIUM CAGE VERSUS SOLELY POSTERIOR FIXATION FOR TRAUMATIC THORACOLUMBAR FRACTURES: A SYSTEMATIC REVIEW AND META-ANALYSIS

Smits AJ
Polack M
Deunk J
Bloemers FW

Abstract

Background

Additional anterior stabilization might prevent posterior implant failure but over time the disadvantages of bone grafts have become evident. The objective of this systematic review was to compare risks and advantages of additional anterior stabilization with a titanium cage to solely posterior fixation for traumatic thoracolumbar fractures.

Methods

An electronic search was performed in the literature from 1980 until March 2016. Studies comparing only posterior with antero-posterior fixation by means of a titanium cage were included. Data extraction and Cochrane risk of bias assessment was done by two independent authors. In addition the PRISMA statement was followed and the GRADE approach was used to present results.

Results

Of the 1584 studies, two RCTs and one retrospective cohort study were included in the meta-analysis. The RCTs reported evidence of high quality that antero-posterior stabilization maintained better kyphosis correction than posterior stabilization alone. However these results were neutralized in the meta-analysis by the cohort study. Implant failure was reported by one study, in the posterior group. No differences in follow-up VAS scores, neurologic improvement and complications were found. Operation time, blood loss and hospital stay all increased in the antero-posterior group.

Conclusions

Patients with a highly comminuted or unstable fracture could benefit from combined antero-posterior stabilization with a titanium cage, for some evidence suggests this prevents loss of correction. However, large randomized studies still lack. There is a risk of cage subsidence and increased perioperative risks have to be considered when choosing the optimal treatment.

Introduction

Posterior short segment fixation is one of the most widely used surgical stabilization techniques for spinal fractures. The posterior spine is relatively easy accessible and posterior fixation provides kyphosis correction, indirect reduction of canal encroachment and stabilization of the fractured vertebra. However in specific fracture types, reported instrumentation failure up to 40% and loss of kyphosis correction^[1-4] indicated the necessity of additional anterior stabilization. In 1994 the load sharing classification (LSC) was introduced^[5] to predict posterior implant failure, and has been used to select patients requiring additional anterior column support.

Different methods to support the anterior column have been developed; bone strut grafts, vertebral body stents, mesh- and expandable cages, with or without additional anterolateral plating. Different techniques have their respective disadvantages. Widely used autologous bone-grafts are associated with post donor site pain^[6, 7], risk of non-union^[8] and increased correction loss^[8, 9]. The anterior approach is surgically invasive, but nowadays minimally-invasive techniques like thoracoscopy^[10-12] are available. Although minimally invasive transpedicular vertebral body stenting is promising^[13], long term results and applicability to traumatic fractures are yet unknown. However, fractures with neurologic deficit and a load sharing score (LSC) of ≥ 7 also have shown acceptable outcomes when treated non-operatively^[14, 15] or with solely short-segment posterior fixation^[16].

Biomechanical studies have shown superior stability of antero-posterior fixation compared to posterior instrumentation alone^[17-20]. While anterior stabilization mainly prevents loss of correction, studies did not show correlation with improved functional outcomes^[9, 15]. In addition, long-term maintenance of correction is possibly affected by fracture type and anterior graft material^[21]. Clinical studies comparing the solely posterior with anterior^[22, 23] or antero-posterior stabilization by means of a titanium cage are scarce or involved mainly bone strut grafts^[3, 24], outdated instruments^[25] or osteoporotic fractures^[26]. The titanium cage might provide additional value while it avoids risks and disadvantages associated with other anterior stabilization techniques.

Yet the exact value of an additional anterior cage remains unclear. The aim of this systematic review is to provide the evidence in the current literature of additional anterior stabilization with a titanium cage compared to solely posterior fixation for traumatic thoracolumbar fractures.

Materials and methods

The systematic review was conducted according to the PRISMA statement^[27]. Electronic searches were performed in Pubmed and Embase from January 1980 until January 2017. Published articles as well as accepted and drafts in English, German, French, Dutch and Chinese were deemed eligible. Authors of articles in languages other than English were contacted for English translation and if they not responded, articles were translated by a medical professional translator. The search consisted of general and Mesh (medical subject heading) terms and variants of "Spinal Fractures", "spine", "vertebrae", "fracture*", "injury", "anterior", "posterior*", "stabilization", "fixation" (Table 1 provides the full electronic search strategy). Furthermore, references of articles retrieved after the first selection were searched for eligible studies. Inclusion criteria consisted of (1) clinical trials or cohort studies involving (2) patients with traumatic thoracolumbar fractures and comparing (3) solely posterior fixation using pedicle screws with (4) combined antero-posterior stabilization by means of a titanium cage and pedicle screws. Excluded were articles comparing treatment for (1) patients with degenerative, pathologic or osteoporotic indication, (2) studies not comparing both treatments, (3) editorials and letters to the editor and (4) articles in languages other than mentioned. Two authors independently selected articles based on title and abstract. Full-texts of the remaining articles were then read, if disagreement existed on inclusion this was solved through discussion or with a tertiary independent author.

Table 1. Full electronic search strategy for at least one database as in concordance with the PRISMA statement. The presented search was used in the PubMed database in the systematic review and meta-analysis.

1	("Spinal Fractures"[Mesh] OR ((spine[tiab] OR spinal[tiab] OR vertebrae[tiab] OR vertebral[tiab]) AND (fractur*[tiab] OR injury[tiab] OR injuries[tiab]))) AND (anterior*[tiab] AND posterior*[tiab] AND (stabil*[tiab] OR fixat*[tiab]))
2	((("Spinal Fractures"[Mesh] OR ((spine[tiab] OR spinal[tiab] OR vertebrae[tiab] OR vertebral[tiab]) AND (fractur*[tiab] OR injury[tiab] OR injuries[tiab]))) AND (anterior*[tiab] AND posterior*[tiab] AND (stabil*[tiab] OR fixat*[tiab]))) AND (cage[tiab] OR synex[tiab] OR obelisk[tiab]))
3	((("Spinal Fractures"[Mesh] OR ((spine[tiab] OR spinal[tiab] OR vertebrae[tiab] OR vertebral[tiab]) AND (fractur*[tiab] OR injury[tiab] OR injuries[tiab]))) AND (anterior*[tiab] AND posterior*[tiab] AND (stabil*[tiab] OR fixat*[tiab]))) AND (randomized controlled trial[pt] OR controlled clinical trial[pt] OR randomized[tiab] OR placebo[tiab] OR drug therapy[sh] OR randomly[tiab] OR trial[tiab] OR groups[tiab])) OR (((("Spinal Fractures"[Mesh] OR ((spine[tiab] OR spinal[tiab] OR vertebrae[tiab] OR vertebral[tiab]) AND (fractur*[tiab] OR injury[tiab] OR injuries[tiab]))) AND (anterior*[tiab] AND posterior*[tiab] AND (stabil*[tiab] OR fixat*[tiab]))) AND (systematic[sb]))

Data collection

Data extraction from included articles was done using a standardized extraction form created for this study. If data was insufficient, authors were contacted for additional information. Data was extracted for (1) patient characteristics (age, sex, fracture type and level), (2) number of patients, (3) surgical techniques, (4) reported outcomes (visual analog scale (VAS), loss of kyphosis correction, neurological improvement, complications, SF-36) and (5) follow-up duration. Loss of kyphosis correction was defined as the difference in kyphosis directly postoperative and at final follow up. One study used the VAS-spine score, 19 questions concerning fracture related back-pain and is rated from 0 to 100 with 100 being no disability/pain. To compare this score to the regular VAS scores of the other studies, it was inverted to 0 being no disability/pain and 100 being maximum pain.

Risk of bias assessment

Risk of bias was assessed at the study level for randomized studies using the Cochrane risk of bias tool. Cohort studies were assessed at the outcome level using the Cochrane risk of bias assessment tool: for non-randomized studies of interventions (Acrobat-NRSI). Outcomes are reported in Tables 2 and 3. Pooled results are presented integrating the risk of bias using the GRADE approach (GRADEpro GDT, McMaster University, 2015).

Statistical analysis

Results from included studies were pooled for a meta-analysis where possible. If not reported, standard deviations were calculated from P-values or confidence intervals, if these were not available the range was used^[28]. Random effects models were used since heterogeneity was suspected. To estimate the total treatment effect, standardized mean differences were calculated for studies using different scoring scales. Mean differences were calculated if studies used the same continuous outcome scale. To compare dichotomous outcomes, risk ratios with 95% confidence intervals were calculated to estimate total effect. To test whether observed differences in results could be due to chance alone, a χ^2 -test was used (with $p < 0,1$ considered significant). The I^2 -test was used to estimate the percentage of variability in effect estimates that is due to heterogeneity, with a value of $>70\%$ considered as substantial heterogeneity. Using a funnel plot to determine publication bias was not feasible due to the small amount of included studies.

Analyses were performed using Review Manager 5.3 (Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014).

Table 2. Cochrane risk of bias assessment for cohort studies, concerning the study of Weiner et al.(31)

Domain	Correction Loss	Post-op VAS score
Confounding bias	Serious	Serious
Selection bias	Moderate	Moderate
Bias in measurements of intervention	Low	Low
Bias due to departures from intended interventions	Low	Moderate
Bias due to missing data	Moderate	Moderate
Bias in measurement of outcomes	Low	Moderate
Bias in selection of reported result	Moderate	Moderate
Overall	Serious	Serious

Table 3. Cochrane risk of bias assessment for randomized controlled studies


	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Korovessis et al.	?	?	-	?	+	-	+
Wang et al.	+	?	-	+	+	-	+
Weiner et al.	Not applicable						

Results

Study selection

The initial search resulted in 1584 articles after duplicates were removed. After selection based on title and abstract nine articles seemed eligible for inclusion based on title and abstract. After full-text was assessed, two RCTs and one retrospective cohort study remained eligible for qualitative analysis and could [29-31] be included in the meta-analysis (Figure 1).

Figure 1. PRISMA flow chart of study selection



Included studies consisted of a total of 134 patients, 69 of which underwent antero-posterior stabilization with pedicle screws and a titanium cage and 65 patients solely posterior stabilization with pedicle screws. All three studies described groups ranging from 20-28 patients. Patient characteristics, generally comparable among studies, are summarized in Table 4. All studies included more males compared to females. Noticeable differences are fracture levels; one study[29] only assessed the mid-lumbar region (L2-L4), whereas other studies assessed the thoracolumbar region (T11-L2).[30, 31] The longest mean follow-up was 70

months.^[30] Wang et al.^[30] also included very severe fractures while Korovessis et al.^[29] used a LSC of 6 as upper boundary for inclusion.

Surgical technique

Two studies used a titanium mesh cage filled with autogenous bone^[29, 30], the other study used a titanium expandable cage^[31]. One study used posterior instrumentation with pedicle screws in the fractured vertebra in both groups^[29], whereas the other two^[30, 31] only used pedicle screws in both groups one level above and below the fractured vertebra. Wang et al.^[30] additionally performed posterolateral fusion with autogenous bone graft. Only Weiner et al.^[31] described removal of posterior implant in a few patients in both groups before final follow up.

Outcomes

All studies reported kyphosis angles and VAS scores, although one study reported the VAS-spine score^[31]. Two studies^[29, 30] reported complication rates for wound infections and deep venous thrombosis, neurologic improvement on Frankel scale, operation time, blood-loss and hospital stay. Two studies^[29, 31] reported different domains of the SF-36 and were therefore not comparable. Outcomes are reported in Table 4.

Quality assessment

Risk of bias of the RCTs are shown in Table 3. Both studies scored a generally low to unclear risk of bias. For both studies it was very unlikely that participants were blinded to intervention. Wang et al.^[30] reported less types of complications compared to Korovessis et al.^[29] In addition Korovessis et al.^[29] did not report method of randomization and blinding, nor did they report all items of the SF-36.

In the cohort study of Weiner et al.^[31] a selection bias may have occurred for both outcomes, as treatment allocation depended on fracture type. Also, start of follow-up and start of intervention did not coincide for most subjects. Of the 46 patients that went missing to follow-up, it was unclear to which group they belonged, and these were neither included in their analysis. The exact time frame of last measurement was unclear. Overall, both correction loss and VAS-spine score scored a serious risk of bias (Table 2).

Meta-analysis

VAS scores

Antero-posterior treatment had a moderate effect on lowering the postoperative VAS score compared to the solely posterior approach (SMD -0,64; 95% CI: -1,69-0,41; $P=0,23$).

Table 4. Summary of study characteristics and outcomes


Author	year	Study type	Outcome*	Inclusion	Exclusion	Approach	n	Age yrs (SD)	M:F	Fract level	FU (months, SD)
Wang et al.	2015	RCT	VAS, Complications, Frankel, Cobb-angle, Surg time & BL	Burst fx with >50% VHL, or >20° kyphosis, or >50% SCE	>1 level fx, concomitant surgical injuries, History of spinal surgery	AP	21	41 (13)	16:5	T12-L2	71 (9)
Korovessis et al.	2006	RCT	VAS, Complications, Frankel, Gardner-angle, Surg time & BL, SF36	AO A3 fx with LSC ≤ 6, fx within 1 week	>1 level fx, polytraumatized, osteoporosis, other spinal disease or surgery	AP	20	39 (19)	16:4	L2-L4	46 (?)
Weiner et al.	2013	Cohort	VAS-spine, Cobb-angle, SF36	Magerl ≥ A3 or looming neu deficit through SCE or kyphosis > 15-20°	Pathologic fx, complete SCI, ≤18 months FU, age <16 > 65, lost to FU	AP	28	45 (?)	6:4	Thoracolumbar	41 (?)

VHL: Vertebral Height Loss, SCE: Spinal Canal Encroachment, AP: anterior-posterior, P: posterior, BL: Blood-loss, fx: fracture, LSC: load sharing classification, SCI: spinal cord injury, IND: intervertebral disk, FU: Follow up

*only outcomes that are used in this review are reported

The substantial heterogeneity ($I^2=88\%$; $X^2: p<0,01$) did not decrease using stratified analysis for RCT's and the cohort study (Figure 2). Quality of evidence for this outcome was graded very low, using all three studies due to risk of bias and imprecision of the cohort study, and indirectness of the RCT's. Using only the RCT's, the quality of evidence was graded moderate due to inconsistency.

Figure 2. Forest plot of visual analog scale scores at follow-up for combined anteroposterior approach with a titanium cage versus solely posterior fixation, stratified for RCT and cohort studies



Radiologic evaluation

Although the antero-posterior group maintained more kyphosis correction on final follow-up compared to the posterior group, this was not significant (MD -2,50; 95% CI: -6,56-1,51; $P=0,22$). With stratified analyses for all studies, heterogeneity decreased from substantial to moderate, I^2 from 84% to 44% and X^2 from $P=0,002$ to $P=0,18$. The RCT group showed significantly more kyphosis maintenance in favor of the antero-posterior group (MD -4,59; 95% CI: -6,95 - -2,22; $P<0,01$) (Figure 3). Using all three studies, the quality of evidence for correction loss was graded very low due to risk of bias and indirectness from inclusion of the cohort study and inconsistency. Excluding the cohort study and including only the RCT's, the quality of evidence was graded High.

Neurologic improvement

Neurologic improvement was graded as at least one grade improvement on Frankel scale. The antero-posterior group had a higher, though not significant, relative risk of neurologic improvement (Risk ratio 1,15; 95% CI: 0,92-1,43; $P=0,22$) (Figure 4). Heterogeneity was not important ($I^2=0\%$, $X^2: P=0,99$). Quality of evidence of neurologic improvement was graded high.

Figure 3. Forest plot of kyphosis correction loss (degrees) measured from postoperative value to final follow-up, for combined anteroposterior approach with a titanium cage vs solely posterior fixation






Figure 4. Forest plot of neurologic improvement on Frankel scale for combined anteroposterior approach with a titanium cage versus solely posterior fixation



Complications

Complications in both RCT's reported were deep venous thrombosis and wound infections. The antero-posterior group showed a higher risk for these complications, however not significant (RR 2,06; 95% CI: 0,56 – 7,61; $P=0,28$) (Figure 5). Heterogeneity was not important ($I^2=0\%$, $X^2: P=0,95$). Korovessis et al.^[29] reported more types of complications than did Wang et al. Quality of evidence concerning complications was rated low due to serious risk of bias and strongly suspected publication bias due to selective reporting.

Figure 5. Forest plot of complications: Wound infections and deep venous thrombosis for combined anteroposterior approach with a titanium cage versus solely posterior fixation



Surgical parameters


Only the RCT's reported surgical parameters. Operation time was significantly shorter in the posterior group compared to the antero-posterior group (MD 141.61 minutes; 95% CI: 125,47 – 157,74; $P < 0,01$) (Figure 6a). Heterogeneity was not important ($I^2 = 0\%$, $X^2: P = 0,69$).

Intraoperative blood loss was significantly less in the posterior group compared to the antero-posterior group (MD 515,97 mL; 95% CI: 297,54-734,41; $P < 0,01$) (Figure 6b). With substantial heterogeneity ($I^2 = 71\%$, $X^2: P = 0,07$).

Hospital stay was significantly shorter in the posterior group compared to the antero-posterior group (MD 6,21 days; 95% CI: 3,41 – 9,02; $P < 0,01$) (Figure 6c), with moderate heterogeneity ($I^2 = 54\%$, $X^2: P = 0,14$).

All quality of evidence concerning surgical parameters was graded high.

Figure 6. Forest plots of (a) Operation time (min), (b) Peroperative blood loss (mL), and (c) Hospital stay (days) for combined anteroposterior approach with a titanium cage versus solely posterior fixation



Grading of evidence

Quality rating of the evidence of each outcome with the Grade approach and additional comments and explanations are presented in Figure 7.

Figure 7. Summary of evidence graded using the GRADE approach

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	№ of participants (studies)	Quality of the evidence (GRADE)	Comments
	Risk with posterior stabilization alone	Risk with Antero-posterior stabilization using a titanium cage				
Neurologic improvement on Frankel scale (Neurologic) assessed with: Frankel scale follow up: mean 59 months	739 per 1.000	850 per 1.000 (680 to 1.000)	RR 1.15 (0.92 to 1.43)	45 (2 RCTs)	⊕⊕⊕⊕ HIGH	
Blood loss assessed with: milliliters	The mean blood loss was 603.5 mL	The mean blood loss in the intervention group was 515,97 mL higher (297,54 higher to 734,41 higher)	-	84 (2 RCTs)	⊕⊕⊕⊕ HIGH	
Operation time assessed with: minutes	The mean operation time was 105 minutes	The mean operation time in the intervention group was 141,61 minutes more (125,47 more to 157,74 more)	-	84 (2 RCTs)	⊕⊕⊕⊕ HIGH	
Hospital stay assessed with: days	The mean hospital stay was 10 days	The mean hospital stay in the intervention group was 6,21 days more (3,41 more to 9,02 more)	-	84 (2 RCTs)	⊕⊕⊕⊕ HIGH	
VAS post op assessed with: Regular VAS and VAS-spine follow up: mean 53 months	-	-	-	134 (3 RCTs) ¹	⊕○○○ VERY LOW 2,3,4,5	Rule of thumb considering SMD: 0.2 represents a small effect, 0.5 a moderate effect, and 0.8 a large effect. Which means that the anteroposterior approach has a moderate effect on lowering VAS score.

Table continues on next page

VAS post op - RCT assessed with: Regular VAS follow up: mean 59 months	-	-	-	84 (2 RCTs)	⊕⊕⊕○ MODERATE 4	Considering only the RCT's included; the anteroposterior approach has a fairly large effect on lowering the VAS score.
Correction loss in degrees follow up: mean 53 months	The mean correction loss in degrees was 6.6 degrees	The mean correction loss in degrees in the intervention group was 2,5 degrees lower (6,56 lower to 1,55 higher)	-	134 (3 RCTs) ¹	⊕○○○ VERY LOW 2,5,6,7	
Correction loss - RCT follow up: mean 59 months	The mean correction loss - RCT was 7 degrees	The mean correction loss - RCT in the intervention group was 4,59 degrees fewer (6,95 fewer to 2,22 fewer)	-	84 (2 RCTs)	⊕⊕⊕⊕ HIGH 8	
Complications: infections and deep venous thrombosis	70 per 1.000	144 per 1.000 (39 to 531)	RR 2.06 (0.56 to 7.61)	84 (2 RCTs)	⊕⊕○○ LOW ⁹	

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio; MD: Mean difference; SMD: Standardised mean difference

GRADE Working Group grades of evidence

High quality: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate quality: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low quality: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low quality: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

- 2 RCT's and 1 Cohort study
- Including one cohort study which scores serious risk of allocation bias because more severely injured patients were allocated to anteroposterior group, overall risk of bias in RCT's is low to unclear with both studies scoring high risk of selective reporting and Wang et al. did not blind participants.
- Includes VAS and VAS-spine
- RCT's show inconsistent or opposite evidence
- All three studies include a wide variety of patients, but the cohort study shows a serious allocation bias allocating only most severe patients to anteroposterior group.
- Cohort study by Weiner et al. shows opposite effect opposed to the RCT's.
- Cohort study of Weiner et al shows a relatively wide confidence interval
- Korovessis et al. used Gardner angle to measure kyphosis while Wang et al. used Cobb-angle, however method of measurement was consistent on both time points so the difference in measurement between time points (correction loss) is comparable.
- High risk of bias due to selective reporting for both studies, Wang et al. report far less complications compared to Korovessis et al. while they do a five year follow-up.

Discussion

The main indication for additional anterior stabilization is to provide support in the anterior column in order to prevent secondary kyphosis and posterior instrumentation failure. Our systematic review shows more persistent kyphosis correction using an additional anterior cage. No difference between groups was seen in pain scores, neurologic improvement, deep venous thrombosis and wound infections. Operation time, blood loss and hospital stay did increase, as expected, in the antero-posterior group.

Kyphosis and implant failure

Independent of fracture location (T12-L2 vs L2-L4), the RCTs reported significant less correction loss using a cage. The cohort study^[31] however contradicted this and reported a high prevalence of cage subsidence. The authors attributed the cage subsidence to small endplate surfaces of their used expendable cages. Resulting in frequent intraoperative damage of vertebral endplates during the cage distraction. Another possible explanation is that no additional anterolateral plating was used which could provide additional anterior support. Cage subsidence is an important concern, although no correlation with quality of life was found^[31]. Observational studies reported low rates of cage subsidence and good kyphosis correction and outcome^[32-37]. We expect consistent kyphosis correction when using a cage with additional plating.

While anterior stabilization is developed to prevent posterior implant failure, only Wang et al.^[30] reported posterior implant failure, in the posterior group. An explanation could be that this was the only study that used solely posterior fixation on fractures with a LSC ≥ 7 . Korovessis et al.^[29] included fractures with a LSC ≤ 6 , which according to the LSC do not need additional anterior stabilization. They, in accordance, concluded that solely posterior fixation was associated with less surgical trauma and provided better clinical outcomes. In addition, they advised to only use an additional cage in the case of high comminution and angulation. Wang et al.^[30] included fractures with a LSC over and below 7 although results were not stratified accordingly. They concluded to only use a combined approach in very comminuted unstable fractures or with posterior column injury. Due to the large selection bias in the study of Weiner et al.^[31] it is not possible to make a recommendation on fracture type based on this study. In conclusion, it seems that a small proportion of fractures with high comminution or instability are indicated for additional anterior stabilization using a cage. While it is not possible to appoint specific fracture types based on these studies, the LSC could be a good indicator.

Pain and neurologic improvement

Pain might be the result of posttraumatic kyphosis, neurologic injury or post-donor site pain, and complications resulting from invasive surgery. Our systematic review shows no significant difference for either group when taking all studies into account, with very low quality of evidence. The randomized studies showed conflicting results. Korovessis et al.^[29] assigned the higher pain score in the antero-posterior group to the more invasive additional surgery. This is likely if the pain score was measured directly postoperative, however the measuring moment remains unclear. The other studies^[30, 31], both reporting less pain for the antero-posterior group, measured pain after 70 and 41 months, respectively. Interestingly, Weiner et al.^[31] report less pain in the antero-posterior group while they report 85% cage subsidence. The antero-posterior group shows a trend towards less pain. Observational studies confirm this, reporting a large improvement of pain scores on long term with the combined approach^[34, 35, 38].

Neurologic damage is reported to be a main cause of long-term persisting pain^[9]. Neurologic improvement is reported after solely posterior fixation^[39] and even after non-operative management^[9]. Our study shows that neurologic improvement is independent of using a cage, so that neurologic injury by itself is not an indication for additional anterior stabilization. However, an anterior approach could be indicated when there is significant ventral bone impingement.

Additional risks of anterior surgery

We found low quality of evidence of a slightly lower risk of wound infection and deep venous thrombosis for the posterior group, although this was not significant. Observational studies reported very few complications after anterior approach^[38] and few posterior wound infections^[35]. While Schnake et al.^[40] reported anterior surgery related complications of 37.5%, 26% was related to the thoracotomy itself and most complications were not clinically significant.

We also found increased perioperative characteristics with additional anterior surgery. However no complications needing re-operation were reported. We therefore think increased maintenance of kyphosis correction is more important in specific fractures at high risk of kyphosis. While it remains important to weight accompanying risks for every patient, minimally invasive but technically demanding thoracoscopic techniques can decrease blood transfusions, pain and hospital stay^[11, 12, 41].

New techniques

Minimally invasive vertebral body stenting (VBS) also provides anterior stability, however the few short term results published are not yet impressive^[13, 42] and applicability to severe

traumatic fractures is unclear. If the stability provided is comparable to a titanium cage it might be an alternative for anterior stabilization.

Functional outcomes

One study^[31] reported no differences between groups on all SF-36 domains, while Korovessis et al.^[29] reported significant improvement on the domains physical and bodily pain in the posterior group. They attributed this to the increased morbidity of anterior surgery, although it is not clear when the SF-36 was assessed. Weiner et al.^[31] reported no difference between groups on functional outcomes of Low Back Outcome score and Oswestry Disability Index.

Strengths and Limitations

There is limited evidence available about the additional value of an anterior cage after posterior stabilization and studies that are available describe small patient groups. Therefore, we included articles in all available languages (including Chinese) and also included cohort studies comparing both groups. While cohort studies may introduce selection bias, stratified forest plots and grading of evidence are reported. Although we think the presented results are not influenced by different fracture locations, ideally results are specified according to fracture level (e.g. thoracolumbar vs lumbar) which was not possible due to the few available studies. It is possible that studies did not report all results in full extent. Kyphosis was measured as Cobb angle and Gardner angle, although these were comparable because they assessed relative kyphosis difference over time. Expandable and titanium cages were used, but literature shows no difference^[43].

In conclusion, evidence suggests that patients with a highly comminuted or unstable fracture benefit from antero-posterior stabilization with a titanium cage due to maintenance of kyphosis correction. Neurologic injury is not a primary indication for an additional anterior cage. There is a risk of cage subsidence and increased perioperative risks have to be considered. Prospective studies focusing on specific patients that are indicated for an additional cage could provide stronger evidence.

References

- McLain RF, Sparling E, Benson DR. Early failure of short-segment pedicle instrumentation for thoracolumbar fractures. A preliminary report. *J Bone Joint Surg Am.* 1993;75(2):162-7.
- Butt MF, Farooq M, Mir B, Dhar AS, Hussain A, Mumtaz M. Management of unstable thoracolumbar spinal injuries by posterior short segment spinal fixation. *Int Orthop.* 2007;31(2):259-64.
- Schnee CL, Ansell LV. Selection criteria and outcome of operative approaches for thoracolumbar burst fractures with and without neurological deficit. *J Neurosurg.* 1997;86(1):48-55.
- Verlaan JJ, Diekerhof CH, Buskens E, van der Tweel I, Verbout AJ, Dhert WJ, et al. Surgical treatment of traumatic fractures of the thoracic and lumbar spine: a systematic review of the literature on techniques, complications, and outcome. *Spine (Phila Pa 1976).* 2004;29(7):803-14.
- McCormack T, Karaikovic E, Gaines RW. The load sharing classification of spine fractures. *Spine (Phila Pa 1976).* 1994;19(15):1741-4.
- Almaiman M, Al-Bargi HH, Manson P. Complication of anterior iliac bone graft harvesting in 372 adult patients from may 2006 to may 2011 and a literature review. *Craniomaxillofac Trauma Reconstr.* 2013;6(4):257-66.
- Qi L, Liu Y, Li H, Zhang Y. Results of 10-year follow-up of the iliac donor site of graft patients. *J Int Med Res.* 2014;42(6):1348-52.
- Pflugmacher R, Schleicher P, Schaefer J, Scholz M, Ludwig K, Khodadadyan-Klostermann C, et al. Biomechanical comparison of expandable cages for vertebral body replacement in the thoracolumbar spine. *Spine (Phila Pa 1976).* 2004;29(13):1413-9.
- Reinhold M, Knop C, Beisse R, Audige L, Kandziora F, Pizanis A, et al. Operative treatment of 733 patients with acute thoracolumbar spinal injuries: comprehensive results from the second, prospective, Internet-based multicenter study of the Spine Study Group of the German Association of Trauma Surgery. *Eur Spine J.* 2010;19(10):1657-76.
- Baulot E, Trouilloud P, Ragois P, Giroux EA, Grammont PM. [Anterior spinal fusion by thoracoscopy. A non-traumatic technique]. *Rev Chir Orthop Reparatrice Appar Mot.* 1997;83(3):203-9.
- Ragel BT, Amini A, Schmidt MH. Thoracoscopic vertebral body replacement with an expandable cage after ventral spinal canal decompression. *Neurosurgery.* 2007;61(5 Suppl 2):317-22; discussion 22-3.
- Ragel BT, Kan P, Schmidt MH. Blood transfusions after thoracoscopic anterior thoracolumbar vertebrectomy. *Acta Neurochir (Wien).* 2010;152(4):597-603.
- Diel P, Roder C, Perler G, Vordemvenne T, Scholz M, Kandziora F, et al. Radiographic and safety details of vertebral body stenting: results from a multicenter chart review. *BMC Musculoskelet Disord.* 2013;14:233.
- Dai LY, Jiang LS, Jiang SD. Conservative treatment of thoracolumbar burst fractures: a long-term follow-up results with special reference to the load sharing classification. *Spine (Phila Pa 1976).* 2008;33(23):2536-44.
- Wood KB, Buttermann GR, Phukan R, Harrod CC, Mehbod A, Shannon B, et al. Operative compared with nonoperative treatment of a thoracolumbar burst fracture without neurological deficit: a prospective randomized study with follow-up at sixteen to twenty-two years. *J Bone Joint Surg Am.* 2015;97(1):3-9.
- Aono H, Tobimatsu H, Ariga K, Kuroda M, Nagamoto Y, Takenaka S, et al. Surgical outcomes of temporary short-segment instrumentation without augmentation for thoracolumbar burst fractures. *Injury.* 2016.
- Kallemeier PM, Beaubien BP, Buttermann GR, Polga DJ, Wood KB. In vitro analysis of anterior and posterior fixation in an experimental unstable burst fracture model. *J Spinal Disord Tech.* 2008;21(3):216-24.
- Wilke HJ, Kemmerich V, Claes LE, Arand M. Combined anteroposterior spinal fixation provides superior stabilisation to a single anterior or posterior procedure. *J Bone Joint Surg Br.* 2001;83(4):609-17.
- Bence T, Schreiber U, Grupp T, Steinhauser E, Mittelmeier W. Two column lesions in the thoracolumbar junction: anterior, posterior or combined approach? A comparative biomechanical in vitro investigation. *Eur Spine J.* 2007;16(6):813-20.
- Cripton PA, Jain GM, Wittenberg RH, Nolte LP. Load-sharing characteristics of stabilized lumbar spine segments. *Spine (Phila Pa 1976).* 2000;25(2):170-9.
- Ringel F, Stoffel M, Stuer C, Totzek S, Meyer B. Endoscopy-assisted approaches for anterior column reconstruction after pedicle screw fixation of acute traumatic thoracic and lumbar fractures. *Neurosurgery.* 2008;62(5 Suppl 2):ONS445-52; discussion ONS52-3.
- Xu GJ, Li ZJ, Ma JX, Zhang T, Fu X, Ma XL. Anterior versus posterior approach for treatment of thoracolumbar burst fractures: a meta-analysis. *Eur Spine J.* 2013;22(10):2176-83.
- Stancic MF, Gregorovic E, Nozica E, Penezic L. Anterior decompression and fixation versus posterior reposition and semirigid fixation in the treatment of unstable burst thoracolumbar fracture: prospective clinical trial. *Croat Med J.* 2001;42(1):49-53.
- P PO, Tuinebreijer WE, Patka P, den Hartog D. Combined anterior-posterior surgery versus posterior surgery for thoracolumbar burst fractures: a systematic review of the literature. *Open Orthop J.* 2010;4:93-100.
- Been HD, Bouma GJ. Comparison of two types of surgery for thoraco-lumbar burst fractures: combined anterior and posterior stabilisation vs. posterior instrumentation only. *Acta Neurochir (Wien).* 1999;141(4):349-57.
- Nakashima H, Imagama S, Yukawa Y, Kanemura T, Kamiya M, Deguchi M, et al. Comparative study of 2 surgical procedures for osteoporotic delayed vertebral collapse: anterior and posterior combined surgery versus posterior spinal fusion with vertebroplasty. *Spine (Phila Pa 1976).* 2015;40(2):E120-6.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 2009;6(7):e1000100.
- Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol.* 2005;5:13.
- Korovessis P, Baikousis A, Zacharatos S, Petsinis G, Koureas G, Iliopoulos P. Combined anterior plus posterior stabilization versus posterior short-segment instrumentation and fusion for mid-lumbar (L2-L4) burst fractures. *Spine (Phila Pa 1976).* 2006;31(8):859-68.
- Wang J, Liu P. Analysis of surgical approaches for unstable thoracolumbar burst fracture: minimum of five year follow-up. *J Pak Med Assoc.* 2015;65(2):201-5.
- Weiner X, El Saman A, Ruger F, Laurer H, Marzi I. [Impact of surgical strategy on quality of life and radiological outcome in traumatic fractures of the thoracolumbar spine]. *Z Orthop Unfall.* 2013;151(3):264-71.
- Theologis AA, Tabaraee E, Toogood P, Kennedy A, Birk H, McClellan RT, et al. Anterior corpectomy via the mini-open, extreme lateral, transpoas approach combined with short-segment posterior fixation for single-level traumatic lumbar burst fractures: analysis of health-related quality of life outcomes and patient satisfaction. *J Neurosurg Spine.* 2016;24(1):60-8.
- Kalicinski M, Szczesniak A, Kalisz J, Tesiorowski M. Anterior fixation of thoracolumbar traumatic spinal injuries. *Ortop Traumatol Rehabil.* 2015;17(1):7-20.
- Choi JI, Kim BJ, Ha SK, Kim SD, Lim DJ, Kim SH. Single-Stage Transpedicular Vertebrectomy and Expandable Cage Placement for Treatment of Unstable Mid and Lower Lumbar Burst Fractures. *J Spinal Disord Tech.* 2014.

35. Graillon T, Rakotozanany P, Blondel B, Adetchessi T, Dufour H, Fuentes S. Circumferential management of unstable thoracolumbar fractures using an anterior expandable cage, as an alternative to an iliac crest graft, combined with a posterior screw fixation: results of a series of 85 patients. *Neurosurg Focus*. 2014;37(1):E10.
36. Ray WZ, Krisht KM, Dailey AT, Schmidt MH. Clinical outcomes of unstable thoracolumbar junction burst fractures: combined posterior short-segment correction followed by thoracoscopic corpectomy and fusion. *Acta Neurochir (Wien)*. 2013;155(7):1179-86.
37. Shawky A, Al-Sabroun AM, El-Meshtawy M, Hasan KM, Boehm H. Thoracoscopically assisted corpectomy and percutaneous transpedicular instrumentation in management of burst thoracic and thoracolumbar fractures. *Eur Spine J*. 2013;22(10):2211-8.
38. Payer M. Unstable burst fractures of the thoraco-lumbar junction: treatment by posterior bisegmental correction/fixation and staged anterior corpectomy and titanium cage implantation. *Acta Neurochir (Wien)*. 2006;148(3):299-306; discussion
39. Khare S, Sharma V. Surgical outcome of posterior short segment trans-pedicle screw fixation for thoracolumbar fractures. *J Orthop*. 2013;10(4):162-7.
40. Schnake KJ, Stavridis SI, Kandziora F. Five-year clinical and radiological results of combined anteroposterior stabilization of thoracolumbar fractures. *J Neurosurg Spine*. 2014;20(5):497-504.
41. Kim DH, Jahng TA, Balabhadra RS, Potulski M, Beisse R. Thoracoscopic transdiaphragmatic approach to thoracolumbar junction fractures. *Spine J*. 2004;4(3):317-28.
42. Werner CM, Osterhoff G, Schlickeiser J, Jenni R, Wanner GA, Ossendorf C, et al. Vertebral body stenting versus kyphoplasty for the treatment of osteoporotic vertebral compression fractures: a randomized trial. *J Bone Joint Surg Am*. 2013;95(7):577-84.
43. Eleraky MA, Duong HT, Esp E, Kim KD. Expandable versus nonexpandable cages for thoracolumbar burst fracture. *World Neurosurg*. 2011;75(1):149-54.