

VU Research Portal

Kids in Action: is engaging children as co-researchers key to promoting healthy physical activity and dietary behaviors?

Anselma, M.

2020

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Anselma, M. (2020). *Kids in Action: is engaging children as co-researchers key to promoting healthy physical activity and dietary behaviors?*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


E-mail address:

vuresearchportal.ub@vu.nl



Chapter 3

Kids in Action: the protocol of a youth participatory action research project to promote physical activity and dietary behavior



Manou Anselma, Teatske M. Altenburg, and Mai J. M. Chin A Paw

Based on the publication in BMJ Open (2019), 9.

Abstract

Introduction In this study, researchers collaborate with children from a deprived neighborhood in Amsterdam in developing, implementing and evaluating interventions targeting their health behaviors. This Youth Participatory Action Research (YPAR) project focuses on the promotion of physical activity and healthy dietary behavior.

Methods and analysis This study is a controlled trial using participatory methods to develop interventions together with 9-12-year-old children. At four primary schools in a deprived neighborhood in Amsterdam an 'Action Team' is installed: a group of six to eight children who actively participate as co-researchers in developing, implementing and evaluating interventions. An academic researcher facilitates the participatory process. Four control schools, also located in deprived areas in and around Amsterdam, continue with their regular curriculum and do not participate in the participatory process. For the effect evaluation, physical activity and sedentary behavior is assessed using: accelerometers and self-reporting; dietary behavior using self-reporting; and motor fitness (strength, flexibility, coordination, speed, endurance) using the MOPER fitness test. Effectiveness of the interventions is evaluated by multi-level regression analysis. The process of co-creating interventions and the implemented interventions is continually evaluated during meetings of the Action Teams and with children participating in the interventions. Empowerment of children is evaluated during focus groups. Summaries and transcripts of meetings are coded and analyzed to enrich children's findings.

Ethics and dissemination The Medical Ethics Committee of the VU Medical Center approved the study protocol (2016.366).

Protocol registration The study protocol has been registered at the Dutch trial registration www.trialregister.nl under number TC=6604.

3.1. Introduction

The number of children with overweight or obesity is growing worldwide and this public health problem is high on municipal and governmental agendas. This is no different in the Netherlands, where in 2016 on average there were 10.7% of the children between 8 and 12 years old with overweight/obesity[1]. In urban areas such as Amsterdam, the rates exceed the country's average, with prevalence rates of overweight/obesity of 12.8% among 5-year-olds and 20.9% among 10-year olds[2]. Even though the overweight numbers are stabilizing, health inequalities still exist[3]: children with overweight or obesity are not only disproportionately divided geographically, but also across income and ethnic groups[4-6]. Looking at race/ethnicity, children from minority groups show higher overweight/obesity rates than children from a majority group[4, 7, 8]. For example, in Amsterdam 10.4% of 5-year-old children with a Dutch ethnicity have overweight while this is almost 30% in 10-year-old children with a non-Western background³. In relation to income groups, in the Netherlands in the age category 4-25-year-olds, 11.2% of the highest income group had overweight, versus 18.0% of the lowest income group[9]. Similarly, in 2017 in Amsterdam 30.1% of the 10-year-old children with a very low socioeconomic status (SES) had overweight, versus 9.8% of the 10-year old children with a very high SES[3]. Importantly, children with overweight are at high risk of remaining overweight and are therefore also at higher risk for chronic illnesses during childhood and in their adult life[10]. This is why prevention of overweight in children is a priority for many health organizations, municipalities and ministries[11, 12].

Many interventions have been developed and implemented to prevent childhood obesity, but most show disappointing effects[13, 14]. Pivotal in childhood obesity prevention is improving dietary behavior, physical activity and sedentary behavior[15, 16], but this is challenging[17-19]. Strikingly, the most affected group of children – i.e. from families with a low SES and from non-Western backgrounds – is most difficult to reach through interventions[20], thereby maintaining or even widening health inequalities[21, 22]. One reason why these interventions show low participation and effectiveness in this target group could be because the target group is seldom involved in the

development of the interventions[23]. Involving the target group is essential to connect to their needs and interests[24], as this influences the reach and effectiveness of the intervention. Therefore, in the current research project – 'Kids in Action' – children from a deprived neighborhood are engaged as co-researchers, i.e. applying Youth Participatory Action Research (YPAR). Children not only co-create interventions to improve their lifestyle and that of their peers and family members, but also collaborate in the implementation and evaluation of these interventions. To structure this process, the systematic Intervention Mapping (IM) methodology is applied alongside YPAR. Through six iterative steps the IM protocol guides health promoters in the development of evidence-based interventions to change behavior[25, 26]. Combining IM and YPAR ensures that the co-created interventions are appropriate to the interests and needs of the children, but also build on existing evidence. The application of IM alongside YPAR is a novel approach which we iteratively shape during this study.

Aims and objectives

The overall aim of the 'Kids in Action' study (April 2016–November 2019) is to develop, implement and evaluate interventions that stimulate a healthy lifestyle to reduce health inequalities in children from a low SES neighborhood in collaboration with the children themselves. This study builds on a participatory needs assessment that was conducted in the same neighborhood[27]. From this needs assessment, two main needs were identified: to improve physical activity and a healthy diet. The organized activities should be offered at a low price and at a nearby location, the education concerning a healthy diet should be organized in a fun and practical manner.

The primary objective of this study is to evaluate whether designing interventions in collaboration with children can lead to interventions that are more effective in improving children's physical activity and dietary behavior.

The secondary objective of this study is to evaluate the process of combining YPAR with IM. This includes evaluating the effects of participating in the YPAR process on the empowerment of children and the judgement of children and other stakeholders of interventions that were co-developed

by their peers.

3.2. Methods

The Medical Ethics Committee of the VU University Medical Center approved the study protocol (2016.366).

3.2.1. Participatory Action Research

Participatory Action Research (PAR) aims to 'improve health and reduce health inequities' by working together with the community and consequently empowering the community by getting them to improve their own health[28]. Throughout the entire process of developing, implementing and evaluating interventions, community members are involved as co-researchers and highly valued as experts of their own lives and experiences. At the same time, the community is empowered and experiences more ownership over their lives and livelihood.

This study specifically works together with children in the PAR process. YPAR engages youth as co-researchers in the research process. In this process, children identify problems in their living environment and become empowered to do something about it[29-32]. Children learn research skills so they can participate in research and have shared power over the research- and decision-making processes[30, 32, 33].

3.2.2. Patient and public involvement

This study is initiated by academic researchers and a community organization. The municipality advises on the selection of the intervention neighborhood, to recruit a neighborhood with high health needs that can benefit from the project. As this study is informed by a participatory needs assessment (see section 1.1)[27], the objectives and outcome measures of this study are determined in collaboration with children, parents and professionals working with children in the neighborhood. The design of the study and recruitment procedures are decided by the academic researchers. The conduct of the study, the development of interventions and the dissemination of the

results to the study participants and other relevant stakeholders, is decided together with the children.

3.2.3. Participants

The four intervention schools are all situated in one deprived neighborhood in Amsterdam, where in 2015-2016 over 50% of the residents had a non-Western background, 27% of the 10-year-olds were overweight/obese and in 2014 31% of the children under 18 years old grew up in a household defined as low-income[34-36]. Potential control schools are selected from different neighborhoods but with similar characteristics regarding overweight/obesity rates, household income and cultural background. Participants in this study are children from four intervention schools and four control schools in deprived neighborhoods in Amsterdam, the Netherlands. The intervention schools participate in the YPAR process, including implementing and evaluating the developed interventions. The control schools only participate in the measurements for the effect evaluation.

Recruitment


Following selection of a neighborhood with high health needs, the intervention schools are contacted by the municipality to inform them about the project and to ask them if they are willing to participate. After the schools agree to participate, the main researcher contacts the schools to give them more information about the project. Control schools in the area of Amsterdam are contacted by the main researcher in a random order via e-mail or telephone until four schools are found that are willing to participate as control schools. Control schools are offered a presentation about the research results after the study is finished.

All 9-12-year-old children (i.e. children of the three highest grades in primary school) of the four intervention and four control schools are eligible to participate in the effect measurements of the project. For the YPAR process, children from intervention schools are invited to collaborate with academic researchers in co-researcher groups, named 'Action Teams'. At each of the four intervention schools one Action Team is formed. For both

the effect measurements and the Action Teams, every year new children can participate as the highest grade leaves the school and new children enter the third-highest grade. All children receive an information letter for themselves and for their parents about the measurements and the Action Team. Attached to the information letters for parents is an informed consent letter that at least one of the parents has to sign if they agree to the participation of their child. At all schools, the researcher explains the project in all classes and encourages children to participate, before handing out the information letters. Children who participate in the measurements and/or in the Action Team receive a small gift.

3.2.4. Procedures

This section describes the five phases of the 'Kids in Action' project. See Figure 1 for an outline.



Continuous activities




Figure 1: Outline of the 'Kids in Action' project

Phase 1: Creating partnerships

The first phase consists of creating partnerships with the schools and other stakeholders in the area, such as social workers, organizers of after-school activities and the community centers. Together with these stakeholders, a project group is started that meets every three months to discuss running projects in the neighborhood and how partners can collaborate. In this phase, an IM expert group is also formed, to advise on how YPAR and IM should be combined throughout all phases of the study.

Phase 2: Formation of Action Teams

In the second phase of the project, the Action Teams are formed. Each of the Action Teams consists of six to eight children, an academic researcher and a research assistant. Meetings with the principals of the four intervention schools are planned to decide upon recruitment methods for the Action Teams and to schedule the meetings. All interested 9-12-year-old children can sign up for the Action Teams. This approach may lead to bias as only children interested in health may sign up, but limits bias that would occur if teachers select the children for the Action Teams (i.e. only the high-performers might be selected). Subsequently, the Action Teams are formed and a general outline of the meetings is developed. In this phase, the baseline effect measurement (T₀) is executed.

Phase 3: Intervention development

In the third phase, the meetings of the Action Teams take place. The meetings with the Action Teams are ideally held biweekly during school hours for one hour. Despite not all schools agreeing to this in the needs assessment, the researchers try to schedule meetings during school hours to raise the children's motivation for participation[27]. If the schools do not agree with this, meetings are held weekly for 45 minutes, followed by a 45-minute sports session[27].

The first three to four meetings are used to verify the data that was gathered in a participatory needs assessment and to decide on determinants that the interventions need to focus on[27]. In the rest of the meetings

(approximately 10 per year) we develop interventions together with the children targeting children's physical activity and healthy dietary habits. The type of the interventions (e.g. environmental changes, organizational changes, or educational approaches) is dependent on this collaborative process. Throughout these meetings, capacity building takes place to help the children through the process of intervention development. Children learn for example about formulating a research question, different kinds of research methodologies, how to analyze qualitative data, how to translate this data into intervention ideas and practical steps that need to be taken when developing intervention plans. At the end of phase 3, pilots of the first intervention activities are carried out. The Action Teams are also asked to identify 'Champions', i.e. people who can help them with the development and implementation of the pilots. A champion is a well-known community member such as a teacher, sports coach or family member. Children discuss who they think is suitable to assist them with a specific intervention and subsequently ask the champions to fulfil this task. The results of this phase (i.e. the needs assessment, the intervention ideas and results of the pilots) are discussed with the stakeholders in the project group to make sure the interventions become a joint and sustainable effort. At the end of the year, a focus group with the Action Teams and their peers is held to discuss the feeling of empowerment that the children of the neighborhood experience as part of the process evaluation.

Phase 4: Implementation and evaluation of interventions

In the beginning of phase 4, new Action Teams are recruited/formed. Children who were in the Action Teams of the previous year can still participate and are approached first. With the new Action Teams, meetings are planned monthly. Champions are involved and asked to participate in the meetings when appropriate. Together with stakeholders from the project group and the Action Teams, the implementation plans are finalized and subsequently the interventions are implemented. In order to offer sustainable interventions we looked for partners within the community whose job description aligns with providing the intervention. Depending on the type of intervention, implementers could be dieticians, sports coaches

or supermarkets in the community. Once the interventions are implemented, the meetings are used to evaluate the interventions. If the Action Teams feel the interventions are going well, they are encouraged to develop and implement additional intervention activities that focus on other determinants or a different subgroup[37]. At the end of the year, focus groups are held focusing on empowering children and evaluating interventions. The first effect measurement (T1) is also executed in this phase.

Phase 5: Gradual transfer of responsibilities

In phase 5, responsibilities are gradually transferred to the identified champions. Specific plans are made together with the champions and other stakeholders to continue the interventions and participatory process after this project has ended. The meetings with the (new) Action Teams continue to take place every month, and are used to evaluate and, if necessary, adapt the interventions and discuss new ideas for interventions. The post-intervention effect measurement (T2) is executed in this phase. The study ends in November 2019.

3.2.5. Measurements

Effect evaluation

The primary outcomes of this study include measures of dietary behavior (consumption of snacks and sugar-sweetened beverages), physical activity (total MVPA time, time spent playing outside, time spent participating in sports), sedentary behavior (total sedentary time and screen time), self-rated health, and physical fitness. Dietary behavior, physical activity and screen behavior is measured by self-report. Additionally, physical activity and sedentary behavior is measured using an accelerometer. Motor fitness is measured using the MOPER fitness test. In the first school year (T0), questionnaire and accelerometer data are gathered in the period September-October 2016. The fitness tests take place in March-April 2017, 2018 and 2019.

Questionnaire

A questionnaire is developed containing questions on: the number of small (e.g. crisps, nuts, chocolate) and large (e.g. hamburger, fries, pizza) snacks children eat; the number of sugar-sweetened beverages they drink; their sports and outdoor play participation; their attitude towards sports and outdoor play; their screen behavior; and their self-rated health. The questionnaire is based on validated items from the ENERGY child questionnaire[38], the DOiT questionnaire[39], and the EuroQol[40]. Table 1 presents the questionnaire items, and the validity and reliability of the original items.

The children fill in the questionnaire during school hours, in the presence of a researcher who explains the procedure of completing the questionnaire before handing out the questionnaires. The children are requested to go through the questionnaire section by section, with the researcher giving a short explanation about each section before the participants fill in that specific section. In this way, examples can be given, for example by showing different sizes of soda cans, and all participants finish at the same time. The questionnaire takes approximately 40 minutes to complete.

Data entry of the multiple-choice questions is done through digital scanning and transferred into SPSS by an independent organization. Qualitative data are manually entered in SPSS.

Accelerometer

Physical activity and sedentary behavior is objectively assessed by ActiGraph GT3X+ accelerometers. The children receive instructions and the accelerometers from an academic researcher after filling in the questionnaire. Children are asked to wear the small and light-weight (4.6 x 3.3 x 1.5 cm; 19 grams) accelerometer on the right hip for eight consecutive days during all waking hours except for water-based activities.

The children receive the accelerometer after completing the questionnaire. The children also receive a diary in which the instructions are summarized and they can write down when and why they did not wear the accelerometer,

if applicable. Additionally, they are asked to write down the time they went to bed. All children who participate in the questionnaire- and accelerometer measurements receive a small present after returning the accelerometer. Additionally, at each school there is one prize for a participant who wore the accelerometer properly (seven days, at least ten hours) and recorded their data correctly in their diary.

Data are downloaded from the accelerometers into the ActiLife program in 15 second epochs. Accelerometer data is analyzed using a customized software program developed in R. We select a cut point of 100 counts per minute (cpm) for sedentary behavior[41, 42] and a cut point of 3000 cpm for MVPA[43]. Non-wear time is defined as a period of ≥ 60 minutes of consecutive zeros[44]. For inclusion in the data analysis, each participant needs a minimum of six days with at least eight valid hours, including at least one weekend day[44]. Data is analyzed on total time spent in MVPA and sedentary, and time in bouts spent in MVPA and sedentary.

MOPER

Children's motor fitness is measured using the Motor Performance Test (MOPER). The MOPER tests speed, flexibility, endurance, coordination and strength by means of eight tests[45]. For practical reasons, the arm pull and 12-minute endurance test have been replaced, leading to the following tests: 1) hang as long as possible on a horizontal bar with flexed arms; 2) jump as high as possible from a standing position; 3) run 10x5 meters as fast as possible; 4) reach as far as possible from a sitting position; 5) hand grip strength measured using a dynamometer[46, 47] (instead of arm pull); 6) lie on their back and lift their extended legs ten times as fast as possible; 7) tap two plates which are 75 cm apart with the preferred hand 50 times as fast as possible; and 8) shuttle run test[48, 49] (instead of 12-minute endurance test). Children can do tests one and eight once. Tests two and five are executed twice, but when the difference between one and two is more than 10%, a third try is performed. The highest score is used. The other tests are performed twice and the highest score is used. The first seven activities of the MOPER test are executed during one Physical Education (PE) class by the PE teacher together with five or six research assistants. The

PE teacher conducts the shuttle run test in a separate PE class. All research assistants and PE teachers are trained by an academic researcher on how the tests should be executed. At the end of the study, or when children from the highest grade leave the school, the PE teacher anonymously shares the results of the test. Parents receive an information letter with a passive consent form, which should be signed by at least one of the parents if they object to anonymously sharing the fitness test results of their child with the researchers.

Process evaluation

The process evaluation includes the description of the process of co-creating interventions, combining IM and YPAR, and empowerment. The PAR process is continually evaluated in the Action Team meetings, and meetings are optimized in accordance with the evaluation[28]. The academic researcher and research assistant who are part of the Action Teams evaluate after every meeting, using a reflection form consisting of a summary of the meeting, what the setting was like, the group process and a personal reflection[50-52].

The interventions are developed by combining the YPAR and IM methodologies in an iterative process and are continuously evaluated during the meetings of the Action Teams and with the children participating in the interventions. In collaboration with the Action Teams, it is determined how to evaluate the experiences of children with the interventions. The Action Teams can for example interview peers or develop a questionnaire. The goal of these evaluations is to see how their peers perceive the interventions and whether quick adaptations need to be made. At the end of each school year, focus groups are organized with children from both the Action Teams and their peers, as well as champions to reflect upon the implementation of ongoing interventions and on the empowerment process. Empowerment consists of a combination of individual, organizational, and community empowerment[53]. In our research, we mostly focus on the empowerment of children (individual), but this cannot be evaluated without taking the organizational (school) and community empowerment into account[54].

Table 1: Questionnaire items, their origin, and reliability and validity [38-40]

Questionnaire item	Question derived from	Reliability (ICC/k)/ Validity (ICC/k)
1.How many days a week do you drink sugar-sweetened beverages?	ENERGY child questionnaire	0.71/0.59
2.On a day you drink sugar-sweetened beverages, how many glasses/small bottles (250ml), cans (330ml) or big bottles (500ml) do you drink?	Combination ENERGY child questionnaire, DOiT questionnaire	ENERGY Glasses/small bottles (250ml) 0.59/0.24 Cans (330ml) 0.53/0.44 Big bottles (500ml) 0.58/-0.01 DOiT Cartons/small bottles (200ml) 0.74/0.12 Glasses (200ml) 0.45/0.47 Cans (330ml) 0.61/0.24 Bottles (500ml) 0.28/0.17
3.How many days a week do you drink energy drinks or sports drinks?	Added based on Q1	
4.On a day you drink energy drinks or sports drinks, how many small cans/bottles (250ml) or big cans/bottles (500ml) do you drink?	Added based on Q2	
5.How many school days per week do you eat sweets?	DOiT questionnaire (adapted) Original: 'How many days a week do you eat sweets?'	0.66/0.60
6.When you eat candy on a school day, how much sweets do you eat?	DOiT questionnaire	0.71/0.21
7.How many days in the weekend (Saturday/Sunday) do you eat sweets?	DOiT questionnaire (adapted) Original: 'How many days a week do you eat sweets?'	0.66/0.60

Questionnaire item	Question derived from	Reliability (ICC/k)/ Validity (ICC/k)
8. When you eat sweets on a day in the weekend, how much candy do you eat?	DOiT questionnaire	0.73/0.07
9. How many schooldays per week do you eat snacks?	DOiT questionnaire (adapted) Original: 'How many days a week do you eat snacks?'	0.50/-0.11
10. When you eat snacks on a school day, how many small and large snacks do you eat?	DOiT questionnaire	Small snacks 0.62/0.13 Large snacks 0.58/-0.08
11. How many days in the weekend (Saturday/Sunday) do you eat snacks?	DOiT questionnaire (adapted) Original: 'How many days a week do you eat snacks?'	0.50/-0.11
12. When you eat snacks in the weekend (Saturday/Sunday), how many small and large snacks do you eat?	DOiT questionnaire	Small snacks 0.53/0.44 Large snacks 0.64/0.08
13. How do you usually travel to school?	DOiT questionnaire	Not in test-retest study
14. How long does it take you to get from home to school?	DOiT questionnaire + ENERGY child questionnaire (adapted) Original: 'If you walk/bike to school, how long does it take you?'	DOiT Walking 0.65/ zero variance Biking 0.91/0.68 ENERGY Walking 0.70/0.59 Biking 0.81/0.66
15. What do you usually do when you play outside at school?	ENERGY child questionnaire	0.80/0.65
16. I like playing outside	ENERGY child questionnaire - Adapted from Q20	
17. I play outside never/1-2 times a week/3-4 times a week/5-6 times a week/every day	Added	
18. When you play outside after school, what do you do?	ENERGY child questionnaire - Adapted from Q15	

Questionnaire item	Question derived from	Reliability (ICC/k)/ Validity (ICC/k)
19. When you play outside after school, how long do you play? (fill in the number of hours per day in table)	Added	
20. I like playing sports	ENERGY child questionnaire	0.64/0.09
21. I play sports often/sometimes/never	Added	
22a. Do you participate in sports in your free time?	DOiT questionnaire (adapted)	
22b. How many times per week do you do this sport?	Original:	1. 0.98/0.86
22c. How many hours per day do you do this sport? (fill in all sports that you do, the number of times and number of hours per week in the table)	1. 'Do you participate in a sport at a sports club?' 2. 'How many hours a week do you do this sport?' 3. 'Do you participate in a second sport at a sports club?' 4. 'How many hours a week do you do this second sport?' 5. 'Do you participate in sports outside a sports club?' 6. 'How many hours a week do you do these sports?'	2. 0.94/0.78 3. 0.79/0.69 4. 0.76/0.96 5. 0.64/0.33 6. 0.64/0.45
23. About how many hours a day do you usually watch television/DVDs/movies on the tablet or iPad in your free time? (fill in the number of hours per day in table)	ENERGY child questionnaire (adapted) Original: 'About how many hours a day do you usually watch television in your free time?' (weekdays and weekend days)	Weekdays 0.67/0.63 Weekend days 0.68/0.56

Questionnaire item	Question derived from	Reliability (ICC/k)/ Validity (ICC/k)
24.About how many hours a day do you usually play games on your game computer, iPad, smartphone or surfing on the internet in your free time? (fill in the number of hours per day in table, weekdays and weekend days)	ENERGY child questionnaire (adapted) Original: 'About how many hours a day do you usually play games on a computer, or use your computer in your free time?' (weekdays and weekend days)	Weekdays 0.67/0.35 Weekend days 0.67/0.65
25.How do you rate your health today?	Euroqol EQ-5D-Y Dutch	0.83/-0.51

The focus groups consist of two exercises. The first exercise is mainly focused on individual empowerment, evaluating what children have learnt about the process of intervention development, how they see their role, and competences[55, 56]. The children can choose an intervention idea which has not been further developed yet. For this intervention they have to make a timeline with all the steps they need to take from coming up with the idea through to implementation. The researchers guide them through questions, for example: in which order do the steps need to be written down?; do they think they can execute this step by themselves?; if not, do they know where they can get help?[55]. The second exercise evaluates the organizational and community empowerment. In pairs, the children first indicate which changes happened at school or in the community; then they indicate whether children had any influence on the changes; finally, the findings are discussed in a plenary session. Again the researchers ask questions, for example: how do you feel when you have influence on changes in the community/school?; do you think children have enough influence?; would different changes have been made if children had had more influence? The findings of this focus group provide critical understanding of the environment, what children have learnt, to what extent children participate in the organizational setting and community, and what collective action has already been taken[54-56].

Of all hard-copy research data gathered in the PAR meetings, identifiable

information is removed and the data are stored in a locked cabinet at the research location until the study is completed. All online data are coded and stored on the VUmc protected drive until five years after the completion date of the study; data from the questionnaires, accelerometers, MOPER and personal data are saved with encryption. Hard-copies of the questionnaires and the audio-recordings are also stored at the VUmc until five years after the study is completed. The three researchers on this project, who are also the authors of this paper, are the only ones who have full access to the trial data. Research assistants have limited and temporary access to copies of the data.

Sample size calculation

Using a significance level of 0.05 and a power of 0.80, 180 children per group are needed to detect a difference of 0.15SD in the primary outcome variables. Taking into account dropout and clustering of data within schools we aim to include 240 children per group.

Data analysis

Effect evaluation

To test for baseline differences in the dependent variables between control and intervention groups, t-test for continuous variables and chi-square tests for categorical variables are used. Effectiveness of the interventions on dietary behavior, physical activity, sedentary behavior, physical fitness and self-rated health is evaluated using multi-level regression analysis with a 3-level structure (i.e. student, class, school) to adjust for clustering of observations. Analyses are adjusted for age, gender, ethnicity and baseline levels. Data are analyzed according to the intention-to-treat principle. All statistical analyses are performed in SPSS, using a significance level of $P < 0.05$.

Process evaluation

Evaluation of the PAR process and its meetings are mainly performed by the Action Teams themselves. The academic researcher stimulates

the children to find patterns and relations in the findings of their own research and assists in interpretation[57]. Children can for example look at the pictures they have taken and write down why they took the picture and what they want to say with the picture. Children can also write down the key issues that come up in the interviews they have conducted and see if they can identify a pattern. By giving children this role in qualitative data analysis, less misinterpretation of data occurs (than would be the case with adults trying to interpret the children's findings). In addition, all meetings are summarized and include field notes, and key meetings are fully transcribed[50]. The academic researcher analyses these transcripts to enrich the children's findings. When, for example, the children discuss the pictures they have taken, these discussions may also contain valuable information in addition to the pictures and conclusions of the children. All summaries and transcripts are coded in ATLAS.ti by two researchers to improve the reliability of the study. For the entire process evaluation, an elaborate coding scheme is produced through open coding[57]. For specific aspects like the evaluation of an intervention, coding is done separately resulting in its own coding scheme. For evaluations relating to empowerment, closed coding is used as this will be linked to a conceptual model.

3.3. Discussion

In the Kids in Action project, children are involved throughout the entire research process. This YPAR approach has previously shown promising results for communities in need with respect to researchers' understanding of the community, lowering health disparities, increasing children's skills (e.g. research skills, life skills), critical awareness, involvement and empowerment concerning community action[58-60]. In the Kids in Action project, children are not involved in the first phase of this study, in which partnerships with other stakeholders in the community have to be set up. This is because creating partnerships can be time-consuming and not very interesting for children, and we do not want to lower their spirits[37]. The partnerships are important in YPAR for creating support in the community for the study[58, 61] and are beneficial in the rest of the research process and outcomes.

A difference between this study and most YPAR studies is that 9-12-year-old children are involved as co-researchers, whereas most YPAR studies collaborate with adolescents older than twelve[62]. Younger children can be more easily distracted, have a limited attention span and might need more 'play', all of which should be taken into account when designing the meetings. Meetings should not be too long, should contain fun and playful exercises, and wording should be suitable for the children, while retaining key principles of YPAR. These principles include: sharing power between researchers and children; training children to participate in research and identify needs in their community; teaching children how to become advocates; creating ownership over the process; and creating involvement in establishing change in their community[63]. When all of this is done with care, children between nine and twelve years old are capable of joining in YPAR research[64-66].

One implication of working with 9-12-year-old children is that you often have to collaborate intensively with the schools. This could mean that changes in the planning have to be made beforehand or during the project, based on the schools' preferences, holidays and other reasons for cancelling meetings[37]. Also, the approval and assistance of schools and other community organizations are likely to be needed for implementing the interventions. Because this is a community project, all primary schools in the neighborhood are included in the intervention and randomization of schools is not possible. However, the inclusion of comparable control schools is a strength of this study as this is seldom included in PAR[67]. Another strength of this study is the combination of YPAR with IM, which makes sure that evidence-based strategies are being applied. As far as we know, this has not been done before. A challenge for all intervention studies in real life is that other initiatives can also take place in the neighborhood. This is part of usual care and can take place both in the intervention school and the control school neighborhoods, and may dilute intervention effects.

3.4. References

1. CBS, (2016). Ervaren gezondheid, zorggebruik en leefstijl bij kinderen tot 12 jaar. [Available from: <http://statline.cbs.nl/Statweb/publication/?DM=SLNL&PA=83716NED&D1=89&D2=0&D3=l&D4=0&D5=2&VW=T> (accessed 12-12 2017).
2. GGD, (2016). Gezondheid in beeld: GGD Amsterdam. [Available from: <https://amsterdam.ggdgezondheidinbeeld.nl/jeugd/> (accessed 12-12 2017).
3. Gemeente Amsterdam, (2017). Staat van gezond gewicht en leefstijl van Amsterdamse kinderen: 1-46.
4. Chi, D.L., Luu, M., and Chu, F., (2017). A scoping review of epidemiologic risk factors for pediatric obesity: Implications for future childhood obesity and dental caries prevention research. *J Public Health Dent*, 77(1): S8-S31.
5. Kumanyika, S.K., Whitt-Glover, M.C., and Haire-Joshu, D., (2014). What works for obesity prevention and treatment in black Americans? Research directions. *Obes Rev*, 15(4): 204-212.
6. Wang, Y., and Lim, H., (2012). The global childhood obesity epidemic and the association between socio-economic status and childhood obesity. *Int Rev Psychiatry*, 24(3): 176-188.
7. Taveras, E.M., Gillman, M.W., Kleinman, K., Rich-Edwards, J.W., and Rifas-Shiman, S.L., (2010). Racial/ethnic differences in early-life risk factors for childhood obesity. *Pediatrics*, 125(4): 686-695.
8. Wang, Y., and Beydoun, M.A., (2007). The obesity epidemic in the United States--gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiol Rev*, 29: 6-28.
9. CBS, (2016). Leefstijl en preventie; geslacht, leeftijd, persoonskenmerken [Available from: <http://statline.cbs.nl/Statweb/publication/?DM=SLNL&PA=83385ned&D1=11-15&D2=a&D3=a&D4=5-9&D5=0&D6=l&VW=T> (accessed 12-12 2017).
10. Singh, A.S., Mulder, C., Twisk, J.W., van Mechelen, W., Chinapaw, M.J., (2008). Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes Rev*, 9(5): 474-488.
11. Hirasing, R.A., Fredriks, A.M., van Buuren, S., Verloove-Vanhorick, S.P., and Wit, J.M., (2001). [Increased prevalence of overweight and obesity in Dutch children, and the detection of overweight and obesity using international criteria and new reference diagrams]. *Ned Tijdschr Geneeskd*, 145(27): 1303-1308.
12. Schonbeck, Y., Talma, H., van Dommelen, P., Bakker, B., Buitendijk, S.E., Hirasing, R.A., and van Buuren, S., (2011). Increase in prevalence of overweight in Dutch children and adolescents: a comparison of nationwide growth studies in 1980, 1997 and 2009. *PLoS One*, 6(11): e27608.
13. Bahia, L., Schaan, C.W., Sparrenberger, K., Abreu, G.A., Barufaldi, L.A., Coutinho, W., and Schaan, B.D., (2018). Overview of meta-analysis on prevention and treatment of childhood obesity. *J Pediatr (Rio J)*.
14. Wang, Y., Cai, L., Wu, Y., Wilson, R.F., Weston, C., Fawole, O., Bleich, S.N., Cheskin, L.J., Showell, N.N., Lau, B.D., Chiu, D.T., Zhang, A., and Segal, L., (2015). What childhood obesity prevention programmes work? A systematic review and meta-analysis. *Obes Rev*,

- 16(7): 547-565.
15. Pandita, A., Sharma, D., Pandita, D., Pawar, S., Tariq, M., and Kaul, A., (2016). Childhood obesity: prevention is better than cure. *Diabetes Metab Syndr Obes*, 9: 83-89.
 16. Davison, K.K., and Birch, L.L., (2001). Childhood overweight: a contextual model and recommendations for future research. *Obes Rev*, 2(3): 159-171.
 17. Metcalf, B., Henley, W., and Wilkin, T., (2012). Effectiveness of intervention on physical activity of children: systematic review and meta-analysis of controlled trials with objectively measured outcomes (EarlyBird 54). *BMJ*, 345: e5888.
 18. Cushing, C.C., Brannon, E.E., Suorsa, K.I., and Wilson, D.K., (2014). Systematic review and meta-analysis of health promotion interventions for children and adolescents using an ecological framework. *J Pediatr Psychol*, 39(8): 949-962.
 19. Kamath, C.C., Vickers, K.S., Ehrlich, A., McGovern, L., Johnson, J., Singhal, V., Paulo, R., Hettlinger, A., Erwin, P.J., and Montori, V.M., (2008). Clinical review: behavioral interventions to prevent childhood obesity: a systematic review and metaanalyses of randomized trials. *J Clin Endocrinol Metab*, 93(12): 4606-4615.
 20. Lorenc, T., Petticrew, M., Welch, V., and Tugwell, P., (2013). What types of interventions generate inequalities? Evidence from systematic reviews. *J Epidemiol Community Health*, 67(2): 190-193.
 21. Chung, A., Backholer, K., Wong, E., Palermo, C., Keating C., and Peeters, A., (2016). Trends in child and adolescent obesity prevalence in economically advanced countries according to socioeconomic position: a systematic review. *Obes Rev*, 17(3): 276-295.
 22. Stamatakis, E., Wardle, J., and Cole, T.J., (2010). Childhood obesity and overweight prevalence trends in England: evidence for growing socioeconomic disparities. *Int J Obes (Lond)*, 34(1): 41-47.
 23. Gittelsohn, J., Anderson Steeves, E., Mui, Y., Kharmats, A.Y., Hopkins, L.C., and Dennis, D., (2014). B'More Healthy Communities for Kids: design of a multi-level intervention for obesity prevention for low-income African American children. *BMC Public Health*, 14: 942.
 24. Brown, H.E., Atkin, A.J., Panter, J., Wong, G., Chinapaw, M.J., and van Sluijs, E.M., (2016). Family-based interventions to increase physical activity in children: a systematic review, meta-analysis and realist synthesis. *Obes Rev*, 17(4): 345-360.
 25. Kok, G., Peters, L.W.H., and Ruiter, R.A.C., (2017). Planning theory- and evidence-based behavior change interventions: a conceptual review of the intervention mapping protocol. *Psicologia: Reflexão e Crítica*, 30(1): 19.
 26. Bartholomew Eldredge, L.K., Markham, C.M., Ruiter, R.A.C., Fernández, M.E., Kok, G., and Parcel, G.S., (2016). *Planning Health Promotion Programs: an Intervention Mapping Approach*. Fourth ed: John Wiley & Sons.
 27. Anselma, M., Chinapaw, M.J.M., and Altenburg, T.M., (2018). Determinants of Child Health Behaviors in a Disadvantaged Area from a Community Perspective: A Participatory Needs Assessment. *Int J Environ Res Public Health*, 15(644).
 28. Baum, F., MacDougall, C., and Smith, D., (2006). Participatory action research. *J Epidemiol Community Health*, 60(10): 854-857.
 29. Rodriguez, L.F., and Brown, T.M., (2009). From voice to agency: guiding principles for participatory action research with youth. *New Dir Youth Dev*, 2009(123): 19-34.
 30. Ozer, E.J., (2016). Youth-Led Participatory Action Research: Developmental and Equity

- Perspectives. *Adv Child Dev Behav*, 50: 189-207.
31. Cammarota, J., and Fine, M., (2008). Youth Participatory Action Research. In: Cammarota, J., and Fine, M., ed. Revolutionizing education. New York, United States of America: Routledge: 1-11.
 32. London, J.K., Zimmerman, K., and Erbstein, N., (2003). Youth-Led Research and Evaluation: Tools for Youth, Organizational, and Community Development. *NEW DIRECTIONS FOR EVALUATION*, (98): 33-45.
 33. Suleiman, A.B., Soleimanpour, S., and London, J. Youth Action for Health Through Youth-Led Research. *Journal of Community Practice* 2006;14(1-2): 125-145.
 34. Municipality of Amsterdam. Bevolking buurten naar herkomstgroepering, (2017) [Available from: <https://www.ois.amsterdam.nl/feiten-en-cijfers/buurten> (accessed 05-02 2018).
 35. GGD Amsterdam. Gezondheid in beeld [cited 2018 05-02]. Available from: <https://amsterdam.ggdgezondheidinbeeld.nl/>.
 36. Municipality of Amsterdam. Gebiedsanalyse 2017 Noord West Stadsdeel Noord, (2017) [Available from: https://www.ois.amsterdam.nl/pdf/2017_gebiedsanalyse_17.pdf (accessed 05-02 2018).
 37. Suleiman, A.B., Soleimanpour, S., and London, J., (2006). Youth Action for Health Through Youth-Led Research. *Journal of Community Practice*, 14(1-2): 125-145.
 38. Singh, A.S., Vik, F.N., Chinapaw, M.J., Uijtdewilligen, L., Verloigne, M., Fernández-Alvira, J.M., Stomfai, S., Manios, Y., Martens, M., and Brug, J., (2011). Test-retest reliability and construct validity of the ENERGY-child questionnaire on energy balance-related behaviours and their potential determinants: the ENERGY-project. *Int J Behav Nutr Phys Act*, 8: 136.
 39. Janssen, E.H., Singh, A.S., van Nassau, F., Brug, J., van Mechelen, W., and Chinapaw, M.J., (2014). Test-retest reliability and construct validity of the DOiT (Dutch Obesity Intervention in Teenagers) questionnaire: measuring energy balance-related behaviours in Dutch adolescents. *Public Health Nutr*, 17(2): 277-286.
 40. Ravens-Sieberer, U., Wille, N., Badia, X., Bonsel, G., Burström, K., Cavrini, G., Devlin, N., Egmar, A.C., Gusi, N., Herdman, M., Jelsma, J., Kind, P., Olivares, P.R., Scalone, L., and Greiner, W., (2010). Feasibility, reliability, and validity of the EQ-5D-Y: results from a multinational study. *Qual Life Res*, 19(6): 887-897.
 41. Trost, S.G., Loprinzi, P.D., Moore, R., and Pfeiffer, K.A., (2011). Comparison of accelerometer cut points for predicting activity intensity in youth. *Med Sci Sports Exerc*, 43(7): 1360-1368.
 42. Fischer, C., Yildirim, M., Salmon, J., and Chinapaw, M.J.M., (2012). Comparing different accelerometer cut-points for sedentary time in children. *Pediatr Exerc Sci*, 24(2): 220-228.
 43. Evenson, K.R., Catellier, D.J., Gill, K., Ondrak, K.S., and McMurray, R.G., (2008). Calibration of two objective measures of physical activity for children. *J Sports Sci*, 26(14): 1557-1565.
 44. Chinapaw, M.J., de Niet, M., Verloigne, M., de Bourdeaudhuij, I., Brug, J., and Altenburg, T.M., (2014). From sedentary time to sedentary patterns: accelerometer data reduction decisions in youth. *PLoS One*, 9(11): e111205.
 45. Leyten, C., Kemper, H.C.G., and Verschuur, R., (1982). De moper fitheidstest: handleiding en prestatieschalen 9 t/m 11 jarigen. Haarlem: BV Uitgeverij De Vrieseborch.

46. Molenaar, H.M., Zuidam, J.M., Selles, R.W., Stam, H.J., and Hovius, S.E., (2008). Age-specific reliability of two grip-strength dynamometers when used by children. *J Bone Joint Surg Am*, 90(5): 1053-1059.
47. Hamilton, G.F., McDonald, C., and Chenier, T.C., (1992). Measurement of grip strength: validity and reliability of the sphygmomanometer and jamar grip dynamometer. *J Orthop Sports Phys Ther*, 16(5): 215-219.
48. Matsuzaka, A., Takahashi, Y., Yamazoe, M., Kumakura, N., Ikeda, A., Wilk, B., and Bar-Or, O., (2004). Validity of the Multistage 20-M Shuttle-Run Test for Japanese Children, Adolescents, and Adults. *Pediatric exercise science*, 16(2): 113-125.
49. Liu, N.Y.S., Plowman, S.A., and Looney, M.A., (1992). The Reliability and Validity of the 20-Meter Shuttle Test in American Students 12 to 15 Years Old. *Research Quarterly for Exercise and Sport*, 63(4): 360-365.
50. Halcomb, E.J., Gholizadeh, L., DiGiacomo, M., Phillips, J., Davidson, P.M., (2007). Literature review: considerations in undertaking focus group research with culturally and linguistically diverse groups. *Journal of Clinical Nursing*, 16(6): 1000-1011.
51. Rabiee F., (2004). Focus-group interview and data analysis. *Proc Nutr Soc*, 63(4):655-60.
52. Marshall, J., and Mead, G., (2005). Editorial: Self-reflective practice and first-person action research. *Action research*, 3(3): 235-244.
53. Israel, B.A., Checkoway, B., Schulz, A., and Zimmerman, M., (1994). Health education and community empowerment: conceptualizing and measuring perceptions of individual, organizational, and community control. *Health Educ Q*, 21(2): 149-170.
54. Zimmerman, M., (2012). Empowerment Theory: Psychological, Organizational and Community Levels of Analysis.
55. Perkins, D.D., and Zimmerman, M.A., (1995). Empowerment theory, research, and application. *Am J Community Psychol*, 23(5): 569-579.
56. Zimmerman, M.A., Israel, B.A., Schulz, A., and Checkoway, B., (1992). Further Explorations in Empowerment Theory: An Empirical Analysis of Psychological Empowerment. *American Journal of Community Psychology*, 20(6): 707-727.
57. Corbin, J., and Strauss, A., (1990). Grounded Theory Research: Procedures, Canons, and Evaluative Criteria. *Qualitative Sociology*, 13(1): 3-21.
58. Berg, M., Coman, E., and Schensul, J.J., (2009). Youth Action Research for Prevention: a multi-level intervention designed to increase efficacy and empowerment among urban youth. *Am J Community Psychol*, 43(3-4): 345-359.
59. Foster-Fishman, P.G., Law, K.M., Lichty, L.F., and Aoun, C., (2010). Youth ReACT for Social Change: a method for youth participatory action research. *Am J Community Psychol*, 46(1-2): 67-83.
60. Checkoway, B., Allison, T., and Montoya, C., (2005). Youth participation in public policy at the municipal level. *Children and Youth Services Review*, 27: 1149-1162.
61. Oen, G., and Stormark, K.M., (2013). Participatory action research in the implementing process of evidence-based intervention to prevent childhood obesity: project design of the "Healthy Future" study. *J Obes*, 2013: 437206.
62. Kim, J., (2016). Youth Involvement in Participatory Action Research (PAR): Challenges and Barriers. *Critical Social Work*, 17(1): 38-53.
63. Ozer, E.J., Ritterman, M.L., and Wanis, M.G., (2010). Participatory action research (PAR) in

- middle school: opportunities, constraints, and key processes. *Am J Community Psychol*, 46(1-2): 152-166.
64. Caro, H.E., Altenburg, T.M., Dedding, C., and Chinapaw, M.J.M., (2016). Dutch Primary Schoolchildren's Perspectives of Activity-Friendly School Playgrounds: A Participatory Study. *Int J Environ Res Public Health*, 13(6).
 65. Wang, C.C., (2006). Youth Participation in Photovoice as a Strategy for Community Change. *Journal of Community Practice*, 14(1-2): 147-161.
 66. Wilson, N., Dasho, S., Martin, A.C., Wallerstein, N., Wang, C.C., and Minkler, M., (2007). Engaging Young Adolescents in Social Action Through Photovoice. *Journal of Early Adolescence*, 27(2): 241-261.
 67. Leykum, L.K., Pugh, J.A., Lanham, H.J., Harmon, J., and McDaniel, R.R. Jr., (2009). Implementation research design: integrating participatory action research into randomized controlled trials. *Implement Sci*, 4:69.