

VU Research Portal

Modulators of proteostasis: therapeutic targets and diagnostic markers to halt and reverse atrial fibrillation

Marion, D.M.S.

2021

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Marion, D. M. S. (2021). *Modulators of proteostasis: therapeutic targets and diagnostic markers to halt and reverse atrial fibrillation: Modulating proteostasis to halt and reverse AF*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Chapter 4

The future of atrial fibrillation therapy: intervention on heat shock proteins influencing electropathology is the next in line

Eva A. H. Lanthers¹, Denise M. S. van Marion², Herman Steen³, Natasja M. S. de Groot^{1#}, Bianca J. J. M. Brundel^{2#}

¹Unit Translational Electrophysiology, Department of Cardiology, Erasmus Medical Center, Rotterdam, The Netherlands

²Department of Clinical Pharmacy and Pharmacology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

³Nyken Therapeutics BV, Groningen, The Netherlands

Netherlands Heart Journal. 2015 Jun; 23(6):327-33.

Keywords: atrial fibrillation, mapping, electropathology, heat shock proteins, geranylgeranylacetone

Abstract

Atrial fibrillation (AF) is the most common age-related cardiac arrhythmia accounting for one-third of hospitalizations. Treatment of AF is difficult, which is rooted in the progressive nature of electrical and structural remodeling, called electropathology, which makes the atria more vulnerable for AF. Importantly, structural damage of the myocardium is already present when AF is diagnosed for the first time. Currently, no effective therapy is known that can resolve this damage.

Previously, we observed that exhaustion of cardio-protective heat shock proteins (HSPs) contributes to structural damage in AF patients. Also, boosting of HSPs, by the heat shock factor 1 (HSF1) activator geranylgeranylacetone (GGA), halted AF initiation and progression in experimental cardiomyocyte and dog models for AF. However, it is still unclear whether induction of HSPs also prolongs the arrhythmia-free interval after, for example, cardioversion of AF.

In this review, we discuss the role of HSPs in the pathophysiology of AF and give an outline of the HALT&REVERSE project, initiated by the HALT&REVERSE Consortium and the AF Innovation Platform. This project will elucidate whether HSPs 1) reverse cardiomyocyte electropathology and thereby halt AF initiation and progression and 2) represent novel biomarkers which predict the outcome of AF conversion and/or occurrence of post-operative AF.

Introduction

Atrial fibrillation (AF) represents the arrhythmia with the highest prevalence, accounting for one-third of hospitalizations related to cardiac rhythm disturbances [1]. The main risk factors to develop AF include age, cardiac surgery, valvular heart disease, congestive heart disease, ischemic cardiomyopathy, obesity, hypertension and diabetes mellitus. These risk factors cause atrial stretch and dilation and condition the heart for AF [1]. Patients with AF are frequently encountered in daily clinical practice. The goal of therapy in AF is, ideally, to abolish AF episodes and to restore sinus rhythm. This in turn re-establishes atrioventricular synchrony and improves atrial function. Despite extensive research, treatment of AF remains difficult, which is rooted in the persistent and progressive nature of this arrhythmia. An initiator, trigger and substrate are mandatory for development of AF. In this manuscript we will focus on the substrate underlying AF.

AF persistence is caused by progressive changes in the electrophysiology and structure of atrial cardiomyocytes [2], as demonstrated in a goat model for AF. In this model, AF resulted in shortening of the atrial effective refractory period and reversion of the physiological rate adaptation (shortening of the atrial refractory period at slower heart rates), facilitating induction and stability of AF. Similar findings have been reported in humans. The goat model of AF also revealed that longstanding AF is associated with changes in myocardial structure, including dedifferentiation, increase in cell size, perinuclear accumulation of glycogen, central loss of sarcomeres, changes in mitochondrial shape, fragmentation of the sarcoplasmic reticulum and disorganization of fiber orientation [3]. These structural changes were observed in human AF as well [4]. Importantly, structural changes are sustainable and impair electrical coupling and the functional recovery to sinus rhythm by pharmacological and electrical cardioversion. In addition, it was found that alterations of the myocardial structure underlie changes in electrophysiology as observed in AF [2, 5]. This phenomenon is commonly defined as electropathology. It should be noted that electropathology is already present when a patient enters the clinic for the first time with an episode of AF. The currently available

pharmacological therapy is directed at alleviation of electrical changes (rhythm control) and has limited effect on patient outcome [1]. Therapeutic approaches that halt the mechanisms conveying the AF-induced structural remodeling, may offer superior therapeutic perspectives. Thus, from science to patients: reversal of electropathology of cardiomyocytes represents a key target to accomplish and maintain cardiac sinus rhythm after AF.

Discontinuous propagation in the atria

It is generally assumed that transition of short-lasting paroxysms of AF to long-lasting episodes, that no longer terminate spontaneously over time, correspond to a progression from a trigger-driven to a substrate mediated arrhythmia due to a combination of electrical, structural and contractile atrial remodeling (Figure 1). Previous studies have demonstrated that alterations in the atrial microscopic structure impair intra-atrial conduction [6]. Spach et al. were the first to provide evidence that *'two-dimensional propagation is discontinuous in nature at a microscopic level'* (Figure 2) [7, 8]. In their experiments, variations in shape and amplitude of extracellular potential waveforms were studied when the direction of conduction was changed from longitudinal to transverse in relation to the fiber axis. They observed that in uniform anisotropic myocardium, the shape of the upstroke of the transmembrane action potential depended on the direction of propagation in relation to the fiber orientation. Fast conduction in the longitudinal direction was associated with a slow upstroke and a long τ -foot (time constant of the action potential representing voltage decay) and slow conduction in the transverse direction with a fast upstroke and a short τ -foot. Differences were explained by the anisotropic distribution of intercellular connections (gap junctions, connexions). This was confirmed in a goat model for sustained (two months) AF, where persistence of AF was associated with a lower density and inhomogeneous distribution of gap junctions [9, 10]. Gap junctions along the fiber axis provide a low resistance to current flow that increases the space constant and hence the electrotonic current flow. Subsequently, the upstroke of the transmembrane potential in this direction decreases

and the τ -foot increases. Due to the high resistance barriers in the transverse direction, the space constant is reduced and the electrotonic current is therefore smaller. This results in an increase in the upstroke of the transmembrane potential and a decrease in the τ -foot. Consequently, as the safety factor is lower during longitudinal conduction than during transverse conduction, conduction block is more likely to occur during longitudinal conduction [11].




Figure 1. Persistence of atrial fibrillation

Surface electrograms of paroxysms (upper panel) and persistent (lower panel) atrial fibrillation. Wave-maps constructed during these different types of maps are also shown. A wave-map shows individual fibrillation waves represented by colors according to their sequence of appearance. A previously described mapping algorithm was used to classify them into peripheral waves (entering the mapping area from outside the electrode array), breakthrough waves (appearing at the epicardial surface inside the mapping area) or discontinuous conduction waves (fibrillation waves starting with a delay of 13–40 ms from boundaries of other waves). Sites of epicardial breakthroughs are indicated by white asterisks; white arrows indicate direction(s) of expansion of epicardial breakthrough or discontinuous fibrillation waves. Peripheral fibrillation waves are indicated by black arrows.

Next to gap junctional remodeling, also ion channel remodeling has been demonstrated to play a role in AF progression. A marked reduction in ion (preferably L-type Ca^{2+} and transient outward K^+) channel protein expression, associated with a shortening of atrial effective refractory period, was observed in patients with AF [12]. This decrease in protein expression might be explained by increased (calcium

dependent) calpain I activity, causing enhanced degradation of ion-channel proteins and contractile proteins in the myocardium [4, 13].

Modifications of the atrial architecture affecting cardiomyocyte geometry (size and shape), gap junctions (distribution and conductivity) or the interstitial space (size and distribution) result in non-uniform anisotropic tissue. Muscle bundles become electrically dissociated due to strands of fibrotic tissue giving rise to fractionation of extracellular potentials [14]. These areas of local conduction abnormalities may facilitate genesis and perpetuation of AF.




Figure 2. Discontinuous conduction

Electrograms recorded in the longitudinal and transversal conduction direction in relation to fiber orientation in uniform anisotropic (A) and non-uniform anisotropic tissue (B). The electrogram in the transverse conduction direction in non-uniform anisotropic tissue consists of multiple deflections indicating discontinuous conduction. These alterations may be caused by changes in myocardial structure, including increase in cell size, perinuclear accumulation of glycogen, central loss of sarcomeres (myolysis), changes in mitochondrial shape, fragmentation of the sarcoplasmic reticulum and disorganization of fiber orientation as (partly) observed in panel (B) and (D). (C) Atrial myocardium from a patient in sinus rhythm (SR) and (D) from a patient with permanent AF (PeAF) stained with toluidine blue. SR patients show normal structural cardiomyocytes, whereas the patient with PeAF reveals cardiomyocytes with extensive perinuclear myolysis (red asterisk).

Fractionation of extracellular potentials

Fractionated electrograms are extracellular waveforms containing multiple, distinct deflections reflecting local asynchronous activation of myocardium surrounding the recording electrode [14]. Fractionation of fibrillation potentials can be functional and/or structural in nature as local asynchronous activation can be the result of a spatial dispersion in refractory periods (functional) or non-uniform tissue anisotropy (structural, as explained above). During AF, multiple, simultaneously circulating wavelets excite the atria. A fibrillation wave may encounter atrial tissue that is partially or totally refractory due to previous excitation by another wave. This results in slowing of conduction, conduction block and turning of fibrillation waves around lines of conduction block. All these conduction abnormalities give rise to local asynchronous activation of the myocardium and thus to multiple deflections in the extracellular potentials.

High density epicardial mapping during electrically induced AF in humans without structural heart disease showed that there is a relation between the degree of fractionation in the right atrium and specific spatial patterns of activation. Double potentials were recorded along lines of conduction block (Figure 3) whereas electrograms with 3 or more deflections were recorded at pivot points or areas of slow conduction [15, 16].




Figure 3. Double potentials recorded along the lines of conduction block

The color-coded activation map (interelectrode distance, 2 mm) shows two fibrillation waves (its trajectories indicated by thick black lines) separated by areas of conduction block. Electrograms consisting of two deflections are shown outside the activation map. They are recorded along the line of conduction block reflecting activation on either side.

In older patients with structural heart disease and longstanding persistent AF, it is likely that fractionation is more structurally determined due to dissociation of muscle bundles by fibrosis or fatty degeneration. Recent studies suggested that ablation of areas with fractionated electrograms may eliminate AF suggesting that fractionated electrograms are indeed indicative for arrhythmogenic regions perpetuating AF [17].

Counteracting electropathology by heat shock proteins

Previous research has unequivocally demonstrated that the induction of the heat shock response provides protection against various cardiac diseases [18]. Sensing of cellular stress activates the heat shock transcription factor 1 (HSF1) and consequently results in the expression of heat shock proteins (HSPs). In general, HSPs function as intra-cellular chaperones for other proteins. They play an important role in protein-protein interactions such as folding and assisting in the establishment of proper protein conformation (shape) and prevention of unwanted protein aggregation, which occur during ageing of the cell and (cardiac) stress [18]. Because of their essential role in protein maintenance, some members of the HSP family are expressed at low to moderate levels in all organisms. In cardiomyocytes, HSPs bind to damaged proteins, including sarcomeric proteins, and thereby prevent cardiomyocyte toxicity and further structural remodeling and functional loss [19, 20]. HSPs consist of five HSP families, including the HSPA1 and HSPB1 family, which both have been suggested to convey cardio-protective effects [19]. Therefore, impairment of HSP expression, especially HSPA1 and HSPB1, has been related to the development of various cardiac diseases, including AF [20], but also metabolic diseases, such as diabetes and obesity [21], which trigger the induction of AF [1].

These findings indicate that HSPs represent an interesting target to treat cardiac diseases such as AF. Indeed, experimental and human studies revealed that HSPs provide protection against AF initiation.

In dogs, pre-treatment with the HSP inducer, geranylgeranylacetone (GGA), attenuates changes in electrophysiology and conduction velocity of atrial cardiomyocytes and prevents initiation of AF in (acute) ischemia [22]. Also, protective effect of HSPs were observed in a rabbit model for AF induced by heart failure [23]. A further indication for the protective effect of HSPs is obtained from two studies in patients undergoing cardiac surgery. In both studies an inverse correlation between higher levels of HSPA1A expression in atrial tissue and lower incidence of post-operative AF was observed [24, 25]. In addition, one study described a correlation between HSPA1A and HSPB1 levels in tissue and improved restoration of sinus rhythm after mitral valve surgery [26]. In addition, high HSPB1 levels in blood predict sinus rhythm maintenance after catheter ablation in patients with paroxysmal AF [26]. In an experimental study in mice, it was observed that induction of HSPA1B prevents angiotensin II-induced AF and limits atrial fibrosis formation [27]. Together, these results indicate that induction of HSPs mitigates the development of the atrial substrate for the induction of AF.

In addition to the cardio-protective effect of HSPs against the induction of AF, it is expected that induced expression of HSPs also protects against AF progression. Such a cardio-protective effect is supported by several studies. For example, general HSP levels were induced by a heat shock or GGA treatment in tachypaced HL-1 atrial cardiomyocytes and these cardiomyocytes were compared to normal tachypaced HL-1 cardiomyocytes [20, 28]. In these studies, tachypaced cardiomyocytes with induced HSP expression revealed improved contractile function and structural integrity. Similar protection by general HSP induction was obtained in a dog model for AF [28]. Dogs pre-treated with GGA showed attenuation of tachypacing-induced AF promotion and limited recurrence of AF after cardioversion, most likely by limiting the electrophysiological changes including shortening of action potential duration and maintenance of L-type Ca^{2+} current [28]. Further studies revealed that specific members of the family of small HSPs play a key role in cardio-protection against AF. Induction of HSPB1 was found sufficient to prevent tachypacing-induced structural remodeling, while overexpressing HSPA1 was not protective [28]. Other specific small HSP family members also show protection in cellular models, including HSPB6, HSPB7 and HSPB8, but not the other small HSP family members [29].

Importantly, selective knockdown of HSPB1 fully attenuates the protective effect of GGA, demonstrating a main role for HSPB1 in cardio-protection. A similar protective effect of the small HSP members as observed in HL-1 cardiomyocytes has been found in *Drosophila melanogaster* (fruit fly) model for tachypacing-induced contractile dysfunction. In the transparent fruit fly pupae, the heart wall can be visualized and subjected to tachypacing [30]. This model also allows for general induction of HSPs, either by heat shock, or by HSP-inducing compounds such as GGA, or specific induction of HSPs via genetic manipulation specifically in the heart. Indeed, induction of HSPs, especially the small HSPs, by these strategies prevents tachypacing-induced contractile dysfunction and structural remodeling in *Drosophila* pupae [30]. Also, in human AF there are indications that HSPB1 may protect against cardiac remodeling and AF progression. Two independent studies report higher atrial expression levels of HSPB1 to relate to a shorter duration of AF and less extensive structural damage [20, 31]. This suggests that in short-duration AF the HSP response gets activated, while it diminishes over time when AF persists. As HSPB1 conserves the cardiomyocyte structure, lower levels of HSPB1 may trigger the progression of structural remodeling, paving the way to longstanding and permanent AF. Therefore, securing HSP levels at an adequate level, for example, by treatment with HSP inducers, may limit the expansion of the AF substrate during paroxysmal and short-term AF. In agreement with this hypothesis, restoration of sinus rhythm in patients with permanent AF after mitral valve surgery is related to the HSF1 activity and induced HSPB1 levels [32].

Taken together, induction of HSPs attenuates electropathology of cardiomyocytes in AF. In the project “Hsf1 Activators Lower cardiomyocyte damage: Towards a novel therapeutic approach to REVERSE atrial fibrillation” (HALT & REVERSE), we will focus on the electropathological substrate of AF and dissect the mechanism of protection.

Novel HSP-inducing agents as a therapeutic approach in AF

Pharmacological approaches to prevent mechanisms underlying atrial electropathology are being studied, with the hope that they might be useful therapeutic agents in treating AF [1]. It has been recognized that the efficacy of commonly used drugs on remodeling is limited [1]. As the protective action of HSPs depends on their temporary induction, drugs that boost the endogenous heat shock responses are of particular interest in prevention of first onset, recurrence and progression of AF. A drug often used to boost HSP expression is GGA. GGA is originally used as an anti-ulcer agent and is a nontoxic acyclic isoprenoid compound with a retinoid skeleton that induces HSP synthesis in various tissues, including gastric mucosa, intestine, liver, heart, retina, and central nervous system [33, 34]. GGA induces HSP expression through the activation of HSF1. The protective effect of GGA-induced HSP expression on electropathology caused by tachycardia has been observed in various experimental models, including the dog model for AF [28]. These findings suggest that the induction of HSPs by GGA might have a potential value for clinical AF. In *in vivo* tachypaced *Drosophila*, GGA treatment protected against contractile dysfunction of the heart wall and structural remodeling [30]. Also in the dog model for (acute) atrial ischemia and tachypacing-induced AF promotion, an HSP-inducing GGA treatment revealed protective effects against cardiomyocyte remodeling, and consequently occurrence and recurrence of AF [22]. A direct protective effect of HSP on the earlier mentioned gap junctional and ion channel remodeling has not been described (yet). However it has been shown that GGA treatment attenuates the proteolytic activity of calpain [30] and prevents ion channel remodeling in AF [28], suggesting a direct protective role of HSPs.

Although protective effects with GGA were observed, an important disadvantage of GGA is its high LogP value, which is around 9, and therefore generally high dosages are required, as found in the dog studies for AF [22, 28]. To overcome this disadvantage various derivatives of GGA have been synthesized, with improved pharmaco-chemical and HSP-boosting properties (Figure 4A). Some of these derivatives also showed cardio-protective effects in tachypaced HL-1 cardiomyocytes (Figure

4B). In the HALT & REVERSE project, we will test whether these derivatives reveal improved cardio-protective effects compared with GGA and elucidate their mode of action. In addition, one of the selected GGA derivatives will be further tested in an animal model for AF. Finally, GGA has also been tested in patients undergoing cardiac surgery and atrial tissue will be used to test whether GGA induces HSP expression in patients and protects against post-operative AF.

Expected findings of HALT & REVERSE

Ultimately, by inducing cardio-protective HSP expression, the HALT & REVERSE Consortium wants to halt or even reverse the structure of cardiomyocytes and therefore the electropathology in patients with AF (Figure 4C). Hereby, we expect to significantly contribute to novel rational approaches, to enhance successful recovery of heart function after electrical cardioversion (ECV) and/or cardiac surgery. This approach will affect the substrate underlying AF, not the trigger. For the first time, we exploit HSP levels to predict the electropathological outcome in patients, as measured by our unique mapping approaches. In addition, at the end of this project, we will know whether HSPs are suitable biomarkers for AF. Our unique approach will ultimately result in reduced hospital time, induced quality of life for the patient and lower costs for the society. The findings will be of benefit to other (cardiac, kidney) diseases, as many of those are characterized by sustainable structural remodeling. Thus, the project is of major importance from a scientific, clinical and economical perspective.




Figure 4. Geranylgeranylacetone (GGA) derivatives induce HSPA1 expression and reveal cardio-protective effects

(A) Western blot showing induction of HSPA1 levels in HL-1 cardiomyocytes treated with GGA and some of the GGA derivatives (Nyk compounds). (B) Representative calcium transient tracers of non-treated and normally paced (1 Hz) HL-1 cardiomyocytes (Control) or tachypaced cardiomyocytes (TP, 5 Hz) without or with treatment of 10 μ M GGA or GGA derivatives (Nyk compounds) as indicated. (C) Overview of the central concept of HALT & REVERSE. Atrial fibrillation (AF) naturally progresses in time (red line), which is rooted in the underlying electropathology. By pharmacological induction of HSP levels (blue arrow), we aim to halt or even reverse electropathology and consequently AF progression (green line).

Acknowledgements This work was supported by the LSH-Impulse grant (40-43100-98-008) and the Dutch Heart Foundation (2013T088, 2013T144, 2013T096 and 2011T046).

Conflict of interest Dr. H. Steen is the CEO of Nyken Therapeutics, which holds IP on the use of HSP-inducing compounds.

References

1. Dobrev D, et al. Novel molecular targets for atrial fibrillation therapy. *Nat Rev Drug Discov* 2012;11(4):275-291.
2. Wijffels MC, et al. Atrial fibrillation begets atrial fibrillation. A study in awake chronically instrumented goats. *Circulation* 1995;92(7):1954-1968.
3. Ausma J, et al. Structural changes of atrial myocardium due to sustained atrial fibrillation in the goat. *Circulation* 1997;96(9):3157-3163.
4. Brundel BJM, et al. Activation of Proteolysis by Calpains and Structural Changes in Human Paroxysmal and Persistent Atrial Fibrillation. *Cardiovasc Res* 2002;54:380-389.
5. De Groot N, et al. Electropathological substrate of longstanding persistent atrial fibrillation in patients with structural heart disease: epicardial breakthrough. *Circulation* 2010;122:1674-1682.
6. Allesie M, et al. Electrical, contractile and structural remodeling during atrial fibrillation. *Cardiovasc Res* 2002;54:230-246.
7. Spach MS, et al. Electrophysiological effects of remodeling cardiac gap junctions and cell size: experimental and model studies of normal cardiac growth. *Circ Res* 2000;86(3):302-311.
8. Spach MS, Dolber PC. Relating extracellular potentials and their derivatives to anisotropic propagation at a microscopic level in human cardiac muscle. Evidence for electrical uncoupling of side-to-side fiber connections with increasing age. *Circ Res* 1986;58(3):356-371.
9. van der Velden HM, et al. Gap junctional remodeling in relation to stabilization of atrial fibrillation in the goat. *Cardiovasc Res* 2000;46(3):476-86.
10. Van der Velden HMW, et al. Altered pattern of connexin40 distribution in persistent atrial fibrillation in the goat. *J Cardiovasc Electrophysiol* 1998;9:596-607.
11. Koura T, et al. Anisotropic conduction properties in canine atria analyzed by high-resolution optical mapping: preferential direction of conduction block changes from longitudinal to transverse with increasing age. *Circulation* 2002;105(17):2092-2098.
12. Brundel BJM, et al. Ion channel remodeling is related to intra-operative atrial refractory periods in patients with paroxysmal and persistent atrial fibrillation. *Circulation* 2001;103:684-690.
13. Ke L, et al. Calpain mediates cardiac troponin degradation and contractile dysfunction in atrial fibrillation. *J Mol Cell Cardiol* 2008;45:685-693.
14. Spach MS. Anisotropy of cardiac tissue: a major determinant of conduction? *J Cardiovasc Electrophysiol* 1999;10(6):887-890.
15. Konings KT, et al. Configuration of unipolar atrial electrograms during electrically induced atrial fibrillation in humans. *Circulation* 1997;95(5):1231-1241.
16. Yaksh A, et al. Atrial fibrillation: to map or not to map? *Neth Heart J* 2014;22(6):259-266.

17. Nademanee K. Trials and travails of electrogram-guided ablation of chronic atrial fibrillation. *Circulation* 2007;115(20):2592-2594.
18. Willis MS, Patterson C. Proteotoxicity and cardiac dysfunction. *N Engl J Med* 2013;368(18):1755.
19. Brundel BJM, et al. Heat shock proteins as molecular targets for intervention in atrial fibrillation. *Cardiovasc Res* 2008;78:422-428.
20. Brundel BJM, et al. Heat shock protein upregulation protects against pacing-induced myolysis in HL-1 atrial myocytes and in human atrial fibrillation. *J Mol Cell Cardiol* 2006;41:555-562.
21. Chung J, et al. HSP72 protects against obesity-induced insulin resistance. *Proc Natl Acad Sci* 2008;105:1739-1744.
22. Sakabe M, et al. Effects of heat shock protein induction on Atrial Fibrillation caused by acute atrial ischemia. *Cardiovasc Res* 2008;78:63-70.
23. Chang SL, et al. Heat shock protein inducer modifies arrhythmogenic substrate and inhibits atrial fibrillation in the failing heart. *Int J Cardiol* 2013 Oct 9;168(4):4019-4026.
24. Mandal K, et al. Association of high intracellular, but not serum, heat shock protein 70 with postoperative atrial fibrillation. *Ann Thorac Surg.* 2005;79:865-871.
25. St Rammos K, et al. Low preoperative HSP70 atrial myocardial levels correlate significantly with high incidence of postoperative atrial fibrillation after cardiac surgery. *Cardiovasc Surg.* 2002;10:228-232.
26. Hu YF, et al. Electrophysiological correlation and prognostic impact of heat shock protein 27 in atrial fibrillation. *Circ Arrhythm Electrophysiol* 2012 Apr;5(2):334-340.
27. Wakisaka O, et al. Hyperthermia treatment prevents angiotensin II-mediated atrial fibrosis and fibrillation via induction of heat-shock protein 72. *J Mol Cell Cardiol.* 2007;436:616-626.
28. Brundel BJM, et al. Induction of heat-shock response protects the heart against atrial fibrillation. *Circ Res* 2006;99:1394-1402.
29. Ke L, et al. HSPB1, HSPB6, HSPB7 and HSPB8 protect against RhoA GTPase-induced remodeling in tachypaced HL-1 atrial myocytes. *PlosOne* 2011;6:e20395.
30. Zhang D, et al. Effects of different small HSPB members on contractile dysfunction and structural changes in a *Drosophila melanogaster* model for Atrial Fibrillation. *J Mol Cell Cardiol* 2011;51:381-389.
31. Yang M, et al. Expression of heat shock proteins in myocardium of patients with atrial fibrillation. *Cell Stress & Chaperones* 2007;12:142-150.
32. Cao H, et al. Heat shock proteins in stabilization of spontaneously restored sinus rhythm in permanent atrial fibrillation patients after mitral valve surgery. *Cell Stress & Chaperones* 2011; 16(5):517-28.
33. Ooie T, et al. Single oral dose of geranylgeranylacetone induces heat-shock protein 72 and renders protection against ischemia/reperfusion injury in rat heart. *Circulation* 2001;104:1837-1843.
34. Katsuno M, et al. Pharmacological induction of heat-shock proteins alleviates polyglutamine-mediated motor neuron disease. *Proc Natl Acad Sci USA* 2005;102:16801-16806.