

VU Research Portal

Blended CBT for Depression

Kooistra, L.C.

2021

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Kooistra, L. C. (2021). *Blended CBT for Depression*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

REFERENCES

References

- Aaronson, N. K., Muller, M., Cohen, P. D. A., Essink-Bot, M.-L., Fekkes, M., Sanderman, R., Sprangers, M. A. G., te Velde, A., & Verrips, E. (1998). Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *Journal of Clinical Epidemiology*, 51(11), 1055–1068.
[https://doi.org/10.1016/S0895-4356\(98\)00097-3](https://doi.org/10.1016/S0895-4356(98)00097-3)
- Allison, P. D. (1999). Logistic regression using the SAS system: Theory and application. SAS Institute Inc. and John Wiley and Sons.
- Alonso, J., Angermeyer, M. C., Bernert, S., Bruffaerts, R., Brugha, T. S., Bryson, H., de Girolamo, G., Graaf, R., Demyttenaere, K., Gasquet, I., Haro, J. M., Katz, S. J., Kessler, R. C., Kovess, V., Lépine, J. P., Ormel, J., Polidori, G., Russo, L. J., Vilagut, G., ... Vollebergh, W. A. M. (2004). Use of mental health services in Europe: Results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatrica Scandinavica. Supplementum*, 109(420), 47–54. <https://doi.org/10.1111/j.1600-0047.2004.00330.x>
- American Psychiatric Association (APA). (2000). Diagnostic and statistical manual of mental disorders, 4th edition, text revision (DSM-IV-TR) (4th ed.). American Psychiatric Association.
- American Psychiatric Association (APA). (2013). Diagnostic and statistical manual of mental disorders (DSM-V) (5th ed.). Author.
- Andersson, G. (2010). The promise and pitfalls of the internet for cognitive behavioral therapy. *BMC Medicine*, 8(1), 82. <https://doi.org/10.1186/1741-7015-8-82>
- Andersson, G., & Cuijpers, P. (2008). Pros and cons of online cognitive-behavioural therapy. *The British Journal of Psychiatry: The Journal of Mental Science*, 193(4), 270–271. <https://doi.org/10.1192/bj.pbp.108.054080>
- Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized psychological treatments for adult depression: A meta-analysis. *Cognitive Behaviour Therapy*, 38(4), 196–205. <https://doi.org/10.1080/16506070903318960>
- Andersson, G., Cuijpers, P., Carlbring, P., Riper, H., & Hedman, E. (2014). Guided Internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: A systematic review and meta-analysis. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)*, 13(3), 288–295. <https://doi.org/10.1002/wps.20151>
- Andersson, G., Rozental, A., Shafran, R., & Carlbring, P. (2017). Long-term effects of internet-supported cognitive behaviour therapy. *Expert Review of Neurotherapeutics*, 18(1), 1–8. <https://doi.org/10.1080/14737175.2018.1400381>
- Andersson, G., Wagner, B., & Cuijpers, P. (2016). ICBT for depression. In N. Lindefors & G. Andersson (Eds.), *Guided internet-based treatments in psychiatry* (pp. 17–32). Springer International Publishing. https://doi.org/10.1007/978-3-319-06083-5_2
- Andrews, G., Basu, A., Cuijpers, P., Craske, M. G., McEvoy, P., English, C. L., & Newby, J. M. (2018). Computer therapy for the anxiety and depression disorders is effective, acceptable and practical health care: An updated meta-analysis. *Journal of Anxiety Disorders*, 55(10), 70–78. <https://doi.org/10.1016/j.janxdis.2018.01.001>

- Andrews, G., Cuijpers, P., Craske, M. G., McEvoy, P., & Titov, N. (2010). Computer therapy for the anxiety and depressive disorders is effective, acceptable and practical health care: A meta-analysis. *PLOS ONE*, 5(10), e13196. <https://doi.org/10.1371/journal.pone.0013196>
- Andrews, G., Henderson, S., & Hall, W. (2001). Prevalence, comorbidity, disability and service utilisation. Overview of the Australian National Mental Health Survey. *The British Journal of Psychiatry*, 178, 145–153. <https://doi.org/10.1192/bjp.178.2.145>
- Andrews, G., & Williams, A. D. (2014). Up-scaling clinician assisted internet cognitive behavioural therapy (iCBT) for depression: A model for dissemination into primary care. *Clinical Psychology Review*, 41. <https://doi.org/10.1016/j.cpr.2014.05.006>
- Araya, R., Zitko, P., Markkula, N., Rai, D., & Jones, K. (2018). Determinants of access to health care for depression in 49 countries: A multilevel analysis. *Journal of Affective Disorders*, 234, 80–88. <https://doi.org/10.1016/j.jad.2018.02.092>
- Bangor, A., Kortum, P., & Miller, J. (2009). Determining what individual SUS scores mean: Adding an adjective rating scale. *Journal of Usability Studies*, 4(3), 114–123. <https://doi.org/10.1080/1554549.2009.1073405>
- Bangor, A., Kortum, P. T., & Miller, J. T. (2008). An empirical evaluation of the System Usability Scale. *International Journal of Human-Computer Interaction*, 24(6), 574–594. <https://doi.org/10.1080/10447310802205776>
- Barth, J., Munder, T., Gerger, H., Nüesch, E., Trelle, S., Znoj, H., Jüni, P., & Cuijpers, P. (2013). Comparative efficacy of seven psychotherapeutic interventions for patients with depression: A network meta-analysis. *PLOS Medicine*, 10(5). <https://doi.org/10.1371/journal.pmed.1001454>
- Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., Dai, B., & Eigen, C. (2017). Linear mixed-effects models using “Eigen” and S4 (1.1-13).
- Bech, P., Olsen, L. R., Kjoller, M., & Rasmussen, N. K. (2003). Measuring well-being rather than the absence of distress symptoms: A comparison of the SF-36 Mental Health subscale and the WHO-Five Well-Being Scale. *International Journal of Methods in Psychiatric Research*, 12(2), 85–91. <https://doi.org/10.1002/mpr.145>
- Beck, A., Brown, G., Steer, R. A., Eidelberg, J. I., & Riskind, J. H. (1987). Differentiating anxiety and depression: A test of the cognitive content-specificity hypothesis. *Journal of Abnormal Psychology*, 96(3), 179–183. <https://doi.org/10.1037/0021-843X.96.3.179>
- Beck, A., Epstein, N., Brown, G., & Steer, R. A. (1988). An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting and Clinical Psychology*, 56(6), 893–897. <https://doi.org/10.1037/0022-006X.56.6.893>
- Beck, A. T., Kovacs, M., & Weissman, A. (1979). Assessment of suicidal intention: The Scale for Suicide Ideation. *Journal of Consulting and Clinical Psychology*, 47(2), 343–352. <https://doi.org/10.1037/0022-006X.47.2.343>
- Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). Cognitive therapy of depression. Guilford.
- Beck, A. T., & Steer, R. A. (1993). Beck Anxiety Inventory: Manual. The Psychological Corporation.

References

- Beck, A. T., Steer, R., & Brown, G. K. (1996). Beck Depression Inventory II: Manual. The Psychological Corporation.
- Beck, J. S. (2011). Cognitive behavior therapy: Basics and beyond (Second edi). The Guilford Press.
- Becker, E. M., & Jensen-Doss, A. (2013). Computer-assisted therapies: Examination of therapist-level barriers to their use. *Behavior Therapy*, 44(4), 614-624.
<https://doi.org/10.1016/j.beth.2013.05.002>
- Berger, T. (2017). The therapeutic alliance in internet interventions: A narrative review and suggestions for future research. *Psychotherapy Research*, 27(5), 511-524.
<https://doi.org/10.1080/10503307.2015.1119908>
- Berger, T., Krieger, T., Sude, K., Meyer, B., & Maercker, A. (2018). Evaluating an e-mental health program ("deprexis") as adjunctive treatment tool in psychotherapy for depression: Results of a pragmatic randomized controlled trial. *Journal of Affective Disorders*, 227, 455-462. <https://doi.org/10.1016/j.jad.2017.11.021>
- Bijl, R. V., & Ravelli, A. (2000). Psychiatric morbidity, service use, and need for care in the general population: Results of the Netherlands Mental Health Survey and Incidence Study. *Am J Public Health*, 90(4), 602-607. <https://doi.org/10.2105/ajph.90.4.602>
- Bijl, R. V., van Zessen, G., Ravelli, A., de Rijk, C., & Langendoen, Y. (1998). The Netherlands Mental Health Survey and Incidence Study (NEMESIS): Objectives and design. *Social Psychiatry and Psychiatric Epidemiology*, 33(12), 581-586.
<http://www.ncbi.nlm.nih.gov/pubmed/9857790>
- Bockting, C. L. H., & Huibers, M. J. H. (2011). Protocollaire behandeling van patienten met een depressieve stoornis. In G. P. J. Keijsers, A. van Minnen, & C. A. L. Hoogduin (Eds.), *Protocollaire behandelingen voor volwassenen met psychische klachten 1* (pp. 251-288). Uitgeverij Boom.
- Bordin, E. S. (1979). The generalizability of the psychoanalytic concept of the working alliance. *Psychotherapy: Theory, Research & Practice*, 16(3), 252-260.
<https://doi.org/10.1037/h0085885>
- Bosmans, J. E., de Bruijne, M. C., van Hout, H. P. J., Hermens, M. L. M., Adèr, H. J., & van Tulder, M. W. (2008). Practical guidelines for economic evaluations alongside equivalence trials. *Value in Health: The Journal of the International Society for Pharmacoeconomics and Outcomes Research*, 11(2), 251-258. <https://doi.org/10.1111/j.1524-4733.2007.00245.x>
- Bower, P., & Gilbody, S. (2005). Stepped care in psychological therapies: Access, effectiveness and efficiency. Narrative literature review. *The British Journal of Psychiatry*, 186, 11-17.
<https://doi.org/10.1192/bjp.186.1.11>
- Brazier, J., Roberts, J., & Deverill, M. (2002). The estimation of a preference-based measure of health from the SF-36. *Journal of Health Economics*, 21(2), 271-292.
[https://doi.org/10.1016/S0167-6296\(01\)00130-8](https://doi.org/10.1016/S0167-6296(01)00130-8)
- Bremmer, F., & van Es, M. (2013). Een analyse van de verwachte kosten en baten van eHealth: Blended behandelen en begeleiden.
- Briggs, A. H., Claxton, K., & Sculpher, M. J. (2006). Decision modelling for health economic evaluation. Oxford university press.

- Brooke, J. (1996). SUS-a quick and dirty usability scale. In P. W. Jordan, B. Thomas, B. A. Weerdmeester, & I. L. McClelland (Eds.), Usability evaluation in industry (Vol. 189, pp. 189–194). Taylor & Francis Ltd.
- Brooks, R. (1996). EuroQoL: The current state of play. *Health Policy* (Amsterdam, Netherlands), 37(1), 53–72. <http://www.ncbi.nlm.nih.gov/pubmed/10158943>
- Bruijniks, S. J. E., Franx, G., & Huibers, M. J. H. (2018). The implementation and adherence to evidence-based protocols for psychotherapy for depression: The perspective of therapists in Dutch specialized mental healthcare. *BMC Psychiatry*, 18(1), 1–10.
<https://doi.org/10.1186/s12888-018-1768-x>
- Bruijniks, S. J. E., Lemmens, L. H. J. M., Hollon, S. D., Peeters, F. P. M. L., Cuijpers, P., Arntz, A., Dingemanse, P., Willems, L., van Oppen, P., Twisk, J. W. R., van den Boogaard, M., Spijker, J., Bosmans, J., & Huibers, M. J. H. (2020). The effects of once- versus twice-weekly sessions on psychotherapy outcomes in depressed patients. *The British Journal of Psychiatry : The Journal of Mental Science*, 216(4), 222–230.
<https://doi.org/10.1192/bjp.2019.265>
- Butler, A. C., Chapman, J. E., Forman, E. M., & Beck, A. T. (2006). The empirical status of cognitive-behavioral therapy: A review of meta-analyses. *Clinical Psychology Review*, 26(1), 17–31. <https://doi.org/10.1016/j.cpr.2005.07.003>
- Cameron, S. K., Rodgers, J., & Dagnan, D. (2018). The relationship between the therapeutic alliance and clinical outcomes in cognitive behaviour therapy for adults with depression: A meta-analytic review. *Clinical Psychology & Psychotherapy*, 25(3), 446–456.
<https://doi.org/10.1002/cpp.2180>
- Carlbring, P., Andersson, G., Cuijpers, P., Riper, H., & Hedman-Lagerlöf, E. (2018). Internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: An updated systematic review and meta-analysis. *Cognitive Behaviour Therapy*, 47(1), 1–18. <https://doi.org/10.1080/16506073.2017.1401115>
- Cohen, J. (1988). Statistical power analysis for the behavioral sciences (revised edition). Lawrence Erlbaum Associates, Inc.
- Costa, P., & McCrae, R. R. (1995). Inventory domains and facets: Hierarchical personality assessment using the Revised NEO Personality Inventory. *Journal of Personality Assessment*, 64(1), 21–50. <https://doi.org/10.1207/s15327752jpa6401>
- Cuijpers, P. (2017). Four decades of outcome research on psychotherapies for adult depression: An overview of a series of meta-analyses. *Canadian Psychology*, 58(1), 7–19.
<https://doi.org/10.1037/cap0000096>
- Cuijpers, P., Donker, T., Johansson, R., Mohr, D. C., van Straten, A., & Andersson, G. (2011). Self-guided psychological treatment for depressive symptoms: A meta-analysis. *PLOS ONE*, 6(6), 1–7. <https://doi.org/10.1371/journal.pone.0021274>
- Cuijpers, P., Donker, T., van Straten, A., Li, J., & Andersson, G. (2010). Is guided self-help as effective as face-to-face psychotherapy for depression and anxiety disorders? A systematic review and meta-analysis of comparative outcome studies. *Psychological Medicine*, 40(12), 1943–1957. <https://doi.org/10.1017/S0033291710000772>

References

- Cuijpers, P., Huibers, M. J. H., Ebert, D. D., Koole, S. L., & Andersson, G. (2013). How much psychotherapy is needed to treat depression? A metaregression analysis. *Journal of Affective Disorders*, 149(1-3), 1-13. <https://doi.org/10.1016/j.jad.2013.02.030>
- Cuijpers, P., & Riper, H. (2014). Internet Interventions for depressive disorders: An overview. *Revista de Psicopatología y Psicología Clínica*, 19(2), 209-216. <https://doi.org/http://dx.doi.org/10.5944/rppc.vol.19.num.3.2014.13902>
- Cuijpers, P., Riper, H., & Andersson, G. (2015). Internet-based treatment of depression. *Current Opinion in Psychology*, 4, 131-135. <https://doi.org/10.1016/j.copsyc.2014.12.026>
- Cuijpers, P., Smit, F., Oostenbrink, J., de Graaf, R., ten Have, M., & Beekman, A. (2007). Economic costs of minor depression: A population-based study. *Acta Psychiatrica Scandinavica*, 115(3), 229-236. <https://doi.org/10.1111/j.1600-0447.2006.00851.x>
- Cuijpers, P., van Straten, A., van Oppen, P., & Andersson, G. (2008). Are psychological and pharmacologic interventions equally effective in the treatment of adult depressive disorders? A meta-analysis of comparative studies. *Journal of Clinical Psychiatry*, 69(11), 1675-1685; quiz 1839-1841. <https://doi.org/10.4088/jcp.v69n1102>
- Cuijpers, P., Berking, M., Andersson, G., Quigley, L., Kleiboer, A., & Dobson, K. S. (2013). A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *The Canadian Journal of Psychiatry*, 58(7), 376-385. <https://doi.org/10.1177/070674371305800702>
- Cuijpers, P., Karyotaki, E., de Wit, L., & Ebert, D. D. (2020). The effects of fifteen evidence-supported therapies for adult depression: A meta-analytic review. *Psychotherapy Research: Journal of the Society for Psychotherapy Research*, 30(3), 279-293. <https://doi.org/10.1080/10503307.2019.1649732>
- Cusin, C., Yang, H., Yeung, A., & Fava, M. (2010). Rating Scales for Depression. In L. Baer & M. A. Blais (Eds.), *Handbook of Clinical Rating Scales and Assessment in Psychiatry and Mental Health*. Humana Press. https://doi.org/10.1007/978-1-59745-387-5_2
- de Brey, H. (1983). A cross-national validation of the Client Satisfaction Questionnaire: The Dutch experience. *Evaluation and Program Planning*, 6(3-4), 395-400. [https://doi.org/10.1016/0149-7189\(83\)90018-6](https://doi.org/10.1016/0149-7189(83)90018-6)
- Demyttenaere, K., Bruffaerts, R., Posada-Villa, J., Gasquet, I., Kovess, V., Lepine, J. P., Angermeyer, M. C., Bernert, S., de Girolamo, G., Morosini, P., Polidori, G., Kikkawa, T., Kawakami, N., Ono, Y., Takeshima, T., Uda, H., Karam, E. G., Fayyad, J. A., Karam, A. N., ... WHO World Mental Health Survey Consortium. (2004). Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. *JAMA*, 291(21), 2581-2590. <https://doi.org/10.1001/jama.291.21.2581>
- Denicoff, K. D., Smith-Jackson, E. E., Disney, E. R., Suddath, R. L., Leverich, G. S., & Post, R. M. (1997). Preliminary evidence of the reliability and validity of the prospective life-chart methodology (LCM-p). *Journal of Psychiatric Research*, 31(5), 593-603. [https://doi.org/10.1016/S0022-3956\(96\)00027-1](https://doi.org/10.1016/S0022-3956(96)00027-1)
- Devilly, G. J., & Borkovec, T. D. (2000). Psychometric properties of the credibility/expectancy questionnaire. *Journal of Behavior Therapy and Experimental Psychiatry*, 31(2), 73-86. [https://doi.org/10.1016/S0005-7916\(00\)00012-4](https://doi.org/10.1016/S0005-7916(00)00012-4)

- Driessen, E., Van, H. L., Don, F. J., Peen, J., Kool, S., Westra, D., Hendriksen, M., Schoevers, R. A., Cuijpers, P., Twisk, J. W. R., & Dekker, J. J. M. (2013). The efficacy of cognitive-behavioral therapy and psychodynamic therapy in the outpatient treatment of major depression: A randomized clinical trial. *American Journal of Psychiatry*, 170(9), 1041–1050. <https://doi.org/10.1177/0003065114525237>
- Drummond, M. F., & Sculpher, M. (2005). Common methodological flaws in economic evaluations. *Medical Care*, 43(7 Suppl), 5-14. <http://www.ncbi.nlm.nih.gov/pubmed/16056003>
- Drummond, M. F., Sculpher, M. J., Claxton, K., Stoddart, G. L., & Torrance, G. W. (2015). Methods for the economic evaluation of health care programmes (4th ed.). Oxford University Press.
- Drummond, M. F., Sculpher, M. J., Torrance, G. W., O'Brien, B. J., & Stoddart, G. (2005). Cost methods for the economic evaluation of health care programmes. In *Methods for the Economic Evaluation of Health Care Programmes* (3rd ed.). Oxford university press.
- Emmelkamp, P. M. G., David, D., Beckers, T., Muris, P., Cuijpers, P., Lutz, W., Andersson, G., Araya, R., Banos Rivera, R. M., Barkham, M., Berking, M., Berger, T., Botella, C., Carlbring, P., Colom, F., Essau, C., Hermans, D., Hofmann, S. G., Knappe, S., ... Vervliet, B. (2014). Advancing psychotherapy and evidence-based psychological interventions. *International Journal of Methods in Psychiatric Research*, 23(Supp. 1), 58–91. <https://doi.org/10.1002/mpr.1411>
- Erbe, D., Eichert, H.-C., Riper, H., & Ebert, D. D. (2017). Blending face-to-face and Internet-based interventions for the treatment of mental disorders in adults: Systematic review. *Journal of Medical Internet Research*, 19(9), e306. <https://doi.org/10.2196/jmir.6588>
- EuroQol Group. (1990). EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy*, 16(3), 199–208. <http://www.ncbi.nlm.nih.gov/pubmed/10109801>
- Fekadu, A., Wooderson, S. C., Markopoulou, K., Donaldson, C., Papadopoulos, A., & Cleare, A. J. (2009). What happens to patients with treatment-resistant depression? A systematic review of medium to long term outcome studies. *Journal of Affective Disorders*, 116(1–2), 4–11. <https://doi.org/10.1016/j.jad.2008.10.014>
- Fenwick, E., Claxton, K., & Sculpher, M. (2001). Representing uncertainty: The role of cost-effectiveness acceptability curves. *Health Economics*, 10(8), 779–787. <http://www.ncbi.nlm.nih.gov/pubmed/11747057>
- Fenwick, E., O'Brien, B. J., & Briggs, A. (2004). Cost-effectiveness acceptability curves-facts, fallacies and frequently asked questions. *Health Economics*, 13(5), 405–415. <https://doi.org/10.1002/hec.903>
- Flückiger, C., Del, A. C., Wampold, B. E., & Horvath, A. O. (2018). The alliance in adult psychotherapy: A meta-analytic synthesis. *Psychotherapy*, 55(4), 316–340. <https://doi.org/10.1037/pst0000172>

References

- Gerhards, S. A. H., de Graaf, L. E., Jacobs, L. E., Severens, J. L., Huibers, M. J. H., Arntz, A., Riper, H., Widdershoven, G., Metsemakers, J. F. M., & Evers, S. M. A. A. (2010). Economic evaluation of online computerised cognitive-behavioural therapy without support for depression in primary care: Randomised trial. *The British Journal of Psychiatry*, 196(4), 310–318. <https://doi.org/10.1192/bjp.bp.109.065748>
- Government of the Netherlands. (2019). Where can I get help for mental health problems? <https://www.government.nl/topics/mental-health-services/question-and-answer/help-for-mental-health-problems>
- Gyani, A., Shafran, R., Myles, P., & Rose, S. (2014). The gap between science and practice: How therapists make their clinical decisions. *Behavior Therapy*, 45(2), 199–211. <https://doi.org/10.1016/j.beth.2013.10.004>
- Hakkaart-van Roijen, L., Tan, S. S., & Bouwmans, C. A. M. (2010). Handleiding voor kostenonderzoek - Methoden en standaard kostprijszen voor economische evaluaties in de gezondheidszorg. Instituut voor Medical Technology Assessment Erasmus Universiteit Rotterdam.
- Hakkaart-van Roijen, L., van der Linden, N., Bouwmans, C., Kanters, T., & Tan, S. S. (2016). Kostenhandleiding: Methodologie van kostenonderzoek en referentieprijzen voor economische evaluaties in de gezondheidszorg.
- Hakkaart-van Roijen, L., van Straten, A., Tiemens, B., & Donker, M. C. H. (2002). Trimbos/iMTA questionnaire for costs associated with psychiatric illness (TIC-P). Institute of Medical Technology Assessment (iMTA).
- Hamilton, M. (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery & Psychiatry*, 23(1), 56–62. <https://doi.org/10.1136/jnnp.23.1.56>
- Harrell, F. E. (2018). Regression modeling strategies (5.1-2). <http://biostat.mc.vanderbilt.edu/rms>
- Harris, M. G., Baxter, A. J., Reavley, N., Diminic, S., Pirkis, J., & Whiteford, H. A. (2015). Gender-related patterns and determinants of recent help-seeking for past-year affective, anxiety and substance use disorders: Findings from a national epidemiological survey. *Epidemiology and Psychiatric Sciences*, 2016, 1–14. <https://doi.org/10.1017/S2045796015000876>
- Harvey, A. G., & Gumpert, N. B. (2015). Evidence-based psychological treatments for mental disorders: Modifiable barriers to access and possible solutions. *Behaviour Research and Therapy*, 68, 1–12. <https://doi.org/10.1016/j.brat.2015.02.004>
- Hatcher, R. L., & Gillaspy, J. A. (2006). Development and validation of a revised short version of the working alliance inventory. *Psychotherapy Research*, 16(1), 12–25. <https://doi.org/10.1080/10503300500352500>
- Hedman, E., Ljótsson, B., Kaldo, V., Hesser, H., El Alaoui, S., Kraepelien, M., Andersson, E., Rück, C., Svanborg, C., Andersson, G., & Lindefors, N. (2014). Effectiveness of Internet-based cognitive behaviour therapy for depression in routine psychiatric care. *Journal of Affective Disorders*, 155(1), 49–58. <https://doi.org/10.1016/j.jad.2013.10.023>

- Hesser, H. (2015). Modeling individual differences in randomized experiments using growth models: Recommendations for design, statistical analysis and reporting of results of internet interventions. *Internet Interventions*, 2(2), 110–120.
<https://doi.org/10.1016/j.invent.2015.02.003>
- Hickie, I. B., Davenport, T. A., Luscombe, G. M., Moore, M., Griffiths, K. M., & Christensen, H. (2010). Practitioner-supported delivery of internet-based cognitive behaviour therapy: Evaluation of the feasibility of conducting a cluster randomised trial. *The Medical Journal of Australia*, 192(11 Supp.), S31-35.
<http://www.ncbi.nlm.nih.gov/pubmed/20528705>
- Higgins, C., Walter, M., Heft, H., Soulodre, C., Laing, A., Kaulback, K., Mitchell, A., Thota, A., Ng, V., Sikich, N., & Dhalla, I. (2017). Psychotherapy for major depressive disorder and generalized anxiety disorder: A health technology assessment. *Ontario Health Technology Assessment Series*, 17(15), 1–167. <http://www.hqontario.ca/evidence-to-improve-care/journal-ontario-health-technology-assessment-series>
- Høifødt, R. S., Lillevoll, K. R., Griffiths, K. M., Wilsgaard, T., Eisemann, M., Waterloo, K., & Kolstrup, N. (2013). The clinical effectiveness of web-based cognitive behavioral therapy with face-to-face therapist support for depressed primary care patients: Randomized controlled trial. *Journal of Medical Internet Research*, 15(8), e153.
<https://doi.org/10.2196/jmir.2714>
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6(2), 65–70. <https://doi.org/10.2307/4615733>
- Honig, A., Hendriks, C. H., Akkerhuis, G. W., & Nolen, W. A. (2001). Usefulness of the retrospective Life-Chart method manual in outpatients with a mood disorder: A feasibility study. *Patient Education and Counseling*, 43(1), 43–48.
[https://doi.org/10.1016/S0738-3991\(00\)00144-0](https://doi.org/10.1016/S0738-3991(00)00144-0)
- Horvath, A. O., Del Re, A. C., Flückiger, C., & Symonds, D. (2011). Alliance in individual psychotherapy. *Psychotherapy*, 48(1), 9–16. <https://doi.org/10.1037/a0022186>
- Horvath, A. O., & Greenberg, L. S. (1989). Development and validation of the Working Alliance Inventory. *Journal of Counseling Psychology*, 36(2), 223–233.
<https://doi.org/10.1037//0022-0167.36.2.223>
- Husereau, D., Drummond, M., Petrou, S., Carswell, C., Moher, D., Greenberg, D., Augustovski, F., Briggs, A. H., Mauskopf, J., & Loder, E. (2013). Consolidated Health Economic Evaluation Reporting Standards (CHEERS) - explanation and elaboration: a report of the ISPOR Health Economic Evaluation Publication Guidelines Good Reporting Practices Task Force. *Value in Health: The Journal of the International Society for Pharmacoeconomics and Outcomes Research*, 16(2), 231–250.
<https://doi.org/10.1016/j.jval.2013.02.002>
- Jacobson, N. S., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59(1), 12–19. <https://doi.org/10.1037/0022-006X.59.1.12>

References

- Johansson, R., & Andersson, G. (2012). Internet-based psychological treatments for depression. *Expert Review of Neurotherapeutics*, 12(7), 861–869; quiz 870. <https://doi.org/10.1586/ern.12.63>
- Joling, K. J., van Marwijk, H. W. J., Piek, E., van der Horst, H. E., Penninx, B. W. J. H., Verhaak, P., & van Hout, H. P. J. (2011). Do GPs' medical records demonstrate a good recognition of depression? A new perspective on case extraction. *Journal of Affective Disorders*, 133(3), 522–527. <https://doi.org/10.1016/j.jad.2011.05.001>
- Karyotaki, E., Ebert, D. D., Donkin, L., Riper, H., Twisk, J., Burger, S., Rozental, A., Lange, A., Williams, A. D., Zarski, A. C., Geraedts, A., van Straten, A., Kleiboer, A., Meyer, B., Ünlü Ince, B. B., Buntrock, C., Lehr, D., Snoek, F. J., Andrews, G., ... Cuijpers, P. (2018). Do guided internet-based interventions result in clinically relevant changes for patients with depression? An individual participant data meta-analysis. *Clinical Psychology Review*, 63(June), 80–92. <https://doi.org/10.1016/j.cpr.2018.06.007>
- Karyotaki, E., Riper, H., Twisk, J., Hoogendoorn, A., Kleiboer, A., Mira, A., MacKinnon, A., Meyer, B., Botella, C., Littlewood, E., Andersson, G., Christensen, H., Klein, J. P., Schröder, J., Bretón-López, J., Scheider, J., Griffiths, K., Farrer, L., Huibers, M. J. H., ... Cuijpers, P. (2017). Efficacy of self-guided internet-based cognitive behavioral therapy in the treatment of depressive symptoms: A meta-analysis of individual participant data. *JAMA Psychiatry*, 74(4), 351–359. <https://doi.org/10.1001/jamapsychiatry.2017.0044>
- Karyotaki, E., Smit, Y., Holdt HenningSEN, K., Huibers, M. J. H., Robays, J., De Beurs, D., & Cuijpers, P. (2016). Combining pharmacotherapy and psychotherapy or monotherapy for major depression? A meta-analysis on the long-term effects. *Journal of Affective Disorders*, 194(1), 144–152. <https://doi.org/10.1016/j.jad.2016.01.036>
- Kelders, S. M., Bohlmeijer, E. T., Pots, W. T. M., & van Gemert-Pijnen, J. E. W. C. (2015). Comparing human and automated support for depression: Fractional factorial randomized controlled trial. *Behaviour Research and Therapy*, 72, 72–80. <https://doi.org/10.1016/j.brat.2015.06.014>
- Keller, M. B. (2003). Past, present, and future directions for defining optimal treatment outcome in depression: Remission and beyond. *JAMA*, 289(23), 3152–3160. <https://doi.org/10.1001/jama.289.23.3152>
- Kemmeren, L. L., van Schaik, D. J. F., Riper, H., Kleiboer, A. M., Bosmans, J. E., & Smit, J. H. (2016). Effectiveness of blended depression treatment for adults in specialised mental healthcare: Study protocol for a randomised controlled trial. *BMC Psychiatry*, 16, 113. <https://doi.org/10.1186/s12888-016-0818-5>
- Kenter, R. M. F., van de Ven, P. M., Cuijpers, P., Koole, G., Niamat, S., Gerrits, R. S., Willems, M., & van Straten, A. (2015). Costs and effects of Internet cognitive behavioral treatment blended with face-to-face treatment: Results from a naturalistic study. *Internet Interventions*, 2(1), 77–83. <https://doi.org/10.1016/j.invent.2015.01.001>
- Kessler, D., Lewis, G., Kaur, S., Wiles, N., King, M., Weich, S., Sharp, D. J., Araya, R., Hollinghurst, S., & Peters, T. J. (2009). Therapist-delivered internet psychotherapy for depression in primary care: A randomised controlled trial. *The Lancet*, 374(9690), 628–634. [https://doi.org/10.1016/S0140-6736\(09\)61257-5](https://doi.org/10.1016/S0140-6736(09)61257-5)

- Kessler, R. C., Chiu, W. T., Demler, O., Merikangas, K. R., & Walters, E. E. (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62(6), 617–627.
<https://doi.org/10.1001/archpsyc.62.6.617>
- Kessler, R. C., Sampson, N. A., Berglund, P., Gruber, M. J., Al-Hamzawi, A., Andrade, L., Bunting, B., Demyttenaere, K., Florescu, S., de Girolamo, G., Gureje, O., He, Y., Hu, C., Huang, Y., Karam, E., Kovess-Masfety, V., Lee, S., Levinson, D., Medina Mora, M. E., ... Wilcox, M. A. (2015). Anxious and non-anxious major depressive disorder in the World Health Organization World Mental Health Surveys. *Epidemiology and Psychiatric Sciences*, 24(3), 210–226. <https://doi.org/10.1017/S2045796015000189>
- Kleiboer, A., Smit, J., Bosmans, J., Ruwaard, J., Andersson, G., Topooco, N., Berger, T., Krieger, T., Botella, C., Baños, R., Chevreul, K., Araya, R., Cerga-Pashoja, A., Cieślak, R., Rogala, A., Vis, C., Draisma, S., van Schaik, A., Kemmeren, L., ... Moher, D. (2016). European comparative effectiveness research on blended Depression treatment versus treatment-as-usual (E-COMPARED): Study protocol for a randomized controlled, non-inferiority trial in eight European countries. *Trials*, 17(1), 387. <https://doi.org/10.1186/s13063-016-1511-1>
- Köhler, S., Chrysanthou, S., Guhn, A., & Sterzer, P. (2019). Differences between chronic and nonchronic depression: Systematic review and implications for treatment. *Depression and Anxiety*, 36(1), 18–30. <https://doi.org/10.1002/da.22835>
- Kohn, R., Saxena, S., Levav, I., & Saraceno, B. (2004). The treatment gap in mental health care. *Bulletin of the World Health Organization*, 82(11), 858–866. <https://doi.org/10.1596/9789241562004001100011>
- Kolovos, S., van Dongen, J. M., Riper, H., Buntrock, C., Cuijpers, P., Ebert, D. D., Geraedts, A. S., Kenter, R. M., Nobis, S., Smith, A., Warmerdam, L., Hayden, J. A., van Tulder, M. W., & Bosmans, J. E. (2018). Cost effectiveness of guided Internet-based interventions for depression in comparison with control conditions: An individual-participant data meta-analysis. *Depression and Anxiety*, 35(3), 209–219. <https://doi.org/10.1002/da.22714>
- Kooistra, L. C., Ruwaard, J., Wiersma, J. E., van Oppen, P., van der Vaart, R., van Gemert-Pijnen, J. E. W. C., & Riper, H. (2016). Development and initial evaluation of blended cognitive behavioural treatment for major depression in routine specialized mental health care. *Internet Interventions*, 4, 61–71. <https://doi.org/10.1016/j.invent.2016.01.003>
- Kooistra, L. C., Wiersma, J. E., Ruwaard, J. J., Riper, H., Penninx, B. W. J. H., & van Oppen, P. (2018). Six-year healthcare trajectories of adults with anxiety and depressive disorders: Determinants of transition to specialised mental healthcare. *Journal of Affective Disorders*, 241(April), 226–234. <https://doi.org/10.1016/j.jad.2018.07.072>
- Kooistra, L. C., Wiersma, J. E., Ruwaard, J., Neijenhuijs, K., Lokkerbol, J., van Oppen, P., Smit, F., & Riper, H. (2019). Costs and effectiveness of blended versus standard cognitive behavioral therapy for depressed outpatients in routine specialized mental health care: Pilot randomized controlled trial. *Journal of Medical Internet Research*, 21(10), e14261. <https://doi.org/10.2196/14261>

References

- Kooistra, L. C., Wiersma, J. E., Ruwaard, J., van Oppen, P., Smit, F., Lokkerbol, J., Cuijpers, P., & Riper, H. (2014). Blended vs. face-to-face cognitive behavioural treatment for major depression in specialized mental health care: Study protocol of a randomized controlled cost-effectiveness trial. *BMC Psychiatry*, 14(1), 290. <https://doi.org/10.1186/s12888-014-0290-z>
- Lamers, L. M., McDonnell, J., Stalmeier, P. F. M., Krabbe, P. F. M., & Busschbach, J. J. V. (2006). The Dutch tariff: Results and arguments for an effective design for national EQ-5D valuation studies. *Health Economics*, 15(10), 1121–1132. <https://doi.org/10.1002/hec.1124>
- Larsen, D. L., Attkisson, C. C., Hargreaves, W. A., & Nguyen, T. D. (1979). Assessment of client/patient satisfaction: development of a general scale. *Evaluation and Program Planning*, 2(3), 197–207. [https://doi.org/10.1016/0149-7189\(79\)90094-6](https://doi.org/10.1016/0149-7189(79)90094-6)
- Laws, H. B., Constantino, M. J., Sayer, A. G., Klein, D. N., Kocsis, J. H., Manber, R., Markowitz, J. C., Rothbaum, B. O., Steidtmann, D., Thase, M. E., & Arnow, B. A. (2017). Convergence in patient-therapist therapeutic alliance ratings and its relation to outcome in chronic depression treatment. *Psychotherapy Research : Journal of the Society for Psychotherapy Research*, 27(4), 410–424. <https://doi.org/10.1080/10503307.2015.1114687>
- Leon, A. C., Davis, L. L., & Kraemer, H. C. (2011). The role and interpretation of pilot studies in clinical research. *Journal of Psychiatric Research*, 45(5), 626–629.
<https://doi.org/10.1016/j.jpsychires.2010.10.008>
- Lépine, J.-P., & Briley, M. (2011). The increasing burden of depression. *Neuropsychiatric Disease and Treatment*, 7(Suppl 1), 3–7. <https://doi.org/10.2147/NDT.S19617>
- Lokkerbol, J., Adema, D., Cuijpers, P., Reynolds, C. F., Schulz, R., Weehuizen, R., & Smit, F. (2014). Improving the cost-effectiveness of a healthcare system for depressive disorders by implementing telemedicine: A health economic modeling study. *The American Journal of Geriatric Psychiatry : Official Journal of the American Association for Geriatric Psychiatry*, 22(3), 253–262. <https://doi.org/10.1016/j.jagp.2013.01.058>
- Lorenzo-Luaces, L., Driessen, E., DeRubeis, R. J., Van, H. L., Keefe, J. R., Hendriksen, M., & Dekker, J. (2017). Moderation of the alliance-outcome association by prior depressive episodes: Differential effects in cognitive-behavioral therapy and short-term psychodynamic supportive psychotherapy. *Behavior Therapy*, 48(5), 581–595.
<https://doi.org/10.1016/j.beth.2016.11.011>
- Lovell, K., & Richards, D. (2000). Multiple Access Points & Levels of Entry (MAPLE): Ensuring choice, accessibility & equity for CBT services. *Behavioural and Cognitive Psychotherapy*, 28, 379–391.
<http://linkinghub.elsevier.com/retrieve/pii/S221478291400030X>
- Ly, K. H., Janni, E., Wrede, R., Sedem, M., Donker, T., Carlbring, P., & Andersson, G. (2015). Experiences of a guided smartphone-based behavioral activation therapy for depression: A qualitative study. *Internet Interventions*, 2(1), 60–68.
<https://doi.org/10.1016/j.invent.2014.12.002>
- Lyketsos, C. G., Nestadt, G., Cwi, J., & Heithoff, K. (1994). The Life Chart Interview: A standardized method to describe the course of psychopathology. *International Journal of Methods in Psychiatric Research*, 4(3), 143–155.

- Måansson, K. N. T., Skagius Ruiz, E., Gervind, E., Dahlin, M., & Andersson, G. (2013). Development and initial evaluation of an Internet-based support system for face-to-face cognitive behavior therapy: A proof of concept study. *Journal of Medical Internet Research*, 15(12), e280. <https://doi.org/10.2196/jmir.3031>
- Mathers, C. D., & Loncar, D. (2006). Projections of global mortality and burden of disease from 2002 to 2030. *PLOS Medicine*, 3(11), e442. <https://doi.org/10.1371/journal.pmed.0030442>
- Mauskopf, J. A., Sullivan, S. D., Annemans, L., Caro, J., Mullins, C. D., Nuijten, M., Orlewska, E., Watkins, J., & Trueman, P. (2007). Principles of good practice for budget impact analysis: report of the ISPOR Task Force on good research practices-budget impact analysis. *Value in Health: The Journal of the International Society for Pharmacoeconomics and Outcomes Research*, 10(5), 336–347. <https://doi.org/10.1111/j.1524-4733.2007.00187.x>
- McCrone, P. (2004). Cost-effectiveness of computerised cognitive-behavioural therapy for anxiety and depression in primary care: Randomised controlled trial. *The British Journal of Psychiatry*, 185(1), 55–62. <https://doi.org/10.1192/bjp.185.1.55>
- McCullough, J. P. (1999). Treatment for chronic depression: Cognitive behavioral analysis system of psychotherapy. Guilford Press.
- Meadows, G., Harvey, C., Fossey, E., & Burgess, P. (2000). Assessing perceived need for mental health care in a community survey: Development of the Perceived Need for Care Questionnaire (PNCQ). *Social Psychiatry and Psychiatric Epidemiology*, 35(9), 427–435. <http://www.ncbi.nlm.nih.gov/pubmed/11089671>
- Meyer, B., Berger, T., Caspar, F., Beevers, C. G., Andersson, G., & Weiss, M. (2009). Effectiveness of a novel integrative online treatment for depression (Deprexis): Randomized controlled trial. *Journal of Medical Internet Research*, 11(2), e15. <https://doi.org/10.2196/jmir.1151>
- Mickey, R. M., & Greenland, S. (1989). The impact of confounder selection criteria on effect estimation. *American Journal of Epidemiology*, 129(1), 125–137. <http://www.ncbi.nlm.nih.gov/pubmed/2910056>
- Moher, D., Hopewell, S., Schulz, K. F., Montori, V., Gøtzsche, P. C., Devereaux, P. J., Elbourne, D., Egger, M., & Altman, D. G. (2010). CONSORT 2010 explanation and elaboration: Updated guidelines for reporting parallel group randomized trials. *Journal of Clinical Epidemiology*, 63(8), e1–37. <https://doi.org/10.1016/j.jclinepi.2010.03.004>
- Moher, D., Schulz, K. F., & Altman, D. (2001). The CONSORT statement: Revised recommendations for improving the quality of reports of parallel-group randomized trials. *JAMA*, 285(15), 1987–1991. <https://doi.org/10.1001/jama.285.15.1987>
- Mohr, D. C., Hart, S. L., Julian, L., Catledge, C., Honos-Webb, L., Vella, L., & Tasch, E. T. (2005). Telephone-administered psychotherapy for depression. *Archives of General Psychiatry*, 62(9), 1007–1014. <https://doi.org/10.1001/archpsyc.62.9.1007>
- Mohr, D. C., Lyon, A. R., Lattie, E. G., Reddy, M., & Schueller, S. M. (2017). Accelerating digital mental health research from early design and creation to successful implementation and sustainment. *Journal of Medical Internet Research*, 19(5), e153. <https://doi.org/10.2196/jmir.7725>

References

- Mol, M., Dozeman, E., Provoost, S., Van Schaik, A., Riper, H., & Smit, J. H. (2018). Behind the scenes of online therapeutic feedback in blended therapy for depression: Mixed-methods observational study. *Journal of Medical Internet Research*, 20(5), 1–11. <https://doi.org/10.2196/jmir.9890>
- Mol, M., Dozeman, E., van Schaik, D. J. F., Vis, C. P. C. D., Riper, H., & Smit, J. H. (2016). The therapist's role in the implementation of internet-based cognitive behavioural therapy for patients with depression: Study protocol. *BMC Psychiatry*, 16(1), 338. <https://doi.org/10.1186/s12888-016-1045-9>
- Munder, T., Wilmers, F., Leonhart, R., Linster, H. W., & Barth, J. (2010). Working alliance inventory-short revised (WAI-SR): Psychometric properties in outpatients and inpatients. *Clinical Psychology and Psychotherapy*, 17(3), 231–239. <https://doi.org/10.1002/cpp.658>
- Murphy, J. A., & Byrne, G. J. (2012). Prevalence and correlates of the proposed DSM-5 diagnosis of chronic depressive disorder. *Journal of Affective Disorders*, 139(2), 172–180. <https://doi.org/10.1016/j.jad.2012.01.033>
- National Institute for Health and Care Excellence (NICE). (2011). Common mental health problems: Identification and pathways to care. May. <https://www.nice.org.uk/guidance/cg123>
- Paykel, E. S., Ramana, R., Cooper, Z., Hayhurst, H., Kerr, J., & Barocka, A. (1995). Residual symptoms after partial remission: An important outcome in depression. *Psychological Medicine*, 25(6), 1171–1180. <https://doi.org/10.1017/S0033291700033146>
- Pearlin, L. I., & Schooler, C. (1978). The structure of coping. *Journal of Health and Social Behavior*, 19(1), 2–21. <https://doi.org/10.2307/2136319>
- Peduzzi, P., Concato, J., Feinstein, A. R., & Holford, T. R. (1995). Importance of events per independent variable in proportional hazards regression analysis. II. Accuracy and precision of regression estimates. *Journal of Clinical Epidemiology*, 48(12), 1503–1510. [https://doi.org/10.1016/0895-4356\(95\)00048-8](https://doi.org/10.1016/0895-4356(95)00048-8)
- Penninx, B. W. J. H., Beekman, A. T. F., Smit, J. H., Zitman, F. G., Nolen, W. A., Spinhoven, P., Cuijpers, P., de Jong, P. J., van Marwijk, H. W. J., Assendelft, W. J. J., van der Meer, K., Verhaak, P., Wensing, M., de Graaf, R., Hoogendoijk, W. J., Ormel, J., van Dyck, R., & NESDA Research Consortium. (2008). The Netherlands Study of Depression and Anxiety (NESDA): Rationale, objectives and methods. *International Journal of Methods in Psychiatric Research*, 17(3), 121–140. <https://doi.org/10.1002/mpr.256>
- Piek, E., van der Meer, K., Hoogendoijk, W. J. G., Penninx, B. W. J. H., & Nolen, W. A. (2011). Most antidepressant use in primary care is justified; results of the Netherlands study of depression and anxiety. *PLOS ONE*, 6(3), 1–8. <https://doi.org/10.1371/journal.pone.0014784>
- Piek, E., van der Meer, K., Penninx, B. W. J. H., Verhaak, P. F. M., & Nolen, W. a. (2011). Referral of patients with depression to mental health care by Dutch general practitioners: An observational study. *BMC Family Practice*, 12(1), 41. <https://doi.org/10.1186/1471-2296-12-41>

- Preschl, B., Maercker, A., & Wagner, B. (2011). The working alliance in a randomized controlled trial comparing online with face-to-face cognitive-behavioral therapy for depression. *BMC Psychiatry*, 11(1), 189. <https://doi.org/10.1186/1471-244X-11-189>
- Prins, M. A., Verhaak, P. F., Hilbink-Smolders, M., Spreeuwenberg, P., Laurant, M. G., van der Meer, K., van Marwijk, H. W., Penninx, B. W., & Bensing, J. M. (2011). Outcomes for depression and anxiety in primary care and details of treatment: A naturalistic longitudinal study. *BMC Psychiatry*, 11, 180. <https://doi.org/10.1186/1471-244X-11-180>
- Prins, M., Meadows, G., Bobevski, I., Graham, A., Verhaak, P., Van derMeer, K., Penninx, B. W. J. H., & Bensing, J. (2011). Perceived need for mental health care and barriers to care in the Netherlands and Australia. *Social Psychiatry and Psychiatric Epidemiology*, 46(10), 1033–1044. <https://doi.org/10.1007/s00127-010-0266-3>
- R Core Team. (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing.
- Ramsey, S., Willke, R., Briggs, A., Brown, R., Buxton, M., Chawla, A., Cook, J., Glick, H., Liljas, B., Petitti, D., & Reed, S. (2005). Good research practices for cost-effectiveness analysis alongside clinical trials: The ISPOR RCT-CEA Task Force report. *Value in Health: The Journal of the International Society for Pharmacoeconomics and Outcomes Research*, 8(5), 521–533. <https://doi.org/10.1111/j.1524-4733.2005.00045.x>
- Revelle, W. (2018). Psych: Procedures for psychological, psychometric, and personality research (R package version 1.8.4). Northwestern University.
- Richards, D., & Richardson, T. (2012). Computer-based psychological treatments for depression: A systematic review and meta-analysis. *Clinical Psychology Review*, 32(4), 329–342. <https://doi.org/10.1016/j.cpr.2012.02.004>
- Riper, H., van Ballegooijen, W., Kooistra, L., de Wit, J., & Donker, T. (2013). Preventie & eMental-health. Vrije Universiteit commisioned by ZonMw.
- Robertson, L., Smith, M., Castle, D., & Tannenbaum, D. (2006). Using the Internet to enhance the treatment of depression. *Australasian Psychiatry: Bulletin of Royal Australian and New Zealand College of Psychiatrists*, 14(4), 413–417. <https://doi.org/10.1111/j.1440-1665.2006.02315.x>
- Robinson, R. (1993). Cost-utility analysis. *BMJ (Clinical Research Ed.)*, 307(6908), 859–862. <https://doi.org/10.1136/bmj.307.6908.859>
- Romijn, G., Riper, H., Kok, R., Donker, T., Goorden, M., van Roijen, L. H., Kooistra, L., van Balkom, A., & Koning, J. (2015). Cost-effectiveness of blended vs. face-to-face cognitive behavioural therapy for severe anxiety disorders: Study protocol of a randomized controlled trial. *BMC Psychiatry*, 15, 311. <https://doi.org/10.1186/s12888-015-0697-1>
- Rush, A. J., Carmody, T., & Reimitz, P. (2000). The Inventory of Depressive Symptomatology (IDS): Clinician (IDS-C) and Self-Report (IDS-SR) ratings of depressive symptoms. *International Journal of Methods in Psychiatric Research*, 9(2), 45–59. <https://doi.org/10.1002/mpr.79>
- Rush, A. J., Giles, D. E., Schlessier, M. A., Fulton, C. L., Weissenburger, J., & Burns, C. (1986). The Inventory for Depressive Symptomatology (IDS): Preliminary findings. *Psychiatry Research*, 18(1), 65–87. [https://doi.org/10.1016/0165-1781\(86\)90060-0](https://doi.org/10.1016/0165-1781(86)90060-0)

References

- Rush, A. J., Gullion, C. M., Basco, M. R., Jarrett, R. B., & Trivedi, M. H. (1996). The Inventory of Depressive Symptomatology (IDS): Psychometric properties. *Psychological Medicine*, 26(3), 477-486. <http://www.ncbi.nlm.nih.gov/pubmed/8733206>
- Rush, A. J., Trivedi, M. H., Ibrahim, H. M., Carmody, T. J., Arnow, B., Klein, D. N., Markowitz, J. C., Ninan, P. T., Kornstein, S., Manber, R., Thase, M. E., Kocsis, J. H., & Keller, M. B. (2003). The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): A psychometric evaluation in patients with chronic major depression. *Biological Psychiatry*, 54(5), 573-583.
[https://doi.org/10.1016/S0006-3223\(02\)01866-8](https://doi.org/10.1016/S0006-3223(02)01866-8)
- Ruwaard, J., & Kok, R. N. (2015). Wild West eHealth: Time to hold our horses? *European Health Psychologist*, 17(1), 45-49.
- Ruwaard, J., Lange, A., Schrieken, B., Dolan, C. V., & Emmelkamp, P. (2012). The effectiveness of online cognitive behavioral treatment in routine clinical practice. *PLOS ONE*, 7(7). <https://doi.org/10.1371/journal.pone.0040089>
- Sauro, J. (2011). Measuring usability with the system usability scale (SUS). <http://www.measuringusability.com/sus.php>
- Schröder, J., Berger, T., Westermann, S., Klein, J. P., & Moritz, S. (2016). Internet interventions for depression: new developments. *Dialogues in Clinical Neuroscience*, 18(2), 203-212. <https://doi.org/10.1136/bmj.1.3087.311-a>
- Schueller, S. M., Tomasino, K. N., & Mohr, D. C. (2017). Integrating human support into behavioral intervention technologies: The efficiency model of support. *Clinical Psychology: Science and Practice*, 24(1), 27-45. <https://doi.org/10.1111/cpsp.12173>
- Schulz, K. F., Altman, D. G., & Moher, D. (2011). CONSORT 2010 statement: Updated guidelines for reporting parallel group randomised trials. *International Journal of Surgery (London, England)*, 9(8), 672-677. <https://doi.org/10.1016/j.ijsu.2011.09.004>
- Schuster, R., Leitner, I., Carlbring, P., & Laireiter, A. R. (2017). Exploring blended group interventions for depression: Randomised controlled feasibility study of a blended computer- and multimedia-supported psychoeducational group intervention for adults with depressive symptoms. *Internet Interventions*, 8(December 2016), 63-71. <https://doi.org/10.1016/j.invent.2017.04.001>
- Shapiro, S. S., & Wilk, M. B. (1965). An analysis of variance test for normality (complete samples). *Biometrika*, 52(3), 591-611.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59 Suppl 2, 22-33;quiz 34-57. <http://www.ncbi.nlm.nih.gov/pubmed/9881538>
- Smit, F., Cuijpers, P., Oostenbrink, J., Batelaan, N., de Graaf, R., & Beekman, A. (2006). Costs of nine common mental disorders: Implications for curative and preventive psychiatry. *The Journal of Mental Health Policy and Economics*, 9(4), 193-200.
<http://www.ncbi.nlm.nih.gov/pubmed/17200596>

- Smit, F., Willemse, G., Meulenbeek, P., Koopmanschap, M., van Balkom, A., Spinhoven, P., & Cuijpers, P. (2009). Preventing panic disorder: Cost-effectiveness analysis alongside a pragmatic randomised trial. *Cost Effectiveness and Resource Allocation*, 7, 8. <https://doi.org/10.1186/1478-7547-7-8>
- Smolders, M., Laurant, M., Verhaak, P., Prins, M., van Marwijk, H., Penninx, B. W. J. H., Wensing, M., & Grol, R. (2010). Which physician and practice characteristics are associated with adherence to evidence-based guidelines for depressive and anxiety disorders? *Medical Care*, 48(3), 240-248. <https://doi.org/10.1097/MLR.0b013e3181ca27f6>
- Spijker, J., Bockting, C. L. H., Meeuwissen, J. A. C., van Vliet, I. M., Emmelkamp, P. M. G., Hermens, M. L. M., & van Balkom, A. J. L. M. (2013). Multidisciplinaire richtlijn depressie. Richtlijn voor de diagnostiek, behandeling en begeleiding van volwassen patienten met een depressieve stoornis. (3rd ed.). Trimbos-instituut.
- http://www.ggzrichtlijnen.nl/index.php?pagina=/richtlijn/item/pagina.php&richtlijn_id=88
- Steer, R. A., Beck, A. T., Clark, D. A., & Beck, J. S. (1994). Psychometric properties of the Cognition Checklist with psychiatric outpatients and university students. *Psychological Assessment*, 6(1), 67-70.
- Stinckens, N., Ulburghs, A., & Claes, L. (2009). De werkalliantie als sleutelelement in het therapiergebeuren: Meting met behulp van de WAV-12, de Nederlandstalige versie. *Tijdschrift Voor Klinische Psychologie*, 39, 44-60.
- Stringer, B., van Meijel, B., Eikelenboom, M., Koekkoek, B., Verhaak, P. F. M., Kerkhof, A. J. M. F., Penninx, B. W. J. H., & Beekman, A. T. F. (2013). Perceived need for care and health care utilization among depressed and anxious patients with and without suicidal ideation. *Crisis*, 34(3), 192-199. <https://doi.org/10.1027/0227-5910/a000182>
- Tan, S., Bouwmans-Frijters, C. A. M., & Roijen, L. (2012). Handleiding voor kostenonderzoek: Methoden en referentieprijs voor economische evaluaties in de gezondheidszorg. *Tijdschrift Voor Gezondheidswetenschappen*, 90(6), 367-372. <https://doi.org/10.1007/s12508-012-0128-3>
- ten Have, M., Nuyen, J., Beekman, A., & de Graaf, R. (2013). Common mental disorder severity and its association with treatment contact and treatment intensity for mental health problems. *Psychological Medicine*, 43(10), 2203-2213. <https://doi.org/10.1017/S0033291713000135>
- ten Have, M., de Graaf, R., van Dorsselaer, S., & Beekman, A. (2013). Lifetime treatment contact and delay in treatment seeking after first onset of a mental disorder. *Psychiatric Services (Washington, D.C.)*, 64(10), 981-989. <https://doi.org/10.1176/appi.ps.201200454>
- Thase, M. E., Wright, J. H., Eells, T. D., Barrett, M. S., Wisniewski, S. R., Balasubramani, G. K., McCrone, P., & Brown, G. K. (2018). Improving the efficiency of psychotherapy for depression: Computer-assisted versus standard CBT. *The American Journal of Psychiatry*, 175(3), 242-250. <https://doi.org/10.1176/appi.ajp.2017.17010089>
- Therneau, T. M. (2017). Survival Analysis (2.41-3). <https://cran.r-project.org/package=survival%3E>

References

- Thompson, S. C., & Schlehofer, M. M. (2008). The many sides of control motivation. Motives for high, low and illusory control. In J. Y. Shah & W. L. Gardner (Eds.), *Handbook of Motivation Science* (pp. 41-56). Guilford Press.
- Titov, N. (2011). Internet-delivered psychotherapy for depression in adults. *Current Opinion in Psychiatry*, 24(1), 18-23. <https://doi.org/10.1097/YCO.0b013e32833ed18f>
- Trivedi, M. H., Rush, A. J., Ibrahim, H. M., Carmody, T. J., Biggs, M. M., Suppes, T., Crismon, M. L., Shores-Wilson, K., Toprac, M. G., Dennehy, E. B., Witte, B., & Kashner, T. M. (2004). The Inventory of Depressive Symptomatology, Clinician Rating (IDS-C) and Self-Report (IDS-SR), and the Quick Inventory of Depressive Symptomatology, Clinician Rating (QIDS-C) and Self-Report (QIDS-SR) in public sector patients with mood disorders. *Psychological Medicine*, 34(1), 73-82. <https://doi.org/10.1017/S0033291703001107>
- Trivedi, M. H., Rush, A. J., Wisniewski, S. R., Nierenberg, A. A., Warden, D., Ritz, L., Norquist, G., Howland, R. H., Lebowitz, B., McGrath, P. J., Shores-Wilson, K., Biggs, M. M., Balasubramani, G. K., & Fava, M. (2006). Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: Implications for clinical practice. *American Journal of Psychiatry*, 163(1), 28-40. <https://doi.org/10.1176/appi.ajp.163.1.28>
- van 't Hof, E., Cuijpers, P., & Stein, D. J. (2009). Self-help and Internet-guided interventions in depression and anxiety disorders: A systematic review of meta-analyses. *CNS Spectrums*, 14(2 Suppl 3), 34-40. <https://doi.org/10.1017/S1092852900027279>
- van Balkom, A. J. L. M., van Vliet, I. M., Emmelkamp, P. M. G., Bockting, C. L. H., Spijker, J., Hermens, M. L. M., & Meeuwissen, J. A. C. (2013). Multidisciplinaire richtlijn angststoornissen. Richtlijn voor de diagnostiek, behandeling en begeleiding van volwassen patiënten met een angststoornis (3rd ed.). Trimbos-instituut.
- http://www.ggzrichtlijnen.nl/index.php?pagina=/richtlijn/item/pagina.php&richtlijn_id=35
- van Ballegooijen, W., Cuijpers, P., van Straten, A., Karyotaki, E., Andersson, G., Smit, J. H., & Riper, H. (2014). Adherence to Internet-based and face-to-face cognitive behavioural therapy for depression: A meta-analysis. *PLOS ONE*, 9(7), e100674. <https://doi.org/10.1371/journal.pone.0100674>
- van Beljouw, I. M. J., Verhaak, P. F. M., Cuijpers, P., van Marwijk, H. W. J., & Penninx, B. W. J. H. (2010). The course of untreated anxiety and depression, and determinants of poor one-year outcome: A one-year cohort study. *BMC Psychiatry*, 10(86), 86. <https://doi.org/10.1186/1471-244X-10-86>
- van den Brink, M., van den Hout, W. B., Stigkelbout, A. M., Putter, H., van de Velde, C. J. H., & Kievit, J. (2005). Self-reports of health-care utilization: Diary or questionnaire? *International Journal of Technology Assessment in Health Care*, 21(3), 298-304. <https://doi.org/10.1017/S0266462305050397>
- van der Vaart, R., Witting, M., Riper, H., Kooistra, L., Bohlmeijer, E. T., & van Gemert-Pijnen, L. J. E. W. C. (2014). Blending online therapy into regular face-to-face therapy for depression: Content, ratio and preconditions according to patients and therapists using a Delphi study. *BMC Psychiatry*, 14, 355. <https://doi.org/10.1186/s12888-014-0355-z>

- van der Zanden, R., Galindo-Garre, F., Curie, K., Kramer, J., & Cuijpers, P. (2013). Online cognitive-based intervention for depression: Exploring possible circularity in mechanisms of change. *Psychological Medicine*, 44, 1159–1170.
<https://doi.org/10.1017/S003329171300175X>
- van Gemert-Pijnen, J. E., Kelders, S. M., & Bohlmeijer, E. T. (2014). Understanding the usage of content in a mental health intervention for depression: An analysis of log data. *Journal of Medical Internet Research*, 16(1), e27. <https://doi.org/10.2196/jmir.2991>
- van Hemert, A. M., Kerkhof, A. J. F. M., De Keijser, J., Verwey, B., van Boven, C., Hummelen, J. W., De Groot, M. H., Lucassen, P., Meerdinkveldboom, J., Steendam, M., Stringer, B., Verlinde, A. A., & van de Glind, G. (2012). Multidisciplinaire richtlijn diagnostiek en behandeling van suïcidaal gedrag. 1–454.
- van Roijen, L. H., van Straten, A., Al, M., Rutten, F., & Donker, M. (2006). Cost-utility of brief psychological treatment for depression and anxiety. *The British Journal of Psychiatry*, 188, 323–329. <https://doi.org/10.1192/bjp.188.4.323>
- van Vliet, I. M., & de Beurs, E. (2007). Het Mini Internationaal Neuropsychiatrisch Interview (MINI). Een kort gestructureerd diagnostisch psychiatrisch interview voor DSM-IV en ICD-10-stoornissen. *Tijdschrift Voor Psychiatrie*, 49(6), 393–397.
- Verhaak, P. F. M., Prins, M. A., Spreeuwenberg, P., Draisma, S., van Balkom, A. J. L. M., Bensing, J. M., Laurant, M. G. H., van Marwijk, H. W. J., van der Meer, K., & Penninx, B. W. J. H. (2009). Receiving treatment for common mental disorders. *General Hospital Psychiatry*, 31(1), 46–55. <https://doi.org/10.1016/j.genhosppsych.2008.09.011>
- Vernmark, K., Hesser, H., Topooco, N., Berger, T., Riper, H., Luuk, L., Backlund, L., Carlbring, P., & Andersson, G. (2019). Working alliance as a predictor of change in depression during blended cognitive behaviour therapy. *Cognitive Behaviour Therapy*, 48(4), 285–299. <https://doi.org/10.1080/16506073.2018.1533577>
- Vis, C., Kleiboer, A., Prior, R., Bønes, E., Cavallo, M., Clark, S. A., Dozeman, E., Ebert, D., Etzelmueller, A., Favaretto, G., Zabala, A. F., Kolstrup, N., Mancin, S., Mathiassen, K., Myrbakk, V. N., Mol, M., Jimenez, J. P., Power, K., van Schaik, A., ... Riper, H. (2015). Implementing and up-scaling evidence-based eMental health in Europe: The study protocol for the MasterMind project. *Internet Interventions*, 2(4), 399–409. <https://doi.org/10.1016/j.invent.2015.10.002>
- Waller, G. (2009). Evidence-based treatment and therapist drift. *Behaviour Research and Therapy*, 47(2), 119–127. <https://doi.org/10.1016/j.brat.2008.10.018>
- Wampold, B. E. (2015). How important are the common factors in psychotherapy? An update. *World Psychiatry*, 14(3), 270–277. <https://doi.org/10.1002/wps.20238>
- Wang, P. S., Aguilar-Gaxiola, S., Alonso, J., Angermeyer, M. C., Borges, G., Bromet, E. J., Bruffaerts, R., de Girolamo, G., de Graaf, R., Gureje, O., Haro, J. M., Karam, E. G., Kessler, R. C., Kovess, V., Lane, M. C., Lee, S., Levinson, D., Ono, Y., Petukhova, M., ... Wells, J. E. (2007). Worldwide use of mental health services for anxiety, mood, and substance disorders: Results from 17 countries in the WHO world mental health surveys. *The Lancet*, 370(9590), 841–850. [https://doi.org/10.1016/S0140-6736\(07\)61414-7](https://doi.org/10.1016/S0140-6736(07)61414-7). Worldwide

References

- Wang, P. S., Angermeyer, M., Borges, G., Bruffaerts, R., Tat Chiu, W., DE Girolamo, G., Fayyad, J., Gureje, O., Haro, J. M., Huang, Y., Kessler, R. C., Kovess, V., Levinson, D., Nakane, Y., Oakley Brown, M. A., Ormel, J. H., Posada-Villa, J., Aguilar-Gaxiola, S., Alonso, J., ... Ustün, T. B. (2007). Delay and failure in treatment seeking after first onset of mental disorders in the World Health Organization's World Mental Health Survey Initiative. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)*, 6(3), 177–185. <http://www.ncbi.nlm.nih.gov/pubmed/2174579>
- Ware, J. E. (2000). SF-36 health survey update. *Spine*, 25(24), 3130–3139 <http://www.ncbi.nlm.nih.gov/pubmed/11124729>
- Ware, J. E. J., Snow, K. K., Kosinski, M., & Gandek, B. (1993). SF-36 Health Survey: Manual and interpretation guide. New England Medical Center, Health Institute.
- Warmerdam, L., Smit, F., van Straten, A., Riper, H., & Cuijpers, P. (2010). Cost-utility and cost-effectiveness of internet-based treatment for adults with depressive symptoms: Randomized trial. *Journal of Medical Internet Research*, 12(5), e53. <https://doi.org/10.2196/jmir.1436>
- Watkins, K. E., Burnam, M. A., Orlando, M., Escarce, J. J., Huskamp, H. a, & Goldman, H. H. (2009). The health value and cost of care for major depression. *Value in Health: The Journal of the International Society for Pharmacoeconomics and Outcomes Research*, 12(1), 65–72. <https://doi.org/10.1111/j.1524-4733.2008.00388.x>
- Webb, C. A., Rosso, I. M., & Rauch, S. L. (2017). Internet-based cognitive-behavioral therapy for depression: Current progress and future directions. *Harvard Review of Psychiatry*, 25(3), 114–122. <https://doi.org/10.1097/HRP.oooooooooooo00000139>
- Weitz, E. S., Hollon, S. D., Twisk, J., van Straten, A., Huibers, M. J. H., David, D., DeRubeis, R. J., Dimidjian, S., Dunlop, B. W., Cristea, I. A., Faramarzi, M., Hegerl, U., Jarrett, R. B., Kheirkhah, F., Kennedy, S. H., Mergl, R., Miranda, J., Mohr, D. C., Rush, A. J., ... Cuijpers, P. (2015). Baseline depression severity as moderator of depression outcomes between cognitive behavioral therapy vs pharmacotherapy: An individual patient data meta-analysis. *JAMA Psychiatry*, 72(11), 1102–1109. <https://doi.org/10.1001/jamapsychiatry.2015.1516>
- Wentzel, J., van der Vaart, R., Bohlmeijer, E. T., & van Gemert-Pijnen, J. E. W. C. (2016). Mixing online and face-to-face therapy: How to benefit From blended care in mental health care. *JMIR Mental Health*, 3(1), e9. <https://doi.org/10.2196/mental.4534>
- White, I. R., Carpenter, J., & Horton, N. J. (2012). Including all individuals is not enough: Lessons for intention-to-treat analysis. *Clinical Trials*, 9(4), 396–407. <https://doi.org/10.1177/1740774512450098>
- Whiteford, H. A., Degenhardt, L., Rehm, J., Baxter, A. J., Ferrari, A. J., Erskine, H. E., Charlson, F. J., Norman, R. E., Flaxman, A. D., Johns, N., Burstein, R., Murray, C. J. L., & Vos, T. (2013). Global burden of disease attributable to mental and substance use disorders: Findings from the Global Burden of Disease Study 2010. *Lancet*, 382(9904), 1575–1586. [https://doi.org/10.1016/S0140-6736\(13\)61611-6](https://doi.org/10.1016/S0140-6736(13)61611-6)

- Wilhelmsen, M., Lillevoll, K., Risør, M. B., Høifødt, R., Johansen, M.-L., Waterloo, K., Eisemann, M., & Kolstrup, N. (2013). Motivation to persist with internet-based cognitive behavioural treatment using blended care: a qualitative study. *BMC Psychiatry*, 13(1), 296–304. <https://doi.org/10.1186/1471-244X-13-296>
- Williams, A. D., & Andrews, G. (2013). The effectiveness of internet cognitive behavioural therapy (iCBT) for depression in primary care: A quality assurance study. *PLOS ONE*, 8(2). <https://doi.org/10.1371/journal.pone.0057447>
- Wiltsey Stirman, S., Gutner, C. A., Crits-Christoph, P., Edmunds, J., Evans, A. C., & Beidas, R. S. (2015). Relationships between clinician-level attributes and fidelity-consistent and fidelity-inconsistent modifications to an evidence-based psychotherapy. *Implementation Science*, 10(1), 1–10. <https://doi.org/10.1186/s13012-015-0308-z>
- Wittchen, H. U., Jacobi, F., Rehm, J., Gustavsson, A., Svensson, M., Jönsson, B., Olesen, J., Allgulander, C., Alonso, J., Faravelli, C., Fratiglioni, L., Jennum, P., Lieb, R., Maercker, A., van Os, J., Preisig, M., Salvador-Carulla, L., Simon, R., & Steinhausen, H. C. (2011). The size and burden of mental disorders and other disorders of the brain in Europe 2010. *European Neuropsychopharmacology*, 21(9), 655–679. <https://doi.org/10.1016/j.euroneuro.2011.07.018>
- World Health Organization (WHO). (1997). Composite International Diagnostic Interview, Core version 2.1: Interviewer's Manual. World Health Organisation.
- World Health Organization (WHO). (2018). Factsheet depression.
- World Health Organization Collaboration Centre for Drug Statistics Methodology. (2016). World Health Organization (WHO) Collaborating Centre for Drug Statistics Methodology. The Anatomical Therapeutic Chemical (ATC) classification system. <http://www.whocc.no/>
- Wright, J. H., Wright, A. S., Albano, A. M., Basco, M. R., Goldsmith, L. J., Raffield, T., & Otto, M. W. (2005). Computer-assisted cognitive therapy for depression: maintaining efficacy while reducing therapist time. *American Journal of Psychiatry*, 162(6), 1158–1164. <https://doi.org/10.1176/appi.ajp.162.6.1158>
- Zilcha-Mano, S. (2017). Is the alliance really therapeutic? Revisiting this question in light of recent methodological advances. *American Psychologist*, 72(4), 311–325. <https://doi.org/10.1037/a0040435>
- Zwerenz, R., Becker, J., Knickenberg, R. J., Siepmann, M., Hagen, K., & Beutel, M. E. (2017). Online self-help as an add-on to inpatient psychotherapy: Efficacy of a new blended treatment approach. *Psychotherapy and Psychosomatics*, 86(6), 341–350. <https://doi.org/10.1159/000481177>