

# VU Research Portal

## Urbanisation as a risk indicator for complex psychiatric disorders and forced admissions

Schoevers, R.; Peen, J.; Dekker, J.J.M.

### **published in**

BMC Psychiatry  
2007

### **DOI (link to publisher)**

[10.1186/1471-244X-7-S1-S148](https://doi.org/10.1186/1471-244X-7-S1-S148)

### **document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

### **citation for published version (APA)**

Schoevers, R., Peen, J., & Dekker, J. J. M. (2007). Urbanisation as a risk indicator for complex psychiatric disorders and forced admissions. *BMC Psychiatry*, 7(S1), S148. <https://doi.org/10.1186/1471-244X-7-S1-S148>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

Oral presentation

Open Access

## Urbanisation as a risk indicator for complex psychiatric disorders and forced admissions

Robert Schoevers\*, Jaap Peen and Jack Dekker

Address: Mentrum Mental Health Care Amsterdam, 2e Constantijn Huyjgensstraat 37, 1054 AG Amsterdam, Netherlands

\* Corresponding author

from WPA Thematic Conference. Coercive Treatment in Psychiatry: A Comprehensive Review  
Dresden, Germany. 6–8 June 2007

Published: 19 December 2007

BMC Psychiatry 2007, 7(Suppl 1):S148 doi:10.1186/1471-244X-7-S1-S148

This abstract is available from: <http://www.biomedcentral.com/1471-244X/7/S1/S148>

© 2007 Schoevers et al; licensee BioMed Central Ltd.

### Background

To determine both prevalence and complexity of psychiatric disorders in the Amsterdam area in relation to other major cities and less urbanized areas in the Netherlands, and to assess whether this is related to higher levels of (coercive) admissions.

### Methods

These associations were explored in a nationwide epidemiological study and the national admission register, and in a local study of the Amsterdam region examining health care use patterns.

### Results

The admission rate for the whole of the Netherlands was twice as high in the group of most highly urbanized municipalities as in the group of least urbanized municipalities. The urban/rural variations in admission rates in the Netherlands are reflected in true psychiatric morbidity rates. The authors found an urban/rural difference in total annual prevalence figures for psychiatric disorders in the population. The difference was also found for the separate disorders, mood disorders and substance-induced disorders, but not for anxiety disorders. Both prevalence and complexity of psychopathology in terms of comorbidity and severity were significantly higher in Amsterdam compared to other larger cities in the Netherlands, as were the number of coercive admissions.

### Conclusion

There is evidence regarding a link between urbanisation, the development of complex psychiatric disorders and the number of (forced) admissions to PICU's [1].

### References

1. Peen J, Dekker J: **Is urbanicity an environmental risk-factor for psychiatric disorders?** *Lancet* 2004, **363**:2013-2014.