

VU Research Portal

Treating obsessions with Competitive Memory Training: A pilot study.

Korrelboom, K.; van der Gaag, M.; Hendriks, V.M.; Huijbrechts, I.; Berretty, E.W.

published in

The Behavior Therapist
2008

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Korrelboom, K., van der Gaag, M., Hendriks, V. M., Huijbrechts, I., & Berretty, E. W. (2008). Treating obsessions with Competitive Memory Training: A pilot study. *The Behavior Therapist*, 31(2), 29-35.
<http://www.abct.org/Members/?m=mMembers&fa=JournalsPeriodicals#sec3>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

the Behavior Therapist

Contents

Clinical Forum

Kees Korrelboom, Mark van der Gaag, Vincent M. Hendriks, Irma Huijbrechts, and Ed W. Berretty
Treating Obsessions With Competitive Memory Training:
A Pilot Study 29

Student Forum

Eric L. Sullivan
Selecting a Theoretical Orientation:
Tips and Interdisciplinary Considerations 36

Letter to the Editor

Milton Spett 40

At ABCT

Katherine Martinez
Find a Therapist ... Who Takes Your Insurance,
Has Saturday Appointments, Is Located Nearby,
and More ... 40

- Call for Award Nominations / Student Awards Program 42
- Call for Papers for the 42nd Annual Meeting 44

*Coming directly to your mailbox soon—
your ABCT membership card!*

FEATURING . . .

- Your unique ABCT membership number
- Important dates
- Useful URLs

Keep this card handy in your wallet. Please be sure we have your correct mailing address! If we don't, please call the central office (212-647-1890).

Clinical Forum

Treating Obsessions With Competitive Memory Training: A Pilot Study

Kees Korrelboom, Mark van der Gaag,
Vincent M. Hendriks, Irma Huijbrechts,
and Ed W. Berretty, *PsyQ Haaglanden,
Parnassigroup, The Hague*

Since the 1980s, in vivo exposure combined with response prevention (ERP) has been considered the psychological intervention of choice in the treatment of obsessive-compulsive disorder (OCD). Exposing patients to their feared stimuli while preventing them from performing neutralizing rituals not only reduces the compulsive behaviors of the patient, but also the frequency and emotional impact of accompanying obsessions (Emmelkamp, Van Oppen, & Van Balkom, 2002).

In OCD, “normal” cognitive intrusions, which are occasionally experienced by everybody, develop into pathological obsessions because OCD patients catastrophically misinterpret the personal meaning of these intrusive thoughts and, hence, become anxious (Salkovskis, 1985). Anxiousness and dysfunctional attempts to neutralize anxiety-provoking intrusions broaden the range of potentially dangerous stimuli and increase the number of intrusive thoughts (Rachman, 1997, 1998). While earlier cognitive interventions such as Meichenbaum’s Self-Instruction Training (Emmelkamp et al., 1980) and Ellis’ Rational Emotive Therapy (Emmelkamp & Beens, 1991; Emmelkamp, Visser, & Hoekstra, 1988) did not enhance the effectiveness of ERP, recent theorizing in OCD has stimulated the development of new cognitive interventions. These interventions have been ap-

the Behavior Therapist

Published by the Association for
Behavioral and Cognitive Therapies
305 Seventh Avenue - 16th Floor
New York, NY 10001-6008
(212) 647-1890/Fax: (212) 647-1865
www.abct.org

EDITOR *Drew Anderson*
Editorial Assistant *Melissa Them*
Behavior Assessment *Timothy R. Stickle*
Book Reviews *C. Alix Timko*
Clinical Forum *John P. Forsyth*
Clinical Dialogues *Brian P. Marx*
Institutional
Settings. *David Penn*
Tamara Penix Sbraga
Lighter Side *Elizabeth Moore*
News and Notes. *David DiLillo*
Laura E. Dreer
James W. Sturges
Public Health Issues. *Jennifer Lundgren*
Research-Practice
Links. *David J. Hansen*
Research-Training
Links. *Gayle Y. Iwamasa*
Science Forum. *Jeffrey M. Lohr*
Special Interest
Groups *Andrea Seidner Burling*
Technology Update. *James A. Carter*

ABCT President *Anne Marie Albano*
Executive Director *Mary Jane Eimer*
Director of Education &
Meeting Services *Mary Ellen Brown*
Director of Communications *David Teisler*
Managing Editor *Stephanie Schwartz*

Copyright © 2008 by the Association for Behavioral and Cognitive Therapies. All rights reserved. No part of this publication may be reproduced or transmitted in any form, or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the copyright owner.

Subscription information: *the Behavior Therapist* is published in 8 issues per year. It is provided free to ABCT members. Nonmember subscriptions are available at \$40.00 per year (+\$32.00 airmail postage outside North America).

Change of address: 6 to 8 weeks are required for address changes. Send both old and new addresses to the ABCT office.

ABCT is committed to a policy of equal opportunity in all of its activities, including employment. ABCT does not discriminate on the basis of race, color, creed, religion, national or ethnic origin, sex, sexual orientation, gender identity or expression, age, disability, or veteran status.

All items published in *the Behavior Therapist*, including advertisements, are for the information of our readers, and publication does not imply endorsement by the Association.

PRESIDENTIAL PANELS

Dissemination and the Public Face of CBT

Moderator: Frank Andrasik
Past Presidents: David Barlow, K. Daniel O'Leary, Steven D. Hollon,
Antonette M. Zeiss, & Linda C. Sobell

The Evolution of CBT and Assessment

Moderator: Anne Marie Albano
Past Presidents: Thomas H. Ollendick, Richard G. Heimberg,
G. Terence Wilson, & G. Alan Marlatt

The Life and Times of AABT/ABCT: An Organizational Memoir

Moderator: Raymond DiGiuseppe
Past Presidents: Michael W. Otto, Richard M. Suinn, & Patricia Resick


Evolution of Evidence-Based Approaches

Moderator: Dianne L. Chambless
Past Presidents: Steven C. Hayes, Marsha M. Linehan, & W. Edward Craighead

The Professional Practice of CBT: Promises and Pitfalls

Moderator: Debra A. Hope
Past Presidents: Jacqueline B. Persons, J. Gayle Beck, Arthur M. Nezu,
Arthur Freeman, & Dennis C. Russo

*David Barlow • K. Daniel O'Leary •
Steven D. Hollon • Antonette M. Zeiss •
Linda C. Sobell • Thomas H. Ollendick •
Richard G. Heimberg • G. Terence Wilson
• G. Alan Marlatt • Michael W. Otto •
Richard M. Suinn • Patricia Resick •
Steven C. Hayes • Marsha M. Linehan •
W. Edward Craighead • Jacqueline B.
Persons • J. Gayle Beck • Arthur M. Nezu
• Arthur Freeman • Dennis C. Russo*



ORDER DVDS ON-LINE
www.abct.org

INSTRUCTIONS for AUTHORS

The Association for Behavioral and Cognitive Therapies publishes *the Behavior Therapist* as a service to its membership. Eight issues are published annually. The purpose is to provide a vehicle for the rapid dissemination of news, recent advances, and innovative applications in behavior therapy.

- Feature articles that are approximately 16 double-spaced manuscript pages may be submitted.
- Brief articles, approximately 6 to 12 double-spaced manuscript pages, are preferred.
- Feature articles and brief articles should be accompanied by a 75- to 100-word abstract.
- Letters to the Editor may be used to respond to articles published in *the Behavior Therapist* or to voice a professional opinion. Letters should be limited to approximately 3 double-spaced manuscript pages.

Submissions must be accompanied by a **Copyright Transfer Form** (a form is printed on p. 24 of the January 2008 issue of *tBT*, or contact the ABCT central office): *submissions will not be reviewed without a copyright transfer form*. Prior to publication authors will be asked to submit a final electronic version of their manuscript. Authors submitting materials to *tBT* do so with the understanding that the copyright of the published materials shall be assigned exclusively to ABCT. Submissions via e-mail are preferred and should be sent to the editor at drewa@albany.edu. Please include the phrase *tBT submission* in the subject line of your e-mail. Include the first author's e-mail address on the cover page of the manuscript attachment. By conventional mail, please send manuscripts to:

Drew A. Anderson, Ph.D.
SUNY-Albany
Dept. of Psychology/SS369
1400 Washington Ave.
Albany, NY 12222

plied to OCD patients who present without overt compulsive behaviors. It has been estimated that between 9% (Foa & Kozak, 1995) and 25% (Freeston et al., 1997) of patients diagnosed with OCD experience obsessions only. The sole use of ERP with these patients has produced ambiguous results, compared to those in patients with both obsessions and compulsions (Craske, 1999; Stanley & Averill, 1998). Supplementing ERP with cognitive interventions was found to be beneficial for primarily obsessional patients (Freeston et al., 1997).

Moreover, although ERP is considered an effective treatment for OCD patients with manifest compulsions, only a minority of these patients achieves a nonsymptomatic status after treatment (Abramowitz, 1998). While several studies have shown that obsessive thinking in these patients reduced after ERP alone, it is suggested that adding cognitive interventions to ERP might further enhance its efficacy (Van Oppen et al., 1995).

While cognitive behavioral therapy in general is considered to be an effective intervention for a range of disorders, debate continues about the mechanisms involved in instigating change. According to Brewin (2006), cognitive therapy does not directly modify negative information in memory, but rather influences the relative retrievability of the different meanings of emotional concepts stored in memory. Strengthening positive representations that are in "retrieval competition" with dysfunctional negative representations is considered to be the common target of various cognitive and behavioral procedures.

Prior to Brewin's (2006) publication, but largely consistent with his suggestions, a series of cognitive behavioral interventions had been developed for patients who failed to develop a positive change in feelings, despite adequate changes in cognition after traditional Socratic challenging and behavioral testing of their dysfunctional thoughts. These interventions were based on Lang's concept of cognitive emotional networks (Lang, 1985, 1994) and were considered to be a modern, cognitive variant of counterconditioning. Assuming that the negative feeling state of these patients was directed by Lang's cognitive emotional networks, counterconditioning—or Competitive Memory Training (COMET)—aims to change these networks. According to Lang, cognitive emotional networks consist of cognitive representations of emotionally relevant stimuli, responses to these stimuli, and the (implicit and/or explicit) thoughts

and interpretations about the relationship between both. Activation of an emotional network is a function of the number of matches between perceptions of the real world and the information already stored in long-term memory.

COMET is a stepwise cognitive-behavioral intervention based on this network theory of emotional representation. Basically, two variants of COMET protocols have been developed. One variant is aimed at changing the (felt) content of dysfunctional thoughts; the other is concerned with changing the attitude toward one's thoughts and thinking style. In the content variant, first the personal meaning and emotional theme of the dysfunctional emotional network are identified (e.g., intrusive thoughts about causing another person's death could mean that one is a "bad person"); next, a more functional but realistic alternative is formulated (e.g., "I am a good person"). Thereafter, this alternative is made more perceptible and retrievable for the patient by having him describe and imagine examples of positive situations in which these more functional alternatives were current (enhancing positive stimulus representations), by instructing him to (sub)vocalize positive self-statements in line with the imagination (enhancing positive meaning representations), and by asking him to accentuate and physically enact the positive images and statements by way a congruent posture and facial expression (stimulating positive response representations). Thus, several times a day the patient practices the strengthening of this competitive emotional network by imagining autobiographical situations in which he is, for instance, acting as a good and responsible adult, being helpful to others, or adequately performing responsible tasks. Implementation of these exercises is relatively straightforward, practiced in sets of no more than a couple of minutes each. The patient is simply asked to close his eyes and imagine himself being in the "positive situation." During these imaging practices, he sits with a straight back, strong shoulders, and a comforting facial expression, (sub)vocalizing, for example, "I am always there when I am needed." Then the exercise is stopped and the patient is asked whether he could experience at least "just a little bit of the positive feeling." If so, the exercise is repeated a couple of times. If not, the patient is helped to get a clearer image by giving him more guidance during his imagination, or by instructing him to be more expressive in his posture and facial expression. The patient is told that he has to become a "world

champion in activating good images and feelings" by practicing 6 times a day for 5 minutes each time. Once a strong and competitive new emotional network has been formed, the patient imagines problematic situations (negative stimulus representations) from the old dysfunctional network, meanwhile assuming the positive posture and facial expression (positive response representations) and (sub)vocalizing the positive self-verbalizations (positive meaning representations) of the newly formed network. So, while activating obsessive thoughts about, for example, the death of his children, he keeps his "good person" posture and expression and keeps telling himself that he "is always there when he is needed." It is thought that in this way the old rigid dysfunctional emotional network is transformed into a more functional one. Although the procedure of COMET resembles that of systematic desensitization in several aspects, the supposed mechanisms behind this procedure are quite different.

An example of an "attitude variant" of COMET pertains to worry. In this protocol the patient learns to distance himself from his worrisome preoccupations by changing stimulus, response, and meaning representations. After explanation of the rationale ("Every person has the capacity to cling to issues of personal importance as well as to let go of things that have lost this personal importance—this protocol helps you to identify your capacities of letting go and to use these capacities in instances where you can't, but should let go"), a patient chooses his or her worry themes. Then, the patient learns, in a stepwise fashion, first to change the images of his worry themes ("zooming out" or "blurring" the picture as can be done with a camera, or picturing his worries on a stage and watching them from a distance like the audience in a theatre). Then, while imaging, he takes on an accepting or uninvolved posture and facial expression, (sub)vocalizing, for example, "These thoughts are boring, I won't allow them to interfere with my life," or other thoughts and attitudes that have previously been helpful in other situations where the patient could indeed take a distance or was able to let things go.

In their discussion on the mechanisms of exposure therapy, Foa and Kozak also refer to Lang's theory. They suggest a two-stage process. First, the automatic process of habituation reduces the patient's fear responses *within* exposure sessions. Second, the experience that fear is reduced automatically during exposure makes the patient less anxious at the beginning of the next ex-

posure session. This is referred to as fear reduction *between* exposure sessions. The combined action of fear reduction within sessions and between sessions is considered to be the working mechanism of exposure therapy (Foa & Kozak, 1986). One of the main differences between COMET and Foa and Kozak's account seems to pertain to the deliberateness of the therapeutic steps in COMET. Whereas for Foa and Kozak, both processes of fear reduction proceed automatically, patients in COMET should willfully and deliberately activate their positive as well as their negative beliefs and fears. In this respect, COMET resembles cognitive training in adjusting cognitive biases, as suggested by Mackintosh et al. (2006).

Until now, COMET protocols for several emotional problems have been developed and applied, among them self-esteem, worry, panic, and frustration (Korrelboom & ten Broeke, 2004). One of the COMET protocols pertains to the treatment of obsessions in OCD. This article describes the first empirical test of this COMET protocol for obsessions. It concerns an uncontrolled naturalistic study in a routine outpatient mental health setting.

Method

Patients

Patients were recruited from the Department of Anxiety Disorders of PsyQ, Parnassia, a large public mental health organization in the Netherlands. The study sample consisted of 17 outpatients with marked obsessions (a score greater than 7 on the obsessions subscale of the Y-BOCS) and a diagnosis of current OCD (*DSM-IV*; American Psychiatric Association, 1994), who had responded insufficiently after at least 7 sessions of regular outpatient treatment for OCD (i.e., "treatment-resistant" patients), consisting of ERP and pharmacotherapy, as recommended in the Dutch Multidisciplinary Guidelines for Anxiety Disorders (LSMR, 2003). On average, these patients had received 30 weeks ($SD = 21$ weeks) of therapy prior to the start of the current intervention. At the start of COMET, mean Y-BOCS scores were 23.47 and mean BDI scores were 16.43. These are in the same range as in several other clinical samples, with mean Y-BOCS and BDI scores of, respectively, 26.79 and 18.44 (Franklin et al., 2000), and 24.9 and 15.7 (Vogel, Stiles, & Götestam, 2004) at the start of therapy.

Patients were predominantly female ($n = 11$), of Western cultural background ($n = 17$), and with a relatively high education

level (i.e., at least 13 years of education; $n = 11$). The mean age of the group was 33.1 years ($SD = 10.4$ years).

Instruments

The following assessments were conducted at baseline and posttreatment.

The Yale-Brown Obsessive Compulsive Scale–Dutch version (Y-BOCS; Goodman, Price, Rasmussen, Mazure, Delgado, et al., 1989; Goodman, Price, Rasmussen, Mazure, Fleishmann, et al., 1989). The Y-BOCS is a semistructured interview, designed to assess the severity of obsessive-compulsive symptoms. It yields a total score (Y-BOCS-tot) as well as two separate subscores: one for obsessions (Y-BOCS-obs) and one for compulsions (Y-BOCS-comp). The Y-BOCS is considered to be reliable and valid in assessing OCD symptomatology and is sensitive to treatment effects (Hiss, Foa, & Kozak, 1994). Y-BOCS ratings were made by the therapists themselves, so they were not independent. All therapists were trained in conducting Y-BOCS ratings.

The Beck Anxiety Inventory–Dutch version (BAI; Beck, Epstein, Brown, & Steer, 1988). The BAI is a 21-item self-administered questionnaire. Each item is scored on a 4-point Likert scale. The BAI is considered to be reliable and valid in measuring state anxiety (Osman et al., 1997).

The Beck Depression Inventory–Dutch translation (BDI; Beck et al., 1961). The BDI is a 21-item self-administered questionnaire, with a 4-point Likert scale for each item. The BDI has been demonstrated to be reliable (Bouman et al., 1985) and valid (Bouman, 1989) in measuring depression in Dutch populations.

Therapists

Patients were treated by four psychologists, all experienced in treating anxiety-disordered patients with CBT. All therapists had received two half days of training in conducting the COMET protocol. During the treatments, they were supervised by the first author for a total of 4 hours.

Procedure

All patients were informed by their therapists about the rationale, procedures, and treatment goal of COMET, and were asked to provide informed consent. After establishing the eligibility of each patient, the baseline assessments were conducted. The patients subsequently received 7 COMET sessions of 45 minutes each. Patients re-

ceived COMET as an add-on to their ongoing routine therapy.

During the first session, all patients received a comprehensive treatment manual, in which the treatment rationale of COMET was emphasized (i.e., "Obsessions are nothing but strange thoughts which reveal nothing important about the person, his future, and the world he lives in"). To achieve this, the obsession and its personal meaning to the patient were determined. For instance: "Having intrusive thoughts about sexually abusing my child means that I am an immoral person." Next, a credible alternative description of the personality of the patient (or his world or his future) was formulated. For instance: "I am a person of high moral standards" (Session 1). Then, this alternative was elaborated in such a way that its meaning was not only intellectually *understood* by the patient but, in particular, more emotionally *felt* by him (Sessions 2 to 4). This was achieved by helping the patient to identify examples of himself being "a person of high moral standards" and by having him write down such examples during homework assignments.

In the next three steps, this process of "making the patient feel what he already knows" was intensified by having the patient (in therapy sessions and during homework) *imagine* these examples, meanwhile accompanying these imaginations by (*sub*)*vocalizing* adequate words and sentences ("I am honest and reliable") and compatible *posture and facial expressions*. It has been demonstrated that emotional experience and attitude can be influenced by manipulating writing, imaging, self-verbalizations, and posture and facial expression (Camras, Holland, & Patterson, 1993; Laird, 1974; Lang, 1985; Lange et al., 1998; Schnall & Laird, 2003; Segal, Gemar, & Williams, 1999). From a theoretical point of view, writing and imaging are considered to activate stimulus representations, while posture and facial expression are thought to be ways of triggering response representations. Finally, self-verbalizations are supposed to be examples of meaning representations.

In Sessions 5 and 6 and subsequent homework practices, the patient attempted to connect the newly achieved positive feeling state to his obsessions. This was realized by asking the patient to activate his obsession (stimulus representation of the dysfunctional network) while, at the same time, activating his incompatible response representation (facial expression and posture) and meaning representations (self-verbalizations) of the alternative, functional

Table 1. Data of Completers on Symptomatology Before and After COMET for OCD

	Pretreatment	Posttreatment	<i>t</i> value	<i>df</i>	Significance	Cohen's <i>d</i>
Y-BOCS-tot	23.47 (5.34)	15.60 (5.70)	6.24	14	.000	1.42
Y-BOCS-obs	12.80 (3.41)	8.33 (2.16)	7.75	14	.000	1.57
Y-BOCS-comp	10.67 (2.94)	7.27 (4.42)	3.34	14	.005	0.91
BAI	18.79 (5.42)	12.43 (5.05)	3.67	13	.003	1.21
BDI	16.43 (7.73)	10.36 (6.20)	3.50	13	.004	0.87

Note. Y-BOCS = Yale-Brown Obsessive-Compulsive Scale (tot = total score; obs = obsessions subscore; comp = compulsions subscore); BAI = Beck Anxiety Inventory; BDI = Beck Depression Inventory.

Table 2. Data of Intention to Treat on Symptomatology Before and After COMET for OCD

	Pretreatment	Posttreatment	<i>t</i> value	<i>df</i>	Significance	Cohen's <i>d</i>
Y-BOCS-tot	22.24 (6.13)	15.29 (5.44)	5.44	16	.000	1.20
Y-BOCS-obs	12.53 (3.29)	8.59 (2.15)	6.35	16	.000	1.42
Y-BOCS-comp	9.71 (3.97)	6.71 (4.51)	3.21	16	.005	0.71
BAI	19.69 (8.93)	14.13 (9.58)	3.46	15	.003	0.60
BDI	17.06 (8.51)	11.75 (8.09)	3.29	15	.005	0.64

Note. Y-BOCS = Yale-Brown Obsessive-Compulsive Scale (tot = total score; obs = obsessions subscore; comp = compulsions subscore); BAI = Beck Anxiety Inventory; BDI = Beck Depression Inventory.

network. It is thought that the patient could then *experience* that his obsession is “nothing more than a strange thought, revealing nothing important about the patient, his future, or the world he lives in.”

In the last part of COMET (Sessions 6 and 7), the patient learned to distance himself from his obsession. Again, he was asked to activate his (now, supposedly, emotionally weakened) obsessive thought. Again, he had to activate incompatible motor and meaning responses. This time the alternative motor responses were “looking bored” or “looking amused” or “surprised,” assuming a physical posture congruent with this feeling state and, at the same time, (sub)vocalizing “bored,” “amused,” or “surprised” words and sentences. Also the patient was asked, if possible, to change the image of his

obsessions by “placing it at a distance in his mind (zooming out) or by making the picture of it less focused (blurring).”

Results

Of the 17 participating patients, 15 patients completed all seven COMET sessions. Both dropouts were female and they were slightly younger than the completers (23.5 versus 34.4 years). Also, dropouts had received less ERP therapy before COMET started (mean: 14 weeks versus 32 weeks), were more depressive (mean BDI: 21.50 versus 16.40), more anxious (mean BAI: 26.00 versus 18.79), and less obsessive-compulsive (mean YBOCS-tot: 13.00 versus 23.47) at the start of COMET. As far as significance can mean anything in this situation, the only significant difference was

found on the Y-BOCS-tot. As no follow-up data could be obtained from the treatment dropouts, pre- to posttreatment patients' changes were primarily analyzed in the treatment completers (see Table 1). However, we also did an intention-to-treat analysis by substituting the pretreatment scores of the two dropouts as their posttreatment scores (see Table 2).

As summarized in Table 1, all measures showed significant reductions of symptomatology from baseline to follow-up. These reductions would remain significant after Bonferroni correction for multiple testing. Effect sizes (Cohen's *d*: Cohen, 1988) were large (> .80) for all anxiety measures as well as for depression. In the intention-to-treat analysis (Table 2), all results remained significant, while effect sizes for compulsions,

general anxiety, and depression decreased from large to moderate (between .40 and .80), but stayed large for obsessions and OCD.

To estimate the number of patients that showed a clinically significant improvement, we followed the procedure of Jacobson and Truax (1991). In accordance with Steketee, Frost, and Bogert (1996) and Vogel et al. (2004), we calculated the percentage of patients that fell within the non-patient range by having a posttreatment Y-BOCS-tot < 16 (47%) as well as the percentage of patients with an Reliable Change Index (RCI) based on the Y-BOCS test-retest reliability of .79 (53%). Patients who had both an RCI and a posttreatment score on Y-BOCS-tot within the normal range were considered to have realized a clinically significant improvement during COMET (27%).

Discussion

This pilot study represents the first empirical investigation of COMET as an adjunct to ongoing routine treatment in OCD patients, specifically aimed at obsessive symptomatology. The results suggest that COMET may be effective in reducing both obsessive-compulsive symptoms and co-occurring depressive and anxiety symptoms in this population. In addition, the procedures used in COMET could be transferred relatively easily to therapists and were well accepted by the patients. These results are in line with findings from other preliminary studies in which variants of the COMET procedures were tested in other clinical conditions. In three uncontrolled studies (Olij et al., 2006; Peeters, Korrelboom, Voermans, & Huijbrechts, 2005; Van der Gaag, personal communication), patients with various psychiatric disorders (i.e., panic disorder, depression, eating disorder, PTSD, schizophrenia, and personality disorder) reacted positively to variants of COMET for treating low self-esteem, auditory hallucinations, and panic.

As a first pilot study in OCD, this study clearly has limitations. These include the small sample size, absence of a (preferably randomized) comparison condition, as well as the lack of a long-term follow-up and adequate control of the therapeutic procedures. In addition, COMET was provided as an adjunct to the patients' ongoing routine therapy. Therefore, the positive results found in this study cannot be attributed unequivocally to COMET per se. Nevertheless, patients had received their routine therapy for an average of about 6 months

prior to the start of COMET, but still exhibited symptom levels for OCD and depression that were comparable to baseline levels found in other studies in OCD patients (e.g., Foa et al., 2005; Vogel et al., 2004). Hence, whereas patients were still symptomatic after their routine therapy, the addition of COMET resulted in substantial reductions of their symptoms in a short period of time, and with effect sizes comparable to that of other studies in OCD (e.g., Foa et al., 2005; Van Balkom et al., 1994; Vogel et al., 2004). Based on the therapists' clinical impression, longer duration of COMET might have been even more beneficial for some patients, as might have a more deliberate tuning of COMET with ERP.

To summarize, COMET may be an effective (adjunctive) intervention for OCD. The promising results found in the current study merit further investigation of this intervention in a well-defined population, using a better controlled design, sufficiently large samples, and a sufficiently long follow-up period.

References

- Abramowitz, J. S. (1998). Does cognitive-behavioral therapy cure obsessive-compulsive disorder? A meta-analytic evaluation of clinical significance. *Behavior Therapy, 29*, 339-355.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Beck, A. T., Epstein, N., Brown, G., & Steer, R. A. (1988). An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting & Clinical Psychology, 56*, 893-897.
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J. E., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry, 18*, 561-571.
- Bouman, T. K. (1989). Assessment van stemmingsstoornissen (Assessment of mood disorders). In F. A. Albersnagel, P. M. G. Emmelkamp, & R. H. van den Hoofdakker (Eds.), *Depressie: Theorie, diagnostiek en behandeling* (Depression: Theory, diagnostics and treatment). Deventer: Van Loghum Slaterus.
- Bouman, T. K., Luteijn, F., Albersnagel, F. A., & van der Ploeg, F. A. E. (1985). Enige ervaringen met de Beck Depression Inventory (Some experiences with the Beck Depression Inventory). *Gedrag-Tijdschrift voor Psychologie, 13*, 13-24.
- Brewin, C. R. (2006). Understanding cognitive behaviour therapy: A retrieval competition account. *Behaviour Research & Therapy, 44*, 765-784.
- Camras, L. A., Holland, E. A., & Patterson, M. J. (1993). Facial expression. In M. Lewis & J. M. Haviland (Eds.), *Handbook of emotions*. New York: The Guilford Press.
- Cohen, J. (1988). *Statistical power analysis for the behavioural sciences* (2nd ed.). Hillsdale, NJ: Lawrence Erlbaum.
- Craske, M. G. (1999). *Anxiety disorders: Psychological approaches to theory and treatment*. Boulder, CO: Westview Press.
- Emmelkamp, P. M. G., & Beens, H. (1991). Cognitive therapy with obsessive-compulsive disorder: A comparative evaluation. *Behaviour Research & Therapy, 29*, 293-300.
- Emmelkamp, P. M. G., van der Helm, M., van Zanten, B. L., & Plogch, I. (1980). Treatment of obsessive-compulsive patients: The contribution of self-instructional training to the effectiveness of exposure. *Behaviour Research & Therapy, 18*, 61-66.
- Emmelkamp, P. M. G., van Oppen, P., & van Balkom, A. (2002). Cognitive changes in patients with obsessive-compulsive rituals treated with exposure in vivo and response prevention. In R. O. Frost & G. Steketee (Eds.), *Cognitive approaches to obsessions and compulsions: Theory, assessment, and treatment*. Oxford: Elsevier.
- Emmelkamp, P. M. G., Visser, S., & Hoekstra, R. J. (1988). Cognitive therapy vs. exposure in vivo in the treatment of obsessive-compulsives. *Cognitive Therapy & Research, 12*, 103-114.
- Foa, E. B., & Kozak, M. J. (1986). Emotional processing of fear: Exposure to corrective information. *Psychological Bulletin, 99*, 20-35.
- Foa, E. B., & Kozak, M. J. (1995). DSM-IV field trial: Obsessive-compulsive disorder. *American Journal of Psychiatry, 152*, 90-96.
- Foa, E. B., Liebowitz, M. R., Kozak, M. J., Davies, S., Campeas, R., Franklin, M. E., Huppert, J. D., Kjernisted, K., Rowan, V., Schmidt, A. B., Simpson, H. B., & Tu, X. (2005). Randomized, placebo-controlled trial of exposure and ritual prevention, clomipramine, and their combination in the treatment of obsessive-compulsive disorder. *American Journal of Psychiatry, 162*, 151-161.
- Franklin, M. E., Abramowitz, J. S., Kozak, M. J., Levitt, J. T., & Foa, E. B. (2000). Effectiveness of exposure and ritual prevention for obsessive-compulsive disorder: Randomized compared with non-randomized samples. *Journal of Consulting and Clinical Psychology, 68*, 594-602.
- Freeston, M. H., Ladouceur, R., Gagnon, F., Thibodeau, N., Rheume, J., Letarte, H., & Bujold, A. (1997). Cognitive-behavioral treatment of obsessive thoughts: A controlled study. *Journal of Consulting and Clinical Psychology, 65*, 405-413.
- Goodman, W. K., Price, L. H., Rasmussen, S. A., Mazure, C., Delgado, P., Henninger, G. R., & Charney, D. S. (1989). The Yale-Brown Obsessive-Compulsive Scale: II. Validity. *Archives of General Psychiatry, 46*, 1012-1016.
- Goodman, W. K., Price, L. H., Rasmussen, S. A., Mazure, C., Fleishmann, R. L., Hill, C. L., Henninger, G. R., & Charney, D. S. (1989).

The Yale-Brown Obsessive-Compulsive Scale: I. Development, use, and reliability. *Archives of General Psychiatry*, 46, 1006-1011.

Hiss, H., Foa, E. B., & Kozak, M. J. (1994). A relapse prevention program for obsessive-compulsive disorder. *Journal of Consulting and Clinical Psychology*, 62, 801-808.

Jacobson, N. S., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12-19.

Korrelboom, C. W., & ten Broeke, E. (2004). *Geïntegreerde cognitieve gedragstherapie: Handboek voor theorie en praktijk (Integrated cognitive behaviour therapy: Handbook for theory and practice)*. Bussum: Coutinho.

Laird, J. D. (1974). Self-attribution of emotion: The effects of expressive behavior on the quality of emotional experience. *Journal of Personality and Social Psychology*, 29, 475-486.

Landelijke Stuurgroep Multidisciplinaire Richtlijnontwikkeling in de GGZ (National Committee for the Development of Multidisciplinary Guidelines in Mental Health). (2003). *Multidisciplinaire richtlijn angststoornissen: Richtlijn voor de diagnostiek, behandeling en begeleiding van volwassen cliënten met een angststoornis (Multidisciplinary guideline for anxiety disorders: Guideline for diagnosis, treatment and counselling of adults with an anxiety disorder)*. Utrecht: Trimbos-Instituut.

Lang, P. J. (1985). The cognitive psychophysiology of emotion: Fear and anxiety. In A. H. Tuma & J. Maser (Eds.), *Anxiety and the anxiety disorders*. Hillsdale, NJ: Lawrence Erlbaum.

Lang, P. J. (1994). The motivational organization of emotions. In S. van Goozen, N. E. van de Poll, & J. A. Sergeant (Eds.), *Emotions: Essays on emotion theory*. Hillsdale, NJ: Lawrence Erlbaum.

Lange, A., Richard, R., Gest, A., de Vries, M., & Lodder, L. (1998). The effects of positive self-instruction: A controlled trial. *Cognitive Therapy & Research*, 22, 225-236.

Mackintosh, B., Mathews, A., Yiend, J., Ridgeway, V., & Cook, E. (2006). Induced biases in emotional interpretation influences stress vulnerability and endure despite changes in context. *Behavior Therapy*, 37, 209-222.

Olij, R. J. B., Korrelboom C. W., Huijbrechts, I. P. A. M., de Jong, M., Cloin, P. A., Maarsingh, M., & Paumen, B. N. W. (2006). De module zelfbeeld in een groep: werkwijze en eerste bevindingen (Treating low self-esteem in a group: procedure and first results). *Directieve Therapie*, 26, 307-325.

Osman, A., Kopper, B. A., Barrios, F. X., Osman, J. R., & Wade, T. (1997). The Beck Anxiety Inventory: Re-examination of factor structure and psychometric properties. *Journal of Clinical Psychology*, 53, 7-14.

International Association for Cognitive Psychotherapy

Facilitate the utilization and
growth of CT by joining IACP.

IACP serves as a resource and information
center for members to enhance
their professional and research activities.

2008 Membership Benefits include:

- ★ Free subscription to *The International Journal of Cognitive Therapy*, our new official journal published by Guilford Press
- ★ Free subscription to the *Cognitive Therapy Newsletter*
- ★ Access to state-of-the-art audio workshops on the web
- ★ Listing in the yearly membership directory
- ★ Optional listing in the online referral directory
- ★ Access to the IACP Listserve
- ★ Various discounts to conferences and book publishers

JOIN US NOW & SAVE \$140 on the
International Congress on Cognitive Psychotherapy (ICCP)
6th Convention, Rome 2008



To apply online, go to: www.the-iacp.com
Or contact the IACP Membership Office at:
iacpmembership@yahoo.com

Peeters, S., Korrelboom, C. W., Voermans, M., & Huijbrechts, I. P. A. M. (2005). Paniekmanagement revisited: Ervaringen met een nieuwe groepsbehandeling (Training in panic control revisited: Experiences with a new group therapy). *Directieve Therapie*, 25, 396-408.

Rachman, S. (1997). A cognitive theory of obsessions. *Behaviour Research & Therapy*, 35, 793-502.

Rachman, S. (1998). A cognitive theory of obsessions: Elaborations. *Behaviour Research & Therapy*, 36, 385-401.

Salkovskis, P. M. (1985). Obsessional-compulsive problems: A cognitive-behavioural analysis. *Behaviour Research & Therapy*, 23, 571-583.

Schnall, S., & Laird, J. D. (2003). Keep smiling: Enduring effects of facial expressions and postures on emotional experience and memory. *Cognition & Emotion*, 17, 787-797.

Segal, Z. V., Gemar, M., & Williams, S. (1999). Differential cognitive response to a mood challenge following successful cognitive therapy or pharmacotherapy for depression. *Journal of Abnormal Psychology*, 108, 3-10.

Stanley, M. A., & Averill, P. M. (1998). Psychosocial treatments for obsessive-compulsive disorder: Clinical applications. In R. P. Swinson, M. M. Antony, S. Rachman & M. A. Richter (Eds.), *Obsessive-compulsive disorder: Theory, research, and treatment*. New York: The Guilford Press.

Steketee, G., Frost, R., & Bogert, K. (1996). The Yale-Brown Obsessive-Compulsive Scale: Interview versus self-report. *Behaviour Research & Therapy*, 34, 675-684.

van Balkom, A. J. L. M., van Oppen, P., Vermeulen, A. W. A., van Dyck, R., Nauta, M. C. E., & Vorst H. C. M. (1994). A meta-analysis on the treatment of obsessive-compulsive disorder: A comparison of antidepressants, behavior and cognitive therapy. *Clinical Psychological Review*, 14, 359-381.

van Oppen, P., de Haan, E., van Balkom, A. J. L. M., Spinhoven, P., Hoogduin, C. A. L., & van Dyck, R. (1995). Cognitive therapy and exposure in vivo in the treatment of obsessive-compulsive disorder. *Behaviour Research & Therapy*, 33, 379-390.

Vogel, P. A., Stiles, T. C., & Götestam, K. G. (2004). Adding cognitive therapy elements to exposure therapy for obsessive-compulsive disorder: A controlled study. *Behavioural and Cognitive Psychotherapy*, 32, 275-290.

Address correspondence to C. W. Korrelboom, Ph.D., Head of Research & Innovation, PsyQ Haaglanden, Parnassia group, Mangostraat 15, 2552 KS, The Hague, Netherlands
email: k.korrelboom@psyq.nl



<http://www.abct.org>

Journals & Publications ↩

the Behavior Therapist ↩

Past issues ↩

Student Forum

Selecting a Theoretical Orientation: Tips and Interdisciplinary Considerations

Eric L. Sullivan, *Suffolk University*

Many students of clinical psychology, as part of their professional development, select a theoretical orientation from which to conceptualize, research, and treat psychopathology. Theoretical orientation is a useful heuristic that guides psychopathology conceptualization and treatment, making it faster and more efficient. An orientation provides a pre-packaged explanation of clinical problems and, usually, a method for changing them. Choosing an orientation, however, can be a daunting task, as there are many from which to select and compelling reasons to select any number of them. A nonexhaustive list of theoretical orientations and therapies includes behaviorism, cognitive therapy, cognitive behavioral therapy (CBT), humanism, psychodynamic therapies, gestalt therapy, existential therapy, and many more. In addition, many recent theorists and practitioners have elected to maintain an eclectic treatment approach, applying a variety of therapeutic technologies from across orientations and/or adopting their principles as part of a larger pastiche orientation.

Despite the oppressive number of possibilities from which to choose, there are steps that students can take to facilitate the selection process. These appear at the end of this

article. In addition, students should consider the theories and research findings of the various nonclinical fields within psychology as well as that of other disciplines. Fusing interdisciplinary knowledge with the theory, research, and practice of clinical psychology can augment, broaden, and deepen the student's understanding and treatment of psychopathology. This idea receives discussion in this article, using the theory of cognitive development as an example of how students might enhance their understanding of clinical phenomena. But first, it is important to consider the reason for selecting a theoretical orientation in the first place.

Why Select an Orientation?

During practicum and internship, many students often struggle with conceptualizing clinical problems and selecting treatments to address them. Many feel overwhelmed by the multitude of possible theoretical models and treatment packages from which to select. In this author's experience, treating clients in practicum raised many doubts and concerns. The questions, "Have I selected the *right* treatment to effectively address this individual's problems?" or even "Am I doing the client any good at all?" often arose. Working within a

theoretical orientation (such as CBT) can ease such concerns by offering a framework within which to understand client problems and a host of treatments that effectively address them. By adopting an orientation, the emerging clinician does not have to flounder with "reinventing the wheel." When using empirically supported treatments and theories, students can increase the likelihood that their case conceptualizations are accurate and their treatments effective because the wisdom and experience of a community of professionals and a large body of empirical support (in the case of CBT) predicated them.

Identifying with a theoretical orientation also provides an opportunity to enter a professional community, which most readers of this newsletter have done by joining ABCT. Professional organizations can provide support for students through e-mail listserves and conferences. Listserves and conferences are important and useful venues for students to explore and learn from the experience, wisdom, and scientific research of professionals. In such forums, professionals and students can exchange and evaluate ideas. These forums are also useful for networking with influential psychologists to build contacts and make friends. Conferences provide opportunities for students to hear, firsthand, the philosophy and science of the various treatments and concepts within an orientation.

An additional benefit of selecting a theoretical orientation is that it promotes consistency in research and clinical practice. This is useful for two reasons. First, it allows one to hone one's understanding of psychopathology and clinical skills to a finer degree. An orientation narrows the focus of conceptualization and treatment, which

provides more time to practice skills and evaluate theoretical conceptions. The example of sports illustrates this notion. That is, one may elect to play many sports or to focus on one sport. The more one elects to participate in multiple sports, the less likely he or she will be able to become expert at any one due to limitations in time and energy. By contrast, electing to focus on one sport allows a person to devote more time and energy to honing his or her abilities in that sport. If a basketball player practices foul shots daily, the amount of foul shots the player successfully makes will increase steadily. The same is true for clinical practice and research. For example, the more one practices thought monitoring (a component of cognitive therapy), the more effective one will become in skillfully leading clients in building awareness of automatic thoughts, identifying emotions linked to those thoughts, and eliciting alternative thoughts.

Second, the conceptual and clinical consistency that accompanies selecting a theoretical orientation also provides an anchor for exploring outside perspectives and ideas. When evaluating outside perspectives, these perspectives can be considered in contrast to a deep understanding of the theoretical framework within which one identifies. While contrasting ideas does not necessitate such an approach, it is useful from a practical standpoint: It prevents the confusion and bewilderment that could result from considering multiple orientations without a point of reference. It focuses exploration to make evaluating other ideas easier. Thus, while some may consider selecting a theoretical orientation confining, it may actually provide a platform from which one may explore other ideas and disciplines.

Selecting a theoretical orientation within which to ground conceptualization and treatment of psychopathology provides multiple benefits. Once one selects a theoretical orientation, however, it is easy to lose sight of the diversity of ideas and insight that nonclinical psychology research and disciplines outside of psychology (e.g., medicine) have to offer. In the following section, the theory of cognitive development serves as an example of how other branches of psychology (and outside disciplines) can augment understanding and treatment of psychopathology. Most readers of *the Behavior Therapist* have adopted a CBT orientation, which tends to emphasize the present and often deemphasizes development and experience. (However, this perspective does not discount their impor-

Looking to relocate to the beautiful Northwest?

FOR SALE

Psychology Practice Specializing in Anxiety Disorders

Lake Oswego, Oregon

The Anxiety Disorders Clinic (TADC) specializes in the evaluation and treatment of anxiety disorders in adults, adolescents, and children. Anxiety disorders affect 13 million people, surpassing substance abuse (10 million people) and depression (9.4 million people) as the number one mental health problem in the country.

TADC was established in 1985 by Ricks Warren, Ph.D., ABPP to provide the most current, evidence-based treatments for each of the anxiety disorders and related conditions. Clinic staff, in conjunction with the School for Professional Psychology at Pacific University, has pioneered in conducting effectiveness studies in a private practice setting, resulting in a rewarding national and international reputation. Studies have included panic disorder, OCD, PTSD, and social anxiety disorder.

TADC currently has six staff members: three psychologists, two LCSWs and a marriage and family therapist. Psychopharmacological evaluations and medication management are provided by a consulting psychiatric nurse practitioner certified in psychopharmacology and other consulting psychiatrists. TADC has an extensive referral network of primary care physicians, pediatricians, and other medical specialists, as well as friends, family, and work associates of former clients.

TADC is located at 4550 S.W. Kruse Way in Lake Oswego, Oregon. The office building is located in one of the most desirable professional office areas in the Portland Metropolitan area, and each office space in the suite has a beautiful view of lush Northwest vegetation. Dr. Warren is selling TADC so he can move closer to extended family.

For more information:

- see the TADC website at www.anxietydisordersclinic.com
- contact Dr. Warren at (503) 635-8710 #1
- or email rickswarren@comcast.net

<http://www.anxietydisordersclinic.com>

tance.) The theory and research of cognitive development can complement CBT conceptualizations by offering insight into the development of psychopathology.

Interdisciplinary Considerations in Selecting Theoretical Orientation

Richardson's (1998) classification system of cognitive development theories provides a useful framework within which to categorize conceptualizations of clinical issues. In Richardson's model, there are four basic schools of cognitive development: nativism, associationism, constructivism, and sociocognitive. These same divisions also underlie some theoretical differences between cognitive-behaviorists. These divisions are not independent, but rather, represent domains of theoretical emphasis or levels of analysis. Many clinical problems have biopsychosocial elements. Nonetheless, understanding these divisions will broaden student understanding of clinical phenomena and personal orientation and improve their ability to understand and change problems.

Nativism

Nativists explain cognitive development through genetic and biological factors. This perspective can also extend to behavior (e.g., skills, abilities, etc.). Nativism in its most extreme form would involve identifying genetic, inborn causes for every cognitive process and behavior. Most nativists, however, are not so extreme, emphasizing inborn and genetic causes to varying degrees. Noam Chomsky is a contemporary example of a nativist. According to his theory of language development (1980), humans are born with the ability to speak, including an innate understanding of grammar, which shapes one's acquisition and use of language.

In CBT, nativists view psychopathology in a similar vein. Nativist CBT therapists seek inherited or biological deficits. Some disorders have known biological underpinnings (such as schizophrenia and bipolar disorder). Therapists working with such clients might "work around" these inherited deficits by emphasizing existing skills, teaching skills through psychoeducation, and restructuring problematic cognition. One such approach is psychiatric rehabilitation (Pratt, Gill, Barrett, & Roberts, 2002).

Many psychiatrists also conceptualize clinical problems from a nativist perspective. They consider the acquired or genetic biological underpinnings of psychopathology and treat them by altering chemical

and other physiology. Interventions may include restoring neurotransmitter concentration balance via prescription medicine, changing the electric behavior of the nervous system via electroconvulsive therapy, or altering physical structures via psychosurgery. Recently, the nativist perspective has surged in popularity via mass media marketing of psychotropic drugs.

Associationism

Acquisition of information about the world and reinforcement of behavior via interaction with the environment forms the underpinnings of associationist explanations of cognition and behavior. Learning, in its simplest form, is the acquisition of associations, or connections between objects and events or behavior and its consequences. Classical (Pavlovian) and operant conditioning are two of the main processes through which people learn. B. F. Skinner is among the most famous of the behaviorists (i.e., associationists). Behavior therapy emerged, in large part, because of his work. He theorized and empirically demonstrated that contingencies in one's environment shape one's behavior (i.e., operant conditioning; Skinner, 1953). While not denying the existence of cognition (Skinner often spoke of "verbal behavior"), associationism traditionally has not emphasized cognition due to the inability to observe it.

Many past CBT therapies emphasized the conceptualization of cognitions and behaviors as learned associations formed via interaction between the individual and his or her environment. In these conceptualizations, psychopathology arises through reinforcement of problematic thoughts and behaviors. Early behavior therapists used reinforcement principles to shape adaptive behaviors or change old, maladaptive ones. In its simplest form, psychiatric hospitals have used token economies to shape the behavior of patients by reinforcing desirable behavior with reward and ignoring or punishing undesirable behavior. Behavior therapists also utilize psychoeducation about behavior and strive to create and reinforce new, adaptive behaviors, such as coping skills (see Wilson, 1995, for a review of behavior therapy). Current third-wave behavior therapies, such as Acceptance and Commitment Therapy, also emphasize action, for example, citing the role of experiential avoidance in problem formation and maintenance (Hayes, Strosahl, & Wilson, 1999). These therapies use behavioral exercises to reshape experience and alter client perspectives.

Associationism extends to an explanation of cognition as well. In part, cognitive schemas are simply a collection of learned associations. The study of schemas is the main content of theory and research in the associationist cognitive development literature. This model does not allow for meaning making to originate within the individual, however.

Constructivism

Constructivists strike a balance between the nativist and associationist perspectives. They propose that individuals genetically inherit core cognitive abilities or structures. These abilities and structures are necessary for cognitive development. Using these structures and abilities, individuals both make meaning of their environment and are shaped by it. People also possess the ability to make novel cognitive constructions. The most famous of the constructivist cognitive development theorists is Jean Piaget.

Many CBT theories and therapies fit most easily into this model. Contemporary theorists view the individual as an active agent of meaning making and change. In support of this model, consider that the way one individual makes sense of events often differs from the manner in which others make sense of those same events. Different events have different meanings for different people. These dissimilarities in processing events arise from a number of sources, including biology, learning, and individual differences. This perspective, therefore, also makes room for environmental shaping. CBT practiced from this perspective takes individual differences in cognition into account and uses cognitive and environmental realities unique to the individual to formulate alternative perspectives that are less harmful to mood. Traditional cognitive therapy takes this approach by seeking to change depressed mood by correcting faulty information processing and distorted thinking (Beck, 1967; Burns, 1980). This often involves assimilation and accommodation of information into existing, individual cognitive associative networks (i.e., schemas). Schema therapy, for example, aims to change maladaptive schemas that have developed because of environmental shaping and personal construction (Young, Klosko, & Weishaar, 2003). At the same time, a constructivist might additionally work to modify maladaptive behaviors or build adaptive ones (e.g., coping skills).

Sociocognitive

This perspective of cognitive development emphasizes the role of society in shaping individual cognition. Vygotsky and other social constructivists argue that, unlike Piaget and pure constructivists, individuals do not simply act upon their world. Rather, larger social networks inculcate that individual with that society's worldview and knowledge. While the child acts upon its environment, making meaning of it, the knowledge learned by that child is reciprocally shaped by that child's social context. This occurs in a way that is congruent with that society's worldview. This view is similar to associationism and behaviorism in its emphasis on environmental influence on behavior, but different in that it speculates about the shaping influence of a particular society's worldview.

In its simplest form, social constructivist psychotherapy would equate to clinics and hospitals shaping behavior, playing on the strength of institutions to shape their members. It could also take the form of a directive therapist who makes suggestions and offers direction to clients. As a social representative of moral authority, a priest, reverend, or spiritual leader could offer guidance to a client seeking help. Considered by many as authorities on the values of a given society, they would be well suited to offer ethical advice and guidance and promote that society's values. *Walden Two* (Skinner, 1976) illustrates the potential of applying theories of operant conditioning in shaping the members of society to create a modern utopia.

Tips for Selecting an Orientation

In addition to considering the theory and research of other fields of psychology and disciplines, here are some general tips for selecting a theoretical orientation.

Supplement Your Reading About Therapies and Theoretical Systems With Supervision From Mentors and Seasoned Researchers and Clinicians

From their experience and knowledge, they can offer a hindsight view of the successes and pitfalls that accompany selecting and utilizing the approaches of theoretical orientations. Similarly, seek opportunities to observe the founders of an approach in action. Read classic case studies. Watch recordings of the founders practicing their therapies. When possible, attend conferences like ABCT for a more interactive experience. At ABCT conferences, one can

observe leading CBT figures in action, ask them questions, and seek their advice.

Integrate theory and research from other fields of psychology and professional disciplines (e.g., medicine) into your search. It will improve the breadth and scope of your understanding of cognition and behavior, as well as improve your ability to predict and change them. While this article presents the schools of cognitive development and their application to CBT, many ideas from other branches of psychology are useful as well. Other disciplines and schools can broaden your understanding of the world. Remain open to the body of knowledge, experience, and wisdom that other disciplines and perspectives can offer.

Consider Your Personal Strengths, Weaknesses, Predilections, and Values When Selecting an Orientation

Does a certain orientation more naturally fit your strengths and way of thinking? Go with what feels right. Much like selecting a career, it will be easier to adopt an approach or perspective that fits with your existing values, proclivities, and strengths. It will facilitate research and understanding, as well as increase clinical effectiveness. Similarly, consider also the theoretical inclination, experiences, and abilities of the faculty and mentors in your program. Benefit from their knowledge and experience. They can offer more of a bird's-eye view of the development and implementation of theories and practices than is possible with a student's limited experience and knowledge.

Consider the Findings of Scientific Research

Empirical inquiry provides the most objective forum in which to test ideas. While proclivity and values are useful guides and can serve as strengths, they can also serve as biases that cloud one's view. While an approach may feel natural or easy, it may not necessarily be effective in addressing clinical problems. Knowing that scientific studies support one's research program or clinical work offers additional confidence in the utility of one's professional activities.

Outside of the empirical literature, it is also useful to think as a scientist in one's daily professional activity. When you have selected an orientation, use practical methods and personal reflection to test its worth. Can you use the theories and research of your orientation to understand a wide range of behavior? Can you use the theory and research of your orientation to understand behavior? If not, why? This will also sharpen

your ability to conceptualize cases. Be open to change. If new scientific findings conflict with theoretical aspects of your orientation, it may be an indication that the perspective is limited and can improve. Perhaps you could even research the anomaly to enhance scientific knowledge in the area.

Selecting and exploring an orientation can be an exciting experience. One can explore a large body of philosophy, ideas, and techniques, within both the entire discipline of psychology and others. Approaching the selection process in this manner can keep the search interesting, as well as provide the benefits of both selecting a theoretical orientation and considering diverse perspectives and wisdom. Have fun with the process and best wishes!

References

- Beck, A. T. (1967). *Depression: Clinical, experimental, and theoretical aspects*. New York: Hoeber.
- Burns, D. M. (1980). *Feeling good: The new mood therapy*. New York: Avon.
- Chomsky, N. (1980). *Rules and representations*. Oxford, UK: Blackwell.
- Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (1999). *Acceptance and commitment therapy: An experiential approach to behavior change*. New York: The Guilford Press.
- Pratt, C. W., Gill, K. J., Barrett, N. M., & Roberts, M. M. (2002). *Psychiatric rehabilitation*. San Diego: Academic Press.
- Skinner, B. F. (1953). *Science and human behavior*. New York: Macmillan.
- Skinner, B. F. (1976). *Walden two*. Oxford, UK: Macmillan.
- Richardson, K. (1998). *Models of cognitive development*. East Sussex, UK: Psychology Press.
- Wilson, G. T. (1995). Behavior therapy. In R. J. Corsini & D. Wedding (Eds.), *Current psychotherapies* (5th ed.). Itasca, IL: F.E. Peacock.
- Young, J. E., Klosko, J. S., & Weishaar, M. E. (2003). *Schema therapy: A practitioner's guide*. New York: The Guilford Press.

Special thanks to Dr. Elisabeth Sandberg for her suggestions and guidance.

Address correspondence to Eric L Sullivan, M.A., Psychology Department, Suffolk University, Boston e-mail: ericsullivan1@gmail.com

Find-a-Therapist . . .

Who Takes Your Insurance, Has Saturday Appointments, Is Located Nearby, and More . . .

Katherine Martinez, *Committee on Clinical Directory and Referral Issues*

One of the first questions a potential client may ask is, "Do you take my insurance?" ABCT's expanded Find-a-Therapist Practice Particulars can resolve this and many other questions asked by clients and referral sources. Giving your valuable time to talk on the phone with a potential client can become burdensome when you realize midway through the call that the client has insurance that you do not take, or requests an evening appointment on days you do not work. To prevent this problem, try using the Practice Particulars option in the Find-a-Therapist directory to specify such important details as:

- insurance taken
- languages spoken
- practice philosophy
- populations served
- areas of specialization

Listing these and other details on your Find-a-Therapist record increases the probability of appropriate referrals from either the ABCT membership or the public. Including the Practice Particulars option in

your profile also benefits ABCT members, who can depend upon the Find-a-Therapist directory as a more efficient and comprehensive referral service.

To be listed in the Find-a-Therapist directory and/or to add Practice Particulars to your listing in the referral directory:

- Go to www.abct.org and select MEMBER LOG-IN.
- Log in.
- Select FIND-A-THERAPIST DIRECTORY AND REFERRAL SERVICE, "join now" (or see list of URLs below, specifically: "Update Find-a-Therapist Listing").

Once your request is processed, you can log on to the member's home page at any time to make edits and ensure your information remains current.

New Year's Resolution: *Update your Practice Particulars!* ■

URLs U Will Use

Dues History (including 2008 payments)

<https://abct.org/members/source/DuesHistory.cfm>

Membership Directory—edit contact information

https://abct.org/members/MemberInfo/Update_Profile.cfm

Membership Directory Searches

https://abct.org/members/Directory/Membership_Directory.cfm

Update Find-a-Therapist Listing

<https://abct.org/members/FindATherapistDirectory/>

Update Passwords

https://abct.org/members/MemberInfo/Update_Password.cfm



These links require that your email address be in ABCT's system. If you've never given us an email address or have changed your email you will not be able to retrieve this information.

Letter to the Editor

In "Convention, Tradition, and the New Wave: Assessing Clinician Identity in Behavior Therapy," Storaasli, Kraushaar, Wilson, and Emrick (2007) compared endorsements of various interventions by therapists who utilize Acceptance and Commitment Therapy (ACT), by therapists who utilize CBT, and by psychology students. The authors found that although there was some overlap, there was a clear difference between ACT therapists and CBT therapists.

One theoretical difference between ACT and CBT therapists is that CBT therapists challenge dysfunctional cognitions, while ACT therapists accept dysfunctional cognitions but encourage patients to pursue their valued goals in spite of conflicting dysfunctional cognitions.

But asking patients to behave in ways that conflict with a cognition indirectly suggests that the cognition is dysfunctional. So I would argue that ACT challenges dysfunctional cognitions indirectly and gently, while CBT challenges dysfunctional cognitions directly and forcefully.

As a CBT therapist, I once gave a workshop with a psychodynamic therapist in which we explained how each of us would treat specific patients. At one point I said that I would tell a patient, "Go home and tell your husband ———." My psychodynamic colleague responded that she would ask the patient, "I wonder why you don't go home and tell your husband ———?" So maybe the most important difference between therapists of various orientations is in how directly we challenge dysfunctional cognitions.

Reference

Storaasli, R. D., Kraushaar, B., Wilson, K. G., & Emrick, C. (2007). Convention, tradition, and the new wave: Assessing clinician identity in behavior therapy. *the Behavior Therapist, 30*, 149-155.

— MILTON SPETT
Private Practice, Cranford, NJ
MCSpett@aol.com

ERRATUM

The January issue of *tBT* (Vol. 31, p. 23) incorrectly identified the winner of the 2007 President's New Researcher Award as "Japster A. J. Smits." The correct spelling is Jasper A. J. Smits. We apologize for this error.

ABCTAMBASSADORS

are needed to:

messenger

Articulate ABCT's vision, purpose, and identity to encourage membership

guide

Mentor individuals through the process of presenting at the annual convention or transitioning into leadership positions

signify

Act as the eyes and ears of the association locally

ABCT's Ambassador program is a brand-new initiative promoting leadership, participation, and membership in ABCT.

ABCT Ambassadors are easily recognized at the annual meeting by their special ribbons. They also receive a certificate of recognition and are featured on our website and in tBT.

For more information, contact Lisa Yarde at ABCT's central office (lyarde@abct.org)



Call for Award Nominations

**Nominate
on-line!**

<http://www.abct.org/awards/>

- Career/Lifetime Achievement
- Distinguished Friend to Behavior Therapy
- Outstanding Service to ABCT
- Outstanding Clinical Contributions
- Outstanding Mentor
- Virginia A. Roswell Student Dissertation

NOMINATIONS DEADLINE: March 3, 2008 | Questions? email M. Joann Wright at ABCTAwards@gmail.com

2008

STUDENT AWARDS PROGRAM

President's New Researcher Award

ABCT's President, Anne Marie Albano, Ph.D., invites submissions for the 30th Annual President's New Researcher Award. The winner will receive a certificate and a cash prize of \$500. Submissions will be accepted on any topic relevant to behavior therapy, but submissions consistent with the conference theme emphasizing basic research are particularly encouraged. Eligible papers must (a) be authored by an individual with five years or less posttraining experience (e.g., post-Ph.D. or postresidency); and (b) have been published in the last two years or currently be in press. Submissions can consist of one's own or any eligible candidate's paper. Papers will be judged by a review committee consisting of Anne Marie Albano, Ph.D.; Raymond DiGiuseppe, Ph.D., ABCT's Immediate Past-President; and Robert Leahy, the ABCT President-Elect. Submissions must be received by August 13, 2008, and must include four copies of both the paper and the author's vita. Send submissions to ABCT President's New Researcher Award, 305 Seventh Ave., 16th floor, New York, NY 10001.

Virginia A. Roswell Student Dissertation Award

This award will be given to a student based on his or her doctoral dissertation proposal. The research should be relevant to behavior therapy. Accompanying this honor will be a \$1,000 award to be used in support of research (e.g., to pay participants, to purchase testing equipment) and/or to facilitate travel to the ABCT convention. Eligible candidates for this award should be student members who have already had their dissertation proposal approved and are investigating an area of direct relevance to behavior therapy, broadly defined. A student's dissertation mentor should complete the nomination. Please complete an on-line nomination by visiting www.abct.org, and completing the appropriate application forms. Then, e-mail the completed forms to ABCTAwards@gmail.com. Also, mail a hard copy of your submission to ABCT, Virginia A. Roswell Dissertation Award, 305 Seventh Ave., New York, NY 10001.

Elsie Ramos Memorial Student Poster Awards

These awards will be given to three student poster presenters (student first authors only), member or nonmember, at ABCT's 42nd Annual Convention in Orlando. The winners will each receive a 2009 ABCT Student Membership, a 1-year subscription to an ABCT journal of their choice, and a complimentary general registration at ABCT's 2009 Annual Convention. To be eligible, students must complete the submission for this year's ABCT convention by **March 3, 2008**. The proposal must then pass ABCT's peer review process. ABCT's Awards and Recognition Committee will judge all student posters.

**Awards
will be
presented
at ABCT's
convention
November
13-16,
Orlando**

CALL *for* PAPERS

42nd
Annual
Meeting

Nov. 13–16, 2008, Orlando

Sandra Pimentel, Ph.D., Program Chair



There is widespread and growing interest in the development and implementation of evidence-based psychotherapies. As evidence has accumulated supporting the efficacy of cognitive and behavioral therapies, policymakers and practitioners seek to disseminate and deliver CBT to an ever-expanding array of clinical populations.

The theme of the 42nd Annual ABCT Convention recognizes the pivotal role of CBT in the delivery of mental health care. The convention will emphasize the role of researchers and practitioners in developing and continuously enhancing theoretical knowledge of psychopathology across the lifespan, developing efficacious forms of CBT, and advancing these treatments into clinical practice.

The meeting will focus on the dissemination of CBT to the range of populations, problems, and systems. We welcome submissions for research symposia, clinical sessions, and workshops focused on the application of CBT across stages of development, diagnostic areas, and organizational systems of care. Submissions that highlight models of dissemination, methods for evaluating and maximizing CBT training and skill transfer, and collaborative arrangements between research and service settings are especially encouraged and will receive special consideration.

**SUBMISSION
DEADLINE:
March 15**

●
Submissions may be in the form of symposia, round tables, panel discussions, and posters. Discussants will be encouraged to integrate efficacy, effectiveness, and dissemination research.

Submission information is located on ABCT's web site, www.abct.org.

————— *Taking it to the Streets* —————


Advancing Dissemination of CBT

About Programs That Work

The successful *Treatments That Work™* series has now been expanded to include treatments and preventative interventions for use with child and adolescent clients. Editors-in-Chief: Anne Marie Albano and David H. Barlow.

- Programs are evaluated by a scientific advisory board to ensure they meet the highest standards for evidence-based practice.
- Interventions and treatments that span the disorder spectrum for youths of all ages, from young children to adolescents and young adults
- Guides include everything you need to successfully implement the program – including session agendas, sample dialogues, and homework assignments

Workbooks are specially designed with age-appropriate and engaging activities, easy-to-read psychoeducational material, and interactive assignments




Helping Parents With Challenging Children Therapist Guide

978-0-19-533298-9 paper \$35.00
978-0-19-533299-7 paper \$24.95




When Children Don't Sleep Well Therapist Guide

978-0-19-532947-6 paper \$35.00
978-0-19-532948-3 paper \$24.95



Coping Power Parent Group Program Facilitator's Guide

978-0-19-532788-5 paper \$45.00
978-0-19-532796-0 paper \$95.00



Coping Power Child Group Program Facilitator's Guide

978-0-19-532787-8 paper \$55.00
978-0-19-537081-2 paper \$64.00



When Children Refuse School Therapist Guide

978-0-19-530830-3 paper \$39.95
978-0-19-530829-7 paper \$29.95



Getting Your Child to say Yes to School

978-0-19-530630-9 paper \$19.95

NEW
from the Author

Helping School Refusing Children and Their Parents A Guide for School-based Professionals

978-0-19-532024-4 paper \$24.95

Also of Interest



Improving Children's Mental Health Through Parent Empowerment

Edited by Peter S. Jensen and Kimberly Hoagwood

"Family support, engagement, and empowerment are critical and enduring concepts that are somehow only now being operationalized and supported. It is a watershed event that the field's most esteemed academics are now packaging and presenting this material."—Eric J. Bruns, Ph.D., Assistant Professor, University of Washington, and coordinator of the National Wraparound Initiative

2008 224 pp. \$27.50
978-0-19-532090-9 paper



Effective Practices for Children with Autism

Educational and Behavior Support Interventions that Work

Edited by James K. Luiselli, Dennis C. Russo, Walter P. Christian, and Susan M. Wilczynski

With contributors from a variety of disciplines, *Effective Practices for Children with Autism* brings together multiple perspectives on intervention effectiveness for autism education and behavior support. Written for practitioners, research scientists, and clinicians, the book is an essential framework for evaluating educational and treatment procedures, selecting those that are most effective, and evaluating outcome.

2008 528 pp. \$75.00
978-0-19-531704-6

OXFORD
UNIVERSITY PRESS

Prices are subject to change and apply only in the US. To order, please contact customer service at call 1-866-445-8685, or visit our website at www.oup.com/us

the Behavior Therapist

Association for Behavioral and Cognitive Therapies
305 Seventh Avenue, 16th floor
New York, NY 10001-6008

Tel.: 212-647-1890
www.abct.org

ADDRESS SERVICE REQUESTED

PRSR STD
U.S. POSTAGE
PAID
Hanover, PA
Permit No. 4