

VU Research Portal

A simplicial A-infinity operad acting on R-completions

Bauer, T.; Libman, A.

published in

Homology, Homotopy and Applications
2009

DOI (link to publisher)

[10.4310/HHA.2009.v11.n2.a5](https://doi.org/10.4310/HHA.2009.v11.n2.a5)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Bauer, T., & Libman, A. (2009). A simplicial A-infinity operad acting on R-completions. *Homology, Homotopy and Applications*, 11(2), 55-73. <https://doi.org/10.4310/HHA.2009.v11.n2.a5>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

A SIMPLICIAL A_∞ -OPERAD ACTING ON R -RESOLUTIONS

TILMAN BAUER AND ASSAF LIBMAN

ABSTRACT. We construct a combinatorial model of an A_∞ -operad which acts simplicially on the cobar resolution (not just its total space) of a simplicial set with respect to a ring R .

1. INTRODUCTION

Let K be a commutative ring and X a simplicial set (henceforth, simply called “space”). Bousfield and Kan [BK72] defined the K -resolution of X to be (a variant of) the cosimplicial space

$$R_K(X)^n = \underbrace{K(K(\dots(K(X)\dots))}_{n+1 \text{ times}},$$

where $K(X)$ denotes the degreewise free K -vector space on X . The K -completion of X is derived from the cosimplicial space R_K and can be defined in at least three different ways, taking values in different categories that each come with a specific notion of homotopy.

Completion functor	taking values in	with homotopy defined by
$X_{\hat{K}} = \text{Tot } R_K(X)$	spaces	(simplicial) homotopy
$X_K^{\text{pro}} = \{\text{Tot}^s R_K(X)\}_{s \geq 0}$	towers of spaces	pro-homotopy
R_K	cosimplicial spaces	external homotopy

For more precise information on pro-homotopical structures, see [Isa04, FI07]. There are natural transformations $R_K \rightarrow (-)_K^{\text{pro}} \rightarrow (-)_{\hat{K}}$ under which homotopies map to homotopies, but the three notions of homotopy are strictly contained in one another. To illustrate this, consider the Bousfield (unstable Adams) spectral sequence associated to R_K . If the cosimplicial space $R_K(X)$ is externally homotopy equivalent [Qui67, Bou03] to $R_K(Y)$, the spectral sequences agree from E^2 on. If the towers are pro-weakly equivalent, then for every (p, q) there is an R such that the $E_{p,q}^r$ -terms are isomorphic for $r \geq R$. If we merely know that $X_{\hat{K}} \simeq Y_{\hat{K}}$, even if this is induced by a map $X \rightarrow Y$, the spectral sequences might not be isomorphic anywhere at any term.

The functors $(-)_K^{\text{pro}}$ and R_K can be made into endofunctors by enlarging the source category to towers or cosimplicial spaces, respectively, applying the completion functor levelwise, and taking the diagonal.

In [BK72, I.5.6], Bousfield and Kan claimed that $(-)_K^{\text{pro}}$ is a monad (levelwise, not just pro-). In fact, they try to prove that R_K is a monad. Both claims were not proved and later retracted by the authors. In the companion paper [BL07]

Date: December 8, 2008.

2000 Mathematics Subject Classification. 55P60, 18D50, 18C15 .

Key words and phrases. monad, A_∞ -monad, completion, cobar resolution.

we showed that $(-)\hat{K}$, while not a strict monad, can be made into a monad up to coherent homotopy, an A_∞ -monad. This works in great generality: whenever given a monad (A_∞ or strict), there is a completion functor which is again an A_∞ -monad.

In this paper, we address the problem of realizing this structure on the resolution. While related in its results, the methods we use here are completely different from [BL07], and this paper can thus be read independently.

Our result is slightly more general than outlined above. Let $c\mathcal{E}$ denote the category of cosimplicial objects in a complete and cocomplete monoidal category (\mathcal{E}, \circ, I) . The category $c\mathcal{E}$ is endowed with an “external” simplicial structure (see Section 3) and inherits a levelwise monoidal structure \diamond from \mathcal{E} . Every object $X \in c\mathcal{E}$ gives rise to an operad $\mathcal{O}\text{End}(X)$ (of simplicial sets) whose n th space is $\text{map}_{c\mathcal{E}}(X^{\circ n}, X)$. As usual [May72], an \mathcal{O} -algebra in $(c\mathcal{E}, \circ, I)$ is an object $X \in c\mathcal{E}$ with a morphism of operads $\mathcal{O} \rightarrow \mathcal{O}\text{End}(X)$. For example, if $\mathcal{O}(n) = *$ is the associative operad, X is simply a monoid. When the spaces $\mathcal{O}(n)$ are weakly contractible we call X an A_∞ -monoid.

Now let (K, η, μ) be a monoid in (\mathcal{E}, \circ, I) defined by $\eta: I \rightarrow K$ and $\mu: K \diamond K \rightarrow K$. Let $R_K \in c\mathcal{E}$ be its cobar construction with $R_K^n = K^{\circ n+1}$ and with coface maps $d^i = K^{\circ i} \eta K^{\circ n-i}$ and codegeneracy maps $s^i = K^{\circ i} \mu K^{\circ n-i}$, cf. [BK72, Ch I, §4].

1.1. Definition. A *twist map* for K is a morphism $t: K \diamond K \rightarrow K \diamond K$ such that

- (1) $\mu \circ t = \mu$,
- (2) $t \circ (K \diamond \eta) = \eta \diamond K$ and $t \circ (\eta \diamond K) = K \diamond \eta$,
- (3) $t \circ (\mu \diamond K) = (K \diamond \mu) \circ (t \diamond K) \circ (K \diamond t)$ and $t \circ (K \diamond \mu) = (\mu \diamond K) \circ (K \diamond t) \circ (t \diamond K)$.

1.2. Theorem. *There exists an A_∞ -operad of simplicial sets TW such that for every monoid (K, η, μ) with a twist map t in a complete and cocomplete monoidal category \mathcal{E} , R_K is a TW -algebra in $c\mathcal{E}$, i.e. there is a morphism of operads $\text{TW} \rightarrow \mathcal{O}\text{End}(R_K)$.*

As an application, let \mathcal{E} be the category of endofunctors of simplicial sets with composition as its monoidal structure; one needs to pass to a larger universe in order for \mathcal{E} to make sense. A monad K on the category of spaces is a monoid in \mathcal{E} .

Let K be the free K -module functor on spaces for a ring K , and t the “twist map” defined by

$$t = (K \diamond \mu) - \text{id}_{K \diamond K} + (\mu \diamond K) \quad (\text{cf. [BK72, p. 28]})$$

We obtain as a corollary the motivating example:

1.3. Corollary. *For any ring K , R_K is a TW -algebra and in particular an A_∞ -monad.*

We do not know any interesting example of a monad with a twist map that is not of the free- K -module type. It would be interesting to know if the monad $X \mapsto \Omega^\infty(E \wedge X)$, for an \mathcal{S} -algebra E in the sense of [EKMM97] or [HSS00] can be given a twist map – this would imply that the Bendersky-Thompson resolution [BT00] is an A_∞ -monad. Despite the lack of other examples, we decided to work in this generality as, if anything, it makes the proofs clearer.

2. THE “TWIST MAP OPERAD” TW

In this section we will give an explicit combinatorial construction of the operad TW of spaces. In the next section we will construct its action on the Bousfield-Kan resolution [BK72].

2.1. Informal description. The k -simplices of the n th space of the operad TW are braid-like diagrams as illustrated in Figure 1 below. These braid diagrams have $n(k + 1)$ incoming strands labelled $(0 \dots k 0 \dots k \dots 0 \dots k)$ and $k + 1$ outgoing strands labelled $(0 \dots k)$ in this order from left to right. Two strands can join if they have the same label, or they can cross provided the label of the strand on the left is bigger than the one of the strand on the right. In addition we allow new strands of any label to emerge.

$$(2.1) \quad \text{Cross } (a > b): \begin{array}{c} a \quad b \\ \diagdown \quad / \\ b \quad a \end{array} \quad \text{Join: } \begin{array}{c} a \quad a \\ \diagdown \quad / \\ a \end{array} \quad \text{Emerge: } \begin{array}{c} \circ \\ | \\ a \end{array}$$

Note that a crossing has no orientation, that is, none of the strands is considered passing above or below the other. Figure 1 shows an example of such a braid. Two such diagrams are considered equal if they are isotopic in \mathbb{R}^2 relative to the




FIGURE 1. An example of an element of $\text{TW}(4)_2$

end points; in particular, we do not allow Reidemeister moves. However, we do identify the following diagrams involving the new strand operation:

$$(2.2) \quad \begin{array}{c} \circ \\ \diagup \\ a \end{array} \begin{array}{c} a \\ \diagdown \\ \circ \end{array} \sim \begin{array}{c} a \\ | \\ a \end{array} \sim \begin{array}{c} a \\ \diagdown \\ \circ \end{array} \begin{array}{c} \circ \\ \diagup \\ a \end{array}, \quad \begin{array}{c} b \quad b \\ \diagdown \quad / \\ a \quad a \end{array} \sim \begin{array}{c} b \\ | \\ b \end{array} \begin{array}{c} a \\ | \\ a \end{array}, \quad \begin{array}{c} a \quad a \\ \diagdown \quad / \\ b \quad b \end{array} \sim \begin{array}{c} \circ \\ | \\ b \end{array} \begin{array}{c} a \\ | \\ a \end{array}$$

The sequence of $\{\text{TW}(n)_k\}_{k \geq 0}$ forms a simplicial set, where the i th face map d_i is obtained by erasing the strands of color i , whereas the i th degeneracy map s_i duplicates the i th strand. An illustration of how this works on the “basic building blocks” of the braids is depicted by (here $a > b$)

$$(2.3) \quad s_a \left(\begin{array}{c} a \quad a \\ \diagdown \quad / \\ a \end{array} \right) = \begin{array}{c} a \quad a+1 \quad a \quad a+1 \\ \diagdown \quad / \quad \diagdown \quad / \\ a \quad a+1 \end{array}, \quad s_b \left(\begin{array}{c} a \quad b \\ \diagdown \quad / \\ b \quad a \end{array} \right) = \begin{array}{c} a+1 \quad b \quad b+1 \\ \diagdown \quad / \quad \diagdown \quad / \\ b \quad b+1 \quad a+1 \end{array}$$

thus introducing new crossings.

Composition in the operad TW is obtained by grafting of these braid-like diagrams with matching labels.

We will now give a more rigid combinatorial description of the operad TW without direct reference to the braid-like diagrams. To do this we will construct a simplicial object $\{\mathcal{B}_k\}_k$ in the category of small monoidal categories. The k -simplices of the space $\text{TW}(n)$ are the morphisms between certain objects in \mathcal{B}_k . In fact, each \mathcal{B}_k is a free monoidal category in a sense which we will now make precise.

2.2. Free monoidal categories and the simplicial monoidal category $\{B_k\}_{k \geq 0}$. By a *graph* we mean a pair $(\mathcal{A} \rightrightarrows \mathcal{O})$ of two sets and two maps (codomain, domain) between them. A *unital graph* is a graph with a common section $\mathcal{O} \rightarrow \mathcal{A}$ of domain and codomain. Every small category \mathcal{C} has an underlying unital graph $U_u(\mathcal{C}) = (\text{Mor}\mathcal{C} \rightrightarrows \text{Obj}\mathcal{C})$ by forgetting the composition rule on morphisms. By further forgetting the unit maps, we obtain a graph $U(\mathcal{C})$. Conversely, every (unital) graph $\mathcal{G} = (\mathcal{A} \rightrightarrows \mathcal{O})$ has an associated category $C(\mathcal{G})$. It is the free category generated by \mathcal{G} in the sense that C is left adjoint to the functor U (resp. U_u) from the category of small categories to the category of (unital) graphs. See [ML98, Section II.7, in particular Theorem 1]. This section is about a version of this free construction for monoidal categories.

2.4. Definition. A *vertex-monoidal graph* is a graph $(\mathcal{A} \rightrightarrows \mathcal{O})$, where \mathcal{O} has the structure of a monoid. A *(unital) monoidal graph* is a graph object in the category of (unital) monoids.

2.5. Lemma. *The following forgetful functors all have left adjoints:*

$$\begin{array}{ccc} \{\text{small monoidal categories}\} & \xrightarrow{U_{um}} & \{\text{unital monoidal graphs}\} \\ & & \xrightarrow{U_m} \{\text{monoidal graphs}\} \xrightarrow{U_{vm}} \{\text{vertex-monoidal graphs}\} \end{array}$$

In particular, for every vertex-monoidal graph \mathcal{G} , there exists a small monoidal category $(\mathcal{M}, \sqcup, \mathcal{O})$ with a morphism of vertex-monoidal graphs $\epsilon: \mathcal{G} \rightarrow U(\mathcal{M})$, such that for any other monoidal category $(\mathcal{C}, \otimes, I)$ and vertex-monoidal graph morphism $X: \mathcal{G} \rightarrow U(\mathcal{C})$, there is a unique functor $X^\flat: \mathcal{M} \rightarrow \mathcal{C}$ of monoidal categories such that $\epsilon \circ U(X^\flat) = X$.

We give a proof of this (easy) fact in the appendix.

2.6. Remark. Fix a set Σ , and let Σ_\sqcup be the set of words, i.e. finite sequences in Σ , including the empty sequence $()$. This is a monoid under concatenation, which we denote by \sqcup . If $(\mathcal{A} \rightrightarrows \mathcal{O})$ is a vertex-monoidal graph whose vertices $\mathcal{O} = \Sigma_\sqcup$ form such a free monoid, then a morphism of vertex-monoidal graphs $X: (\mathcal{A} \rightrightarrows \Sigma_\sqcup) \rightarrow U(\mathcal{M})$, where \mathcal{M} is monoidal, is given by a map $X: \Sigma \rightarrow \text{Obj}(\mathcal{M})$ and, for every arrow $a: (\sigma_1, \dots, \sigma_k) \rightarrow (\tau_1, \dots, \tau_l)$ in \mathcal{A} , a morphism $X(\sigma_1) \otimes \dots \otimes X(\sigma_k) \rightarrow X(\tau_1) \otimes \dots \otimes X(\tau_l)$.

2.7. Lemma. *Let Σ and $\mathcal{G} = (\mathcal{A} \rightrightarrows \Sigma_\sqcup)$ be as in Remark 2.6 and let \mathcal{M} denote the free monoidal category generated by \mathcal{G} by Lemma 2.5. Let \sim be an equivalence relation on the morphism set of \mathcal{M} such that if $f \sim f'$ then their domains and codomains are equal.*

Then there exists a monoidal category \mathcal{M}/\sim together with a quotient functor $\mathcal{M} \rightarrow \mathcal{M}/\sim$ which is the identity on the object sets and such that \mathcal{M}/\sim is the free monoidal category on (Σ, \sqcup) subject to the relation \sim . That is, given any monoidal category $(\mathcal{C}, \otimes, I)$ and a morphism of monoidal graphs $X^\#: \mathcal{G} \rightarrow U(\mathcal{C})$ whose associated monoidal functor $X: \mathcal{M} \rightarrow \mathcal{C}$ has the property that $X(f) = X(g)$ if $f \sim g$, there exists a unique monoidal functor $X: \mathcal{M}/\sim \rightarrow \mathcal{C}$ which factors X through the quotient $\mathcal{M} \rightarrow \mathcal{M}/\sim$.

Proof. Consider the equivalence relation \approx on the morphism set of \mathcal{M} which is generated by \sim and which is closed to the property that $(f \sqcup g) \approx (f' \sqcup g')$ and $f \circ g \approx f' \circ g'$ if $f \approx f'$ and $g \approx g'$ (for the composition, whenever it makes sense). One easily checks that the domain and codomain of \approx -equivalent morphisms are equal. We can now apply [ML98, §II.8, Proposition 1] to obtain \mathcal{M}/\approx which is monoidal due to the requirements on \approx . \square

The definition of the category \mathcal{B}_k is modelled on the ideas in 2.1.

2.8. Definition. We use Lemma 2.7 to define \mathcal{B}_k as the free monoidal category generated by the set of symbols $[k] = \{0, \dots, k\}$ and morphisms (cf. (2.1))


$$\begin{aligned} () \xrightarrow{u^a} (a) \quad \text{and} \quad (a, a) \xrightarrow{m^a} (a) \quad \text{for all } a = 0, \dots, k \\ (a, b) \xrightarrow{t^{a,b}} (b, a) \quad \text{for all } 0 \leq b < a \leq k \end{aligned}$$

subject to the relations (cf. 2.2)

$$\begin{aligned} (R1) \quad m^a \circ (\text{id}_{(a)} \sqcup u^a) &= \text{id}_{(a)} = m^a \circ (u^a \sqcup \text{id}_{(a)}) \\ (R2)_i \quad t^{c,d} \circ (\text{id}_{(c)} \sqcup u^d) &= u^d \sqcup \text{id}_{(c)} \\ (R2)_{ii} \quad t^{c,d} \circ (u^c \sqcup \text{id}_{(d)}) &= \text{id}_{(d)} \sqcup u^c \end{aligned}$$

where $0 \leq a, c, d \leq k$ and $d < c$.

For an object $\sigma = (a_1, \dots, a_n)$ and a morphism f in \mathcal{B}_k we will write $\sigma \sqcup f$ for $\text{id}_\sigma \sqcup f$. The morphism in \mathcal{B}_k between sequences $(a_1, \dots, a_s) \rightarrow (b_1, \dots, b_t)$ can be depicted by braid-like diagrams with s incoming strands labelled by the integers a_1, \dots, a_s and r outgoing strands labelled by (b_1, \dots, b_t) . For example, the diagram



describes a morphism $(1232) \rightarrow (123)$ in \mathcal{B}_3 which is the composition

$$[m^1 \sqcup (23)] \circ [(1) \sqcup t^{21} \sqcup (3)] \circ [(1) \sqcup m^2 \sqcup t^{31}] \circ [(12) \sqcup t^{32} \sqcup u^1].$$

Most of this section is devoted to the proof of the following result.

2.9. Proposition. *There are face and degeneracy operators $d_i: \mathcal{B}_k \rightarrow \mathcal{B}_{k-1}$ and $s_i: \mathcal{B}_k \rightarrow \mathcal{B}_{k+1}$ which make $\{\mathcal{B}_k\}_{k \geq 0}$ a simplicial object of small monoidal categories. On objects, d_i and s_i act by deleting (resp. duplicating) the symbol $i \in [k]$.*

A more elaborate description of the simplicial operators is given in Definition 2.14. In what follows we will write $[k]$ for the object $(01 \dots k)$ in \mathcal{B}_k use $[k]^n$ to denote the n -fold concatenation of $[k]$ with itself, i.e. the object $(0 \dots k 0 \dots k \dots)$.

2.10. Definition. Define $\text{TW}(n)_k = \text{Hom}_{\mathcal{B}_k}([k]^n, [k])$. By Proposition 2.9, this collection of sets, for all k , forms a simplicial set.

The composition law in the operad $\text{TW}(n) \times \prod_{i=1}^n \text{TW}(m_i) \rightarrow \text{TW}(\sum_{i=1}^n m_i)$ is defined using the monoidal structure of \mathcal{B}_k :

$$\begin{aligned} \mathcal{B}_k([k]^n, [k]) \times \prod_{i=1}^n \mathcal{B}_k([k]^{m_i}, [k]) &\xrightarrow{\text{id} \times \sqcup} \mathcal{B}_k([k]^n, [k]) \times \mathcal{B}_k([k]^{\sum_i m_i}, [k]^n) \\ &\xrightarrow{\circ} \mathcal{B}_k([k]^{\sum_i m_i}, [k]). \end{aligned}$$

Unitality and associativity of the monoidal structure in \mathcal{B}_k descend to this composition law.

The goal of this section is to prove

2.11. Theorem. *The sequence $\text{TW}(n)$ is a (nonsymmetric) A_∞ -operad of simplicial sets.*

Here are some easy results on the categories \mathcal{B}_k (Definition 2.8.) An object $\sigma \in \mathcal{B}_k$ is a sequence of symbols (s_1, \dots, s_n) and we write u^σ for $u^{s_1} \sqcup \dots \sqcup u^{s_n}$.

2.12. Lemma. Fix an object $\tau = (t_1 < \dots < t_n)$ in \mathcal{B}_k . Then given an object $\sigma \in \mathcal{B}_k$, any morphism $\varphi: \tau \rightarrow \sigma$ has the form

$$(2.13) \quad \varphi = u^{\sigma_0} \sqcup (t_1) \sqcup u^{\sigma_1} \sqcup \dots \sqcup u^{\sigma_{n-1}} \sqcup (t_n) \sqcup u^{\sigma_n}$$

where σ_i are subsequences of σ such that $\sigma = \sigma_0 \sqcup (t_1) \sqcup \sigma_1 \sqcup \dots \sqcup \sigma_{n-1} \sqcup (t_n) \sqcup \sigma_n$. In particular, the object $()$ is initial in \mathcal{B}_k and $\text{End}_{\mathcal{B}_k}(\tau) = \{\text{id}\}$.

Proof. By Lemma 2.7, the morphisms in \mathcal{B}_k are compositions of morphisms of the form $x \sqcup u^a \sqcup y$, $x \sqcup m^a \sqcup y$, and $x \sqcup t^{ab} \sqcup y$, where x, y are objects of \mathcal{B}_k .

Let φ be as in the statement of the lemma. We prove the result by induction on the “generation length” s of φ . If $s = 0$ then $\varphi = \text{id}$. If φ satisfies (2.13) and is followed by some $x \sqcup u^a \sqcup y$, then the resulting morphism still has the form (2.13). If φ is followed by some $x \sqcup t^{ab} \sqcup y$ then (a, b) cannot be contained in τ because $a > b$, hence one of a or b belongs to σ_i and we can apply the relation (R2) to show that the composition is again of the form (2.13). Finally, if φ is followed by $x \sqcup m^a \sqcup y$ then the domain (a, a) of m^a cannot be contained in τ , and relation (R1) guarantees that the composition is of the form (2.13). \square

2.14. Definition. We turn the categories \mathcal{B}_k into a simplicial monoidal category as follows. To do this it is convenient to replace $[k]$ with a totally ordered finite set S of cardinality $k + 1$. Then monotonic functions $S \rightarrow T$ are generated by the following functions

$$\begin{aligned} \eta_a: (S - \{a\}) &\xrightarrow{\text{inclusion}} S && \text{for some } a \in S \\ \epsilon_a: (S - \{a\}) \sqcup \{a', a''\} &\xrightarrow{\text{collapse } \{a', a''\} \rightarrow \{a\}} S \end{aligned}$$

where $a' < a''$ are inserted to S instead of $a \in S$. In this notation the simplicial identities become

$$(2.15) \quad \begin{aligned} \eta_a \circ \eta_b &= \eta_b \circ \eta_a, & \epsilon_b \circ \eta_a &= \eta_a \circ \epsilon_b, & \epsilon_a \circ \epsilon_b &= \epsilon_b \circ \epsilon_a & (a \neq b \in S) \\ \epsilon_a \circ \epsilon_{a'} &= \epsilon_a \circ \epsilon_{a''}, & \epsilon_a \circ \eta_{a'} &= \epsilon_a \circ \eta_{a''} & &= \text{Id} & (a \in S) \end{aligned}$$

We define face operators $d_a = (\eta_a)^*: \mathcal{B}_S \rightarrow \mathcal{B}_T$ where $T = S - \{a\}$ and degeneracy operators $s_a = (\epsilon_a)^*: \mathcal{B}_T \rightarrow \mathcal{B}_S$ where $T = S - \{a\} \sqcup \{a' < a''\}$ as follows. On objects we define

$$d_a((b)) = (b) = s_a((b)) \text{ if } b \neq a; \quad d_a((a)) = (); \quad s_a((a)) = (a', a'')$$

Extend s_a and d_a to all the objects of \mathcal{B}_k by monoidality with respect to \sqcup .

On morphisms we define d_a and s_a by defining them on generators:

$$\begin{aligned} d_a(m^a) &= \text{id}_{()}, & s_a(m^a) &= (m^{a'} \sqcup m^{a''}) \circ ((a') \sqcup t^{a'', a'} \sqcup (a'')) && \text{cf. (2.3)} \\ d_a(m^b) &= m^b, & s_a(m^b) &= m^b && \text{if } b \neq a \\ d_a(u^a) &= \text{id}_{()}, & s_a(u^a) &= u^{a'} \sqcup u^{a''} \\ d_a(u^b) &= u^b, & s_a(u^b) &= u^b && \text{if } b \neq a \\ d_a(t^{cd}) &= t^{cd}, & s_a(t^{cd}) &= t^{cd} && \text{if } c, d \neq a \\ d_a(t^{cd}) &= \text{id}_{(d)}, & s_a(t^{cd}) &= (t^{a'd} \sqcup (a'')) \circ ((a') \sqcup t^{a''d}) && \text{if } c = a \\ d_a(t^{cd}) &= \text{id}_{(c)}, & s_a(t^{cd}) &= ((a') \sqcup t^{ca''}) \circ (t^{ca'} \sqcup (a'')) && \text{if } d = a, \text{ cf. (2.3)} \end{aligned}$$

To see that these assignments define functors we need to check that they respect the relations (R1) and (R2). For (R1), we compute ($a \neq b$):

$$\begin{aligned}
s_b(m^a \circ ((a) \sqcup u^a)) &= m^a \circ ((a) \sqcup u^a) = \text{id}_{(a)} = s_b(\text{id}_{(a)}) \\
s_a(m^a \circ ((a) \sqcup u^a)) &= (s^{a'} \sqcup s^{a''}) \circ ((a') \sqcup t^{a''a'} \sqcup (a'')) \circ ((a'a'') \sqcup u^{a'} \sqcup u^{a''}) \\
&\stackrel{(R2)}{=} (s^{a'} \sqcup s^{a''}) \circ ((a') \sqcup u^{a'} \sqcup (a'') \sqcup u^{a''}) \\
&\stackrel{(R1)}{=} \text{id}_{(a')} \sqcup \text{id}_{(a'')} = \text{id}_{(a'a'')} = s_a(\text{id}_{(a)}).
\end{aligned}$$

Similarly one checks that d_a and d_b carry the relation (R1) to equal morphisms. The second relation of (R1) is checked similarly.

For (R2), assume that $c > d$ and consider some a . If $a \neq c, d$ then it is clear that applying d_a and s_a to the relations (R2) leaves them unaffected. If $a = c$ then applying s_a to the relation (R2)_i gives

$$\begin{aligned}
s_a(t^{ad} \circ ((a) \sqcup u^d)) &= (t^{a'd} \sqcup (a'')) \circ ((a') \sqcup t^{a'd}) \circ ((a'a'') \sqcup u^d) \stackrel{(R2)}{=} \\
& (t^{a'd} \sqcup (a'')) \circ ((a') \sqcup u^d \sqcup (a'')) \stackrel{(R2)}{=} u^d \sqcup \text{id}_{(a'a'')} = s_a(u^d \sqcup \text{id}_{(a)})
\end{aligned}$$

When $a = d$ a similar calculation shows that $s_a(t^{ca} \circ (\text{id}_{(c)} \sqcup u^a)) = s_a(u^a \sqcup \text{id}_{(c)})$. Checking that s_a respects the relation (R2)_{ii} is analogous, and we leave it to the reader to check that d_a respects (R2) as well. \square

Proof of Proposition 2.9. We will use the ‘‘coordinate free’’ notation of Definition 2.14 and prove that the functors d_a and s_a render $\{\mathcal{B}_k\}_{k \geq 0}$ a simplicial monoidal category. The functors d_a and s_a are monoidal, so it only remains to prove that they satisfy the simplicial identities dual to the identities (2.15), i.e.

$$(2.16) \quad d_b d_a = d_a d_b \quad (a \neq b \in S)$$

$$(2.17) \quad d_b s_a = s_a d_b \quad (a \neq b \in S)$$

$$(2.18) \quad s_b s_a = s_a s_b \quad (a \neq b \in S)$$

$$(2.19) \quad s_{a'} s_a = s_{a''} s_a \quad (a \in S)$$

$$(2.20) \quad d_{a'} s_a = d_{a''} s_a = \text{id} \quad (a \in S)$$

This is obviously the case on the object sets and it only remains to check the simplicial identities on the generating morphisms u^c, m^c, t^{cd} . If $c, d \in S - \{a, b\}$ then by definition d_a, d_b as well as s_a and s_b map u^c, m^c and t^{cd} to themselves, hence all the identities hold. Thus, it remains to check the simplicial identities when $\{c, d\}$

and $\{a, b\}$ are not disjoint. For (2.16) and (2.18), the only non-immediate cases are

$$\begin{aligned}
s_b s_a(m^a) &= s_b(m^{a'} \sqcup m^{a''}) \circ s_b((a') \sqcup t^{a'' a'} \sqcup (a'')) \\
&= (m^{a'} \sqcup m^{a''}) \circ ((a') \sqcup t^{a'' a'} \sqcup (a'')) &= s_a s_b(m^a) \\
s_b s_a(t^{ab}) &= s_b(t^{a' b} \sqcup (a'')) \circ s_b((a') \sqcup t^{a'' b}) \\
&= ((b') \sqcup t^{a' b''} \sqcup (a'')) \circ (t^{a' b'} \sqcup (b'') \sqcup (a'')) \\
&\quad \circ ((a') \sqcup (b') \sqcup t^{a'' b''}) \circ ((a') \sqcup t^{a'' b'} \sqcup (b'')) \\
&= ((b') \sqcup t^{a' b''} \sqcup (a'')) \circ (t^{a' b'} \sqcup t^{a'' b''}) \circ ((a') \sqcup t^{a'' b'} \sqcup (b'')) \\
&= ((b') \sqcup t^{a' b''} \sqcup (a'')) \circ ((b') \sqcup (a') \sqcup t^{a'' b''}) \\
&\quad \circ (t^{a' b'} \sqcup (a'') \sqcup (b'')) \circ ((a') \sqcup t^{a'' b'} \sqcup (b'')) \\
&= s_a((b') \sqcup t^{a b''}) \circ s_a(t^{a b'} \sqcup (b'')) &= s_a s_b(t^{ab})
\end{aligned}$$

The remaining cases, for example $d_b d_a(m^a) = d_a d_b(m^a)$ or when $\{a, b\} \neq \{c, d\}$, are straightforward.

For (2.17), we check

$$\begin{aligned}
d_b s_a(m^a) &= d_b(m^{a'} \sqcup m^{a''}) \circ d_b((a') \sqcup t^{a'' a'} \sqcup (a'')) \\
&= (m^{a'} \sqcup m^{a''}) \circ ((a') \sqcup t^{a'' a'} \sqcup (a'')) &= s_a d_b(m^a) \\
d_b s_a(t^{ac}) &= d_b(t^{a' c} \sqcup (a'')) \circ d_b((a') \sqcup t^{a'' c}) \\
&= (t^{a' c} \sqcup (a'')) \circ ((a') \sqcup t^{a'' c}) &= s_a d_b(t^{ac})
\end{aligned}$$

and leave the rest to the reader.

For (2.19), we use the convention that if $a \in S$ then the domain of $\epsilon_a: S' \rightarrow S$ is formed from S by replacing a with $a' < a''$. The morphisms $\epsilon_{a'}, \epsilon_{a''}: S'' \rightarrow S'$ have $S'' = S - \{a\} \sqcup \{a', a'', a'''\}$ and $s_{a'}(a') = (a', a'')$, $s_{a'}(a'') = a'''$ whereas $s_{a''}(a') = a'$, and $s_{a''}(a'') = (a'', a''')$. We now have the following calculation, which exploits the monoidal structure of \mathcal{B}_k :

$$\begin{aligned}
s_{a'} s_a(m^a) &= s_{a'}(m^{a'} \sqcup m^{a''}) \circ s_{a'}((a') \sqcup t^{a'' a'} \sqcup (a'')) \\
&= \{[(m^{a'} \sqcup m^{a''}) \circ ((a') \sqcup t^{a'' a'} \sqcup (a''))] \sqcup m^{a'''}\} \\
&\quad \circ \{(a' a'') \sqcup [(a') \sqcup t^{a'' a''}] \circ (t^{a'' a'} \sqcup (a''))\} \sqcup (a''') \\
&= (m^{a'} \sqcup m^{a''} \sqcup m^{a'''}) \circ ((a') \sqcup t^{a'' a'} \sqcup (a'' a'' a''')) \\
&\quad \circ ((a' a'' a') \sqcup t^{a'' a''} \sqcup (a''')) \circ ((a' a'') \sqcup t^{a'' a'} \sqcup (a'' a''')) \\
&= (m^{a'} \sqcup m^{a''} \sqcup m^{a'''}) \circ ((a' a' a'') \sqcup t^{a'' a''} \sqcup (a''')) \\
&\quad \circ ((a') \sqcup t^{a'' a'} \sqcup (a'' a'' a''')) \circ ((a' a'') \sqcup t^{a'' a'} \sqcup (a'' a''')) \\
&= \{m^{a'} \sqcup [(m^{a''} \sqcup m^{a'''}) \circ ((a'') \sqcup t^{a'' a''} \sqcup (a'''))]\} \\
&\quad \circ \{(a') \sqcup [(t^{a'' a'} \sqcup (a''')) \circ ((a'') \sqcup t^{a'' a''})] \sqcup (a'' a''')\} \\
&= s_{a''}(m^{a'} \sqcup m^{a''}) \circ s_{a''}((a') \sqcup t^{a'' a'} \sqcup (a'')) &= s_{a''} s_a(m^a)
\end{aligned}$$

This is the most subtle calculation in verifying (2.17). The remaining cases, e.g. $s_{a'} s_a(u^a) = s_{a''} s_a(u^a)$, are proved similarly and left to the reader, as is the proof of (2.20). \square

2.21. **Proposition.** *The spaces $\text{TW}(0)$ and $\text{TW}(1)$ are points.*

Proof. This follows immediately from Lemma 2.12. \square

2.22. **Remark.** The categories \mathcal{B}_k also enjoy the structure of a “cofacial monoidal category”, namely a cosimplicial object in the category of small monoidal categories without codegeneracy maps. In the notation of Definition 2.14 it is induced by the inclusion of symbols $T \subseteq S$ where T and S are totally ordered sets. We will not need this structure but we will make use of the operator

$$d^0: \mathcal{B}_k \rightarrow \mathcal{B}_{k+1}$$


which has the effect of shifting indices by 1. That is, $d^0(i) = i + 1$ for all $i \in [k]$ and on morphisms $d^0(u^a) = u^{a+1}$, $d^0(m^a) = m^{a+1}$ and $d^0(t^{ab}) = t^{a+1, b+1}$. It is obvious that the relations (R1) and (R2) are respected by these assignments. The “coface operator” d^0 also has the following relations with the face and degeneracy operators:

$$(2.23) \quad \begin{aligned} d_0 d^0 &= \text{id}, & s_0 d^0 &= d^0 d^0 \\ d_i d^0 &= d^0 d_{i-1}, & s_i d^0 &= d^0 s_{i-1} \quad \text{for all } i > 0. \end{aligned}$$

If σ is some sequence, we use the power notation σ^n for the n -fold concatenation of σ with itself. We observe that $d_a([k+1]^n) = [k]^n = s_a([k-1]^n)$ and $d^0[k] = (1, 2, \dots, k+1)$.

2.24. **Definition.** For every $n \geq 2$ let $b_n \in \mathcal{B}_1$ denote the object (01^n) . Define $\beta_n: [1]^n \rightarrow b_n$ by induction on $n \geq 2$ as follows:


$$\beta_2 = \{m^0 \sqcup (11)\} \circ \{(0) \sqcup t^{10} \sqcup (1)\}, \quad \beta_n = \{\beta_{n-1} \sqcup (1)\} \circ \{[1]^{n-2} \sqcup \beta_2\}$$



2.25. **Lemma.** *The following hold for all $n \geq 2$.*

- (1) $d_0(\beta_n) = \text{id}_{[0]^n} \in \mathcal{B}_1$.
- (2) $s_0(\beta_n) = \{(0) \sqcup d^0(\beta_n)\} \circ s_1 \beta_n$ (in \mathcal{B}_2).
- (3) $s_0 d_1(\beta_n) = \{(0) \sqcup d^0 d_1(\beta_n)\} \circ \beta_n$ (in \mathcal{B}_2).

The content of (2) is displayed as the obvious equivalence of the following “braids”.



The content of (3) is obtained by deleting all the strands labelled with 2.

Proof. (1) By inspection of Definitions 2.14 and 2.24,

$$d_0(\beta_2) = d_0(m^0 \sqcup (11)) \circ d_0((0) \sqcup t^{10} \sqcup (1)) = \text{id}.$$

The result follows easily using the inductive definition of β_n .

(2) Using Definition 2.24 and the pictures given there, we first observe that $\beta_n = \gamma_n \sqcup (1)$ for some $\gamma_n: [1]^{n-1} \rightarrow b_{n-1}$. It now follows, using the monoidal structure in \mathcal{B}_2 , that for every morphism $\varphi: [1]^2 \rightarrow b_2$ in \mathcal{B}_1 we have

$$(2.26) \quad \begin{aligned} & \{s_1 b_{n-1} \sqcup d^0 \varphi\} \circ \{s_1 \beta_n \sqcup (12)\} = \{s_1 b_{n-1} \sqcup d^0 \varphi\} \circ \{s_1 \gamma_n \sqcup (12)^2\} \\ & = \{s_1 \gamma_n \sqcup (122)\} \circ \{[2]^{n-1} \sqcup (0) \sqcup d^0 \varphi\} = \{s_1 \beta_n \sqcup (2)\} \circ \{[2]^{n-1} \sqcup (0) \sqcup d^0 \varphi\}. \end{aligned}$$

To complete the proof of (2) we use induction on n . The hardest part is the base of induction $n = 2$. Using Definition 2.24 and the monoidal structure in \mathcal{B}_2 we find

$$\begin{aligned} & \{(0) \sqcup d^0 \beta_2\} \circ s_1(\beta_2) = \\ & \underbrace{\{(0) \sqcup m^1 \sqcup (22)\} \circ \{(01) \sqcup t^{21} \sqcup (2)\} \circ \{m^0 \sqcup (1212)\}}_{(0) \circ m^0 = m^0 \circ (00)} \circ \underbrace{\{(0) \sqcup s_1(t^{10}) \sqcup (12)\}}_{\text{expand}} = \\ & \{m^0 \sqcup m^1 \sqcup (22)\} \circ \underbrace{\{(001) \sqcup t^{21} \sqcup (2)\} \circ \{(0) \sqcup t^{10} \sqcup (212)\}}_{\text{expand}} \circ \{(01) \sqcup t^{20} \sqcup (12)\} = \\ & \{m^0 \sqcup m^1 \sqcup (22)\} \circ \{(0) \sqcup t^{10} \sqcup (122)\} \circ \{(010) \sqcup t^{21} \sqcup (2)\} \circ \{(01) \sqcup t^{20} \sqcup (12)\} = \\ & s_0(m^0 \sqcup (11)) \circ s_0((0) \sqcup t^{10} \sqcup (1)) = s_0(\beta_2). \end{aligned}$$

Now assume that the formula holds for n and we prove it for β_{n+1} by using its inductive definition. Note that $(0) \sqcup d^0[1]^n = (01212 \dots 12) = s_1(b_n)$.

$$\begin{aligned} & \{(0) \sqcup d^0 \beta_{n+1}\} \circ s_1 \beta_n \\ & = \{(0) \sqcup d^0 \beta_n \sqcup (2)\} \circ \underbrace{\{s_1 b_{n-1} \sqcup d^0 \beta_2\} \circ \{s_1 \beta_n \sqcup (12)\}}_{\text{Formula (2.26)}} \circ \{[2]^{n-1} \sqcup s_1 \beta_2\} \\ & = \underbrace{\{(0) \sqcup d^0 \beta_n \sqcup (2)\} \circ \{s_1 \beta_n \sqcup (2)\}}_{\text{induction}} \circ \underbrace{\{[2]^{n-1} \sqcup (0) \sqcup d^0 \beta_2\} \circ \{[2]^{n-1} \sqcup s_1 \beta_2\}}_{\text{induction}} \\ & = s_0(\beta_n \sqcup (1)) \circ \{[2]^{n-1} \sqcup s_0 \beta_2\} = s_0(\beta_{n+1}). \end{aligned}$$

This completes the proof of (2). Part (3) follows by applying d_2 to (2), using the simplicial identities and (2.23). \square

2.27. Remark. In Proposition 2.21 we saw that $\text{TW}(0) = \text{TW}(1) = *$. One can also check that $\text{TW}(2)$ is a point, whereas all $\text{TW}(n)$ are infinite-dimensional for $n \geq 3$. Since we will not use these facts, we omit the proofs.

Proof of Theorem 2.11. We have to show that $\text{TW}(n)$ is contractible for all $n \geq 0$. The cases $n = 0, 1$ are covered by Proposition 2.21. Consider $n \geq 2$. Use the morphisms β_n , see Definition 2.24, to define a 0-simplex $\tau(n) \in \text{TW}(n)_0 = \mathcal{B}_0([0]^n, [0])$ by

$$(2.28) \quad \tau(n) = d_1(\beta_n).$$

It is depicted by the tree obtained by deleting the strands labelled 1 from the diagram describing β_n in Definition 2.24.

For every $k \geq 1$, observe that $(0) \sqcup d^0[k-1]^n = s_1^{k-1}(b_n)$, and define

$$(2.29) \quad \Theta_{n,k} = s_1^{k-1}(\beta_n) \in \mathcal{B}_k([k]^n, (0) \sqcup d^0[k-1]^n).$$

We picture $\Theta_{n,k}$ by duplicating $k - 1$ times the strand labelled 1 in the figures in Definition 2.24 and labelling the resulting strands with $1, \dots, k$.

Augment TW by $\text{TW}(n)_{-1} = *$. To show that $\text{TW}(n)$ is contractible, it suffices to show that there is a left contraction [DMN89]

$$s_{-1}: \text{TW}(n)_k \rightarrow \text{TW}(n)_{k+1}, \quad k \geq -1$$

which satisfies

$$(2.30) \quad d_0 s_{-1} = \text{id}; \quad d_i s_{-1} = s_{-1} d_{i-1}; \quad s_i s_{-1} = s_{-1} s_{i-1} \quad \text{for all } i \geq 0.$$

We define $s_{-1}: * \rightarrow \text{TW}(n)_0$ by $s_{-1}(*) = \tau(n)$. For any $\varphi \in \text{TW}(n)_k$, $k \geq 0$, we let $s_{-1}(\varphi)$ be the composition depicted in Figure 2; in formulae:

$$[k+1]^n \xrightarrow{\Theta_{n,k+1}} (0) \sqcup d^0[k]^n \xrightarrow{(0) \sqcup d^0(\varphi)} (0) \sqcup d^0[k] = [k+1] \quad (\text{cf. Remark 2.22}).$$

The first identity of (2.30) holds trivially when $k = -1$. For $k \geq 0$, Lemma 2.25(1),




FIGURE 2. The image of B under the simplicial contraction. B' denotes B with all labels increased by 1.

the simplicial identities, and (2.23) imply

$$d_0(\Theta_{n,k+1}) = d_0 s_1^k(\beta_n) = s_{k+1} \dots s_0^k d_0(\beta_n) = \text{id}.$$

It follows that $d_0 s_{-1} = \text{id}$ because for any $\varphi \in \text{TW}(n)_k$ where $k \geq 0$ we have

$$d_0 s_{-1}(\varphi) = d_0((0) \sqcup d^0(\varphi)) \circ d_0(\Theta_{n,k+1}) = d_0 d^0(\varphi) = \varphi.$$

For the second identity in (2.30), note that $k \geq 0$ because $i > 0$. If $k = 0$ then $i = 1$, and $d_1 d^0(\varphi) = \text{id}_{()}$ for any $\varphi \in \mathcal{B}_0$. Therefore

$$d_1 s_{-1}(\varphi) = d_1((0) \sqcup d^0(\varphi)) \circ d_1(\Theta_{n,1}) = d_1(\beta_n) = \tau(n) = s_{-1}(*) = s_{-1} d_0(\varphi).$$

If $k \geq 1$, the simplicial identities imply that

$$d_i \Theta_{n,k+1} = d_i s_1^k(\beta_n) = s_1^{k-1}(\beta_n) = \Theta_{n,k}.$$

Therefore for any $\varphi \in \text{TW}(n)_k$, we have

$$d_i s_{-1}(\varphi) = d_i(\Theta_{n,k+1}) \circ d_i((0) \sqcup d^0(\varphi)) = \Theta_{n,k} \circ \{(0) \sqcup d^0 d_{i-1}(\varphi)\} = s_{-1} d_{i-1}(\varphi).$$

For the third identity of (2.30), we start with the case $i > 0$, which forces $k \geq 0$:

$$s_i(\Theta_{n,k+1}) = s_i s_1^k(\beta_n) = s_1^{k+1}(\beta_n) = \Theta_{n,k+2}.$$

It follows that for any $\varphi \in \text{TW}(n)_k$,

$$s_i s_{-1}(\varphi) = s_i(\Theta_{n,k+1}) \circ s_i((0) \sqcup d^0(\varphi)) = \Theta_{n,k+2} \circ \{(0) \sqcup d^0 s_{i-1}(\varphi)\} = s_{-1} s_{i-1}(\varphi).$$

It remains to check that $s_0s_{-1} = s_{-1}s_{-1}$, which is the most subtle identity. When $k = -1$ we need to prove that $s_{-1}s_{-1}(\ast) = s_0s_{-1}(\ast)$. By the definition of s_{-1} on $\text{TW}(n)_0$ and by Lemma 2.25(3)

$$s_{-1}s_{-1}(\ast) = s_{-1}\tau(n) = \{(0) \sqcup d^0d_1\beta_n\} \circ \beta_n = s_0d_1\beta_n = s_0\tau(n) = s_0s_{-1}(\ast).$$

For $k \geq 0$, consider some $\varphi \in \text{TW}(n)_k$. By construction,

$$s_{-1}(\varphi) = \{(0) \sqcup d^0(\varphi)\} \circ \Theta_{n,k+1},$$

and therefore, by (2.23),

$$\begin{aligned} s_0s_{-1}(\varphi) &= \{(01) \sqcup s_0d^0(\varphi)\} \circ s_0(\Theta_{n,k+1}) = \{(01) \sqcup d^0d^0(\varphi)\} \circ s_0(\Theta_{n,k+1}) \\ s_{-1}s_{-1}(\varphi) &= \{(0) \sqcup (1) \sqcup d^0d^0(\varphi)\} \circ \{(0) \sqcup d^0(\Theta_{n,k+1})\} \circ \Theta_{n,k+2}. \end{aligned}$$

We are left to show that

$$s_0(\Theta_{n,k+1}) = \{(0) \sqcup d^0(\Theta_{n,k+1})\} \circ \Theta_{n,k+2}.$$

This follows by applying $s_{k+1} \dots s_2$ to Lemma 2.25(2) because $\Theta_{n,k+2} = s_1^{k+1}(\beta_n)$, and because (2.23) implies

$$\begin{aligned} s_0(\Theta_{n,k+1}) &= s_0s_1^k(\beta_n) = s_2^k s_0(\beta_n); \\ d^0(\Theta_{n,k+1}) &= d^0s_1^k(\beta_n) = s_2^k d^0(\beta_n). \end{aligned}$$

□

3. THE OPERAD ACTION ON COMPLETION COSIMPLICIAL OBJECTS

Let $(\mathcal{E}, \diamond, I)$ be a complete and cocomplete monoidal category. In particular, it is tensored and cotensored over sets by

$$E \otimes_{\mathcal{E}} K = \coprod_K E \quad \text{and} \quad [K, E] = \prod_K E.$$

3.1. The simplicial category of cosimplicial objects. Let $c\mathcal{E}$ denote the category of cosimplicial objects in \mathcal{E} . The monoidal structure in \mathcal{E} gives rise to a levelwise monoidal structure \diamond in $c\mathcal{E}$.

The category $c\mathcal{E}$ has a simplicial structure [GJ99, II.2] even if \mathcal{E} does not have one, see [Qui67, II.1.7ff] or [Bou03, §2.10]. This structure is called the *external simplicial structure* and is given by

$$\begin{aligned} [K, X] &= \delta^*[K_\bullet, X^\bullet] && (K \in s\mathbf{Sets}, X \in c\mathcal{E}) \\ X \otimes_{c\mathcal{E}} K &= (\text{LKan}_\delta X) \otimes_\Delta K && (K \in s\mathbf{Sets}, X \in c\mathcal{E}) \\ \text{map}_{c\mathcal{E}}(X, Y) &= \text{Hom}_{c\mathcal{E}}(\text{LKan}_\delta X, Y) && (X, Y \in c\mathcal{E}), \end{aligned}$$

where $\delta: \Delta \rightarrow \Delta \times \Delta$ is the diagonal map, LKan_δ is the left Kan extension along δ^{op} and \otimes_Δ denotes the coend.

If $\Delta[k]$ denotes the standard k -simplex, the usual adjunctions imply

$$(3.1) \quad \text{map}_{c\mathcal{E}}(X, Y)_k = \text{Hom}_{c\mathcal{E}}(X \otimes_{c\mathcal{E}} \Delta[k], Y) = \text{Hom}_{c\mathcal{E}}(X, [\Delta[k], Y]),$$

so we think of the k -simplices of $\text{map}(X, Y)$ as cosimplicial maps $X \rightarrow [\Delta[k], Y]$.

3.2. Proposition. *If $(\mathcal{E}, \diamond, I)$ is a complete and cocomplete monoidal category then the definitions above make $(c\mathcal{E}, \diamond, I)$ a simplicial monoidal category.*

Proof. Quillen shows in [Qui67, II.1] that $c\mathcal{E}$ is tensored and cotensored over $s\mathbf{Sets}$.

The monoidal structure in $c\mathcal{E}$ also gives rise to

$$(3.3) \quad \text{map}_{c\mathcal{E}}(X_1, Y_1) \times \text{map}_{c\mathcal{E}}(X_2, Y_2) \xrightarrow{\circ} \text{map}_{c\mathcal{E}}(X_1 \diamond X_2, Y_1 \diamond Y_2)$$

as follows. Given k -simplices $\varphi: X_1 \rightarrow [\Delta[k], Y_1]$ and $\psi: X_2 \rightarrow [\Delta[k], Y_2]$ we define $\varphi \circ \psi: X_1 \diamond X_2 \rightarrow [\Delta[k], Y_1 \diamond Y_2]$ in cosimplicial degree j as the composition

$$X_1^j \diamond X_2^j \xrightarrow{\prod_{\sigma \in \Delta[k]_j} \varphi^j(\sigma) \circ \psi^j(\sigma)} \prod_{\sigma \in \Delta[k]_j} Y_1^j \diamond Y_2^j.$$

One easily checks that $\varphi \circ \psi$ is a cosimplicial map and that the assignment $(\varphi, \psi) \mapsto \varphi \circ \psi$ gives rise to an associative simplicial map (3.3) which has the constant cosimplicial object I as a unit.

This monoidal product in $c\mathcal{E}$ distributes over the composition. That is, given k simplices $\varphi_i \in \text{map}(X_i, Y_i)_k$ and $\psi_i \in \text{map}(Y_i, Z_i)$ ($i = 1, 2$), we have

$$(\psi_1 \circ \psi_2) \circ (\varphi_1 \circ \varphi_2) = (\psi_1 \circ \varphi_1) \circ (\psi_2 \circ \varphi_2). \quad \square$$

Thus, as mentioned in the introduction, every object $X \in c\mathcal{E}$ gives rise to an operad $\mathcal{O}\text{End}(X)$ whose n th space is $\text{map}_{c\mathcal{E}}(X^{\circ n}, X)$. Its composition law is obtained from the composition and monoidal structure in $c\mathcal{E}$.

3.2. Monoids and twist maps. Before proving Theorem 1.2, we show how to derive Corollary 1.3 from it.

Proof of Corollary 1.3. We need to see that $t: K \diamond K \rightarrow K \diamond K$, given by

$$t = (K \diamond \mu) - \text{id}_{K \diamond K} + (\mu \diamond K),$$

is a twist map (Definition 1.1). It is convenient to write $t = d^0 s^0 - \text{id}_{K \diamond K} + d^1 s^0$ where d^0, d^1 and s^0 are the coface and codegeneracy maps between $K \diamond K$ and K in the cosimplicial object R_K .

We compute

$$\begin{aligned} \mu \circ t &= s^0 \circ (d^0 s^0 - \text{id} + d^1 s^0) = s^0 - s^0 - s^0 = s^0 = \mu, \\ t \circ (K \diamond \eta) &= (d^0 s^0 - \text{id} + d^1 s^0) \circ d^0 = d^0 - d^0 + d^1 = d^1 = \eta \diamond K \\ (K \diamond \mu) \circ (t \diamond K) \circ (K \diamond t) &= s^0 \circ (d^1 s^1 - \text{id}_{K \diamond K} + d^2 s^1) \circ (d^0 s^0 - \text{id}_{K \diamond K} + d^1 s^0) \\ &= d^0 s^0 s^0 - s^1 + d^1 s^0 s^0 = t \circ s^1 = t \circ (\mu \diamond K). \end{aligned}$$

The identities $t \circ (\eta \diamond K) = K \diamond \eta$ and $t \circ (K \diamond \mu) = (\mu \diamond K) \circ (K \diamond t) \circ (t \diamond K)$ follow by symmetry. \square

Proof of Theorem 1.2. Let (K, η, μ) be a monoid in a complete and cocomplete monoidal category $(\mathcal{E}, \diamond, I)$, $R_K \in c\mathcal{E}$ its cobar construction with coface maps d_K^i and codegeneracy maps s_K^i . These extend by monoidality to coface and codegeneracy maps on $R_K^{\circ n}$.

It follows from Lemma 2.7 and Definition 2.8 of \mathcal{B}_k that there are unique monoidal functors $\Phi_k: \mathcal{B}_k \rightarrow \mathcal{E}$ for all $k \geq 0$ defined by

$$\begin{aligned} \Phi_k((a)) &= K, \\ \Phi_k(u^a) &= \eta, \quad \Phi_k(m^a) = \mu, \quad \Phi_k(t^{cd}) = t. \end{aligned}$$

The relations (R1) of Definition 2.8 hold due to the fact that (K, η, μ) is a monoid, and the relations (R2) hold by Property (2) of the twist map. On objects, we thus have $\Phi_k([l]) = (R_K)^l = K^{\circ(l+1)}$.

In Definition 2.8 it was convenient to define the categories \mathcal{B}_S where S is any finite ordered set. In this way, given $a \in S$ we defined functors d_a and s_a in Definition 2.14 which render the categories \mathcal{B}_k a simplicial monoidal category.

We now construct two natural transformations D^a and S^a of functors $\mathcal{B}_S \rightarrow \mathcal{E}$.

3.4. Lemma. *For $a \in S$, there is a natural transformation $D^a: \Phi_{d_a(S)} \circ d_a \rightarrow \Phi_S$ with*

$$D^a: (b) \mapsto \begin{cases} I \xrightarrow{\eta} K, & \text{if } b = a \\ K \xrightarrow{\text{id}} K, & \text{if } b \neq a, \end{cases}$$

and extended to all objects by monoidality. Here $d_a(S) = S - \{a\}$ as in Definition 2.14.

Proof. By construction, \mathcal{B}_k defined in 2.8 is monoidally generated by the objects (b) and the morphisms u^b, m^b and t^{cd} . To prove that D^a is a natural transformation, it suffices by Lemma 2.7 to show that for all $a, b, c, d \in S$,

$$\begin{aligned} \Phi_S(u^b) \circ D^a((b)) &= D^a((b)) \circ \Phi_{d_a(S)}(d_a(u^b)), \\ \Phi_S(m^b) \circ D^a((bb)) &= D^a((bb)) \circ \Phi_{d_a(S)}(d_a(m^b)), \\ \Phi_S(t^{cd}) \circ D^a((cd)) &= D^a((cd)) \circ \Phi_{d_a(S)}(d_a(t^{cd})). \end{aligned}$$

The first equality hold if $a = b$ because $d_a(u^a) = \text{id}$. If $a \neq b$ then $D^a((b)) = \text{id} = D^a((b))$ and the equality holds again since $d^a(u^b) = u^b$.

The second equality holds if $a \neq b$ because $D^a((bb)) = \text{id} = D^a((b))$ and $d^a(m^b) = m^b$. If $a = b$ then $d_a(m^a) = \text{id}_{()}$ so by the relations between η and μ ,

$$\Phi_S(m^a) \circ D^a((aa)) = \mu \circ (\eta \circ \eta) = \eta = D^a((a)).$$

The third equality holds when $a \neq c, d$ because $D^a((cd)) = \text{id}_{K \circ K} = D^a((dc))$ and $d_a(t^{cd}) = t^{cd}$. If $a = c$ then $d_a(t^{cd}) = \text{id}$ (cf. Definition 2.8) so

$$\Phi_S(t^{cd}) \circ D^a((cd)) = t \circ (\mu \circ K) = K \circ \eta = D^a((dc)) \circ \Phi_{d_a(S)}(d_a(t^{cd})).$$

A similar calculation applies when $a = d$. □

3.5. Lemma. *For $a \in S$, there is a natural transformation $S^a: \Phi_{s_a(S)} \circ s_a \rightarrow \Phi_S$ with*

$$S^a: (b) \mapsto \begin{cases} K \circ K \xrightarrow{\mu} K, & \text{if } b = a \\ K \xrightarrow{\text{id}} K, & \text{if } b \neq a, \end{cases}$$

and extended to all objects by monoidality. Here $s_a(S) = S - \{a\} \sqcup \{a', a''\}$ as in Definition 2.14.

Proof. As in the proof of Lemma 3.4, in order to prove the naturality of S^a we have to show that

$$\begin{aligned} \Phi_S(u^b) \circ S^a((b)) &= S^a((b)) \circ \Phi_{s_a(S)}(s_a(u^b)), \\ \Phi_S(m^b) \circ S^a((bb)) &= S^a((bb)) \circ \Phi_{s_a(S)}(s_a(m^b)), \\ \Phi_S(t^{cd}) \circ S^a((cd)) &= S^a((cd)) \circ \Phi_{s_a(S)}(s_a(t^{cd})). \end{aligned}$$

The first and second equalities are easy to prove when $a \neq b$ because $S^a(bb) = \text{id} = S^a(b)$ and $s^a(m^b) = m^b$. When $a = b$ we see from Definition 2.8 that

$$\Phi_S(u^b) \circ S^a() = u = \mu \circ (\eta \diamond \eta) = S^a(a) \circ \Phi_{s_a S}(s_a(u^a)).$$

For the second equality ($a = b$), we use the associativity of μ and Definitions 1.1(1) and 2.14 to deduce that

$$\begin{aligned} S^a(a) \circ \Phi_{s_a S}(s^a(m^a)) &= \mu \circ \Phi_{s_a S}(m^{a'} \sqcup m^{a''}) \circ \Phi_{s_a S}((a') \sqcup t^{a'' a'} \sqcup (a'')) \\ &= \mu \circ (\mu \diamond \mu) \circ (K \diamond t \diamond K) = \mu \circ (\mu \diamond \mu) = \Phi_S(m^a) \circ S^a(aa). \end{aligned}$$

The third equality is again straightforward when $a \neq c, d$. If $a = c$ then by Definitions 2.8, 2.14 and 1.1(3),

$$\begin{aligned} S^a(da) \circ \Phi_{s_a S}(s_a(t^{ad})) &= (K \diamond \mu) \circ \Phi_{s_a S}(t^{a'd} \sqcup (a'')) \circ \Phi_{s_a S}((a') \sqcup t^{a'' d}) \\ &= (K \diamond \mu) \circ (t \diamond K) \circ (K \diamond t) = t \circ (\mu \diamond K) = \Phi_S(t^{ad}) \circ S^a(ad). \end{aligned}$$

A similar calculation applies when $a = d$. \square

When $S = [k]$, we observe that $\Phi_S([k]) = K^{\circ(k+1)} = R_K^k$ and that for any $i \in [k]$ we have $D^i([k]) = d_{R_K}^i$, $S^i([k]) = s_{R_K}^i$, and consequently $D^i([k]^n) = d_{R_K^{\circ n}}^i$ and $S^i([k]^n) = s_{R_K^{\circ n}}^i$.

3.6. Definition. For integers $n, m \geq 0$ let $\text{TW}(n, m)$ denote the simplicial set whose set of k -simplices is $\mathcal{B}_k([k]^n, [k]^m)$. In particular, $\text{TW}(n, 1) = \text{TW}(n)$ (cf. Definition 2.10).

Here we use the fact that $d_i: \mathcal{B}_k \rightarrow \mathcal{B}_{k-1}$ and $s_i: \mathcal{B}_k \rightarrow \mathcal{B}_{k+1}$ are monoidal functors which carry $[k]$ to $[k-1]$ and $[k+1]$ respectively, see Proposition 2.9. Composition of morphisms and the monoidal operation \sqcup in \mathcal{B}_k give rise to ‘‘composition’’ and ‘‘monoidal’’ products of simplicial sets

$$(3.7) \quad \begin{aligned} \text{TW}(n, m) \times \text{TW}(m, l) &\xrightarrow{\circ} \text{TW}(n, l), \\ \text{TW}(n, m) \times \text{TW}(n', m') &\xrightarrow{\sqcup} \text{TW}(n+n', m+m'). \end{aligned}$$

Write $X(n) = R_K^{\circ n}$ and $E(m, n) = \text{map}_{c\mathcal{E}}(X(m), X(n))$. Thus $\Phi_k([k]^n) = X(n)^k$. By Proposition 2.9, \mathcal{B}_\bullet is a simplicial monoidal category and thus a morphism $\alpha \in \Delta([i], [j])$ gives rise to a functor $\alpha^*: \mathcal{B}_j \rightarrow \mathcal{B}_i$.

For $f \in \text{TW}(n, m)_k$ and $j \geq 0$, we define a morphism $X(n)^j \rightarrow [\Delta[k], X(m)]^j$ by

$$X(n)^j \xrightarrow{\Pi_\alpha \Phi_j(\alpha^*(f))} \prod_{\alpha \in \Delta[k]_j} X(m)^j.$$

These maps assemble to a cosimplicial map

$$\rho(n, m)_k(f): X(n) \rightarrow [\Delta[k], X(m)],$$

because by Lemma 3.4, for every injective $\epsilon^i: [j] \rightarrow [j+1]$ in Δ and every $\alpha \in \Delta[k]_{j+1}$,

$$\begin{aligned} d_{X(m)}^i \circ \Phi_j((\alpha \circ \epsilon^i)^*(f)) &= D^i([k]^m) \circ \Phi_{d_i[j+1]}(d_i(\alpha^*(f))) = \\ &= \Phi_{j+1}(\alpha^*(f)) \circ D^i([k]^n) = \Phi_{j+1}(\alpha^*(f)) \circ d_{X(n)}^i. \end{aligned}$$

Similarly, by Lemma 3.5, for any surjective $\eta^i: [j] \rightarrow [j-1]$ and every $\alpha \in \Delta[k]_{j-1}$,

$$\begin{aligned} s_{X(m)}^i \circ \Phi_j((\alpha \circ \eta^i)^*(f)) &= S^i([k]^m) \circ \Phi_{s_i([j-1])}(s_i(\alpha^*(f))) = \\ &= \Phi_{j-1}(\alpha^*(f)) \circ S^i([k]^n) = \Phi_{j-1}(\alpha^*(f)) \circ s_{X(n)}^i. \end{aligned}$$

Thus, $\rho(n, m)_k(f) \in E(n, m)_k$, see (3.1). They assemble to form a simplicial map

$$\rho(n, m): \text{TW}(n, m) \rightarrow E(n, m).$$

because for any $\gamma \in \Delta([\ell], [k])$, $\alpha \in \Delta[\ell]_j$, and $f \in \text{TW}(n, m)_k$, we have

$$\Phi_j((\gamma \circ \alpha)^*(f)) = \Phi_j(\alpha^*(\gamma^*(f))),$$

so $\rho(n, m)_\ell(\gamma^*(f)) = \gamma^*\rho(n, m)_k(f)$.

The following diagrams of spaces commute:

$$\begin{array}{ccc} \text{TW}(n, m) \times \text{TW}(m, l) & \xrightarrow{\circ} & \text{TW}(n, l) \\ \rho(n, m) \times \rho(m, l) \downarrow & & \downarrow \rho(n, l) \\ E(n, m) \times E(m, l) & \xrightarrow{\circ} & E(n, l) \end{array}$$

$$\begin{array}{ccc} \text{TW}(n, m) \times \text{TW}(n', m') & \xrightarrow{\sqcup} & \text{TW}(n+n', m+m') \\ \rho(n, m) \times \rho(n', m') \downarrow & & \downarrow \rho(n+n', m+m') \\ E(n, m) \times E(n', m') & \xrightarrow{(3.3)} & E(n+n', m+m'). \end{array}$$

This follows by verification in every simplicial degree k using the fact that the functors Φ_k are monoidal. For example, given $f \in \text{TW}(n, m)_k$ and $g \in \text{TW}(m, l)_l$, the cosimplicial map $\rho(m, l)_k(g) \circ \rho(n, m)_k(f)$ has the form

$$X(n)^j \xrightarrow{\prod_{\alpha} \Phi_j(\alpha^*(g)) \circ \Phi(\alpha^*(f))} \prod_{\alpha \in \Delta[k]_j} X(l)^j.$$

Now $\Phi_j(\alpha^*(g)) \circ \Phi_j(\alpha^*(f)) = \Phi_j(\alpha^*(g \circ f))$; which assemble to $\rho(n, l)_k(g \circ f)$. A similar argument works for the second diagram.

Since the composition law in TW (Definition 2.10) is derived from (3.7), the commutativity of the diagrams above implies that the maps

$$\rho(n, 1): \text{TW}(n) \rightarrow \text{map}_{c\mathcal{E}}(X(n), X(1))$$

form a morphism of operads $\text{TW} \rightarrow \mathcal{O}\text{End}(R_K)$. This completes the proof of Theorem 1.2. \square

APPENDIX A. THE FREE MONOIDAL CONSTRUCTION

In this appendix we will prove Lemma 2.5. The result is an exercise in free constructions, and probably known to anybody who has worked with small monoidal categories, but we could not find it in the literature.

A.1. Lemma. *The functor $U_{vm}: \{\text{monoidal graphs}\} \rightarrow \{\text{vertex-monoidal graphs}\}$ has a left adjoint.*

Proof. The adjoint is given by $F_{vm}(\mathcal{A} \rightrightarrows \mathcal{O}) = (\mathcal{A}_{\sqcup} \rightrightarrows \mathcal{O})$, where \mathcal{A}_{\sqcup} is the free monoid on \mathcal{A} and the structure maps are extended uniquely to monoidal maps. The adjointness is obvious. \square

A.2. Lemma. *The functor $U_m: \{\text{unital monoidal graphs}\} \rightarrow \{\text{monoidal graphs}\}$ has a left adjoint.*

Proof. Let $\mathcal{A} \rightrightarrows \mathcal{O}$ be a monoidal graph. For every $\sigma \in \mathcal{O}$ we introduce a symbol $\iota(\sigma) \notin \mathcal{A}$ and let $\tilde{\Delta}$ denote $\mathcal{A} \sqcup \{\iota(\sigma)\}_{\sigma \in \mathcal{O}}$. Extend $\partial_0, \partial_1: \mathcal{A} \rightarrow \mathcal{O}$ to $\tilde{\Delta}$ by setting $\partial_0(\iota(\sigma)) = \partial_1(\iota(\sigma)) = \sigma$.

This is not quite a unital monoidal graph because $\iota: \mathcal{O} \rightarrow \tilde{\Delta}$ is not monoidal. We enforce this by setting $\Delta = \tilde{\Delta} / \sim$, where \sim is the equivalence relation generated by

$$\iota(1_{\mathcal{O}}) \sim 1_{\mathcal{A}} \quad \text{and} \quad \iota(\sigma \sqcup \tau) \sim \iota(\sigma) \sqcup \iota(\tau).$$

Since $\partial_i(a) = \partial_i(a')$ if $a \sim a' \in \tilde{\Delta}$, they are well-defined on Δ , and we obtain a unital monoidal graph (Δ, \mathcal{O}) . Again, adjointness is easily checked. \square

A.3. Lemma. *The functor $\{\text{small monoidal categories}\} \xrightarrow{U_m} \{\text{unital monoidal graphs}\}$ has a left adjoint.*

Proof. Let $\mathcal{G} = (\partial_0, \partial_1: \mathcal{A} \rightrightarrows \mathcal{O})$ be a unital monoidal graph. Let $\tilde{\mathcal{M}}$ denote $C(\mathcal{G})$, the free category generated by the graph \mathcal{G} . However, $\tilde{\mathcal{M}}$ is not monoidal yet.

Let \sim be the smallest equivalence relation on the morphism set of $\tilde{\mathcal{M}}$ with

- (i) $(f \sqcup g) \circ (f' \sqcup g') \sim (f \circ f') \sqcup (g \circ g')$ for all composable arrows f, f' and g, g' in $\tilde{\mathcal{M}}$,
- (ii) \sim is closed under \sqcup , i.e. if $f \sim f'$ and $g \sim g'$ then $(f \sqcup g) \sim (f' \sqcup g')$.

Again, the equivalence relation \sim has the property that if $f \sim f'$ then their domains and codomains are equal, and thus we obtain a quotient category $\mathcal{M} = \tilde{\mathcal{M}} / \sim$ [ML98, §II.8, Proposition 1]. The monoidal structure \sqcup descends from $\tilde{\mathcal{M}}$ thanks to (ii). It renders \mathcal{M} a monoidal category by virtue of relation (i). \square

REFERENCES

- [BK72] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Springer-Verlag, Berlin, 1972. Lecture Notes in Mathematics, Vol. 304.
- [BL07] Tilman Bauer and Assaf Libman. A_∞ -monads and completion. Preprint, 2007.
- [Bou03] A. K. Bousfield. Cosimplicial resolutions and homotopy spectral sequences in model categories. *Geom. Topol.*, 7:1001–1053 (electronic), 2003.
- [BT00] Martin Bendersky and Robert D. Thompson. The Bousfield-Kan spectral sequence for periodic homology theories. *Amer. J. Math.*, 122:599–635, 2000.
- [DMN89] William Dwyer, Haynes Miller, and Joseph Neisendorfer. Fibrewise completion and unstable Adams spectral sequences. *Israel J. Math.*, 66(1-3):160–178, 1989.
- [EKMM97] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory*, volume 47 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole.
- [FI07] Halvard Fausk and Daniel C. Isaksen. Model structures on pro-categories. *Homology, Homotopy Appl.*, 9(1):367–398 (electronic), 2007.
- [GJ99] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*. Birkhäuser Verlag, Basel, 1999.
- [HSS00] Mark Hovey, Brooke Shipley, and Jeff Smith. Symmetric spectra. *J. Amer. Math. Soc.*, 13(1):149–208, 2000.
- [Isa04] Daniel C. Isaksen. Strict model structures for pro-categories. In *Categorical decomposition techniques in algebraic topology (Isle of Skye, 2001)*, volume 215 of *Progr. Math.*, pages 179–198. Birkhäuser, Basel, 2004.
- [May72] Peter May. *The Geometry of Iterated Loop Spaces*. Number 271 in LNM. Springer Verlag, 1972.
- [ML98] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.

[Qui67] Daniel Quillen. *Homotopical Algebra*. Number 43 in LNM. Springer Verlag, 1967.

FACULTEIT DER EXACTE WETENSCHAPPEN, VRIJE UNIVERSITEIT AMSTERDAM, DE BOELELAAN
1081A, 1081HV AMSTERDAM, THE NETHERLANDS
E-mail address: `tilman@few.vu.nl`

DEPARTMENT OF MATHEMATICAL SCIENCES, KING'S COLLEGE, UNIVERSITY OF ABERDEEN, AB-
ERDEEN AB24 3UE , SCOTLAND, U.K.
E-mail address: `assaf@maths.abdn.ac.uk`