

VU Research Portal

Single-fluid radiation-hydrodynamic modeling of laser-driven EUV-emitting plasmas

Hemminga, Diko Jelle

2023

DOI (link to publisher)
[10.5463/thesis.423](https://doi.org/10.5463/thesis.423)

document version
Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Hemminga, D. J. (2023). *Single-fluid radiation-hydrodynamic modeling of laser-driven EUV-emitting plasmas*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].
<https://doi.org/10.5463/thesis.423>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:
vuresearchportal.ub@vu.nl

Contents

Introduction	1
1 Plasma	1
2 Extreme ultraviolet lithography	2
3 Laser-produced plasma	4
4 Research methodology	6
5 Thesis outline and summary	7
1 Radiation hydrodynamics as a model for laser-produced plasma	11
1.1 Plasma	13
1.1.1 Laser-produced plasmas	13
1.2 Kinetic theory of plasma	15
1.2.1 Distribution functions and the Boltzmann equation	16
1.2.2 Vlasov equation	17
1.2.3 Macroscopic quantities based on distribution functions	18
1.2.4 Maxwell–Boltzmann distribution and the Boltzmann factor	19
1.3 Fluid approach to plasma modeling	20
1.3.1 From plasma particle distributions to a fluid approach	20
1.3.2 Single-fluid, single-temperature approach	24
1.4 System of equations underlying the fluid simulations	26

1.4.1	Heat conduction	27
1.4.2	Radiation transport	29
1.4.3	Laser propagation and absorption	30
1.5	Equation-of-state	37
1.5.1	Ideal gas law	37
1.5.2	Speed of sound	38
1.5.3	Frankfurt equation-of-state	40
1.6	Analytic formulas for plasma expansion	42
1.6.1	Planar isothermal expansion	43
1.6.2	Planar isentropic expansion	47
1.6.3	Application to laser-produced plasma expansion	47
2	High-energy ions from Nd:YAG laser ablation of tin microdroplets: Comparison between experiment and a single-fluid hydrodynamic model	51
2.1	Introduction	53
2.2	Experimental setup, method and results	56
2.3	Radiation-hydrodynamic simulations and the RALEF-2D code	58
2.3.1	Single-fluid single-temperature radiation hydrodynamics	58
2.3.2	RALEF-2D	59
2.4	Plasma formation and expansion	61
2.4.1	Initial burst of laser-induced ablation	64
2.4.2	Second burst of laser-induced ablation	65
2.5	Ion kinetic energy distributions: experiment and simulation	68
2.6	Conclusion	71
3	Strongly anisotropic ion emission in the expansion of Nd:YAG-laser-produced plasma	73
3.1	Introduction	75
3.2	Experiments	76
3.2.1	Experimental setup	76
3.2.2	Experimental methods	79
3.3	Radiation-hydrodynamic simulations	79
3.3.1	RALEF-2D	79
3.3.2	Application of RALEF-2D	80
3.3.3	Ion distributions from simulation	80
3.4	Results and discussion	82
3.4.1	Ion emission anisotropy	82
3.4.2	Angular distributions of number, momentum, and energy	85

3.4.3	Droplet plasma propulsion	88
3.5	Conclusion	90
	Appendix	91
4	Early-time hydrodynamic response of a tin droplet driven by laser-produced plasma	93
4.1	Introduction	95
4.2	Experiment	98
4.2.1	Experimental setup	98
4.2.2	Experimental results	99
4.3	Simulation	106
4.3.1	Code description	106
4.3.2	Simulation results	108
4.4	Discussion	111
4.5	Conclusion	115
	Appendix I: Scaling of U and \dot{R}_0 with laser energy on droplet	116
	Appendix II: Kinetic energy partition versus velocity ratio	118
5	Simulations of plasmas driven by laser wavelengths in the 1.064 – 10.6 μm range as future extreme ultraviolet light sources	121
5.1	Introduction	123
5.2	Numerical simulations	124
5.2.1	Code description	126
5.2.2	Simulation parameters	126
5.3	Wavelength-dependent plasma dynamics	127
5.3.1	Establishment of steady-state plasma flow	128
5.3.2	Wavelength-dependent power partitioning	130
5.4	Intensity-dependent plasma dynamics	131
5.4.1	Dependence of spatial plasma features on laser intensity	133
5.4.2	Intensity-dependent power partitioning	135
5.5	Summary and conclusion	138
	Conclusion and outlook	139
1	Conclusion	140
2	Outlook	143
	Bibliography	146

List of Publications	163
Acknowledgments	165