

VU Research Portal

SIMONI (smart integrated monitoring) as a novel bioanalytical strategy for water quality assessment: Part I—model design and effect-based trigger values

van der Oost, Ron; Sileno, Giulia; Suárez-Muñoz, Maria; Nguyen, Mai Thao; Besselink, Harrie; Brouwer, Abraham

published in

Environmental Toxicology and Chemistry
2017

DOI (link to publisher)

[10.1002/etc.3836](https://doi.org/10.1002/etc.3836)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

van der Oost, R., Sileno, G., Suárez-Muñoz, M., Nguyen, M. T., Besselink, H., & Brouwer, A. (2017). SIMONI (smart integrated monitoring) as a novel bioanalytical strategy for water quality assessment: Part I—model design and effect-based trigger values. *Environmental Toxicology and Chemistry*, 36(9), 2385-2399. <https://doi.org/10.1002/etc.3836>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

SIMONI (SMART INTEGRATED MONITORING) AS A NOVEL BIOANALYTICAL STRATEGY FOR WATER QUALITY ASSESSMENT: PART I—MODEL DESIGN AND EFFECT-BASED TRIGGER VALUES

RON VAN DER OOST,^{a,*} GIULIA SILENO,^{a,b} MARIA SUÁREZ-MUÑOZ,^b MAI THAO NGUYEN,^c HARRIE BESSELINK,^d and ABRAHAM BROUWER^{d,e}^aDepartment of Technology, Research and Engineering, Waternet Institute for the Urban Water Cycle, Amsterdam, The Netherlands^bInstitute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands^cWaterproof Laboratory, Research & Validation, Edam, The Netherlands^dBioDetection Systems, Amsterdam, The Netherlands^eInstitute for Ecological Sciences, Free University, Amsterdam, The Netherlands

(Submitted 15 July 2016; Returned for Revision 4 November 2016; Accepted 23 February 2017)

Abstract: It is virtually impossible to reliably assess water quality with target chemical analyses only. Therefore, a complementary effect-based risk assessment by bioanalyses on mixtures of bioavailable micropollutants is proposed: the Smart Integrated Monitoring (SIMONI) strategy. The goal of this strategy is to obtain more reliable information on the water quality to select optimum measures for improvement. The SIMONI strategy is 2-tiered. Tier 1 is a bioanalytical hazard identification of sites. A tier 2 ecological risk assessment is carried out only at a limited number of sites where increased hazards are detected in tier 1. Tier 2 will be customized, based on tier 1 evaluation and additional knowledge of the aquatic system. The present study focuses on the tier 1 bioanalytical hazard identification to distinguish “hot spots” of chemical pollution. First, a selection was made of relevant and cost-effective bioanalytical endpoints to cover a wide spectrum of micropollutant modes of action. Specific endpoints may indicate which classes of chemicals might cause adverse effects. Second, effect-based trigger values (EBT) were derived for these bioassays to indicate potential ecological risks. Comparison of EBT with bioassay responses should discriminate sites exhibiting different chemical hazards. Third, a model was designed to estimate the overall risks for aquatic ecosystems. The associated follow-up for risk management is a “toxicity traffic light” system: green, low hazard (no action required); orange, potential risk (further research needed); and red, high risk (mitigation measures). Thanks to cost-effectiveness, flexibility, and relevance, the SIMONI strategy has the potential to become the first bioanalytical tool to be applied in regular water quality monitoring programs. *Environ Toxicol Chem* 2017;36:2385–2399. © 2017 SETAC

Keywords: Bioanalytical monitoring Micropollutant Bioassay Aquatic hazard and risk assessment Effect-based trigger value

INTRODUCTION

Regular water quality monitoring and its limitations

Water quality monitoring should be a control mechanism to assess 1) whether or not measures are needed to improve quality, 2) which measures should be applied, and 3) how successful these measures are in improving the water quality. The European Water Framework Directive tries to integrate biological and chemical information to obtain an overall insight into the quality of individual water bodies. According to the Water Framework Directive, ecological status is determined by monitoring biological quality elements (absence and presence of taxa), hydromorphological elements, and physicochemical elements of the water bodies. The chemical status of a water body is determined by analyzing the concentrations of 45 (groups of) priority substances. A good chemical status is reached when the concentrations of all substances are below the annual average and maximal allowable environmental quality standards defined to protect the environment and human health [1].

Regular chemical monitoring of water quality is almost exclusively performed by targeted chemical analysis of a limited set of compounds. There are, however, some serious limitations related to the use of only chemical analyses of spot samples for monitoring the overall chemical status. First, because only a limited number of target substances are analyzed, the risk of nontarget and unknown substances in the aquatic environment is unclear [2]. At present (December 2016), more than 125 000 000 substances are registered at the Chemical Abstracts Service, whereas already 20 yr ago it was estimated that more than 100 000 chemicals are potential pollutants of aquatic ecosystems [3]. Second, it is obvious that chemicals occur not as single substances in the environment but in complex mixtures with potential synergism or antagonism. Although concentrations of individual chemicals can be below lowest-observed-effect concentrations or detection limits, the entire mixture may still cause adverse effects [4]. Moreover, transformation products of micropollutants may be more toxic and persistent than the parent compounds [5]. These limitations may thus result in an incomplete assessment of the chemical hazards (e.g., Van der Oost et al. [6]), urging alternative approaches to be explored.

Chemical quality monitoring with bioanalytical tools

Effect-based monitoring tools to assess the chemical water quality by measuring effects instead of substances have already been applied for more than 3 decades. Bioanalyses are being

This article includes online-only Supplemental Data.

* Address correspondence to ron.van.der.oost@waternet.nl

Published online 4 May 2017 in Wiley Online Library (wileyonlinelibrary.com).

DOI: 10.1002/etc.3836

performed using 2 approaches, 1) biomarkers in feral or caged organisms that are exposed in the field (e.g., reviews by Stegeman et al. [7] and Van der Oost et al. [6]), and 2) bioassays with laboratory organisms or cell lines that are exposed to environmental samples or extracts (e.g., reviews by Castaño et al. [8] and Durand et al. [9]). Most in vivo assays (whole organisms) measure effects on gross parameters, such as growth, reproduction, feeding activity, and mortality, whereas most in vitro assays (cell lines or unicellular organisms) measure specific biochemical effects of bioactive compounds, such as endocrine disruption and genotoxicity. Three major mechanism-of-action groups can be distinguished regarding the type of interaction between a chemical and its molecular target: nonspecific, specific, and reactive toxicity. "Nonspecific toxicity" or "baseline toxicity" refers to the minimum cytotoxicity that a chemical can exhibit not mediated by specific mechanisms (narcosis). "Specific toxicity" refers to all common mechanisms that involve the selective binding of a chemical to a protein (enzyme or receptor). Mechanisms of action are classified as reactive when covalent bonds are formed between the chemical and its target or when chemical reactions such as oxidative stress are involved [10].

The added value of these effect-based tools for ecological risk assessment has been demonstrated in numerous studies (reviewed by Van der Oost et al. [6]). Firstly, in vivo assays respond to the presence of all pollutants in the water sample as well as their bioavailability and physical transfer into the test organism. Secondly, all bioanalytical tools give a more holistic assessment of biologically active chemicals present in the water because they are able to detect mixture toxicity and the effects of metabolites and unknowns. For bioassays indicative of integrative effects (i.e., cytotoxicity, reactive and adaptive stress responses) typically only a very small fraction of the effect (often <1%) can be explained by known and identified chemicals [11,12]. For other endpoints (e.g., estrogenic activity) it is easier to identify the compounds that cause the effects [13]. Water contaminants can elicit effects by interacting with critical cellular targets such as receptors, proteins, DNA, or phospholipids that trigger a range of cellular events like the activation of genes, production of proteins, and altered protein signaling. Therefore, a series of pathways can be activated by contaminant exposure [14]. The concept of toxicity pathways is put into a wider ecotoxicological perspective as adverse outcome pathways, linking the toxicity pathway at the cellular level, via responses at organ and organism levels, and ultimately to the response at the population level [15].

Requirements for a bioanalytical hazard identification

There are European Union regulations that allow for bioanalytical methods in screening of feed and food for dioxin-like chemicals [16]. It is conceivable to adopt the bioanalytical equivalency concept (BEQ) also for the development of bioanalytical trigger values for water quality assessment [11]. Yet, several conditions have to be met. A well-designed bioassay battery has the potential to provide a cost-effective assessment of the environmental risks caused by thousands of micropollutants. Several preconditions are important for the design of a monitoring strategy that is aimed at assessing the potential ecological risks from chemical pollution. The 6 criteria we considered most important for an effect-based monitoring strategy to be widely applied for regular water quality monitoring are as follows. First, identification of a broad spectrum of chemical pollutants. A hazard-identification strategy based on effect monitoring should

be able to demonstrate the overall hazard of a wide range of chemical pollutants and their transformation products. The design of a good bioassay panel should thus cover the various types of toxic action, that is, nonspecific (various trophic levels), specific, and reactive toxicity. Second, discrimination of sites with potential ecological health risks. The hazard assessment should use effect-based trigger (EBT) values to prioritize the sites where the highest ecological risks can be expected, but not all sites should be classified as hazardous. Third, cost-effectiveness of the panel of bioassays in terms of equipment and consumables. Important for the acceptance of an alternative monitoring strategy is an attractive alternative that provides better ecological health-based information for the same budget or less. Fourth, good performance of bioassays, preferably with International Organization for Standardization or related validation level. The selected toxicological endpoints have to be measured in bioassays that meet certain performance quality standards, such as selectivity, accuracy, reproducibility, robustness, sensitivity, speed, and potential high-throughput capacity. Fifth, easy implementation and applicability of bioanalytical techniques by routine laboratories. The bioassays should be able to analyze environmental samples (complex mixtures) without high-tech laboratory requirements or specialist knowledge. Sixth, relevant and effective water sampling. Snapshot grab sampling is unreliable when water concentrations of micropollutants are varying. An alternative method is time-integrated sampling with passive samplers that are able to concentrate bioavailable micropollutants on site and may be a good reflection of the micropollutants that accumulate in tissues of aquatic organisms [17,18]. There are, however, certain pitfalls when combining passive sampling and bioassays [19] that will be discussed in a follow-up study [20]. Although online bioassays for real-time effects-based monitoring would be ideal for this purpose, current technologies are not sensitive enough to analyze the risks of low concentrations of micropollutants.

Objectives of the present study

Because of the shortcomings of regular chemical monitoring, there is an urgent need for more holistic evaluations of chemical water quality. In addition, monitoring should provide a more concrete quantitative risk classification, instead of the one-out-all-out principle of chemical environmental quality standard. The aim of the present study is to develop a strategy using complementary toxicological and chemical approaches, providing holistic risk quantification for chemical micropollutants to the aquatic environment. This strategy not only is appropriate for research projects but can be applied in regular water quality monitoring. The present study describes the design of the SIMONI (Smart Integrated Monitoring) strategy that meets the requirements outlined in the *Requirements for a bioanalytical hazard identification* section. This alternative for regular chemical monitoring does not advocate revolutionary new bioanalytical tools but aims to find the optimal cost-effective combination of validated in vivo and in vitro endpoints to estimate environmental hazards by applying EBT values.

The primary objectives of the present study were to describe the selection of relevant toxicological endpoints and the derivation of EBT values for a suite of in vivo and in vitro bioassays, to be used in the SIMONI strategy for water quality assessment. This strategy is based on a bioanalytical monitoring battery that is suitable to identify a wide range of chemical hazards. The EBT values for environmental risks were derived and used to interpret and classify the observed bioassay responses. In addition, a simple model is described that

integrates all bioassay responses into a quantitative SIMONI score for hot spot assessment of ecological risks. The SIMONI strategy for environmental hazard assessment will allow regulators to link the bioassay results to potential adverse effects on the aquatic ecosystem. The strategy aims to establish a framework for the most cost-effective bioassay panel that reliably indicates the broad-spectrum chemical hazards for both invertebrate and vertebrate aquatic organisms.

SIMONI DESIGN AND BIOANALYTICAL ENDPOINTS

Design of the SIMONI strategy

A 2-tiered strategy was designed, based on the combination of field-exposed passive samplers and laboratory bioassay measurements (Figure 1). The first tier is hazard identification, and the second tier is risk assessment. The strategy can also be applied for bioanalytical examination of concentrated large-volume water samples to more accurately quantify the results (e.g., for tier 2 risk assessment).

The first tier of the strategy, hazard identification, is applied to assess the potential risks of a broad-spectrum mixture of chemical micropollutants. The main objective of this screening phase is to identify the “hot spots” of chemical water pollution. Hazards of organic micropollutants are characterized by evaluating the responses of a suite of validated bioassays, using EBT values. In this tier, chemical analyses are only performed on metals and ammonium. Concentrations of inorganic compounds can be directly compared with environmental quality standard values for ecological risk assessment. A limitation of the strategy described in the present study is that mixture effects determined with bioassays do not account for potential mixture interaction between organic and inorganic substances, except for a field *Daphnia* test that is exposed to surface water. Only a limited number of sites, where bioassay responses exceed EBT and indicate potential ecological risks, should be examined by tier 2, a more expensive second phase, for the actual risk assessment. All EBT exceedances indicate environmental hazards. However, if only a slight EBT exceedance is observed in a bioassay, this does not necessarily indicate an increased ecological risk. Therefore, a model is designed (see *SIMONI Model for Overall Micropollutant Risks*) which adds up EBT excesses in all bioassays and calculates an

overall SIMONI score that should be indicative for the ecological risks. The tier 2 risk-assessment phase should be customized, based on tier 1 data and knowledge of the water system. If the tier 1 classification indicates low chemical hazards, there is no need to apply more advanced and expensive chemical analyses (e.g., Water Framework Directive priority pollutants) and bioanalytical methods (e.g., fish bioassays and biomarkers). These analyses have to be carried out at sites with increased chemical hazards, where they are most relevant. If tier 2 chemical analyses do not give the answers needed to explain the effects observed in tier 1, a chemical–toxicological effect-directed analysis can be performed using (in vitro) bioassays with the most pronounced responses to identify the unknown compounds that cause the observed toxicity. Results of the risk assessment can be verified with ecological observations, such as the occurrence of species that are sensitive to certain micropollutants.

Selection of bioanalytical endpoints

The selection of the bioanalytical endpoints for the SIMONI effect-based strategy was based on the requirements listed (see *Requirements for a bioanalytical hazard identification*). For the detection of toxic responses of a broad spectrum of micropollutants it is important to select some nonspecific in vivo assays, covering different trophic levels of aquatic organisms. In addition, the most relevant specific and reactive in vitro endpoints for water quality assessment have to be selected. A major advantage of specific in vitro responses is that these may indicate which classes of chemicals and which types of effects may cause the main problem for aquatic organisms. The final selection of toxic endpoints was based on the literature [21–24] as well as our own research [20].

Willemsen et al. [21] evaluated the use of approximately 30 aquatic bioassays for nonspecific toxicity, teratogenicity, and genotoxicity. Selection criteria used for the composition of a test battery were 1) acute tests with whole organisms, 2) a small test volume, 3) tests available in kits, and 4) no specifically trained personnel or extensive laboratory facilities needed. Various methods were compared in terms of documentation, reproducibility, sensitivity, exposure time, standardization, technical simplicity, and costs. A test battery was proposed using bacteria (Microtox[®]), algae (microplate assay), crustaceans (Thamnotoxkit FTM), and a *Daphnia* (IQ Toxicity TestTM) test.

Van der Linden et al. [22] investigated the possible adverse effects of endocrine-disrupting compounds (EDCs) with a panel of in vitro bioassays. The bioassays for estrogen (ER), androgen (AR), and glucocorticoid (GR) receptor activities were applied to extracts of municipal wastewater-treatment plant (WWTP) effluents. Different types of hormone receptor activity were detected and used to quantify the presence of EDCs.

In a study by Macova et al. [23] a bioanalytical test panel was used for monitoring organic micropollutants across the complete water cycle from sewage to drinking water. Six endpoints targeting groups of chemicals with modes of toxic action relevant for human and environmental health were included in the evaluation: genotoxicity, endocrine disruption, neurotoxicity, phytotoxicity, xenobiotic metabolism, and nonspecific cell toxicity. All selected toxicity endpoints appeared to be relevant to evaluate the water cycle quality and showed the highest responses in WWTP influents. The effects in all 6 selected bioassays decreased across 7 water-treatment barriers.

In an extensive interlaboratory study by Escher et al. [24] a representative set of water samples was analyzed for a broad




Figure 1. Schematic presentation of the Smart Integrated Monitoring (SIMONI) effect-based monitoring strategy. EBT = effect-based trigger value; EDA = effect directed analysis; EQS = environmental quality standard; msPAF = multiple substances potentially affected fraction of species; TIE = toxicity identification and evaluation.

range of toxicological effects by using 103 unique in vitro bioassays. Ten samples of WWTP effluent, recycled waters, storm water, surface water, and drinking water were examined. Each water type had a characteristic bioanalytical profile. The most relevant modes of toxic action identified in that study were related to nonspecific toxicity, xenobiotic metabolism (activation of aryl hydrocarbon receptor [AhR] and pregnane X receptor [PXR]), hormone-mediated mechanism of action (estrogenic, antiandrogenic, and glucocorticoid activities), and reactive mechanism of action (genotoxicity and oxidative stress).

Several bioassays were performed on surface water extracts of the Amsterdam region in The Netherlands [20]. Less than 50% of the polar extracts responded in the nonspecific in vivo toxicity assays. Main in vitro responses of the polar extracts were estrogenic activity, antiandrogenic activity, and genotoxicity. The nonpolar extracts showed a higher percentage of detectable responses in the nonspecific toxicity assays, especially in the Microtox and *Daphnia* assays. Very high detectable response percentages were obtained in the in vitro assays for dioxin- and polycyclic aromatic hydrocarbon (PAH)-like effects, oxidative stress (nuclear factor erythroid 2-related factor 2 [Nrf2]), pregnane X response, and estrogenic and antiandrogenic activities.

The selection of toxic endpoints for the SIMONI model is presented in Table 1. This endpoint selection aims to cover a broad range of micropollutants, multiple modes of action (nonspecific, specific, and reactive), and multiple biological levels (in vitro and in vivo). It is emphasized that this is a

selection of bioanalytical endpoints, not bioassays. Bioassays that were used in the present study can be replaced by cost-effective alternatives that measure comparable endpoints and meet the criteria mentioned (see *Requirements for a bioanalytical hazard identification*). Nonspecific in vivo assays were included in the panel because they are responsive to the broadest range of micropollutants. Because of varying sensitivity for different organisms to different kinds of pollutants, assays with organisms at different trophic levels were chosen (bacteria [Microtox], algae [Algaetoxkit], and crustaceans [Daphniatoxkit]). In addition to the bioassay battery that was applied on concentrated water samples, a field *Daphnia magna* assay that assesses mortality after 1-wk exposure to surface water was added to the nonspecific endpoints. In vitro Chemical Activated Luciferase Gene Expression (CALUX) controls for cytotoxicity were also applied as apical endpoints. A fish embryo bioassay would fit very well into this panel because it would be a whole-organism response measured in an in vitro assay. Because of the high costs, however, it did not meet the requirement of a cost-effective approach.

Specific in vitro responses were selected because these are generally much more sensitive than in vivo responses and able to detect specific activities caused by unknown mixtures of compounds with the same mechanisms of action [25]. Endocrine-disruptive effects were most frequently detected by estrogenic, antiandrogenic, and glucocorticoid activities (e.g., ER, anti-AR, and GR CALUX). Another promising endpoint, the (anti) progestagenic activity, was not included in the current selection but will be evaluated in future research. Specific

Table 1. Selection of SIMONI endpoints for effect-based hazard identification of micropollutants, with examples of targeted chemicals

Category	Endpoint (mode of action)	Targeted chemicals	
Nonspecific (in vivo)	Nonspecific toxicity zooplankton, in situ	All chemicals	
	Nonspecific toxicity zooplankton	All extracted chemicals	
	Nonspecific toxicity phytoplankton	All extracted chemicals	
	Nonspecific toxicity bacteria	All extracted chemicals	
	Nonspecific toxicity cytotoxicity	All extracted chemicals	
Specific (in vitro)	Estrogenic activity	Natural and synthetic estrogens, pseudoestrogens, bisphenol A, alkyl phenols, pharmaceuticals, pesticides	
	Antiandrogenic activity	Various pesticides, insecticides, herbicides, brominated flame retardants, (pseudo-)androgens, anabolic steroids, antibiotics, growth promoters, estrogens, PCBs	
	Glucocorticoid activity	Wide range of pharmaceuticals, corticosteroids	
	Metabolism: pregnane X receptor	Pesticides, PAHs, alkyl phenols, triazin pesticides, pharmaceuticals, PCBs, cyanotoxins	
	Metabolism: aryl hydrocarbon receptor (persistent substances)	PCDDs, PCDFs, PCBs, brominated compounds	
	Metabolism: aryl hydrocarbon receptor (degradable substances)	PAHs, nitro-PAHs, halogenated PAHs	
	Lipid metabolism (PPAR)	Organotins, perfluorinated compounds, esters, fatty acid derivatives, retinoic acid	
	Antibiotic activity	Five classes of antibiotics (amidoglycosides, macrolides and β -lactams, sulfonamides, tetracyclines and quinolones), biocides (triclosan)	
	Reactive (in vitro)	Genotoxicity	Chlorinated byproducts, aromatic amines, PAHs
		Adaptive stress response: oxidative stress	General chemical stress, reactive compounds, fungicides, insecticides, phenoles, pharmaceuticals, estrogens

PAH = polycyclic aromatic hydrocarbon; PCB = polychlorinated biphenyl; PCDD = polychlorinated dibenzodioxin; PCDF = polychlorinated dibenzofuran; PPAR = peroxisome proliferator-activated receptor.

assays for activation of AhR responded to many degradable and persistent hydrocarbons (e.g., PAH and dioxin responsive [DR] CALUX). Because lipid metabolism can be disrupted by environmentally relevant compounds, such as perfluorinated chemicals, peroxisome proliferation (e.g., peroxisome proliferator-activated receptor- γ [PPAR γ] CALUX) is considered a relevant endpoint; based on bioactivity profiling of environmental chemicals, the gamma isoform of PPAR is considered the most relevant in environmental monitoring. Oxidative stress (e.g., Nrf2 CALUX) and activation of the PXR (e.g., PXR CALUX) are relevant endpoints that respond to many Water Framework Directive priority substances [26]. Finally, the activity of antibiotics (e.g., RIKILT WaterSCAN assay) is interesting to monitor, both for toxic responses to antibiotics and biocides and for potential increase of resistant bacteria. Although CALUX bioassays were used for EBT derivation, it is important to mention that most selected in vitro endpoints can be tested by alternative technologies for specific toxicity measures (T47DKBluc, MDA-kb2, GeneBLAzer battery, MELN, MVLN, HG5LN battery, LUC battery, etc.). A disadvantage of many assays (including CALUX) is that license fees have to be paid for implementation. If relative effect potencies (REP) of key pollutants are significantly different in the alternative assays, EBT values as derived in the present study have to be adjusted. This selection of SIMONI endpoints is intended to be updated as new insights emerge. New endpoints can be added if sufficient weight of evidence is available for detecting an important group of pollutants that is not covered by the initial selection. For implementation in the tier 1 screening, it should be possible to measure this endpoint with a bioassay that meets the criteria mentioned (see *Requirements for a bioanalytical hazard identification*).

DEVELOPMENT OF ENVIRONMENTAL EBT VALUES

If bioanalytical tools are applied for water quality assessment, it should be decided which bioassay response is considered to indicate an environmental hazard. To this purpose a suite of EBT values have been derived that differentiate between 1) low risk for adverse ecological health effects if bioassay responses are below EBT, and 2) potential risk for adverse ecological health effects if bioassay responses exceed the EBT.

EBT values for nonspecific toxicity

The only bioassay that was applied on-site in nonconcentrated surface water was the in situ *Daphnia* assay. The percentage survival of 20 in situ exposed *D. magna* was monitored after 1 wk of exposure. An observed mortality of 20% was used as the EBT for potential ecological effects because this percentage is used as a blank validity criterion for the chronic *Daphnia* assay [27].

An approach to derive environmental EBT values for apical endpoints is described by Durand et al. [9]. These EBT values were based on the assumption that acute toxicity in a concentrated sample provides an indication of chronic effects in the original sample. Although it is obvious that acute-to-chronic ratios may vary for different chemicals, the applied average acute-to-chronic ratio is 10, which is based on a comparison of the species sensitivity distributions (SSDs) using hundreds of tests for different species and substances [28,29]. Chronic effects are estimated to occur at concentrations 10 times lower than the concentration where acute toxicity is observed (i.e., a detectable acute 50% effect concentration [EC50] in a 10

times concentrated water sample), and negligible effects are expected at concentrations 100 times lower than the observed acute EC50 [9]. Because most substances are not fully recovered by either solid-phase extraction methods or passive sampling, the toxicity of the original sample may be underestimated in concentrated samples. Values of EBT are therefore corrected for limited recovery by an assumed safety factor of 2 (50% recovery), as proposed by Durand et al. [9]. Taking this safety factor into account, effects measured below a relative enrichment factor (REF) of 20 are considered indicative of chronic effects, whereas REFs above 20 indicate a lower risk. If the REF is converted to toxic units (toxic unit = 1/REF), an EBT of 0.05 toxic unit is proposed for potential chronic effects.

EBT values for specific and reactive toxicity

Trigger values for in vitro bioassays should be derived by combining an approach based on toxic equivalents (TEQs) or BEQs of selected substances that trigger the bioassays, together with a benchmark approach using known chemical, toxicological, and biological data [30]. The rationale behind the benchmark approach is that bioassay responses observed at sites with a good ecological status should be considered as a background BEQ level of the bioassay. The BEQ concentration is a measure to express the effect of mixtures of unknown and potentially unidentified chemicals into the concentration of a known reference compound eliciting the same effect [2]. It is impossible to derive practically applicable trigger values that are safe for 100% of the aquatic organisms because most of them would be exceeded even at unpolluted sites [20]. Therefore, a more realistic approach was applied to derive "low-risk" EBT values. These specific EBT values do not protect all aquatic organisms against adverse effects, but exceedances indicate elevated hazards for the aquatic ecosystem attributable to micropollutants.

The novel 3-step approach for EBT derivation of all in vitro endpoints developed in the present study is schematically presented in Figure 2. In the following paragraphs the 3 steps will be explained one by one.

Toxicological database with BEQ and acute-to-chronic conversions

Because bioassays are effect-based tools that measure activities caused by a mixture of compounds, the identity of the compounds that cause the observed effect is unknown. Because using toxicological data for only one reference compound is therefore unreliable for most bioassay EBT derivations, a selection of other compounds that are able to trigger a response was made for all endpoints. Selection of compounds that have high relevance for EBT assessment was based on toxicity, REP compared to reference compounds, available toxicity data, and the range of reported concentrations in water. A complete list of REP values of the selected compounds for each bioassay is provided in Supplemental Data, Table S1. Because chemicals with very low REPs and high toxicity (low EC50) will give extremely low toxic BEQ values, a certain restriction was needed for realistic hazard identification.

Assumption 1. To restrict the REP impact, all EBT procedures were performed on chemicals with REPs > 0.001. The REPs for the CALUX bioassays were calculated from the concentration giving 10% induction with respect to the positive reference compound from different compounds (provided by BioDetection Systems). The REPs for the antibiotic activities were estimated from the detection limits for the selected




Figure 2. Schematic presentation of the approach for the derivation of effect-based trigger values. << means at least 100 times less. BEQ = bioanalytical equivalent; EBT = effect-based trigger; EC50 = 50% effect concentration; HC5 = 5% hazard concentration; LC50 = 50% lethal concentration; LOEC = lowest-observed-effect concentration; NOEC = no-observed-effect concentration; PNEC = predicted-no-effect concentration.

compounds in the RIKILT WaterSCAN (provided by RIKILT) because concentrations that cause a detectable bacterial growth inhibition should correspond to effect potencies.

Aquatic toxicity data (both specific and nonspecific) were collected for all selected substances for a wide range of aquatic species. The toxicity databases were not restricted to population-relevant endpoints (survival, growth, and reproduction) or to specific responses related to the mode of action of the bioassay. Toxic concentrations were converted to BEQs by multiplying them by the REPs of the compound. Toxicity data, such as no-observed-effect concentration (NOEC), lowest-observed-effect concentration (LOEC), EC50, or lethal concentration for 50% of the test animals (LC50), were used to build the data sets for each in vitro bioassay. Because predicted-no-effect (PNEC) concentrations are derived by using assessment factors, they cannot be directly compared with the other toxicity data. However, BEQ transformed PNECs were also used to derive safe BEQ values. The complete data set of all toxicity data that were used for the trigger value designs of all bioassays in the present study is presented in Supplemental Data, Table S2. Using both acute and chronic toxicity data causes a discrepancy in data interpretation. Because chronic toxicity data were considered most relevant for environmental risk assessment, a second assumption was made to obtain a standardized acute-to-chronic data conversion.

Assumption 2. Acute toxicity data were converted to chronic toxicity data by dividing them by an acute-to-chronic ratio of 10 [9]. Because there are no strict definitions for acute/chronic exposure times, it had to be decided to which category the obtained data were allocated, depending on the duration of the life cycles of the test species (estimations listed in Table 2).

Safe BEQ assessment

Safe BEQs should indicate no-risk levels of active compounds to the ecosystem. The lowest chronic BEQ concentrations for each toxicological endpoint (NOEC, LOEC, EC50, and LC50) or the lowest PNECs were selected and divided by an assessment factor, which ranged from 1 to 100

according to the toxic endpoint considered. Values for these assessment factors or safety factors are proposed by the authors in consultation with other Dutch experts on aquatic toxicology (see *Acknowledgment*).

Assumption 3. Assessment factors to estimate safe BEQs by extrapolation of PNECs and 4 different toxicity parameters are listed in Table 3.

The lowest of all observed chronic BEQ data, divided by its respective assessment factor, was considered to be an ecologically “safe BEQ” for the bioassay response. As an illustration of the safe approach, the collected toxicity BEQ data for compounds with a significant estrogenic activity (REP > 0.001) are presented in Figure 3. The safe BEQ determined for estrogenic activity is 0.007 ng estradiol equivalents (EEQs)/L, based on the LOEC for vitellogenin induction in rainbow trout (*Oncorhynchus mykiss*) after chronic exposure to 3.3 ng estrone/L [31]. The safe BEQ is derived after multiplication by the estrone REPs of 0.01 and division by an assessment factor of

Table 2. Criteria applied in the present study to estimate the duration of chronic exposure

Organism	Chronic exposure (days)
Protozoans	≥1
Bacteria	
Fungi	
Polyps	≥4
Algae	
Rotifers	
Crustaceans	
Insects	
Mollusks	
Worms	
Plants	≥7
Amphibians	
Fish	

Table 3. Assessment factors applied in the present study to convert toxicity data to the assumed safe bioanalytical equivalency levels

Endpoint	Assessment factor
Predicted-no-effect concentration	1
No-observed-effect concentration	1
Lowest-observed effect concentration	5
50% effect concentration	10
50% lethal concentration	100

5 for LOEC. The lowest BEQ-converted toxic concentrations found for the compounds with a significant response in each endpoint are listed in Supplemental Data, Table S3. Graphic representations of all collected toxicity data used for the EBT design for all bioassays are presented in Supplemental Data, Appendix S4.

Hazard concentration (5%) BEQ assessment

A more realistic trigger value approach (“low risk” instead of “no risk”) was based on a SSD analysis [32], conforming with Escher et al. [33], who also applied a bioanalytical SSD analysis for recycled water EBT derivation. Species sensitivity distribution curves are usually generated by fitting the distribution of average log-transformed toxicity data of several species for a single compound. The output of the SSD can be used to determine the 5th percentile hazard concentration (HC5), which represents the concentration that will negatively affect 5% of the species. In the present study a standard SSD approach was hampered because toxicity data of all compounds that trigger a bioassay response had to be included, and it is impossible to generate SSDs with data of different substances. To overcome this, all toxicity data were converted to BEQ concentrations of the reference compounds of the bioassay. This conversion allowed us to generate SSD curves with toxicity data for all species and all selected compounds. If different toxicity values were available for the same compound in the same species, the average BEQ values were used. The SSD curves were generated from EC50 BEQ values with the statistical software ETX 2.0 [34]. For DR CALUX and GR CALUX the

number of available EC50 values was insufficient to run a SSD analysis, so additional NOEC and LOEC BEQs (multiplied by factors of 10 and 2, for conversion of NOEC and LOEC to EC50, respectively) were used for these assays. The HC5 BEQ values that were determined by this approach represent the intended upper limits of the low-risk EBT values.

As an illustration of the SSD approach, the SSD curve with collected toxicity data (nanograms EEQ per liter) for estrogenic compounds is presented in Figure 4. The BEQ level that is hazardous to 5% of the organisms can be derived from the SSD curve (affected fraction 0.05). The HC5 BEQ for estrogenic compounds was estimated to be 0.5 ng EEQs/L. Species sensitivity distribution curves of all collected toxicity data that were used for the EBT design for all bioassays are presented in Supplemental Data, Appendix S5.

Background BEQ assessment

Finally, a benchmark study with available field data was crucial in determining a realistic EBT. The rationale behind this approach was that average bioassay responses observed at sites with a good ecological status should be considered as a background level of the bioassay. The reasons for these background responses and the substances causing these effects at clean sites are unknown. Nevertheless, responses below these background BEQ levels are not considered indications for serious environmental hazards from micropollutants because no ecological problems were observed at the investigated healthy ecosystems. This approach was primarily based on the results of a bioassay field survey at 8 Dutch reference sites with a good ecological status, according to Water Framework Directive quality guidelines and/or expert opinion of Dutch ecologists. The background BEQs were determined as the average bioassay response at these ecologically healthy reference sites, as described in Van der Oost et al. [20].

Evaluation of BEQ data and EBT derivation

The background BEQ data should ideally be between the safe BEQ and the HC5 BEQ. If the background BEQ was lower than the HC5 BEQ of the bioassay, then an EBT close to the




Figure 3. Bioanalytical equivalent-converted chronic toxicity data of compounds with estrogenic response. EC50 = 50% effect concentration; EEQ = estradiol equivalent; LC50 = 50% lethal concentration; LOEC = lowest-observed-effect concentration; NOEC = no-observed-effect concentration; PNEC = predicted-no-effect concentration.




Figure 4. Species sensitivity distribution of bioanalytical equivalent-converted toxicity data (nanograms estradiol equivalents per liter of 50% effect concentration) of estrogenic compounds; \log_{10} toxicity data are expressed as nanograms estradiol equivalents per liter.

HC5 BEQ was proposed. If the background BEQ was much lower than the HC5 BEQ of the bioassay, then a multiplication factor on the safe BEQ was proposed, depending on the strength of the data set used to determine safe and HC5 BEQs. The multiplication factor varies between 2 (if toxicity data only for a limited number of substances and species were collected) and 10 (if sufficient information was available). If the background BEQ was close to the HC5 BEQ, then the proposed EBT was above the background BEQ but below the upper limit of the 95% confidence interval of the HC5 BEQ. If the background BEQ was much higher than the HC5 BEQ, this could mean that the reference sites were (slightly) polluted or that the bioassays were triggered by confounding factors. Increased anti-AR responses and oxidative stress, for instance, were observed after exposure to natural organic matter and humic acids of unpolluted soil [35,36]. In these cases, trigger values based on the lower HC5 BEQ would lose their discriminative power because they are exceeded at all sites. Therefore, an EBT based on multiplication of the background BEQ (average of sites with a good ecological status) was proposed. The multiplication factor varies between 2 (if responses at polluted sites were close to the background BEQ) and 5 (if responses at polluted sites were much higher). This case was typically observed for bioassays that are responsive to a wide array of chemicals, such as anti-AR, oxidative stress, and PXR responses. The EBT values derived for these assays are not considered indicative for low micropollutant risks but are indicative for overall chemical stress.

Genotoxicity

Current guidelines for genotoxic substances that may be potentially carcinogenic assume that there is no safe level. The chances of adverse effects decrease at lower exposure levels, but a theoretical risk always remains. When interpreting bioassays for genotoxicity (e.g., Ames, umuC, and p53 CALUX) the risks are assumed to be higher than for chronic toxicity. Because most genotoxicity bioassays are not easily quantifiable, it was hard to use a BEQ approach for this endpoint. In consultation with other Dutch experts on aquatic toxicology (see *Acknowledgment*), the EBT values as derived for the nonspecific bioassays (see *EBT values for nonspecific toxicity*) are further reduced by an assessment factor of 10. The proposed EBT for genotoxicity is 0.005 genotoxic units, that is, a significant genotoxicity in a 200 times concentrated sample. This EBT was slightly exceeded

at 1 of the 8 clean sites (0.0065 toxic unit); at other reference sites no genotoxicity was observed in 400 times concentrated samples [20], supporting the choice made for 0.005 genotoxic unit.

RESULTS OF THE EBT DESIGN FOR 9 IN VITRO BIOASSAYS

The trigger value design of 8 specific CALUX bioassays, as well as a RIKILT bioassay for 5 classes of antibiotics, will be described in this section. Chemical Activated Luciferase Gene Expression assays are in vitro bioassays with genetically modified cell lines that contain specific receptor binding sites controlling expression of luciferase reporter genes. Antibiotic activities were analyzed by the RIKILT WaterSCAN bioassay using agar plates inoculated with 5 species of bacteria that are sensitive to 5 classes of antibiotics with different mechanism of actions [37]. Both specific and nonspecific effect concentrations were considered, and toxicity data were not restricted to endpoints related to the mode of action of the bioassay. Because the BEQ is the effect concentration times the REP, it is possible that the lowest converted BEQ (safe BEQ) is found for a low-REP compound with an entirely different mechanism of action from the one typical for the bioassay. Values of EBT were determined using an evaluation of safe, HC5, and background BEQ values, according to the algorithms described (see *Evaluation of BEQ data and EBT derivation*) and by expert judgment on varying multiplication factors.

Estrogenic activity with ER α CALUX

The toxicity of estrogenic compounds, including biomarker endpoints (e.g., production of vitellogenin and changes in gene expression), has been reported in many studies. Fish were found to be the most sensitive organisms to estrogenic effects. The safe BEQ was derived from a LOEC of 3.3 ng/L for vitellogenin induction in rainbow trout (*O. mykiss*) after chronic exposure to estrone [31]. After conversions, this yielded a chronic safe BEQ of 0.0066 ng EEQ/L. The SSD analysis yielded an HC5 BEQ of 0.5 ng EEQ/L. The average ER CALUX background BEQ was 0.06 ng EEQ/L. Because the background BEQ is lower than the HC5 BEQ, the proposed EBT for ER α CALUX activity is equal to the HC5 BEQ of 0.5 ng EEQ/L.

Antiandrogenic activity with anti-AR CALUX

The group of compounds that can inhibit the human AR and block its action (antiandrogenic response) is very heterogeneous, so many compounds were investigated (see Table S1, Supplemental Data). These included estrogenic compounds (e.g., 17 α -ethinylestradiol and estradiol), pesticides (e.g., alachlor, triclosan, vinclozolin), synthetic materials (e.g., bisphenol A and phthalates), and non-ionic surfactants (alkylphenols). The safe BEQ was derived from the lowest LC50 of 0.016 μ g/L after acute exposure of the copepod *Mesocyclops longisetus* to endosulfan [38]. After conversions, this yielded a chronic safe BEQ of 0.05 ng flutamine equivalents (FluEQs)/L. The SSD analysis yielded an HC5 BEQ value of 0.13 μ g FluEQ/L. The background BEQ at unpolluted sites was 4.55 μ g FluEQ/L. Because the background BEQ is 35 times higher than the HC5 BEQ, the EBT derivation is based on the background BEQ. Responses at polluted sites are clearly elevated, so the background BEQ is multiplied by a factor 5 to get an EBT for anti-AR CALUX activity of 25 μ g FluEQ/L. This EBT is not indicative for low micropollutant risks but can be considered as an indicator of overall chemical stress.

Glucocorticoid activity with GR CALUX

Because toxic effects of glucocorticoids on the aquatic community are poorly investigated, the toxicity data set is limited. Most studies were conducted on fish, whereas information for other trophic levels is scarce or nonexistent. The safe BEQ was derived from a LOEC for an increased gonadal somatic index in male fathead minnow (*Pimephales promelas*) after chronic exposure to 100 ng/L dexamethasone [39]. After conversions, this yielded a chronic safe BEQ of 20 ng dexamethasone equivalents (DxEQs)/L. This safe BEQ is 3 orders of magnitude lower than the prednisolone PNEC of 27 800 ng DxEQ/L, derived with green algae toxicity and a safety factor of 1000 [40]. The SSD analysis yielded an HC5 BEQ of 2145 ng DxEQ/L. The 8 sites with good ecological status did not show any glucocorticoid activity above the detection limit of 1.2 ng DxEQ/L, so 3 orders of magnitude lower than the HC5 BEQ. Because the background BEQ is much lower than the HC5 BEQ, EBT derivation was based on the safe BEQ. This safe BEQ was determined with a limited number of toxicity data, so a multiplication factor of 5 was used to get a proposed EBT for GR CALUX activity of 100 ng DxEQ/L.

Dioxin-like activity with DR CALUX

Most dioxins and dioxin-like compounds are poorly water-soluble. They tend to accumulate in organisms because of bioaccumulation or biomagnification. Most of the studies reported nominal exposure concentrations, which may lead to an underestimation of the risk this group of compounds pose to aquatic organisms. Generally, samples for DR CALUX are cleaned up by acid silica columns to remove easily biodegradable compounds such as PAHs. The safe BEQ was derived from the LOEC for reproduction of rare minnow (*Gobiocypris rarus*) after chronic exposure to 2 pg 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (2,3,7,8-TCDD)/L [41]. After conversions, this yielded a chronic safe BEQ of 0.4 pg TEQ/L (2,3,7,8-TCDD equivalents). The SSD analysis yielded an HC5 BEQ of 137 pg TEQ/L. The average background BEQ of the DR-CALUX response was 13 pg TEQ/L, which is 10 times lower than the HC5 BEQ. The EBT derivation is based on the observed HC5 BEQ; but because of underestimations by nominal concentrations, a lower BEQ was chosen for EBT derivation. The proposed EBT for overall dioxin-like activity is 50 pg TEQ/L, which is approximately 3 times lower than the HC5 BEQ.

PAH toxicity with PAH CALUX

As for dioxins, PAHs are lipophilic compounds that tend to accumulate in soil, organic particulate, and tissues rather than dissolving in water. Therefore, the concentration of this class of pollutants should be measured during the exposure period, but the majority of studies have reported nominal exposure concentrations. The PAH CALUX does not respond exclusively to PAHs, but dioxins, polychlorinated biphenyls, and pharmaceuticals (e.g., cyclophosphamide) also have significant REPs in the assay. With a limited exposure time of 4 h, the PAH CALUX effect will mainly be caused by carcinogenic PAHs [42]. The safe BEQ was derived from a LOEC for reproduction of rare minnow (*G. rarus*) after chronic exposure to 0.002 ng/L 2,3,7,8-TCDD [41]. After conversions, this yielded a chronic safe BEQ of 0.04 ng benzo[*a*]pyrene equivalents (BaP-EQs)/L. The SSD analysis yielded an HC5 BEQ value of 41 ng BaP-EQ/L. The average background BEQ for the PAH CALUX was 63 ng BaP-EQ/L, which was close to

the HC5 BEQ. The EBT derivation is largely based on background BEQs because PAHs are found everywhere, even at remote places, as a result of atmospheric deposition. The proposed EBT for overall PAH activity is 150 ng BaP-EQ/L, that is, more than twice the background BEQ. This value is above the HC5 BEQ but below the upper limit of its 95% confidence interval (Table 4).

Xenobiotic metabolism with PXR CALUX

The PXR CALUX is able to detect many Water Framework Directive priority compounds, including pesticides, PAHs, and alkyl phenols. The safe BEQ was derived from the LOEC for hemoglobin gene expression in *D. magna* after acute exposure to 1 ng/L of the acetyl cholinesterase inhibitor chlorpyrifos-ethyl [43]. This value was converted to the chronic safe BEQ of 0.004 ng nicardipine equivalents (NicEQs)/L. The SSD analysis yielded an HC5 BEQ of 8 ng NicEQ/L. The average background BEQ for PXR CALUX biotransformation activity was 1.71 μ g NicEQ/L, which is 200 times higher than the HC5 BEQ. The EBT derivation was based on the background BEQ. Because responses measured at polluted sites were close to the background BEQ, the proposed EBT for PXR CALUX is 3 μ g NicEQ/L, which is approximately twice the background BEQ. This EBT corresponds to 15 μ g chlorpyrifos-ethyl equivalents/L. As stated earlier, this EBT does not indicate low micropollutant risks and should be considered as an indicator of overall chemical stress.

Lipid metabolism with PPAR γ CALUX

The in vitro PPAR γ bioassays are able to detect compounds that activate the PPAR γ receptor, including several classes of aquatic contaminants, such as organotins and perfluorinated compounds. The safe BEQ was based on the lowest PNEC of 0.14 ng/L for the AhR inducer dibenzo[*a,h*]anthracene [44]. After conversion, this yielded a chronic safe BEQ of 0.00014 ng rosiglitazone equivalents (RosEQs)/L. The SSD analysis yielded an HC5 BEQ of 0.3 ng RosEQ/L. The average background BEQ of PPAR γ CALUX was 4.37 ng RosEQ/L, which is an order of magnitude above the HC5 BEQ. The EBT derivation was based on the background BEQ. Because responses measured at polluted sites were close to the background BEQ, the proposed EBT for peroxisome proliferation is 10 ng RosEQ/L, which is approximately twice the background BEQ. The EBT was just above the upper limit of the HC5 BEQ 95% confidence interval (7 ng RosEQ/L; Table 4) and the highest response observed at the clean sites (9 ng RosEQ/L). This EBT corresponds to 12.5 μ g tributyltin equivalents/L. This EBT is not indicative for low micropollutant risks but can be used to indicate overall chemical stress.

Oxidative stress with Nrf2 CALUX

It was not possible to find any aquatic toxicity data for the reference compound of the Nrf2 CALUX bioassay (i.e., curcumin), but sufficient information was available on the many other compounds that cause oxidative stress to cells and trigger a response in Nrf2 CALUX. The safe BEQ was derived from the lowest NOEC for decreased gene expression of gonadotropin-releasing hormone in transgenic Japanese medaka (*Oryzias latipes*) after acute exposure to 0.001 μ g/L estradiol [45]. After conversions, this yielded a chronic safe BEQ of 0.006 ng curcumin equivalents (CurEQs)/L. The SSD analysis yielded an HC5 BEQ of 0.034 μ g CurEQ/L. The average background BEQ of the Nrf2 CALUX response for

Table 4. Derivation of SIMONI 1.2 environmental effect-based trigger values for in vitro bioassays

Endpoint ^a	Safe BEQ endpoint/compound	HC5 BEQ (95% CI range)	Clean BEQ	EBT
Estrogenic activity	0.0066	0.52	0.06	0.5
ERa CALUX (ng EEQ/L)	LOEC/estrone	(0.019–5.4)		
Antiandrogenic	0.00005	0.13	4.6	25 ^b
anti-AR CALUX (μg Flueq/L)	LC50/endosulfan	(0.05–0.27)		
Dioxin and dioxin-like	0.4	137	13.2	50
DR CALUX (pg TEQ/L)	LOEC/2,3,7,8-TCDD	(15–736)		
Glucocorticoid	20	2145	<LOD	100
GR CALUX (ng DexEQ/L)	LOEC/dexamethasone	(116–143 11)		
PPAR _γ receptor	0.00014	0.3	4.4	10 ^b
PPAR _γ CALUX (ng RosEQ/L)	PNEC/dibenzo[<i>a,h</i>]anthracene	(0.002–6.9)		
Toxic PAHs	0.04	41	63	150
PAH CALUX (ng BaPEQ/L)	LOEC/2,3,7,8-TCDD	(2.5–254)		
Oxidative stress	0.000006	0.034	4.3	10 ^b
Nrf2 CALUX (μg CurEQ/L)	NOEC/estradiol	(0.008–0.11)		
Pregnane X receptor	0.000004	0.008	1.5	3 ^b
PXR CALUX (μg NicEQ/L)	LOEC/chlorpyrifos-ethyl	(0.002–0.024)		
Antibiotic activities				
Aminoglycosides	300	33 222	<LOD	500
RIKILT (ng NeoEQ/L)	PNEC/neomycin	(1546–219 614)		
Macrolides and β-lactams	1.8	98	<LOD	50
RIKILT (ng PenEQ/L)	EC50/tiamulin	(13–470)		
Sulfonamides	10	67 037	4.6	100
RIKILT (ng SulEQ/L)	LOEC/sulfadiazine	(24 675–148 222)		
Tetracyclines	170	27 275	<LOD	250
RIKILT (ng OxyEQ/L)	PNEC/oxytetracycline	(8292–68 544)		
Quinolones	0.53	8759	<LOD	100
RIKILT (ng FlqEQ/L)	EC50/triclosan	(2197–26 050)		

^aExpressed as equivalents of the reference compounds.

^bEBT>HC5 BEQ (+CI); No toxic relevance, indication for chemical stress.

AR = androgen receptor; BaP = benzo[*a*]pyrene; BEQ = bioanalytical equivalency; CI = confidence interval; Cur = curcumin; Dex = dexamethasone; EBT = effect-based trigger; EC50 = 50% effect concentration; E = estradiol; EQ = equivalent; ERa = estrogen receptor alpha; Flq = flumequine; Flu = flutamide; GR = glucocorticoid receptor; HC5 = 5% hazard concentration; LC50 = 50% lethal concentration; <LOD = all below limit of detection; LOEC = lowest-observed-effect concentration; Neo = neomycin; Nic = nicardipine; NOEC = no-observed-effect concentration; Nrf2 = nuclear factor erythroid 2-related factor 2; Oxy = oxytetracycline; PAH = polycyclic aromatic hydrocarbon; Pen = penicillin; PNEC = predicted-no-effect concentration; PPAR = peroxisome proliferator-activated receptor; PXR = pregnane X receptor; Ros = rosiglitazone; Sul = sulfamethoxazole; T = 2,3,7,8-TCDD; TCDD = tetrachlorodibenzo-*p*-dioxin.

oxidative stress was 4.25 μg CurEQ/L. Because of the broad range of toxic concentrations of compounds that cause oxidative stress, the low HC5 BEQ value was exceeded at all reference sites. The EBT derivation was based on the background BEQ. Because responses measured at polluted sites were close to the background BEQ, the proposed EBT for overall oxidative stress activity is 10 μg CurEQ/L, which is approximately twice the background BEQ value. This EBT corresponds to 25 μg carbendazim equivalents/L. As for the anti-AR and PXR endpoints, this EBT does not indicate low micropollutant risks and should be considered as an indicator for overall chemical stress. This mode of action is recommended to be included in any bioanalytical test battery [24].

Antibiotic activities with RIKILT WaterSCAN

According to their mode of action, antibiotics are generally divided into 5 classes: amidoglycosides (A), macrolides and β-lactams (M+B), sulfonamides (S), tetracyclines (T), and quinolones (Q). The RIKILT WaterSCAN (Screening for Antibiotics) is a semiquantitative bioassay, designed to determine activities of all 5 classes of antibiotics. Because the modes of action of these 5 antibiotic classes are different, separate EBTs were developed for each class. Because of their designed specific modes of antibacterial action, antibiotics are most hazardous to microorganisms. Relative effect potencies for the RIKILT assay could only be calculated for a limited number of antibiotic compounds.

Amidoglycosides. The safe BEQ of the A antibiotics was based on the lowest PNEC of 300 ng/L for neomycin [46]. No conversion to a chronic safe BEQ in neomycin equivalents (NeoEQs) was needed. The SSD analysis resulted in a very high HC5 of 33 μg NeoEQ/L. No detectable aminoglycoside activity (>90 ng NeoEQ/L) was found at the 8 clean reference sites. The EBT derivation was based on the safe BEQ. Because of a limited toxicity data set, the proposed EBT for amidoglycosides is 500 ng NeoEQ/L, which is approximately twice the safe BEQ.

Macrolides and β-lactams. The safe BEQ of the M+B antibiotics was based on the lowest EC50 for growth inhibition of *Microcystis aeruginosa* after chronic exposure to 18 ng/L tiamulin [47]. After conversions, a chronic safe BEQ of 1.8 ng penicillin equivalents (PenEQs)/L was derived. The SSD analysis yielded an HC5 BEQ of 98 ng PenEQ/L. No detectable activities of macrolides and β-lactams (>1.4 ng PenEQ/L) were found at the 8 clean reference sites. Large differences were observed between safe BEQ and HC5 BEQ values. The EBT derivation was based on the HC5 BEQ. The proposed EBT for M+B antibiotics is 50 ng PenEQ/L, that is, half of the HC5 BEQ.

Sulfonamides. The safe BEQ for S antibiotics was based on the lowest LOEC found for growth inhibition of zebra fish embryos (*Danio rerio*) after acute exposure to 1000 ng/L sulfadiazine [48]. After conversions, this yielded a chronic safe BEQ of 10 ng sulfamethoxazole equivalents (SulEQs)/L. The SSD analysis yielded a very high HC5 BEQ of 67 μg SulEQ/L.

A sulfonamide activity of 37 ng Suleq/L was found at one of the reference sites, whereas no activity (>2 ng Suleq/L) was found at the other clean sites. The background BEQ was 4.6 ng Suleq/L. The EBT derivation was based on the safe BEQ. Because a sufficient toxicity data set was collected, the proposed EBT for sulfonamides is 100 ng Suleq/L, that is, 10 times the safe BEQ.

Tetracyclines. The safe BEQ of the T antibiotics was based on the lowest PNEC of 170 ng/L derived for oxytetracycline [46]. This PNEC was based on an acute LC50 of the algae *Selenastrum capricornutum* after exposure to 170 µg/L oxytetracycline [49], using a safety factor of 1000. No conversion to chronic safe BEQ in oxytetracycline equivalents (OxyEQs) was needed. The SSD analysis yielded a high HC5 BEQ of 27 µg OxyEQ/L. No detectable tetracycline activity (>22 ng OxyEQ/L) was found at the 8 clean reference sites. The EBT derivation was based on the safe BEQ of a limited data set. The proposed EBT for tetracyclines is 250 ng Suleq/L, which is approximately twice the safe BEQ.

Quinolones. Although it is not used as an antibiotic, toxicity data for triclosan (an antibacterial and antifungal substance) was also investigated because it gives a clear response in the quinolones bioassay. The safe BEQ was based on the lowest EC50 for growth inhibition of the algae *Pseudokirchneriella subcapitata* after acute exposure to 530 ng/L triclosan [50]. After conversions, this yielded a chronic safe BEQ of 0.53 ng flumequine equivalents (FlqEQs)/L. The SSD analysis yielded an HC5 BEQ of 8.8 µg FlqEQ/L. No detectable quinolone activity (>44 ng FlqEQ/L) was found at any of the 8 clean reference sites. Considering the high HC5 BEQ and the low safe BEQ because of triclosan, the proposed EBT for this group of antibiotics is 100 ng/L, which is 2 orders of magnitude above the safe BEQ. An exception of the algorithm that EBT should be approximately 5 times the safe BEQ was made in the present study because the safe BEQ was attributable to only one compound (triclosan), whereas the proposed EBT is still much lower than the HC5 BEQ.

Overall evaluation

The complete set of toxicity data used for the EBT design with literature references, lowest BEQ values, graphs with toxicity data, and SSD-curves for all endpoints is presented in Supplemental Data, Appendices S2–S5. All relevant data for the EBT design, such as lowest observed BEQ values (safe BEQ), hazardous BEQ for 5% of the water organisms (HC5 BEQ with 95% confidence intervals), average background BEQ (observed at sites with good ecological status), and the proposed EBT are summarized in Table 4. The extensive toxicity database that was created for the trigger value development can be applied to derive EBT values for similar in vitro bioassays detecting the same modes of action, possibly after adjusting the REPs of the compounds. The derived EBTs of Table 4 can be applied for bioassays with similar endpoints if REP values are close to those of CALUX and RIKILT assays (Supplemental Data, Table S1).

SIMONI MODEL FOR OVERALL MICROPOLLUTANT RISKS

A final step in the SIMONI hazard-assessment strategy was the development of a model that translates all bioassay responses into potential ecological hazards. This model, SIMONI 1.2, aims to derive an indication of the environmental risks of micropollutants based on the results of individual bioassays, as well as the total battery of bioassays (earlier Ver 1.0 and 1.1 were not published). To obtain a broad and sensitive identification of the chemical risks, the model is based on a

panel of in vivo and in vitro bioassays that are performed on concentrated water extracts. The ecological hazards can be predicted when a substantial subset (or all) of the bioassays shown in Table 1 are applied. It is important to realize that the reliability of the outcome will be highest if the entire bioassay battery is applied.

A “no-risk” approach would require the use of the safe BEQ values for risk assessment. A more realistic approach for the tier 1 hazard identification is to apply the proposed “low-risk” EBT. If only slight exceedances of the EBT are observed in 1 or 2 bioassays, this does not necessarily indicate an increased ecological risk. However, with slight EBT excesses in many bioassays or large EBT exceedances, the hazard indications become stronger. Therefore, the model uses a simple formula that aims to quantify the combined ecological hazards attributable to micropollutants by integrating all individual bioassay responses. All bioassays have been given a weight factor: 2 for apical toxicity endpoints (responsive to all micropollutants) and 1 for specific and reactive toxicity endpoints (responsive to specific groups of micropollutants). These weight factors were introduced to get a better balance between the impact of in vivo and in vitro bioassays. The total weight of the SIMONI bioassay selection is 20, that is, 10 for in vivo and 10 for in vitro bioassays. If the endpoint selection is changed in the future, weight factors have to be adjusted to restore the in vivo versus in vitro balance. As a requirement for a reliable result it is assumed that the total weight of the applied bioassays must be at least 10, and both in vivo and in vitro endpoints should be used. The SIMONI model divides all bioassay responses (toxic unit or BEQ) by their associated EBT and multiplies them by this weight factor. Average relative responses are calculated for the 5 antibiotic assays and for the genotoxicity test with and without metabolic activation. Results are summed for all applied bioassays and divided by a percentage of the total weight of all applied bioassays.

$$\text{SIMONI score} = \frac{\sum_{i=1}^n \left(\frac{\text{bioassay response}_i}{\text{EBT}_i} \right) \times \text{weight}_i}{0.5 \times \text{total bioassay weight}}$$

A total SIMONI score above 1 indicates a potential ecological risk as a result of elevated concentrations of micropollutants. In this SIMONI score model, it is assumed that an increased hazard for the ecosystem occurs when the responses of all bioassays are, on average, more than 50% of the proposed trigger values, that is, the factor 0.5 in the equation. The provisional weight factors and EBT percentage threshold are proposed by the authors in consultation with other Dutch experts on aquatic toxicology (see *Acknowledgment*). The feasibility of the model with these assumptions will be demonstrated in a follow-up study [20].

DISCUSSION

The SIMONI strategy aims to estimate broad-spectrum chemical risks for invertebrate and vertebrate aquatic organisms, by combining validated in vivo and in vitro endpoints. Three aspects are of main importance for successful implementation of the SIMONI system in regular monitoring programs: 1) the choice of the most relevant bioanalytical endpoints, 2) the design of acceptable bioanalytical trigger values for potential risks, and 3) a comparison with regular monitoring strategies that are mainly based on chemical analyses. The first 2 aspects will be discussed in the present study, the third aspect is

discussed in a second study on a field feasibility survey of the SIMONI strategy [20].

Selection of bioanalytical endpoints

For a complete assessment of toxic hazards it is relevant to select a panel of *in vivo* and *in vitro* bioassays representing all mechanism-of-action groups. The suggested SIMONI endpoints (Table 1) combine *in vivo* assays that are most relevant for invertebrate organisms and *in vitro* bioassays with defined adverse outcome pathways that are most relevant for vertebrate organisms. The selected *in vivo* test organisms are ecologically relevant; bacteria are important for nutrient cycling, algae are primary producers, and the invertebrates are primary consumers of algae, as well as food for small fish. Selected *in vitro* endpoints, such as effects on endocrine activities, are known indicators for impaired reproduction (fecundity and spawning behavior); and AhR induction and oxidative stress are related to tissue damage and fish mortality. The initial selection may be adjusted in the future if new knowledge becomes available, new assays are developed, or expensive bioassays are transformed into more cost-effective formats. The impact of other relevant toxicity endpoints, such as immunotoxicity and neurotoxicity that act by many different pathways, is hard to estimate with one simple bioassay [2]. Reproductive toxicity and developmental toxicity are 2 classical examples for which there is currently no comprehensive *in vitro* model [2]. In this respect, fish embryo tests would be a very relevant bioassay for higher-trophic level organisms, but current costs of these assays are too high. The fish embryo test will, however, be implemented in the SIMONI tier 2 risk-assessment phase. Moreover, it is possible to use multiplex polymerase chain reaction to measure fish embryo biomarkers on more than 40 gene expressions (J. Legradi, Free University, Amsterdam, The Netherlands, unpublished data). If a large part of the *in vitro* bioassay screening could be replaced by fish embryo biomarker responses, this would make the assay very interesting for the tier 1 assessment.

EBT values

Routine monitoring with bioassays is still hampered by the lack of reliable interpretation guidelines. The challenge is therefore to provide EBT values that allow regulators to link the test results to possible adverse effects on environmental or human health [2]. The fact that in many studies only a small percentage of the effect observed in a bioassay can be explained by known (chemically determined) substances makes it imperative to derive such EBT values [11,12]. The present study describes a format to derive trigger values for bioassays to distinguish between low and increased ecological risks as a result of organic micropollutants. We realize, however, that it is impossible to make a solid distinction between a “good” and a “bad” chemical status if the identity of compounds causing the bioassay responses is unknown. Therefore, chemical identification on the tier 1 hot spots is performed in tier 2. The discriminative power of the first tier bioanalytical screening should be high enough to indicate potential hazards that have to be verified in a second tier (Figure 1).

Our intention to derive EBTs indicative for low toxic risks between the safe BEQ and the HC5 BEQ is best met for those bioassays that are triggered by a limited amount of specific substances. Increased estrogenic activity, for instance, is mainly the result of exposure to 4 substances: estrone, 17 β -estradiol, estriol, and 17 α -ethinylestradiol [13]. It is hard to derive toxicity-based EBTs for the more “promiscuous” endpoints, such as antiandrogenic activity, pregnane X metabolism, and

oxidative stress, because of the immense number of substances that are able to trigger these bioassays. These endpoints are, however, relevant to be taken up in the SIMONI panel as sensitive indicators of the overall chemical or toxic pressure. The EBT values for these bioassays were mainly based on the benchmark approach using bioassay responses at sites with good ecological status.

A similar bioanalytical strategy as we propose for water quality is already in use for determination of food quality in Europe. Bioassays are being used for high-throughput screening of large amounts of food and feed samples. Only samples with bioassay responses exceeding effect-based action levels (BEQ trigger values) need to be confirmed by chemical analysis. A decision limit was derived for bioanalyses of dioxins, based on a gas chromatography mass spectrometry confirmation and the condition that the chance of false negatives should be <5%. This bioanalytical procedure is laid down in European Union legislation [51]. In addition, for hormones and antibiotics established methods are available to regularly apply bioassays in food quality control (e.g., Bovee et al. [52] and Gizzi et al. [53]).

A limited number of EBT values for water quality assessment can be found in the literature. Most of these, however, are derived for human health risk from drinking water and cannot be compared with the EBT values developed in the present study. Brand et al. [54] derived some human health EBTs for hormonal activity in drinking water. These EBTs were derived for *in vitro* CALUX assays on estrogenic, androgenic, progestagenic, and glucocorticoid activities. An EBT approach by Tang et al. [12] for cytotoxicity measured by bioluminescence inhibition in *Vibrio fischeri* was based on combined effects of mixtures of regulated chemicals, according to the concentration addition model and a quantitative structure-activity relationship model approach for nonspecific baseline toxicity. Escher et al. [11] used a similar strategy to derive an EBT for the oxidative stress response pathway with the AREc32 cell line. Escher et al. [33] proposed a statistical EBT design with a read-across from existing water quality guidelines. This EBT derivation was based on matching the effect concentrations with chemical guidelines and filtering out appropriate chemicals that are responsive in a given bioassay at concentrations near the guideline values. Statistical methods were used to derive specific EBT-BEQ concentrations for 11 bioassays that target receptor-mediated toxicity.

Most comparable to the EBT values of the present study are the environmental thresholds determined by Jarošová et al. [55], Kunz et al. [56], and Leusch et al. [57]. Jarošová et al. [55] derived estrogenic equivalents that are safe regarding major steroid estrogens in municipal WWTP effluents. This EBT was based on the assumption that only steroid estrogens are responsible for *in vitro* estrogenicity, using estrogenic REPs, *in vivo* estrogenic PNECs, and relative contributions to overall estrogenicity of WWTP effluents. Derived EEQs that are safe regarding major steroid estrogens for ER CALUX varied from 0.2 to 0.4 ng EEQs/L for long-term exposure. The European Commission proposed an annual average environmental quality standard of 0.4 ng/L for 17 β -estradiol, based on toxicity SSDs for population-relevant effects. Kunz et al. [56] proposed to use this annual average environmental quality standard as a trigger value for overall estrogenic activity. Leusch et al. [57] proposed threshold levels above which *in vitro* responses were expected to lead to adverse effects in exposed organisms. The threshold for estrogenic activity (0.07 ng EEQ/L) was based on the lowest PNEC \times REP_{E-SCREEN} (relative effect potency in E-Screen

bioassay) value of 4 estrogens (lowest value for estrone). If we applied this approach to the ERa CALUX REPs, this would lead to an EBT of 0.06 ng EEQ/L, which is almost 10 times lower than the “realistic” EBT of 0.5 ng EEQ/L we propose. The EBT derived in the present study was, however, close to the trigger values proposed by Jarošová et al. and Kunz et al.

Leusch et al. [57] also proposed threshold levels for nonspecific toxicity determined with Microtox (1 toxic unit) and genotoxicity determined with the umuC assay (1 genotoxic unit), which were less conservative than those proposed in the present study (0.05 toxic unit and 0.005 genotoxic unit, respectively). These 2 trigger values of Leusch et al. were, however, based on acute toxic effects, whereas those derived in the present study were based on chronic toxicity estimations, with an additional safety factor for genotoxic effects.

Hamers et al. [58] obtained toxicity pathway profiles as toxicological “fingerprints,” using a bioassay battery with different modes of action (genotoxicity, [anti]estrogenic activity, thyroid activity, dioxin-like activity, and nonspecific cytotoxicity). Three potential approaches were described: 1) toxicity profiles were translated into hazard profiles, indicating the relative distance to the desired quality status for each toxic mode of action; 2) toxicity profiles were translated into ecological risk profiles, that is, the ratio between the bioassay responses and those considered safe for environmental health; and 3) use of toxicity and hazard profiles to select hot spot samples for further in-depth effect-directed analysis. A combination of the second and third approaches is most similar to the SIMONI strategy proposed in the present study.

Limitations of the bioanalytical approach

Both overestimations and underestimations of the overall toxic impact of the mixture can be made with the BEQ concept. Moreover, it is hard or impossible to predict adverse in vivo effects with in vitro responses if the impact of absorption, distribution, metabolism, and excretion is unknown. It must be stressed that an EBT value should not be used as a stand-alone value, but a battery of biological endpoints should be combined.

There are several assumptions and limitations connected to the EBT design of the present study that may be optimized in future research. Three assumptions had to be made in the EBT design: acute-to-chronic toxicity ratio and duration of chronic studies, assessment factors to convert toxicity data to safe values, and the REP-based selection of compounds. Limitations with regard to the toxicological database are that only chemicals with a known REP in the bioassay could be considered for the BEQ approach. Reliable REP values are essential for a good conversion of substance concentrations to BEQ, but it was not always possible to assess REPs with EC50 data because of the cytotoxicity in the bioassay at higher concentrations. The CALUX REPs were estimated with 10% of the maximum bioassay response of the reference compound (PC10 values). Future research should be aimed at refining bioassay REP values of the most relevant compounds. The publically available REP database from the Toxicity ForeCaster (ToxCast) project on the US Environmental Protection Agency website could be very useful in this matter. An additional limitation is that there may be large deviations between in vivo and in vitro REP values (e.g., Van Ede et al. [59]).

For some of the bioanalytical endpoints, background BEQs are higher than HC5 BEQ, which resulted in EBTs that are not indicative for low micropollutant risks. Because it might be possible that the background BEQs are different in other regions, it is advised to perform additional background studies on local reference sites.

For several compounds it was hard or impossible to find data on aquatic toxicity, which resulted in limited data sets for some of the bioassays (e.g., GR CALUX and some antibiotics). All toxicity data used for EBT derivation were concentrations in the water that indicated a certain effect to aquatic organisms or safe PNEC values. Laboratory exposures with single compounds, however, differ from the situation in the natural environment as a result of varying physicochemical conditions, such as pH, temperature, and light exposure. In addition, the majority of the studies investigating the toxicity of highly hydrophobic compounds reported nominal concentrations, which may give underestimations of toxicity. This is, however, also the case when bioassays are applied on environmental extracts.

SIMONI model for integrated risk assessment

The holistic approach of effect-based toxicity monitoring has many advantages over the substance-based chemical monitoring of a limited number of target compounds. However, the effect-based methods are hardly used in regular monitoring programs because of the lack of established guidelines for interpretation of the data. The integrated monitoring strategy presented in the present study aims to provide these guidelines.

The in vivo and in vitro bioassay panel that is proposed for hazard identification in water concentrates, together with a field-exposed in situ *Daphnia* assay, should be able to distinguish the chemical quality of sites indicating low or increased ecological hazards. Despite its limitations and uncertainties, the authors believe that the approach proposed in the present study constitutes a better alternative to the current European Union's Water Framework Directive monitoring, that is, chemical analyses of a limited amount of priority substances, whereas the potential impact of more than 100 000 compounds remains unknown. The alternative strategy proposed in the present study represents one of the first attempts to connect multiple bioassay responses with potential negative effects for aquatic organisms, by using EBT values for low ecological risks.

In comparison with regular monitoring programs, the SIMONI strategy can be more cost-effective. A Water Framework Directive chemical surveillance monitoring consists of 12 monthly grab samples that are analyzed for 45 priority substances. A Water Framework Directive chemical campaign for one water body currently (2016) costs approximately €40 000 in The Netherlands. The suggested battery of bioassays in Table 1 (total ~€2000) is less expensive than chemical analysis of 45 Water Framework Directive priority pollutants (total ~€3000). The SIMONI strategy combines passive sampling with bioanalytical methods. Because several weeks of passive sampling will obtain time-weighted average concentrations of bioavailable micropollutants, fewer sampling campaigns are assumed necessary for representative water quality assessment. When the entire panel of bioassays is analyzed, 3 seasonal SIMONI campaigns in one water body would cost approximately €10 000. A more expensive second tier risk assessment, using effect-directed analyses [60], should only be performed at the sites with elevated ecological risks.

The SIMONI model estimates an integrated quantitative measure for potential ecological risks from micropollutants. The outcome of the model indicates which sites are hot spots, relevant for additional chemical-toxicological research. Moreover, if specific or reactive activities are above the EBT value, the model will indicate which class of chemicals may cause the main problem for aquatic organisms. The associated follow-up for risk management is the “toxicity traffic light” system: Green,

SIMONI score <1: low hazard, so no action is required. Orange, SIMONI score ≥ 1 : potential risk, so tier 2 research should be performed. Red, SIMONI score $\geq 1 +$ tier 2 identification toxic substances: high risk, so mitigation measures are needed. SIMONI scores ≥ 2 (responses of all bioassays are, on average, more than 100% of the EBT) are also indicative for high risks, but most appropriate measures can be taken if the (class of) chemicals causing the effects is known.

If the information from the SIMONI hazard assessment is combined with influences of other ecological key factors, a tailor-made plan can be designed for a tier 2 ecological risk assessment. The next steps to make the proposed concept attractive for risk assessors would be to gain experience on the applicability to case studies and to evaluate its robustness for practical use. Field validation studies with this strategy are described in the second paper of the SIMONI series [20]. Because of its low costs and high relevance, this model has the potential to become the first bioanalytical strategy to be applied in regular monitoring of surface water quality. Most Dutch water authorities started feasibility studies with the SIMONI strategy in 2016.

Supplemental Data—The Supplemental Data are available on the Wiley Online Library at DOI: 10.1002/etc.3836.

Acknowledgment—We thank the Waternet Research & Innovation group for funding the main part of the project. L. Moria (Waternet Institute for the Urban Water Cycle, The Netherlands) is acknowledged for her useful input on practical applications of the strategy. We thank the STOWA Foundation for Applied Water Research for funding a study on EBT development and finalizing the SIMONI 1.2 modeling. We thank D. de Zwart and L. Postuma (both National Institute for Public Health and the Environment, The Netherlands), J. Postma (Ecofide Environmental and Ecological Consultancy, The Netherlands), and L. Osté (Deltares, Dutch Institute for Applied Water Research, The Netherlands) for consultancy on the assumptions for trigger value derivation. We thank T. Williams (University of Birmingham, UK) for critically reading and adjusting the manuscript. We especially thank B. Escher (Centre for Environmental Research, University of Tübingen, Germany, and University of Queensland, Australia) for her extensive review and useful comments that significantly improved the quality of the manuscript.

Data Availability—Data, associated metadata, and calculation tools are available on request (ron.van.der.oost@waternet.nl).

REFERENCES

- European Commission. 2012. Proposal for a directive of the European Parliament and of the Council amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy. COM/2011/0876 final. [cited 2017 May 14]. Available from: <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:52011PC0876&rid=3>
- Escher BI, Leusch FDL. 2012. *Bioanalytical Tools in Water Quality Assessment*. IWA, London, UK.
- Depledge MH, Fossi MC. 1994. The role of biomarkers in environmental assessment (2). *Ecotoxicology* 3:161–172.
- Silva E, Rajapakse N, Kortenkamp A. 2002. Something from “nothing”—Eight weak estrogenic chemicals combined at concentrations below NOECs produce significant mixture effects. *Environ Sci Technol* 36:1751–1756.
- Escher BI, Fenner K. 2011. Recent advances in the environmental risk assessment of transformation products. *Environ Sci Technol* 45:3835–3847.
- Van der Oost R, Beyer J, Vermeulen NPE. 2003. Fish bioaccumulation and biomarkers in environmental risk assessment: A review. *Environ Toxicol Phar* 13:57–149.
- Stegeman JJ, Brouwer M, Di Giulio RT, Förlin L, Fowler BA, Sanders BM, Van Veld PA. 1992. Molecular responses to environmental contamination: Enzyme and protein systems as indicators of chemical exposure and effect. In Huggett RJ, Kimerly RA, Mehrle PM Jr, Bergman HL, eds., *Biomarkers: Biochemical, Physiological and Histological Markers of Anthropogenic Stress*. Lewis, Chelsea, MI, USA, pp 235–335.
- Castaño A, Bols N, Braunbeck T, Dierickx P, Halder M, Isomaa B, Kawahara K, Lee LEJ, Mothersill C, Pärt P, Repetto G, Sintes JR, Rufli H, Smith R, Wood C, Segner H. 2003. The use of fish cells in ecotoxicology. The report and recommendations of ECVAM Workshop 47. *ATLA-Altern Lab Anim* 31:317–351.
- Durand AM, Rotteveel S, Collombon MT, Van der Grinten E, Maas JL, Verweij W. 2009. Toxicity measurements in concentrated water samples. Evaluation and validation. Report 607013010/2009. RIVM (National Institute for Public Health and the Environment), Bilthoven, The Netherlands. [cited 2017 May 14]. Available from: http://www.rivm.nl/Documenten_en_publicaties/Wetenschappelijk/Rapporten/2009/mei/Toxicity_measurements_in_concentrated_water_samples_Evaluation_and_validation?s_p=cml2bXE9ZmFsc2U7c2VhcmNoYmFzZT01OTQwMDtyaXZtcTImYWxzZT0=&pagenr=5941
- Escher BI, Hermens JLM. 2002. Modes of action in ecotoxicology: Their role in body burdens, species sensitivity, QSARs, and mixture effects. *Environ Sci Technol* 36:4201–4217.
- Escher BI, Van Daele C, Dutt M, Tang JYM, Altenburger R. 2013. Most oxidative stress response in water samples comes from unknown chemicals: The need for effect-based water quality trigger values. *Environ Sci Technol* 47:7002–7011.
- Tang JYM, McCarty S, Glenn E, Neale PA, Warne MSJ, Escher BI. 2013. Mixture effects of organic micropollutants present in water: Towards the development of effect-based water quality trigger values for baseline toxicity. *Water Res* 47:3300–3314.
- Aerni HR, Kobler B, Rutishauser BV, Wettstein FE, Fischer R, Giger W. 2004. Combined biological and chemical assessment of estrogenic activities in wastewater treatment plant effluents. *Analytical and Bioanalytical Chemistry* 378:688–696.
- Collins FS, Gray GM, Bucher JR. 2008. Transforming environmental health protection. *Science* 319:906–907.
- Ankley GT, Bennet RS, Erickson RJ, Hoff DJ, Hornung MW, Johnson RD, Mount dioux responsive, Nickols DL, Russom CL, Schmieder PK, Serrano JA, Tietge JE, Villeneuve DL. 2010. Adverse outcome pathways: A conceptual framework to support ecotoxicology research and risk assessment. *Environ Toxicol Chem* 29:730–741.
- Hoogenboom LAP, Haedrich J, Eppe G, Goeyens L, Elskens M, Malagocki P, Scippo M, Vanderperren H, Windal I, Kotz A, Denisov M, Malisch R. 2010. Revised EU-criteria for applying bioanalytical methods for screening of feed and food for dioxins and dioxin-like PCBs. *Organohalogen Compounds* 72:1800–1905.
- Smedes F. 2007. Monitoring of chlorinated biphenyls and polycyclic aromatic hydrocarbons by passive sampling in concert with deployed mussels. In Greenwood R, Mills G, Vrana B, eds, *Passive Sampling Techniques in Environmental Monitoring*, Vol 48—Comprehensive Analytical Chemistry. Elsevier, Amsterdam, The Netherlands; Oxford, UK, pp 407–448.
- Li J-Y, Tang JYM, Jin L, Escher BI. 2013. Understanding bioavailability and toxicity of sediment-associated contaminants by combining passive sampling with in vitro bioassays in an urban river catchment. *Environ Toxicol Chem* 32:2888–2896.
- Jahnke A, Mayer P, Schäfer S, Witt G, Haase N, Escher BI. 2016. Strategies for transferring mixtures of organic contaminants from aquatic environments into bioassays. *Environ Sci Technol* 50:5424–5431.
- van der Oost R, Sileno G, Janse T, Nguyen MT, Besselink H, Brouwer A. 2017. SIMONI (Smart Integrated Monitoring) as a novel bioanalytical strategy for water quality assessment: Part II—Field feasibility survey. *Environ Toxicol Chem* 36:xx–xx, (this issue), in press, DOI: 10.1002/etc.3837.
- Willemsen A, Vaal MA, De Zwart D. 1995. Microbiotests as tools for environmental monitoring. Report 607042005. RIVM (National Institute for Public Health and the Environment), Bilthoven, The Netherlands.
- Van der Linden S, Heringa M, Man H-Y, Sonneveld E, Puijker LM, Brouwer A, Van der Burg B. 2008. Detection of multiple hormonal activities in wastewater effluents and surface water, using a panel of steroid receptor CALUX bioassays. *Environ Sci Technol* 42:5814–5820.
- Macova M, Toze S, Hodggers L, Mueller JF, Bartkow M, Escher BI. 2011. Bioanalytical tools for the evaluation of organic micropollutants during sewage treatment, water recycling and drinking water generation. *Water Res* 45:4238–4247.
- Escher BI, Allinson M, Altenburger R, Bain PA, Balaguer P, Busch W, Crago J, Denslow NA, Dopp E, Hilscherova K, Humpage AR, Kumar A, Grimaldi M, Jayasinghe BS, Jarosova B, Jia A, Makarov S, Maruya

- KA, Medvedev A, Mehinto AC, Mendez JE, Poulsen A, Prochazka E, Richard J, Schifferli A, Schlenk D, Scholz S, Shiraishi F, Snyder S, Su G, Tang JYM, Van der Burg B, Van der Linden SC, Werner I, Westerheide SD, Wong CKC, Yang M, Yeung BHY, Zhang X, Leusch FDL. 2014. Benchmarking organic micropollutants in wastewater, recycled water and drinking water with in vitro bioassays. *Environ Sci Technol* 48:1940–1956.
25. Sonneveld E, Jansen HJ, Riteco JAC, Brouwer A, Van der Burg B. 2005. Development of androgen- and estrogen-responsive bioassays, members of a panel of human cell line–based highly selective steroid responsive bioassays. *Toxicol Sci* 83:136–148.
 26. Neale PA, Ait-Aissa S, Brack W, Creusot N, Denison MS, Deutschmann B, Hilscherová K, Hollert H, Krauss M, Novák J, Schulze T, Seiler T-B, Serra H, Shao Y, Escher BI. 2015. Linking in vitro effects and detected organic micropollutants in surface water using mixture-toxicity modeling. *Environ Sci Technol* 49:14614–14624.
 27. Organisation for Economic Co-operation and Development. 2008. Test No. 211: *Daphnia magna* reproduction test. *OECD Guidelines for the Testing of Chemicals*. Paris, France.
 28. De Zwart D, Sterkenburg A. 2002. Toxicity-based assessment of water quality. In Posthuma L, Suter GW II, Traas TP, eds. *Species Sensitivity Distributions in Ecotoxicology*. Lewis, Boca Raton, FL, USA, pp 383–402.
 29. Raimondo S, Montague BJ, Barron MG. 2007. Determinants of variability in acute to chronic ratios for aquatic invertebrates and fish. *Environ Toxicol Chem* 26:2019–2023.
 30. Maas JL, Van de Plassche E, Straetmans A, Vethaak AD, Belfroid AC. 2003. Normstelling voor bioassays. Uitwerking voor oppervlaktewater en waterbodem. Report 2003.005. Rijksinstituut voor Integraal Zoetwaterbeheer en Afvalwaterbehandeling, Lelystad, The Netherlands.
 31. Thorpe KL, Cummings RI, Hutchinson TH, Scholze M, Brighty G, Sumpter JP, Tyler CR. 2003. Relative potencies and combination effects of steroidal estrogens in fish. *Environ Sci Technol* 37:1142–1149.
 32. Posthuma L, Suter GW II, Traas TP. 2002. *Species Sensitivity Distributions in Ecotoxicology*. Lewis, Boca Raton, FL, USA.
 33. Escher BI, Neale PA, Leusch FDL. 2015. Effect-based trigger values for in vitro bioassays: Reading across from existing water quality guideline values. *Water Res* 81:137–148.
 34. Van Vlaardingen PLA, Traas TP, Wintersen AM, Aldenberg T. 2004. ETX 2.0—A Program to Calculate Hazardous Concentrations and Fraction Affected, Based on Normally Distributed Toxicity Data. Report 601501028/2004. RIVM (National Institute for Public Health and the Environment), Bilthoven, The Netherlands.
 35. Bittner M, Saul N, Steinberg CEW. 2012. Antiandrogenic activity of humic substances. *Sci Total Environ* 432:93–96.
 36. Saebelfeld M, Minguez L, Griebel J, Gessner MO, Wolinska J. 2016. Humic dissolved organic carbon drives oxidative stress and severe fitness impairments in *Daphnia*. *Aquat Toxicol* 182:31–38.
 37. Pikkemaat MG, Dijk SO, Schouten J, Rapallini M, van Egmond HJ. 2008. A new microbial screening method for the detection of antimicrobial residues in slaughter animals: The Nouws antibiotic test (NAT-screening). *Food Control* 19:781–789.
 38. Gutierrez MF, Gagneten AM, Paggi JC. 2013. Acute and behavioral sensitivity of *Mesocyclops longisetus* to atrazine and endosulfan formulations under predation pressure. *Water Air Soil Pollut* 224:1375–1383.
 39. LaLone CA, Villeneuve DL, Olmstead AW, Medlock EK, Kahl MD, Jensen KM, Durhan EJ, Makyne EA, Blank CA. 2012. Effects of a glucocorticoid receptor agonist, dexamethasone, on fathead minnow reproduction, growth, and development. *Environ Toxicol Chem* 31:611–622.
 40. Escher BI, Baumgartner B, Koller M, Treyer K, Lienert J, McAndell CS. 2011. Environmental toxicology and risk assessment of pharmaceuticals from hospital wastewaters. *Water Res* 45:75–92.
 41. Wu WZ, Li W, Xu Y, Wang JW. 2001. Long-term toxic impact of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin on the reproduction, sexual differentiation, and development of different life stages of *Gobiocypris rarus* and *Daphnia magna*. *Ecotox Environ Safe* 48:293–300.
 42. Pieterse B, Felzel E, Winter R, Van der Burg B, Brouwer A. 2013. PAH-CALUX, an optimized bioassay for AhR-mediated hazard identification of polycyclic aromatic hydrocarbons (PAHs) as individual compounds and in complex mixtures. *Environ Sci Technol* 47:11651–11659.
 43. Ha MH, Choi J. 2009. Effects of environmental contaminants on hemoglobin gene expression in *Daphnia magna*: A potential biomarker for freshwater quality monitoring. *Arch Environ Con Tox* 57:330–337.
 44. OSPAR. 2014. Establishment of a list of predicted no effect concentrations (PNECs) for naturally occurring substances in produced water. OSPAR Agreement 2014-05. London, UK.
 45. Lee W, Kang CW, Su CK, Okubo K, Nagahama Y. 2012. Screening estrogenic activity of environmental contaminants and water samples using a transgenic medaka embryo bioassay. *Chemosphere* 88:945–952.
 46. Park S, Choi K. 2008. Hazard assessment of commonly used agricultural antibiotics on aquatic ecosystems. *Ecotoxicology* 17: 526–538.
 47. Halling-Sorensen B. 2000. Algal toxicity of antibacterial agents used in intensive farming. *Chemosphere* 40:731–739.
 48. Lin T, Chen YQ, Chen W. 2013. Impact of toxicological properties of sulfonamides on the growth of zebrafish embryos in the water. *Environ Toxicol Phar* 36:1068–1076.
 49. Nunes B, Carvalho F, Guilhermino L. 2005. Acute toxicity of widely used pharmaceuticals in aquatic species: *Gambusia holbrooki*, *Artemia parthenogenetica* and *Tetraselmis chuii*. *Ecotox Environ Safe* 61: 413–419.
 50. Yang LH, Ying GG, Su HC, Stauber JL, Adams MS, Binet MT. 2008. Growth-inhibiting effects of 12 antibacterial agents and their mixtures on the freshwater microalga *Pseudokirchneriella subcapitata*. *Environ Toxicol Chem* 27:1201–1208.
 51. European Union. 2012. Commission Regulation (EU) No 252/2012. Laying down methods of sampling and analysis for the official control of levels of dioxins, dioxin-like PCBs and non-dioxin-like PCBs in certain foodstuffs and repealing Regulation (EC) No 1883/2006. *Official J Eur Union* L84:1–22.
 52. Bovee TFH, Bor G, Heskamp HH, Hoogenboom LAP, Nielen MWF. 2006. Validation and application of a robust yeast estrogen bioassay for the screening of estrogenic activity in animal feed. *Food Addit Contam* 23:556–568.
 53. Gizzi G, Hoogenboom LAP, Von Holst C, Rose M, Anklam E. 2005. Determination of dioxins (PCDDs/PCDFs) and PCBs in food and feed using the DR CALUX[®] bioassay: Results of an international validation study. *Food Addit Contam* 22:472–481.
 54. Brand W, De Jongh CM, Van der Linden SC, Mennes W, Puijker LM, Van Leeuwen CJ, Van Zewel AP, Schriks M, Heringa MB. 2013. Trigger values for investigation of hormonal activity in drinking water and its sources using CALUX bioassays. *Environ Int* 55:109–118.
 55. Jarošová B, Bláha L, Giesy JP, Hilscherová K. 2014. What level of estrogenic activity determined by in vitro assays in municipal waste waters can be considered as safe? *Environ Int* 64:98–109.
 56. Kunz PY, Kienle C, Carere M, Homazava N, Kase R. 2015. In vitro bioassays to screen for endocrine active pharmaceuticals in surface and waste waters. *J Pharmaceut Biomed* 106:107–115.
 57. Leusch FDL, Khan SJ, Gagnon MM, Quayle P, Trinh T, Coleman H, Rawson C, Chapman HF, Blair P, Nice H, Reitsema T. 2014. Assessment of wastewater and recycled water quality: A comparison of lines of evidence from in vitro, in vivo and chemical analyses. *Water Res* 50:420–431.
 58. Hamers T, Leonards PEG, Legler J, Vethaak AD, Schippers CA. 2010. Toxicity profiling: An integrated effect-based tool for site-specific sediment quality assessment. *Integr Environ Assess Manag* 6:761–773.
 59. Van Ede KI, Van Duursen MBM, Van den Berg M. 2016. Evaluation of relative effect potencies (REPs) for dioxin-like compounds to derive systemic or human-specific TEFs to improve human risk assessment. *Arch Toxicol* 90:1293–1305.
 60. Brack W, Ait-Aissa S, Burgess RM, Busch W, Creusot N, Di Paolo C, Escher BI, Hewitt LM, Hilscherova K, Hollender J, Hollert H, Jonker W, Kool J, Lamoree M, Muschket M, Neumann S, Rostkowski P, Ruttkies C, Schollee J, Schymanski EL, Schulze T, Seiler T-B., Tindall AJ, Umbuzeiro GDA, Vrana B, Krauss M. 2016. Effect-directed analysis supporting monitoring of aquatic environments—An in-depth overview. *Sci Total Environ* 544:1073–1118.