

## VU Research Portal

### **Within-Person Pain Variability and Mental Health in Older Adults With Osteoarthritis: An Analysis Across 6 European Cohorts**

de Koning, Elisa J.; Timmermans, E.J.; van Schoor, N.M.; Stubbs, Brendon; van den Kommer, Tessa N.; Dennison, E.M.; Limongi, Federica; Castell, Maria Victoria; Edwards, M.H.; Queipo, Rocio; Cooper, Cyrus; Siviero, Paola; van der Pas, Suzan; Pedersen, N.L.; Sánchez-Martínez, Mercedes; Deeg, D.J.H.; Denkinge, Michael D.; Nikolaus, T.; Denkinge, M.; Peter, R.

**published in**

Journal of Pain

2018

**DOI (link to publisher)**

[10.1016/j.jpain.2018.02.006](https://doi.org/10.1016/j.jpain.2018.02.006)

**document version**

Publisher's PDF, also known as Version of record

**document license**

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

**citation for published version (APA)**

de Koning, E. J., Timmermans, E. J., van Schoor, N. M., Stubbs, B., van den Kommer, T. N., Dennison, E. M., Limongi, F., Castell, M. V., Edwards, M. H., Queipo, R., Cooper, C., Siviero, P., van der Pas, S., Pedersen, N. L., Sánchez-Martínez, M., Deeg, D. J. H., Denkinge, M. D., Nikolaus, T., Denkinge, M., ... Edwards, M. H. (2018). Within-Person Pain Variability and Mental Health in Older Adults With Osteoarthritis: An Analysis Across 6 European Cohorts. *Journal of Pain*, 19(6), 690-698. <https://doi.org/10.1016/j.jpain.2018.02.006>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)



## Within-Person Pain Variability and Mental Health in Older Adults With Osteoarthritis: An Analysis Across 6 European Cohorts

Elisa J. de Koning,<sup>\*</sup> Erik J. Timmermans,<sup>\*</sup> Natasja M. van Schoor,<sup>\*</sup> Brendon Stubbs,<sup>†</sup> Tessa N. van den Kommer,<sup>‡</sup> Elaine M. Dennison,<sup>§</sup> Federica Limongi,<sup>¶</sup> Maria Victoria Castell,<sup>||</sup> Mark H. Edwards,<sup>§</sup> Rocio Queipo,<sup>||</sup> Cyrus Cooper,<sup>§</sup> Paola Siviero,<sup>¶</sup> Suzan van der Pas,<sup>\*</sup> Nancy L. Pedersen,<sup>\*\*</sup> Mercedes Sánchez-Martínez,<sup>||</sup> Dorly J.H. Deeg,<sup>\*</sup> and Michael D. Denkingert<sup>††</sup> and the EPOSA Group

<sup>\*</sup>Department of Epidemiology and Biostatistics / Amsterdam Public Health Research Institute, Vrije Universiteit Medical Center, Amsterdam, The Netherlands.

<sup>†</sup>Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London / Physiotherapy Department, South London and Maudsley National Health Service Foundation Trust, London, United Kingdom.

<sup>‡</sup>Geestelijke Gezondheidszorg inGeest / Vrije Universiteit Medical Center, Amsterdam, The Netherlands.

<sup>§</sup>Medical Research Council Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital / Queen Alexandra Hospital, Cosham, Portsmouth, United Kingdom.

<sup>¶</sup>National Research Council, Aging Branch, Institute of Neuroscience, Padova, Italy.

<sup>||</sup>Department of Preventive Medicine and Public Health, Unit of Primary Care and Family Medicine, Faculty of Medicine, Universidad Autonoma de Madrid, Madrid, Spain.

<sup>\*\*</sup>Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.

<sup>††</sup>AGAPLESION Bethesda Hospital, Geriatric Research Unit / Institute of Epidemiology and Medical Biometry, University of Ulm, Ulm, Germany.

**Abstract:** Pain is a key symptom of osteoarthritis (OA) and has been linked to poor mental health. Pain fluctuates over time within individuals, but a paucity of studies have considered day-to-day fluctuations of joint pain in relation to affective symptoms in older persons with OA. This study investigated the relationship of pain severity as well as within-person pain variability with anxiety and depression symptoms in 832 older adults with OA who participated in the European Project on OsteoArthritis (EPOSA): a 6-country cohort study. Affective symptoms were examined with the Hospital Anxiety and Depression Scale, pain severity was assessed with the Western Ontario and McMaster Universities OA Index and the Australian/Canadian Hand Osteoarthritis Index, and intraindividual pain variability was measured using pain calendars assessed at baseline, 6, and 12 to 18 months. Age-stratified multiple linear regression analyses adjusted for relevant confounders showed that more pain was associated with more affective symptoms in older-old participants (74.1–85 years). Moreover,

Received April 6, 2017; Revised January 16, 2018; Accepted February 13, 2018.

The Indicators for Monitoring COPD and Asthma—Activity and Function in the Elderly in Ulm study (IMCA—ActiFE) was supported by the European Union (2005121) and the Ministry of Science, Baden-Württemberg. The Italian cohort study is part of the National Research Council Project on Aging (PNR). The Longitudinal Aging Study Amsterdam is financially supported by the Dutch Ministry of Health, Welfare and Sports. The Penagrande study was partially supported by the National Fund for Health Research (Fondo de Investigaciones en Salud) of Spain (project numbers FIS PI 05/1898; FIS RETICEF RD06/0013/1013, and FIS PS09/02143). The Swedish Twin Registry is supported in part by the Swedish Ministry of Higher Education. The Hertfordshire Cohort Study is funded by the Medical Research Council of Great Britain, Arthritis Re-

search UK, the British Heart Foundation and the International Osteoporosis Foundation. The funding sources had no role in the study design, the collection, analysis and interpretation of the data, in the writing of the report and in the decision to submit the article for publication.

The authors have no conflicts of interest to declare.

Address reprint requests to Elisa J. de Koning, MSc, Vrije Universiteit Medical Center, Dpt. of Epidemiology & Biostatistics / Amsterdam Public Health Research Institute, De Boelelaan 1089a, 1081 HV Amsterdam, The Netherlands. E-mail: [ej.dekoning@vumc.nl](mailto:ej.dekoning@vumc.nl)

1526-5900/\$36.00

© 2018 by the American Pain Society

<https://doi.org/10.1016/j.jpain.2018.02.006>

older-old participants experienced fewer symptoms of anxiety (ratio = .85, 95% confidence interval [CI], .77–.94), depression (ratio = .90, 95% CI, .82–.98), and total affective symptoms (ratio = .87, 95% CI, .79–.94) if their pain fluctuated more. No such association was evident in younger-old participants (65–74.0 years). These findings imply that stable pain levels are more detrimental to mental health than fluctuating pain levels in older persons.

**Perspective:** *This study showed that more severe and stable joint pain levels were associated with anxiety and depressive symptoms in older persons with OA. These findings emphasize the importance of measuring pain in OA at multiple time points, because joint pain fluctuations may be an indicator for the presence of affective symptoms.*

© 2018 by the American Pain Society

**Key words:** Osteoarthritis, pain variability, depressive symptoms, anxiety symptoms, older persons.

Osteoarthritis (OA) is the most common form of musculoskeletal disorders worldwide<sup>41</sup> and a leading cause of functional limitations and loss of independence in later life. Chronic pain represents one of the key debilitating symptoms of OA.<sup>10</sup> Moreover, OA—and in particular pain experienced by OA patients—has been linked to an increased risk of affective symptoms, such as depression<sup>5,6,20,29</sup> and anxiety.<sup>5,6,29</sup> In turn, affective symptoms may have a detrimental effect on pain perception.<sup>5</sup> A recent systematic review and meta-analysis concluded that approximately 20% of persons with OA experience depressive and/or anxiety symptoms.<sup>35</sup> More insight is needed in the vicious cycle in which pain, depression, and anxiety may continually reinforce each other over time in persons who suffer from OA.

The severity of pain in OA may vary greatly between individuals.<sup>11</sup> Moreover, individuals may experience stable pain levels or fluctuations in pain over time.<sup>11,32</sup> A fluctuating course requires repeated measurements and is difficult to capture with commonly used instruments such as the Western Ontario and McMaster Universities OA Index (WOMAC)<sup>7</sup> and the Australian/Canadian Hand Osteoarthritis Index (AUSCAN).<sup>8</sup>

The association between variability in pain experience and mental health is bidirectional<sup>18,21</sup> and has been rarely studied.<sup>32</sup> Schneider et al<sup>32</sup> showed more day-to-day fluctuations of pain in adult patients with OA or other rheumatic diseases who had higher levels of depressive symptoms. Anxiety symptoms were not predictive of variability in pain scores in their study.

To our knowledge, no study to date has focused on the effect of intraindividual pain variability over time on anxiety and depressive symptoms in persons with OA. The European Project on OsteoArthritis (EPOSA) provides a unique opportunity to examine these associations in older persons with OA from the general population.<sup>40</sup> In this study, we used data of the EPOSA cohort to examine the relationship of pain severity as well as within-person pain variability over time with symptoms of anxiety and depression. Presuming that a lack of predictability of pain is detrimental to mental well-being,<sup>32</sup> it was hypothesized that more severe pain and greater pain variability over time would be associated with more affective symptoms.<sup>5,20,32</sup>

## Methods

### Study Design and Participants

The EPOSA is a multicohort study aimed at investigating the personal and societal consequences of OA and its determinants, using preharmonized data from 6 European population-based cohorts of older adults. Participating countries include Germany, Italy, The Netherlands, Spain, Sweden, and the United Kingdom. Detailed information about the EPOSA design, participants, and procedures of data collection are described elsewhere.<sup>40</sup> In short, 2,942 community-dwelling persons of 65 to 85 years with and without OA were included. Interviews and clinical assessments took place at baseline and after 12 to 18 months. Furthermore, immediately after the baseline and follow-up interview as well as 6 months after baseline, participants were asked to complete a 2-week calendar assessing joint pain. Presence of the 3 most common forms of OA—knee, hip, and hand OA—was assessed with a clinical examination on the basis of the criteria of the American College of Rheumatology,<sup>3</sup> which resulted in a total of 889 participants with clinical OA of the knee, hand, and/or hip. For the current study, we used data from persons with clinical OA who did not report joint replacements or surgery in the past 2 weeks at the site of the clinical OA (N = 832). All 6 cohort studies were approved by the ethical review boards of the respective institutions and all participants gave written informed consent before the start of the study.

## Measurements

### Pain Severity

Pain severity in the past 48 hours was measured with the WOMAC<sup>7</sup> (knee and hip OA) and AUSCAN<sup>8</sup> (hand OA) pain subscales at baseline and follow-up. These commonly used questionnaires assess pain in 5 daily life situations, rated on a 0 (no pain) to 4 (extreme pain) scale. Because the WOMAC and AUSCAN pain subscales are scored on the same scale (score range = 0–20), the raw scores of knee, hip, and hand pain were added, resulting in a total pain severity score (range = 0–60). This score was then transformed to a 0 to 100 scale for comparability with other studies.<sup>7</sup>

## Pain Variability

At baseline and after 6 and 12 to 18 months, participants completed 2-week calendars assessing daily joint pain. On these calendars, participants indicated their amount of joint pain per day, ranging from 0 (no pain) to 10 (extreme pain). In addition, participants were asked to indicate whether they took pain medication, whether they consulted a doctor, and whether they had surgery as a result of their joint pain. No information regarding site and context of the joint pain was collected. Participants from Spain did not complete the 6-month calendar. As a measure of individual day-to-day pain variability over a longer time period, the SD of the scores of all available days of the 3 calendars was calculated for each participant. The more day-to-day pain variability a person reported, the larger the SD. At least 7 of 14 days of at least 1 calendar had to be completed by a participant to be included in the pain variability analyses.

## Affective Symptoms

The Hospital Anxiety and Depression Scale (HADS)<sup>44</sup> was used to measure symptoms of anxiety (HADS-A; 7 items) and depression (HADS-D; 7 items) at baseline and follow-up. The HADS is a widely used, self-rated measure assessing affective symptoms the participant experienced in the past 4 weeks. The total HADS score (HADS-T) is a general measure of total affective symptoms. Item scores range from 0 (rarely or never) to 3 (mostly or always). Total scores for the subscales range from 0 to 21, with a score of  $\geq 8$  being indicative of the presence of a disorder. Total scores for the combined scale range from 0 to 42. The HADS shows good reliability and validity in several medical and general population samples<sup>9</sup> and in people with OA.<sup>6</sup>

## Potential Effect Modifiers and Confounders

We examined whether age<sup>23,27</sup> and gender<sup>25,38</sup> were effect modifiers in the relationship between pain and affective symptoms. The analyses were adjusted for the following confounders: age, gender (if no effect modifier), country, education level (less than elementary school, elementary school, vocational/secondary education, or college/university education), smoking<sup>15</sup> (never, former, or current smoker), body mass index (BMI; calculated as weight in kilograms divided by the height in meters squared),<sup>26</sup> number of chronic diseases (selected from the 7 most common chronic somatic diseases in the older population: chronic nonspecific lung disease, cardiovascular disease, disease of peripheral arteries, stroke, diabetes mellitus, cancer, and osteoporosis), physical activity (in kcal/d, assessed with the Longitudinal Aging Study Amsterdam Physical Activity Questionnaire),<sup>34,36</sup> and use of psychotropic medication (antidepressants, anxiolytics and/or antipsychotics use yes/no) and pain medication (analgesics or anti-inflammatory medication use yes/no).<sup>14,28</sup>

## Statistical Analysis

Descriptive characteristics of the study sample were examined with frequencies and percentages for categorical

variables and medians and interquartile ranges (IQRs) for skewed continuous variables. With the exception of age, gender, and country, all descriptive statistics were weighted to adjust for differences in the distributions of age and gender across the 6 samples. Weights were calculated per gender and per 5-year age category, using the formula:  $\text{weight} = \text{Nexp}/\text{Nobs}$ , where *Nexp* is the number of persons in a specific age/gender category in the European population, and *Nobs* is the number of persons in a specific age/gender category in the cohort.<sup>40</sup> Differences between participants in the analyzed sample and persons who dropped out after baseline were examined with  $\chi^2$  (for dichotomous or categorical variables) or Mann–Whitney (continuous variables) tests in nonresponse analyses. Cronbach  $\alpha$ s were calculated for the HADS and the WOMAC/AUSCAN, at baseline as well as at follow-up, to evaluate the reliability of the scales in the EPOSA sample.

For pain severity, cross-sectional as well as longitudinal multiple linear regression analyses were conducted. For the cross-sectional analysis, the baseline WOMAC/AUSCAN combined pain score was used as predictor and the baseline HADS-A, HADS-D, and HADS-T scores as continuous outcome variables. To examine the relationship between pain severity and affective symptoms over time, the baseline WOMAC/AUSCAN combined pain score was used as predictor, the HADS-A, HADS-D, and HADS-T scores after 12 to 18 months, respectively, as outcomes, and the respective baseline HADS score as an additional covariate to control for baseline levels of affective symptoms.

To study whether intraindividual pain variability over time affects affective symptoms, multiple linear regression analyses were performed with the individual joint pain SDs (calculated from the 3 pain calendars) as predictor and the 3 HADS scores at 12 to 18 months as outcomes. A potential drawback of using the SD of the individual pain scores as a measure of pain variability is that the SD is the same for persons who report the same magnitude of fluctuations, regardless of their mean pain score. For instance, 2 persons with mean pain levels of 0 and 10, respectively, who experience no pain fluctuations, will both have an SD of 0. For this reason, we adjusted all pain variability analyses for the individual mean joint pain score, calculated from all available joint pain scores on the calendars (similar to the calculation of the SD pain variability score).

To study possible effect modification by age and gender, an interaction term of the predictor with the potential effect modifier was created. This interaction term was added to the unadjusted analyses. If the *P* value of the interaction term was  $< .10$ , effect modification was considered to be present and stratified analyses were conducted.

To examine the robustness of the effects, we analyzed 2 models. In model 1, we adjusted for country, age, and gender (if no effect modifier). In model 2, we additionally adjusted for education level, smoking, BMI, number of chronic diseases, physical activity, and use of psychotropic and pain medication.

We conducted preplanned sensitivity analyses for pain severity as well as pain variability omitting persons who

reported taking psychotropic medications, because these drugs may not only decrease affective symptoms, but also alleviate pain.<sup>28</sup>

Because the score distribution of all 3 HADS measures was skewed to the right, the following natural log transformation was conducted on the HADS scores:  $\ln(1 + \text{HADS score})$ . The regression coefficients and confidence intervals from the regression analyses were back-transformed to obtain interpretable ratios. These ratios can easily be converted into a percentage of change in the outcome variable (affective symptoms) per 1 unit change in the predictor (pain). Because a 1-unit increase/decrease on the pain severity scale of 0 to 100 is very small,

a more meaningful 10-unit increase/decrease<sup>4</sup> was derived by calculating ratio<sup>10</sup>. A 2-sided *P* value of .05 was regarded as statistically significant. All analyses were conducted with SPSS version 22 (IBM Corp, Armonk, NY).

## Results

Characteristics of the study population are shown in Table 1. More than two-thirds of the participants were female (70.2%) and the median age was 74.0 years. Of the 832 participants in the present analyses, 67.0% had OA at 1 site, 26.6% at 2 sites, and 6.5% at all 3 sites. The median range of fluctuations in joint pain within 1 cal-

**Table 1. Characteristics of the EPOSA Participants With Clinical OA of the Knee, Hip, and/or Hand\***

	TOTAL GROUP	YOUNGER OLD (65–74.0 YEARS)	OLDER OLD (74.1–85 YEARS)
Presence of clinical OA	N = 832	n = 450	n = 382
Knee OA	523 (63.3)	266 (59.3)	257 (70.2)
Hip OA	162 (19.7)	93 (21.1)	69 (18.2)
Hand OA	468 (56.3)	263 (58.8)	205 (54.8)
Country	N = 832	n = 450	n = 382
Germany	76 (9.1)	47 (10.4)	29 (7.6)
Italy	185 (22.2)	109 (24.2)	76 (19.9)
The Netherlands	132 (15.9)	53 (11.8)	79 (20.7)
Spain	173 (20.8)	76 (16.9)	97 (25.4)
Sweden	157 (18.9)	115 (25.6)	42 (11.0)
United Kingdom	109 (13.1)	50 (11.1)	59 (15.4)
Gender	N = 832	n = 450	n = 382
Female	548 (70.2)	322 (71.6)	262 (68.6)
Male	248 (29.8)	128 (28.4)	120 (31.4)
Age	N = 832	n = 450	n = 382
Median [IQR]	74.0 [70.0–78.0]	70.7 [68.0–72.0]	78.0 [76.0–81.0]
Education level	N = 831	n = 449	n = 382
Elementary school not completed	127 (15.6)	50 (11.3)	77 (21.0)
Elementary school completed	307 (37.3)	152 (34.7)	155 (40.4)
Vocational or general secondary education	263 (31.4)	161 (35.3)	102 (26.4)
College or university education	134 (15.8)	86 (18.7)	48 (12.2)
Smoking	N = 828	n = 449	n = 379
Never	445 (55.0)	239 (53.2)	206 (57.2)
Current	46 (5.9)	34 (8.3)	12 (3.0)
Former	337 (39.1)	176 (38.5)	161 (39.8)
BMI	N = 810	n = 441	n = 369
Median [IQR]	27.8 [24.8–31.2]	27.5 [24.6–30.9]	28.2 [25.2–31.3]
Number of chronic diseases	N = 824	n = 445	n = 379
Median [IQR]	1 [0–2]	1 [0–2]	1 [1–2]
Physical activity, kcal/d	N = 798	n = 437	n = 361
Median [IQR]	686 [428–1,058]	749 [494–1,176]	597 [349–940]
Use of psychotropic medication	N = 832	n = 450	n = 382
Median [IQR]	178 (21.7)	82 (17.5)	96 (27.1)
Use of analgesic/anti-inflammatory medication	N = 831	n = 449	n = 382
Median [IQR]	265 (31.6)	143 (30.5)	122 (32.9)
Pain severity (WOMAC/AUSCAN)†			
Baseline	N = 823	n = 447	n = 376
Median [IQR]	15.0 [10.0–25.0]	15.0 [8.3–23.3]	16.7 [10.0–26.7]
Follow-up	N = 646	n = 367	n = 279
Median [IQR]	15.0 [6.7–26.7]	13.4 [5.0–25.0]	16.7 [6.7–28.3]
Pain variability (from pain calendar)‡	N = 762	n = 428	n = 334
Median [IQR]	1.3 [0.9–1.9]	1.3 [0.9–1.8]	1.3 [0.9–1.9]
HADS			
Baseline	N = 802	n = 437	n = 365
HADS-A	5 [3–9]	6 [3–9]	5 [3–8]
HADS-D	4 [2–7]	4 [1–6]	5 [2–7]
HADS-T	10 [5–15]	9 [5–15]	10 [6–15]
Follow-up	N = 650	n = 370	n = 280
HADS-A	5 [2–7]	4 [2–8]	5 [2–7]
HADS-D	3 [1–6]	3 [1–6]	4 [2–7]
HADS-T	8 [4–13]	8 [4–13]	9 [5–14]

NOTE. Data are presented as n (%) or as median [IQR].

\*With exception of age, gender, and country, descriptive statistics are weighted; the n is unweighted.

†Score range = 0 to 100.

‡Pain variability shown in individual SD units.

endar (2 weeks) was 3 points (IQR = 2–4, range = 0–10) for the baseline pain calendar and 2 (IQR = 1–4, range = 0–10) for the 6 and 12 to 18 months pain calendars.

Cronbach  $\alpha$ s of the HADS-A, HADS-D, and HADS-T at baseline and follow-up varied between .736 and .859, indicating good reliability of the scale. Similarly, the reliability of the WOMAC/AUSCAN pain severity scale was also good (Cronbach  $\alpha$  = .891 and .908 for the baseline and follow-up scales, respectively).

In nonresponse analyses for pain variability, participants in the analyzed sample were compared with persons who dropped out of the study or did not complete at least 7 days on at least 1 pain calendar (n = 69). At baseline, 85.2% of the participants completed  $\geq 7$  days of the calendar. At 6 months, 57.7% completed  $\geq 7$  days (Spain not taken into account), and at 12 to 18 months, 58.5% completed  $\geq 7$  days. At baseline, nonresponders to the pain calendar were older ( $P < .001$ ) and had more depressive symptoms ( $P = .019$ ), compared with participants who did complete at least 7 days. At 6 months, persons who dropped out or did not complete the calendar were older ( $P = .029$ ), lower educated ( $P = .009$ ), had more anxiety ( $P < .001$ ) and depressive symptoms ( $P = .001$ ), more severe pain ( $P = .043$ ), more chronic diseases ( $P = .044$ ), and higher BMI ( $P = .047$ ). At 12 to 18 months, this pattern was similar, with the addition that nonresponders were more often female ( $P = .022$ ) and did not have a higher BMI compared with the responders to the calendar.

### Pain Severity

#### Cross-Sectional Analyses

In the cross-sectional baseline analyses, gender was not an effect modifier. However, age was a significant effect modifier for anxiety ( $P < .001$ ) and total affective symptoms ( $P = .002$ ), but not for depressive symptoms ( $P = .22$ ).

Therefore, analyses were stratified at the median age (74.0 years) for anxiety and total affective symptoms and analyses for depressive symptoms were conducted in the group as a whole (Table 2). The regression analyses showed that, adjusted for relevant confounders, greater pain severity was associated with more anxiety (ratio = 1.011,  $P < .001$ ) and total affective symptoms (ratio = 1.007,  $P = .011$ ) in persons older than 74 years (“older old”), but not in persons between 65 and 74 years (“younger old”; ratio = 1.002,  $P = .43$  and ratio = 1.004,  $P = .12$ , respectively). In the total group, more severe pain was associated with more depressive symptoms (ratio = 1.006,  $P = .001$ ). Accordingly, a 10-unit increase in pain severity corresponded to an 11.6% increase in anxiety symptoms and a 7.2% increase in total affective symptoms in persons older than 74 years, and with a 6.2% increase in depressive symptoms in the total group.

#### Longitudinal Analyses

Contrary to the cross-sectional analyses, age was not an effect modifier in the longitudinal analyses of pain severity. However, sex was a significant effect modifier for anxiety and total affective symptoms ( $P = .002$  and .021, respectively; HADS-D  $P = .21$ ). Hence, analyses for anxiety and total affective symptoms were conducted separately for male and female participants and analyses for depressive symptoms were conducted in the total group (Table 2). Adjusted for confounding, more severe pain at baseline was associated with more affective symptoms at follow-up in men (ratio = 1.009,  $P = .022$ ) but not in women (ratio = .99,  $P = .78$ ). In the total group, more severe pain at baseline was associated with more depressive symptoms at follow-up (ratio = 1.006,  $P = .002$ ). In accordance, a 10-unit increase in baseline pain severity corresponded to a 9.4% increase in total affective symptoms at follow-up in men and a 6.2% increase in depressive symptoms in the total group.

**Table 2. Cross-sectional and Longitudinal Associations Between Pain Severity and Affective Symptoms, Analyzed with Multiple Linear Regression Analysis**

	TOTAL GROUP		YOUNGER OLD (65.0–74.0 YEARS)		OLDER OLD (74.1–85 YEARS)	
	MODEL 1†	MODEL 2‡	MODEL 1†	MODEL 2‡	MODEL 1†	MODEL 2‡
Cross-sectional						
HADS-A			1.04 (.99–1.09)	1.02 (.97–1.07)	1.14* (1.08–1.21)	1.12* (1.05–1.19)
HADS-D	1.08* (1.05–1.13)	1.06** (1.02–1.11)				
HADS-T			1.06*** (1.01–1.12)	1.04 (.99–1.09)	1.12* (1.06–1.17)	1.07*** (1.02–1.13)
	TOTAL GROUP		FEMALE		MALE	
	MODEL 1†	MODEL 2‡	MODEL 1†	MODEL 2‡	MODEL 1†	MODEL 2‡
LONGITUDINAL						
HADS-A			.97 (.93–1.01)	.98 (.94–1.02)	1.10*** (1.02–1.20)	1.08 (1.00–1.17)
HADS-D	1.06** (1.02–1.09)	1.06** (1.02–1.09)				
HADS-T			.99 (.95–1.03)	.99 (.95–1.04)	1.09*** (1.02–1.17)	1.09*** (1.01–1.17)

NOTE. Data are presented as ratio per 10-point difference/change in WOMAC/AUSCAN pain severity (95% confidence interval).

\* $P < .001$ .

\*\* $P < .01$ .

\*\*\* $P < .05$ .

†Model 1 adjusted for sex, age, and country.

‡Model 2 additionally adjusted for education level, smoking, BMI, chronic diseases, physical activity, and use of psychotropic and pain medication.

**Table 3. Longitudinal Associations Between Pain Variability and Affective Symptoms in Persons With OA, Analyzed Using Multiple Linear Regression Analysis**

	YOUNGER OLD (65.0–74.0 YEARS)		OLDER OLD (74.1–85 YEARS)	
	MODEL 1†	MODEL 2‡	MODEL 1†	MODEL 2‡
HADS-A	1.07 (.97–1.18)	1.07 (.97–1.18)	.86* (.77–.95)	.85* (.77–.94)
HADS-D	1.02 (.94–1.11)	1.02 (.94–1.12)	.91 (.83–1.00)	.90** (.82–.98)
HADS-T	1.01 (.92–1.10)	1.04 (.95–1.15)	.90† (.82–.98)	.87* (.79–.94)

NOTE. Data are presented as ratio (95% confidence interval). Pain variability measured with individual SDs of joint pain scores on a maximum of 3 pain calendars.

\* $P < .01$ .

\*\* $P < .05$ .

†Model 1 adjusted for sex, age, and country.

‡Model 2 additionally adjusted for education level, smoking, BMI, chronic diseases, physical activity, and use of psychotropic and pain medication.

### Sensitivity Analyses

The sensitivity analyses without persons who used psychotropic medications ( $n = 178$ ) did not change any of the conclusions (data not shown but available on request from the author).

### Pain Variability

Table 3 shows the results of the pain variability analyses. Gender was not a significant effect modifier. However, age was a significant effect modifier for all 3 HADS outcomes (HADS-A:  $P = .002$ ; HADS-D:  $P = .004$ ; HADS-T:  $P = .001$ ). Therefore, all analyses were conducted separately for the 2 age groups, again stratified at the median age. Greater pain variability over 12 to 18 months was significantly associated with lower anxiety (ratio = .85,  $P = .002$ ), depressive (ratio = .90,  $P = .018$ ), and total affective symptoms (ratio = .87,  $P = .002$ ) after 12 to 18 months in the older-old participants. In accordance, a 1-unit increase in pain variability was associated with a 15% lower anxiety score, a 10% lower depression score, and a 13% lower affective symptoms score in persons older than 74 years with OA. No statistically significant associations between pain variability and affective symptoms were observed in participants between 65 and 74 years of age.

### Sensitivity Analyses

The sensitivity analyses without persons who use psychotropic medications ( $n = 178$ ) did not change any of the conclusions (data not shown but available on request from the author).

### Discussion

Pain is a common and debilitating symptom of OA that can have substantial consequences for daily life. The present study investigated whether pain severity and pain fluctuations over time in older persons with OA from the general European population were associated with symptoms of anxiety and depression. The results showed that more severe pain was associated with more depressive symptoms. In addition, higher pain severity in persons older than 74 years ("older-old" persons) was associated with more anxiety and total affective symptoms.

Over time, depressive symptoms became worse when pain severity increased. In men, total affective symptoms increased with increasing pain severity. Previous studies have also indicated that pain severity is related to symptoms of anxiety and depression.<sup>4,6</sup> The results from the EPOSA study strengthen these findings by showing the association in a large sample of community-dwelling people with OA from multiple countries.

Day-to-day pain variability was assessed with 3 two-week pain calendars over a time course of 12 to 18 months. The calendars showed substantial individual day-to-day fluctuations of pain levels. Contrary to our expectations, more fluctuations in joint pain were associated with less anxiety, depressive, and total affective symptoms in the older-old persons. In the younger-old participants (65–74 years), pain variability was not significantly associated with affective symptoms.

Similar to the study by Schneider et al,<sup>32</sup> we expected to find more affective symptoms in persons with more fluctuating pain levels, possibly because of lack of control over and predictability of pain episodes. However, we found the opposite pattern. There are some differences between our study and Schneider's study that may explain this discrepancy. For instance, the study by Schneider et al recruited patients with OA and other rheumatic diseases from clinics, whereas we investigated a population-based sample with (clinical) OA. Furthermore, Schneider et al assessed pain in general, whereas we specifically asked about joint pain. Finally, the sample size of the study by Schneider et al was substantially smaller ( $N = 300$ ) than the sample size of the present study ( $N = 832$ ).

An explanation for the observed pain variability results might be that more predictable, stable, and chronic pain levels activate feelings of hopelessness and lack of perspective.<sup>37</sup> In addition, persons who also have "good days" may be better able to withstand the "bad days," in contrast to persons who have the same, chronic pain level every day. The latter group may also experience more interference with daily life functioning. Furthermore, evidence suggests that maladaptive plastic brain changes occur in persons with chronic pain, that do not occur in persons with more acute pain. These changes may alter pain perception and are related to depression.<sup>13,16</sup>

The negative association between pain variability and affective symptoms was observed in the age group older

than 74 years only. The cross-sectional pain severity analyses showed a similar pattern. Older persons may have fewer effective coping strategies because of more morbidity and/or frailty, compared with younger persons.<sup>17,19,31</sup> In attempting to explain this age difference further, we investigated differences in pain and affective symptoms between these 2 age groups in our sample. The older-old group did not experience more severe or more fluctuating pain levels, but did report more depressive and total affective symptoms than their younger-old counterparts. This finding may reflect the mutually reinforcing effect of pain and affective symptoms in the older age group.<sup>5,43</sup> Our age-stratified results are in contrast with the study by Sanders et al, who reported that older age did not influence the relationship between pain and depression.<sup>31</sup> However, the study by Sanders et al was conducted with older persons from the general population, not in a specific OA sample.

Because the minimally important difference (MID) in HADS scores has rarely been studied, it is difficult to draw conclusions about whether the observed association between pain severity and variability and affective symptoms is clinically meaningful. One study involving patients with moderate to severe chronic obstructive pulmonary disease showed that the MID for their study sample was 1.5 points.<sup>30</sup> This MID can be taken as an example for the EPOSA study, but it should be kept in mind that the chronic obstructive pulmonary disease sample differed from the EPOSA sample in multiple ways. Because we calculated ratios in our study, it depends on the baseline HADS score whether a change is clinically meaningful. Our analyses showed that a reduction in pain severity or variability was associated with up to 15% lower HADS scores. Hence, for higher baseline HADS scores, this change is clinically meaningful. One could argue that this change is especially relevant for persons with higher baseline HADS scores, because they experience the most affective symptoms.

This study combines several strengths. First, EPOSA is a large study with a preharmonized data set including prospective cohort data from persons with OA from 6 European countries. Presence of OA was determined using standardized assessments methods. The study samples were drawn from the general population, which means that we have data from persons with OA ranging from mild to severe and from persons who did and did not receive care for their OA. This study characteristic increases the generalizability of our results. Because of the large data set, we were able to adjust for many relevant confounders and examine effect modification. Another asset is that we analyzed day-to-day pain variability with multiple joint pain calendars over a large time range, which is a novel and promising way of analyzing pain in OA. To our knowledge, no other study has focused on the effect of pain variability over time on anxiety and depressive symptoms in persons with OA.

This study also has some limitations. First, we did not ask the participants about other types of pain that they may have had at the same time as their joint pain, so we were not able to adjust for this potentially important confounder. Second, the pain calendar at 12 to 18 months was completed after the HADS interview, whereas we

defined the HADS as our dependent variable. However, the time frame between these measurements was rather short (1 day to 2 weeks), because the calendar was started the day after the interview. Leaving the third pain calendar out would not only reduce the power of our analyses, but would also mean that the HADS would be measured 6 months after the last pain calendar, which is a much longer time frame. Third, the information from the pain calendar was not specific enough to determine the site and context of the joint pain. Fourth, the calendars were filled out by hand, without electronic time stamps. Although this paper-and-pen method is preferred by part of the older population, it is prone to "backfilling," which could potentially harm data quality.

A potential methodological issue is the use of the SDs of all available joint pain scores as a measure of pain variability. We combined pain scores from the 3 calendars that were relatively far apart in time. Correlations between pain variability scores of each calendar were moderate in strength (Spearman  $\rho = .409-.571$ , all  $P_s < .001$ ), indicating that the 3 time points are not dissimilar. Moreover, individual SD scores may suffer from relatively low reliability.<sup>42</sup> This potential issue is more pronounced if the number of observations is low. In most cases, the SD was calculated from  $3 \times 14 = 42$  observations. Potentially, this number is not sufficient to reach high reliability, which may have led to an underestimation of the results. Future research should examine whether the use of more observations can create a more reliable pain variability score.

The nonresponse analyses revealed that the persons with the highest pain and affective symptom levels were lost to follow-up, which may have attenuated our results. Previous research, however, indicates that this loss of follow-up data is not likely to be problematic.<sup>39</sup> Finally, despite the longitudinal design, it is still difficult to draw firm conclusions about the causality of pain severity/variability and affective symptoms. Previous research suggests that the relationship may even be bidirectional.<sup>5,33</sup>

The evidence from this study suggests that persons with OA experience substantial day-to-day fluctuations of pain levels. This result is in accordance with another study that also observed that changes in pain were associated with changes in mood in persons with OA.<sup>2,22</sup> These results emphasize the importance of measuring pain variability in persons with OA, because it may shed a different light on experienced pain and its consequences, compared with measuring pain at only 1 time point.<sup>2</sup>

At the moment, treatment for depression and anxiety in persons with OA is suboptimal.<sup>1,43</sup> Pain, anxiety, and depression are risk factors for further functional decline in OA<sup>12</sup> and the disease burden increases if a person experiences affective symptoms.<sup>24</sup> Future research should focus on finding effective interventions for reducing (the burden of) chronic pain to improve mental well-being in older persons with OA.

## Conclusions

The present study showed that more severe and stable joint pain levels are associated with more symptoms of



anxiety and depression in persons older than 74 years of age who suffer from OA. Because of the fluctuating nature of joint pain in OA, assessing pain severity at only 1 time point is not enough to capture the whole story. Therefore, it is important to assess the chronicity of the pain in OA, because it may be an indicator for the presence of anxiety and/or depressive symptoms.

## Acknowledgments

We are very grateful to all participants of the EPOSA study for their valued contributions.

## References

1. Agarwal P, Pan X, Sambamoorthi U: Depression treatment patterns among individuals with osteoarthritis: A cross sectional study. *BMC Psychiatry* 13:121, 2013
2. Allen KD: The value of measuring variability in osteoarthritis pain. *J Rheumatol* 34:2132-2133, 2007
3. Altman RD: Classification of disease: Osteoarthritis. *Semin Arthritis Rheum* 20:40-47, 1991
4. Angst F, Aeschlimann A, Stucki G: Smallest detectable and minimal clinically important differences of rehabilitation intervention with their implications for required sample sizes using WOMAC and SF-36 quality of life measurement instruments in patients with osteoarthritis of the lower extremities. *Arthritis Rheum* 45:384-391, 2001
5. Arola HM, Nicholls E, Mallen C, Thomas E: Self-reported pain interference and symptoms of anxiety and depression in community-dwelling older adults: Can a temporal relationship be determined? *Eur J Pain* 14:966-971, 2010
6. Axford J, Butt A, Heron C, Hammond J, Morgan J, Alavi A, Bolton J, Bland M: Prevalence of anxiety and depression in osteoarthritis: Use of the Hospital Anxiety and Depression Scale as a screening tool. *Clin Rheumatol* 29:1277-1283, 2010
7. Bellamy N: WOMAC Osteoarthritis Index: A User's Guide, London Health Services Centre, Ontario; 1996
8. Bellamy N, Campbell J, Haraoui B, Gerez-Simon E, Buchbinder R, Hobby K, MacDermid JC: Clinimetric properties of the AUSCAN Osteoarthritis Hand Index: An evaluation of reliability, validity and responsiveness. *Osteoarthritis Cartilage* 10:863-869, 2002
9. Bjelland I, Dahl AA, Haug TT, Neckelmann D: The validity of the hospital anxiety and depression scale. An updated literature review. *J Psychosom Res* 52:69-77, 2002
10. Brooks PM: Impact of osteoarthritis on individuals and society: How much disability? Social consequences and health economic implications. *Curr Opin Rheumatol* 14:573-577, 2002
11. Collins JE, Katz JN, Dervan EE, Losina E: Trajectories and risk profiles of pain in persons with radiographic, symptomatic knee osteoarthritis: Data from the osteoarthritis initiative. *Osteoarthritis Cartilage* 22:622-630, 2014
12. Dekker J, van Dijk GM, Veenhof C: Risk factors for functional decline in osteoarthritis of the hip or knee. *Curr Opin Rheumatol* 21:520-524, 2009
13. Doan L, Manders T, Wang J: Neuroplasticity underlying the comorbidity of pain and depression. *Neural Plast* 2015: 504691, 2015
14. Eyre HA, Air T, Proctor S, Rositano S, Baune BT: A critical review of the efficacy of non-steroidal anti-inflammatory drugs in depression. *Prog Neuropsychopharmacol Biol Psychiatry* 57:11-16, 2015
15. Felson DT, Zhang Y: Smoking and osteoarthritis: A review of the evidence and its implications. *Osteoarthritis Cartilage* 23:331-333, 2015
16. Flor H: New developments in the understanding and management of persistent pain. *Curr Opin Psychiatry* 25:109-113, 2012
17. Fulop T, Larbi A, Witkowski JM, McElhane J, Loeb M, Mitnitski A, Pawelec G: Aging, frailty and age-related diseases. *Biogerontology* 11:547-563, 2010
18. Gerrits MM, van Marwijk HW, van Oppen P, van der Horst H, Penninx BW: Longitudinal association between pain, and depression and anxiety over four years. *J Psychosom Res* 78: 64-70, 2015
19. Gibson SJ, Farrell M: A review of age differences in the neurophysiology of nociception and the perceptual experience of pain. *Clin J Pain* 20:227-239, 2004
20. Hawker GA, Gignac MA, Badley E, Davis AM, French MR, Li Y, Perruccio AV, Power JD, Sale J, Lou W: A longitudinal study to explain the pain-depression link in older adults with osteoarthritis. *Arthritis Care Res (Hoboken)* 63:1382-1390, 2011
21. Hilderink PH, Burger H, Deeg DJ, Beekman AT, Oude Voshaar RC: The temporal relation between pain and depression: Results from the Longitudinal Aging Study Amsterdam. *Psychosom Med* 74:945-951, 2012
22. Hutchings A, Calloway M, Choy E, Hooper M, Hunter DJ, Jordan JM, Zhang Y, Baser O, Long S, Palmer L: The Longitudinal Examination of Arthritis Pain (LEAP) study: Relationships between weekly fluctuations in patient-rated joint pain and other health outcomes. *J Rheumatol* 34:2291-2300, 2007
23. Jakobsson U, Hallberg IR: Pain and quality of life among older people with rheumatoid arthritis and/or osteoarthritis: A literature review. *J Clin Nurs* 11:430-443, 2002
24. Katon W, Lin EH, Kroenke K: The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *Gen Hosp Psychiatry* 29:147-155, 2007
25. Keefe FJ, Affleck G, France CR, Emery CF, Waters S, Caldwell DS, Stainbrook D, Hackshaw KV, Fox LC, Wilson K: The EPOSA research group: Germany: T. Nikolaus† (Principal Investigator), M. Denkinger, R. Peter, and F. Herbolsheimer; Italy: S. Maggi (Principal Investigator), S. Zambon, F. Limongi; M. Noale, and P. Siviero; The Netherlands (coordinating center): D.J.H. Deeg (Principal Investigator), S. van der Pas, N.M. van Schoor, L.A. Schaap, E.J. Timmermans, and P. Lips; Spain: Á. Otero (Principal Investigator), M.V. Castell, and M. Sanchez-Martinez; Sweden: N.L. Pedersen (Principal Investigator); United Kingdom: E.M. Dennison (Principal Investigator), C. Cooper, and M.H. Edwards.

Gender differences in pain, coping, and mood in individuals having osteoarthritic knee pain: A within-day analysis. *Pain* 110:571-577, 2004

26. Martin-Rodriguez E, Guillen-Grima F, Marti A, Brugos-Larumbe A: Comorbidity associated with obesity in a large population: The APNA study. *Obes Res Clin Pract* 9:435-447, 2015

27. McIlvane JM, Schiaffino KM, Paget SA: Age differences in the pain-depression link for women with osteoarthritis. Functional impairment and personal control as mediators. *Womens Health Issues* 17:44-51, 2007

28. Patetsos E, Horjales-Araujo E: Treating chronic pain with SSRIs: What do we know? *Pain Res Manag* 2016:2020915, 2016.

29. Penninx BW, Beekman AT, Ormel J, Kriegsman DM, Boeke AJ, van Eijk JT, Deeg DJ: Psychological status among elderly people with chronic diseases: Does type of disease play a part? *J Psychosom Res* 40:521-534, 1996

30. Puhan MA, Frey M, Buchi S, Schunemann HJ: The minimal important difference of the hospital anxiety and depression scale in patients with chronic obstructive pulmonary disease. *Health Qual Life Outcomes* 6:46, 2008

31. Sanders JB, Comijs HC, Bremmer MA, Deeg DJ, Beekman AT: A 13-year prospective cohort study on the effects of aging and frailty on the depression-pain relationship in older adults. *Int J Geriatr Psychiatry* 30:751-757, 2015

32. Schneider S, Junghaenel DU, Keefe FJ, Schwartz JE, Stone AA, Broderick JE: Individual differences in the day-to-day variability of pain, fatigue, and well-being in patients with rheumatic disease: Associations with psychological variables. *Pain* 153:813-822, 2012

33. Somers TJ, Keefe FJ, Godiwala N, Hoyler GH: Psychosocial factors and the pain experience of osteoarthritis patients: New findings and new directions. *Curr Opin Rheumatol* 21: 501-506, 2009

34. Stel VS, Pluijm SM, Deeg DJ, Smit JH, Bouter LM, Lips P: Functional limitations and poor physical performance as independent risk factors for self-reported fractures in older persons. *Osteoporos Int* 15:742-750, 2004

35. Stubbs B, Aluko Y, Myint PK, Smith TO: Prevalence of depressive symptoms and anxiety in osteoarthritis: A systematic review and meta-analysis. *Age Ageing* 45:228-235, 2016

36. Stubbs B, Binnekade TT, Soundy A, Schofield P, Huijnen IP, Eggermont LH: Are older adults with chronic musculoskeletal pain less active than older adults without pain? A systematic review and meta-analysis. *Pain Med* 14:1316-1331, 2013

37. Tang NK, Crane C: Suicidality in chronic pain: A review of the prevalence, risk factors and psychological links. *Psychol Med* 36:575-586, 2006

38. Theis KA, Helmick CG, Hootman JM: Arthritis burden and impact are greater among U.S. women than men: Intervention opportunities. *J Womens Health (Larchmt)* 16:441-453, 2007

39. Twisk J, de Vente W: Attrition in longitudinal studies. How to deal with missing data. *J Clin Epidemiol* 55:329-337, 2002

40. van der Pas S, Castell MV, Cooper C, Denkinger M, Dennison EM, Edwards MH, Herbolzheimer F, Limongi F, Lips P, Maggi S, Nasell H, Nikolaus T, Otero A, Pedersen NL, Peter R, Sanchez-Martinez M, Schaap LA, Zambon S, van Schoor NM, Deeg DJ: European project on osteoarthritis: Design of a six-cohort study on the personal and societal burden of osteoarthritis in an older European population. *BMC Musculoskelet Disord* 14:138, 2013

41. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, Aboyans V, Abraham J, Ackerman I, Aggarwal R, Ahn SY, Ali MK, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Bahalim AN, Barker-Collo S, Barrero LH, Bartels DH, Basanez MG, Baxter A, Bell ML, Benjamin EJ, Bennett D, Bernabe E, Bhalla K, Bhandari B, Bikbov B, Bin Abdulhak A, Birbeck G, Black JA, Blencowe H, Blore JD, Blyth F, Bolliger I, Bonaventure A, Boufous S, Bourne R, Boussinesq M, Braithwaite T, Brayne C, Bridgett L, Brooker S, Brooks P, Brugha TS, Bryan-Hancock C, Bucello C, Buchbinder R, Buckle G, Budke CM, Burch M, Burney P, Burstein R, Calabria B, Campbell B, Canter CE, Carabin H, Carapetis J, Carmona L, Cella C, Charlson F, Chen H, Cheng AT, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, de Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahiya M, Dahodwala N, Damsere-Derry J, Danaei G, Davis A, De Leo D, Degenhardt L, Dellavalle R, Delossantos A, Denenberg J, Derrett S, Des Jarlais DC, Dharmaratne SD, Dherani M, Diaz-Torne C, Dolk H, Dorsey ER, Driscoll T, Duber H, Ebel B, Edmond K, Elbaz A, Ali SE, Erskine H, Erwin PJ, Espindola P, Ewoigbokhan SE, Faragad F, Feigin V, Felson DT, Ferrari A, Ferri CP, Fevre EM, Finucane MM, Flaxman S, Flood L, Foreman K, Forouzanfar MH, Fowkes FG, Franklin R, Fransen M, Freeman MK, Gabbe BJ, Gabriel SE, Gakidou E, Ganatra HA, Garcia B, Gaspari F, Gillum RF, Gmel G, Gosselin R, Grainger R, Groeger J, Guillemin F, Gunnell D, Gupta R, Haagsma J, Hagan H, Halasa YA, Hall W, Haring D, Haro JM, Harrison JE, Havmoeller R, Hay RJ, Higashi H, Hill C, Hoen B, Hoffman H, Hotez PJ, Hoy D, Huang JJ, Ibeanusi SE, Jacobsen KH, James SL, Jarvis D, Jasrasaria R, Jayaraman S, Johns N, Jonas JB, Karthikeyan G, Kassebaum N, Kawakami N, Keren A, Khoo JP, King CH, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Laloo R, Laslett LL, Lathlean T, Leasher JL, Lee YY, Leigh J, Lim SS, Limb E, Lin JK, Lipnick M, Lipshultz SE, Liu W, Loane M, Ohno SL, Lyons R, Ma J, Mabweijano J, Macintyre MF, Malekzadeh R, Mallinger L, Manivannan S, Marcenes W, March L, Margolis DJ, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGill N, McGrath J, Medina-Mora ME, Meltzer M, Mensah GA, Merriman TR, Meyer AC, Miglioli V, Miller M, Miller TR, Mitchell PB, Mocumbi AO, Moffitt TE, Mokdad AA, Monasta L, Montico M, Moradi-Lakeh M, Moran A, Morawska L, Mori R, Murdoch ME, Mwaniki MK, Naidoo K, Nair MN, Naldi L, Narayan KM, Nelson PK, Nelson RG, Nevitt MC, Newton CR, Nolte S, Norman P, Norman R, O'Donnell M, O'Hanlon S, Olives C, Omer SB, Ortblad K, Osborne R, Ozgediz D, Page A, Pahari B, Pandian JD, Rivero AP, Patten SB, Pearce N, Padilla RP, Perez-Ruiz F: Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380:2163-2196, 2012

42. Wang LP, Grimm KJ: Investigating reliabilities of intraindividual variability indicators. *Multivariate Behav Res* 47:771-802, 2012

43. Yohannes AM, Caton S: Management of depression in older people with osteoarthritis: A systematic review. *Aging Ment Health* 14:637-651, 2010

44. Zigmond AS, Snaith RP: The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 67:361-370, 1983