

VU Research Portal

Risk factors for mental disorder development in asylum seekers and refugees resettled in Western Europe and Turkey

Barbui, Corrado; Tedeschi, Federico; Acarturk, Ceren; Anttila, Minna; Au, Teresa; Baumgartner, Josef; Carswell, Ken; Churchill, Rachel; Cuijpers, Pim; Karyotaki, Eirini; Klein, Thomas; Koesters, Markus; Lantta, Tella; Nosè, Michela; Ostuzzi, Giovanni; Pasquini, Massimo; Prina, Eleonora; Sijbrandij, Marit; Tarsitani, Lorenzo; Turrini, Giulia

published in

International Journal of Social Psychiatry
2023

DOI (link to publisher)

[10.1177/00207640221132430](https://doi.org/10.1177/00207640221132430)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Barbui, C., Tedeschi, F., Acarturk, C., Anttila, M., Au, T., Baumgartner, J., Carswell, K., Churchill, R., Cuijpers, P., Karyotaki, E., Klein, T., Koesters, M., Lantta, T., Nosè, M., Ostuzzi, G., Pasquini, M., Prina, E., Sijbrandij, M., Tarsitani, L., ... Purgato, M. (2023). Risk factors for mental disorder development in asylum seekers and refugees resettled in Western Europe and Turkey: Participant-level analysis of two large prevention studies. *International Journal of Social Psychiatry*, 69(3), 664-674. <https://doi.org/10.1177/00207640221132430>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Risk factors for mental disorder development in asylum seekers and refugees resettled in Western Europe and Turkey: Participant-level analysis of two large prevention studies

International Journal of
Social Psychiatry
2023, Vol. 69(3) 664–674
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/00207640221132430
journals.sagepub.com/home/isp


Corrado Barbui¹, Federico Tedeschi¹, Ceren Acarturk²,
Minna Anttila³, Teresa Au⁴, Josef Baumgartner⁵, Ken Carswell⁴,
Rachel Churchill⁶, Pim Cuijpers^{7,8,9}, Eirini Karyotaki^{7,8},
Thomas Klein¹⁰, Markus Koesters¹⁰, Tella Lantta³, Michela Nosè¹,
Giovanni Ostuzzi¹, Massimo Pasquini¹¹, Eleonora Prina¹,
Marit Sijbrandij^{7,8}, Lorenzo Tarsitani¹¹, Giulia Turrini¹,
Ersin Uygun¹², Maritta Välimäki^{3,13}, Lauren Walker¹⁴,
Johannes Wancata⁵, Ross G. White¹⁵ and Marianna Purgato¹

Abstract

Background: In asylum seekers and refugees, the frequency of mental disorders, such as depression, anxiety and post-traumatic stress disorder, is higher than the general population, but there is a lack of data on risk factors for the development of mental disorders in this population.

Aim: This study investigated the risk factors for mental disorder development in a large group of asylum seekers and refugees resettled in high- and middle-income settings.

Methods: Participant-level data from two randomized prevention studies involving asylum seekers and refugees resettled in Western European countries and in Turkey were pooled. The two studies randomized participants with psychological distress, but without a diagnosis of mental disorder, to the Self-Help Plus psychological intervention or enhanced care as usual. At baseline, exposure to potentially traumatic events was measured using the Harvard Trauma Questionnaire-part I, while psychological distress and depressive symptoms were assessed with the General Health Questionnaire and the Patient Health Questionnaire. After 3 and 6 months of follow-up, the proportion of participants who developed a mental disorder was calculated using the Mini International Neuropsychiatric Interview.

Results: A total of 1,101 participants were included in the analysis. At 3- and 6-month follow-up the observed frequency of mental disorders was 13.51% (115/851) and 24.30% (207/852), respectively, while the frequency estimates after missing data imputation were 13.95% and 23.78%, respectively. After controlling for confounders, logistic regression analysis showed that participants with a lower education level ($p=.034$), a shorter duration of journey ($p=.057$) and arriving from countries with war-related contexts ($p=.017$), were more at risk of developing mental disorders. Psychological distress ($p=.004$), depression ($p=.001$) and exposure to potentially traumatic events ($p=.020$) were predictors of mental disorder development.

Conclusions: This study identified several risk factors for the development of mental disorders in asylum seekers and refugees, some of which may be the target of risk reduction policies. The identification of asylum seekers and refugees at increased risk of mental disorders should guide the implementation of focused preventative psychological interventions.

Keywords

Asylum seekers, refugees, migrants, mental health, mental disorders, high-income countries, low-income countries

Introduction

According to the last United Nations Refugee Agency data, despite COVID-related movement restrictions, at the end of 2020 more than 82 million people were forcibly

displaced as a result of persecution, conflict, violence and other human rights violations (United Nations High Commissioner for Refugees [UNHCR], 2022). Of these, more than 30 million were asylum seekers or refugees. While Turkey continued to host the largest number of

refugees, most of whom were Syrian refugees, 10% of all the world's refugees were resettled in Europe at the end of 2020 (UNHCR, 2022). These numbers are expected to further increase, as recent estimates have suggested that over 6.5 million refugees from Ukraine have crossed to Poland, Hungary, Romania, Moldova and other countries (UNHCR, 2022).

Despite heterogeneity between the studies and the populations of refugees and asylum seekers assessed (Giacco, 2019), a large body of evidence consistently showed that the frequency of mental disorders in refugees and asylum seekers is increased as compared with the general population. Existing World Health Organization (WHO) estimates suggest a prevalence of 13.0% for mild forms of depression, anxiety and post-traumatic stress disorder (PTSD), and 4.0% for moderate forms (Charlson et al., 2019). Other reviews of prevalence studies found that adult refugees and asylum seekers have high and persistent rates of depression and PTSD, while the prevalence of anxiety disorders and psychosis are more comparable to findings from general populations (Blackmore et al., 2020; Henkelmann et al., 2020).

The increased frequency of mental disorders in refugees and asylum seekers may be related to the refugee experience, which is characterized by loss of homes, hopes, possessions and disruption of personal, family and professional life projects. In addition, a wide range of physical, psychological and psychosocial problems associated with adversities may occur, such as bombings, threats, captivity, torture, injury, witnessing death or injury of loved ones, discrimination, economic stress and uncertainty about the future (Priebe et al., 2016). Interestingly, while these factors have been studied as determinants of poor mental health outcomes, or as determinants of

psychological distress (Jannesari et al., 2020), to our knowledge they have never been investigated in this population as risk factors for the development of mental disorders considered as full-blown diagnostic entities (Gleeson et al., 2020; Hajak et al., 2021). Knowledge of determinants of the development of mental disorders is particularly important to implement policies aiming at decreasing exposure to such determinants, to improve identification of refugees and asylum seekers at increased risk of developing mental disorders, and to plan the provision of psychological interventions to individuals and communities at risk.

Aiming to test the effect of Self Help Plus (SH+), a low intensity, group-based, self-help psychological intervention recently developed by the WHO (Epping-Jordan et al., 2016; WHO, 2021a), in reducing the frequency of mental disorders, we conducted two large randomized prevention trials in asylum seekers and refugees, one in Western Europe and another in Turkey (Purgato et al., 2019). The studies followed the same research protocol and were conducted in parallel (Acarturk et al., 2022; Purgato et al., 2021). Both the Western European and Turkey studies showed evidence of an effect of SH+ in preventing the onset of mental disorders, but differences were observed between the studies. The effect was much more pronounced for the Turkey study where efficacy (i.e. reducing the frequency of any mental disorder) was observed at 6 months, compared to the Western European study where a preventative effect was only found post-intervention and not after 6 months (Acarturk et al., 2022; Purgato et al., 2021). As these are the only two studies that enrolled asylum seekers and refugees without any mental disorders at baseline, and that assessed the frequency of mental disorders at follow-up as primary

¹WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation, Department of Neuroscience, Biomedicine and Movement Sciences, Section of Psychiatry, University of Verona, Italy

²Department of Psychology, Koc University, Istanbul, Turkey

³University of Turku, Department of Nursing Science, University of Turku, Finland

⁴Department of Mental Health and Substance Use, World Health Organization, Geneva, Switzerland

⁵Clinical Division of Social Psychiatry, Department of Psychiatry and Psychotherapy, Medical University of Vienna, Austria

⁶Centre for Reviews and Dissemination, University of York, UK

⁷Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health research institute, Vrije Universiteit Amsterdam, The Netherlands

⁸WHO Collaborating Centre for Research and Dissemination of Psychological Interventions, Vrije Universiteit Amsterdam, The Netherlands

⁹Babeş-Bolyai University, International Institute for Psychotherapy, Cluj-Napoca, Romania

¹⁰Department of Psychiatry II, Ulm University, Germany

¹¹Department of Human Neurosciences, Sapienza University of Rome, Italy

¹²Trauma and Disaster Mental Health, Istanbul Bilgi University, Turkey

¹³Xiangya Nursing School, Central South University, Xiangya Center for Evidence-Based Practice and Healthcare Innovation, Changsha, China

¹⁴Department of Health Sciences, University of York, UK

¹⁵School of Psychology, Queen's University Belfast, UK

Corresponding author:

Professor Corrado Barbui, WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation, Ospedale Policlinico GB Rossi, Piazzale Scuro 10, Verona 37100, Italy.

Email: corrado.barbui@univr.it

outcome (Papola et al., 2020), they offered a unique opportunity to prospectively investigate the risk factors for the onset of mental disorders in a large group of asylum seekers and refugees.

Methods

Participants and measures

Participant-level data from two randomized prevention trials involving asylum seekers and refugees resettled in Western European countries (Austria, Finland, Germany, Italy and UK) and in Turkey were pooled (Acarturk et al., 2022; Purgato et al., 2021). In both studies, participants were randomly assigned to the SH+ psychological intervention, consisting of SH+ combined with enhanced care as usual (ECAU), or to ECAU only. The two studies were conducted in parallel following the same study design (Purgato et al., 2019). The protocol of the present study was registered within the Open Science Framework (<https://osf.io/37h5n>).

In both studies, participants were included if they met the following criteria: (a) aged 18 years or older; (b) able to speak and understand Arabic, Dari, English or Urdu; (c) being under temporary protection with a refugee or asylum seeker status; (d) experiencing psychological distress, as shown by a score of 3 or more on the 12-item dichotomously-scored General Health Questionnaire (el-Rufaie & Daradkeh, 1996; Goldberg et al., 1998; Kilic et al., 1997); (e) having completed oral and written informed consent to enter the study. Exclusion criteria were: (a) presence of any mental disorder according to the Mini International Neuropsychiatric Interview (M.I.N.I.), a brief structured diagnostic interview for the major psychiatric disorders in DSM-III-R, DSM-IV and DSM-5 and ICD-10 (Kadri et al., 2005; Sheehan et al., 1998); (b) evidence of acute medical conditions contraindicating study participation; (c) evidence of imminent suicide risk, or suicide risk scored as 'moderate or high' on the M.I.N.I.; (d) signs of impaired decision-making capacity emerging from responses during the clinical interview.

The M.I.N.I. was administered before random allocation, in order to exclude participants with a mental disorder, and at 3 and 6 months of follow-up, in order to calculate the proportion of participants who developed a mental disorder. Exposure to potentially traumatic events was measured at baseline using the Harvard Trauma Questionnaire-part I (HTQ; Mollica et al., 1992). The HTQ-part I covers a variety of trauma events that may affect refugee mental health, and the scoring represents the number of different types of traumatic events experienced by the participants (higher score is associated with high number of traumatic events). In addition to using the HTQ total score, we identified subtypes of traumatic events performing a Principal Components analysis (PCA).

Mental health symptoms were measured using the following validated instruments. The GHQ-12 questionnaire was used to measure psychological distress (el-Rufaie & Daradkeh, 1996; Goldberg et al., 1998; Kilic et al., 1997), while PTSD was assessed with the PTSD Checklist for DSM-5 (PCL-5), a 20-item questionnaire that measures overall PTSD symptoms (score zero to 80) and symptoms by cluster (intrusions, avoidance, negative changes in thoughts and mood and changes in arousal), with higher scores indicating higher levels of PTSD symptoms (Blanchard et al., 1996). Levels of self-reported depression symptoms were measured with the Patient Health Questionnaire, nine-item version (PHQ-9) which has a 4-point scale (score 0–27; Kroenke et al., 2001). All measures were collected at baseline before random allocation, and after 3 and 6 months of follow-up. Assessors were trained in the administration of rating scales, instructed on how to perform follow-up assessments while preserving effective masking, and assisted by cultural mediators when needed.

Interventions

The SH+ intervention was developed by the WHO, as described elsewhere (Acarturk et al., 2022; Purgato et al., 2021), and is now publicly available (WHO, 2021a). SH+ consists of a pre-recorded audio course complemented with an illustrated self-help book. The book has been recently updated and published by WHO as *Doing What Matters in Times of Stress* (WHO, 2021b). All SH+ materials were adapted for the cultural groups included in the studies. The SH+ pre-recorded audio material was delivered across five 2-hour sessions to groups of up to 30 people. The audio material imparts key information about stress management and guides participants through individual exercises and small group discussions. To augment the audio recordings, an illustrated self-help book reviews all essential content and concepts. The SH+ intervention was fully delivered in the language of participants by trained facilitators with a migration background, who were native speakers of the target languages. ECAU, provided both to the experimental and control group, consisted of routinely delivered social support and/or care according to local regulations. Participants in the ECAU arm received the same baseline and follow-up assessments of the intervention arm, according to the study schedule (around 3 and 6 months after randomization), information about freely available health and social services and links to community networks providing support to refugees and asylum seekers.

Statistical analysis

Descriptive statistics (M and SD for continuous variables and absolute numbers and percentages for dichotomous variables) were computed on sociodemographic, premigration, migration and postmigration variables at baseline and for clinical variables.

Participants who met criteria for any mental disorder on the M.I.N.I. at 3- or 6-month follow-up were considered cases with a mental disorder. Multiple imputation was adopted to address the issue of missing data in all the variables included in the model. In particular, in case of missing data at the M.I.N.I. in one timepoint, imputation was performed on single M.I.N.I. values at 3 and 6 months. In case of missing values on continuous clinical measures, the imputation was performed on single item scales. Specifically, imputation followed the approach reported by Plumptre and colleagues (Plumptre et al., 2016), that is we used scale totals within prediction equations and, for imputations of responses to individual scale items, we additionally included the responses to the other scale items, using the 'ice' Stata routine (Royston, 2005; Sterne et al., 2009), and considering single-item scores as ordered categorical variables. M.I.N.I. values at the two time-points were used in the prediction equations of regression predictors, and upper and lower bounds were set for continuous variables with missing values as appropriate. The number of imputed samples was determined by following the rule of thumb suggested by I. R. White et al. (2011), that is: 'at least equal to the percentage of incomplete cases'. We rounded such number to the nearest multiple of 10 above.

In order to identify subtypes of traumatic events, we performed a PCA on tetrachoric correlations of HTQ items, with Quartimin oblique rotation to allow for between-factor non-null correlations (Tabachnick, 2007), and summing items with loadings above 0.40 on the same factor to create scores for the regression model (Stevens, 2002). The number of factors was determined by adopting Kaiser's rule (Kaiser, 1960), that is using the cut-off scores of 1 for the eigenvalues.

In order to investigate predictors of the development of mental disorders, unadjusted and adjusted logistic regression analyses were performed, using the frequency of participants with a mental disorder in at least one timepoint as binary dependent variable. The following independent variables were inserted into the model: age (years), gender (men, women), education (years), unemployment (yes/no), country of origin (Syria, Iraq, Nigeria, other countries), length of journey (below 1 month, between 1 and 3 months and above 3 months), study (Western Europe vs. Turkey), length of stay in the resettlement country (months), HTQ (total score at baseline), GHQ-12 (total score at baseline), PCL-5 (total score at baseline), PHQ-9 (total score at baseline), number of SH+ sessions received (0–5). Considering that PTSD and depression are known to be interrelated experiences following trauma (Rytwinski et al., 2013), and considering that the PCL-5 largely overlaps with the HTQ (Patel et al., 2022), and that a substantial overlap exists between the PCL-5 and the PHQ-9 (Dabrowski et al., 2020), we excluded the PCL-5 from the final model, but it was included (total score at baseline) in the imputation model.

In order to check the robustness of the analysis, we re-ran the model after excluding the arm receiving SH+. As a further sensitivity analysis, in order to estimate within-centre effects, we performed a model including a fixed effect for recruiting centre. As a subgroup analysis, we re-ran the logistic regression analysis separately for each study sample (Western Europe vs. Turkey). All analyses were performed using Stata 17 (Statacorp, 2017).

Results

Participants

The two prevention trials randomized a total of 1,101 participants. After 6 months, 249 participants (22.6%) were not available for follow-up assessments, for the reasons reported in Figure 1. Participant characteristics at baseline are presented in Table 1. Almost half were female, with a mean age of 32 ($SD=9.521$) years, and a mean education of 10 years ($SD=4.424$). Slightly less than 40% was unemployed. The migration journey lasted more than 3 months in around one third of participants, in another one third it lasted between 1 and 3 months, while in the remaining 40% the host country was reached in less than 1 month. On average, the mean length of stay in the resettlement country was slightly more than 3 years. Most of the participants were from Syria, Nigeria and Iraq (Table 1). Participants attended a mean number of 1.4 ($SD=2.015$) SH+ sessions. The mean (SD) baseline scores on the measures of interest are reported in Table 1.

Frequency and determinants of mental disorders

At 3- and 6-month follow-up the observed frequency of mental disorders, as measured with the M.I.N.I., was 13.51% (115/851) and 24.30% (207/852), respectively, while the frequency estimates after missing data imputation were 13.95% and 23.78%, respectively. The majority of detected mental disorders were major depressive disorders (9.5% and 20.3% after 3 and 6 months), PTSD (3.2% and 6.7%) and anxiety disorders (2.1% and 4.1%; Figure 2).

The PCA model found a three-factor solution where the set of items was exhaustive and mutually exclusive, with the HTQ items grouped as follows: factor 1 – lack of basic needs: lack of food or water, ill health without access to medical care, lack of shelter; factor 2 – violence and abuse: imprisonment, serious injury, brain washing, rape or sexual abuse, forced isolation from others, forced separation from family members, torture, other (e.g. domestic violence); factor 3 – being close to death: combat situation, being close to death, murder of family or friend, unnatural death of family or friend, murder of stranger(s), lost or kidnapped. The factor loadings of the




Figure 1. Flow-diagram of participant enrolment and follow-up.

PCA-model with Quartimin oblique rotation are reported in the Supplemental Materials.

The results of unadjusted and adjusted logistic regression analyses investigating factors associated with the development of mental disorders are presented in Table 2. Participants with lower education level ($p = .034$), a shorter duration of journey ($p = .057$) and arriving from countries with war-related contexts (Iraq, Syria; $p = .017$), were those more at risk of developing mental disorders. In terms of mental health symptoms and exposure to traumatic events, psychological distress ($p = .004$), depressive symptoms ($p = .001$) and HTQ total score ($p = .020$) at baseline were predictors of mental disorder development, as was the HTQ factor violence and abuse (imprisonment, serious injury, brain washing, rape or sexual abuse, forced isolation from others, forced separation from family members, torture and domestic violence; $p = .008$). In terms of post-migration factors, the number of SH+ sessions ($p < .001$) was inversely associated with the risk of developing mental disorders (Table 2).

Secondary logistic regression analyses including only the ECAU arms of the two studies, and analysing the two studies separately, confirmed the role of mental health symptoms and exposure to traumatic events as risk factors for mental disorders (Supplemental Materials). However, due to lower statistical power, some factors lost significance despite a similar or even higher estimated effect, as for example the HTQ in the model restricted to the Western European study, or years of

education and travel duration in the model restricted to the ECAU sample, and GHQ-12 in the model restricted to the study conducted in Turkey (Supplemental Materials).

Consistently with the results of the two studies, the number of SH+ sessions was inversely associated with the frequency of mental disorders only in the study conducted in Turkey (Supplemental Materials). The inclusion of recruiting centres as predictor variables revealed that, in comparison with participants recruited in Istanbul, participants recruited in Vienna were less likely to develop a mental disorder (Supplemental Materials).

Discussion

To the best of our knowledge, this is the first prospective study that examined risk factors for the onset of mental disorders among asylum seekers and refugees without a mental disorder at baseline. We showed the significant influence of socio-demographic, clinical and contextual factors, including potentially traumatic events, on the development of mental disorders in asylum seekers and refugees resettled in Western European countries and in Turkey.

Overall, one in four refugees and asylum seekers developed a diagnosable mental disorder over a period of 6 months. Interestingly, the most frequently reported diagnostic group was depression, followed by PTSD and anxiety disorders. These figures, derived from two intervention

Table 1. Participants characteristics at baseline.

Variables	M (SD)
Age (years)	31.797 (9.521)
Years of education	9.556 (4.424)
Months in host country	40.636 (32.599)
Host country GDP in \$1,000 (2019)	
Austria	58,091.3
Finland	50,321.6
Germany	55,652.9
Italy	44,334.2
Turkey	27,144.2
UK	49,070.3
Number of SH+ sessions	1.434 (2.015)
GHQ	5.737 (2.230)
PHQ-9	7.232 (5.230)
HTQ total score	5.795 (4.349)
HTQ lack of basic needs	1.277 (1.210)
HTQ violence and abuse	1.753 (2.085)
HTQ being close to death	2.766 (1.947)
Frequencies	n/N (%)
Gender (female)	538/1,101 (48.86)
Unemployed	418/1,099 (38.03)
Country of origin	
Nigeria	114/1,100 (10.36)
Syria	758/1,100 (68.91)
Iraq	94/1,100 (8.55)
Other	134/1,100 (12.18)
Travel duration	
Less than 1 month	277 (40.44)
One to 3 months	203 (29.64)
More than 3 months	205 (29.93)
Study	
Western Europe	459 (41.69)
Turkey	642 (58.31)

Note. M=mean; SD=standard deviation; SH+ =self-help plus; GDP=gross domestic product; GHQ=General Health Questionnaire; PHQ=Patient Health Questionnaire; HTQ=Harvard Trauma Questionnaire.

studies, cannot be compared with those from epidemiological studies, which generally show similar frequencies for depression and PTSD, or slightly higher frequencies for depression, depending on the population surveyed and the study setting (Blackmore et al., 2020; Charlson et al., 2019; Henkelmann et al., 2020; Hoell et al., 2021). In migrants exposed to armed conflict, by contrast, the frequency of PTSD was found to be higher than depression (31% vs. 25%; Mesa-Vieira et al., 2022).

The finding that educational level is inversely associated with the development of mental disorders expands previous data collected in the general population showing that higher educational level seems to have a protective effect against anxiety and depression symptoms, and against common mental disorders in general (Araya et al., 2003; Bjelland et al., 2008). In the general population, poor education has been suggested to be a marker of lack

of opportunities and resources, including material and psychological resources, and a marker of childhood adversity (Araya et al., 2003). In asylum seekers and refugees, these factors may directly contribute to the emergence of mental disorders, or may interfere with the coping skills required to deal with all the adversities associated with the migration and resettlement process, which, in turn, may increase the risk of developing mental disorders (Kirmayer et al., 2011).

Consistently with an extensive literature showing that exposure to potentially traumatic events represents a risk factor for poor mental health and well-being (Gleeson et al., 2020; Hajak et al., 2021; Priebe et al., 2016), the present study adds that exposure to potentially traumatic events is also a risk factor for the development of mental disorders considered as full-blown diagnostic entities. The finding that being displaced from countries with war-related contexts emerged as an additional risk factor for mental disorders further corroborates this association. War-related contexts imply exposure to multiple and serious traumatic events, and the exposure to combat situations creates the risk of witnessing violence and/or death and experiencing physical and psychological violence (Crepet et al., 2017). Extensive literature shows that these situations are associated with poor mental health outcomes, including PTSD, depression, anxiety and somatization symptoms (Jongedijk et al., 2020; Knipscheer et al., 2015; Nickerson et al., 2021).

The present study has limitations and strengths. A first limitation is that the population enrolled in the two trials cannot be considered representative of the general population of asylum seekers and refugees, as we selected participants scoring above a threshold of psychological distress, and we excluded those at suicide risk. During the studies, in addition, around half of the participants was exposed to a psychological intervention aimed at preventing the development of mental disorders. For these reasons, the overall frequency estimate of mental disorders cannot apply to the general population of asylum seekers and refugees resettled in Western European countries or in Turkey. A second issue is that we did not calculate a true incidence of mental disorders, but only frequency figures at two time points with a structured diagnostic interview that might overestimate the true frequency of some mental disorders, such as depression (Wu et al., 2020). Third, the SARS-CoV2 pandemic impacted the study procedures, because in all the recruiting sites follow-up assessments were conducted using online tools instead of face-to-face meetings. Although assessors were trained in the administration of rating scales, instructed on how to perform online follow-up assessments, and assisted by cultural mediators when needed, it is unknown whether this may have impacted the responses of participants to the instruments. Related to this, even though several studies documented that a careful and culturally appropriate use of available instruments is




Figure 2. Frequency of mental disorders, as measured with the M.I.N.I., after 3 and 6 months of follow-up.

feasible and allows a standardization of the screening process and a systematic recognition of psychological distress and psychiatric diagnoses (Acarturk et al., 2021), we acknowledge that formal studies on use of these tools in refugee groups are lacking. Fourth, despite a growing body of literature showed that the duration of the asylum procedure is an important risk factor for mental health conditions (Laban et al., 2004; Winkler et al., 2019), we acknowledge that this information was not collected. However, months in host country, which may be considered a proxy of length of asylum procedure, was included in the model but did not emerge as a significant factor.

Despite these limitations, there are strengths that should be emphasized. The main strength is the exclusion of participants with a mental disorder at baseline, and the choice of a dependent variable that is fully consistent with a preventative design, namely the frequency of mental disorders at follow-up, assessed with validated measures. This design allowed to prospectively investigate the determinants of mental disorders rather than of poor mental health, as previous studies have done in populations who might already be with mental disorders at baseline (Priebe et al., 2016). Another strength is a sample size of over a thousand of asylum seekers and refugees, resettled in a variety of Western European sites and in Turkey. This aspect is of relevance not only in terms of statistical power, but also in terms of generalizability and applicability of study findings to different types of reception settings in high-income and middle-income countries. Lastly, despite an attrition rate of around 20%, a follow-up assessment of 6 months is noteworthy in such a difficult-to-follow population, who is

often moved from one reception site to another, and may not perceive mental health as a priority, having a number of other challenging concerns such as housing, unstable working conditions, management of visa issues, safety of family members, fear of being returned to home country, plans to move to another country or to another location.

The present study has important practical implications. The finding of a positive relationship between exposure to potentially traumatic events and risk of mental disorder development suggests a pressing need of developing policies aiming to decrease exposure to such traumatic experiences after resettlement. Host countries may have opportunities to decrease exposure to situations such as material and economic hardship that could affect integrity, independence, dignity and well-being (financial strain), social hardship due to loss of status (social strain), feelings of inadequacy in relation with specific skills needed in the host-country to successfully function in daily life (competency strain), experiences of unfair treatment on the basis of prejudice (perceived discrimination). This ambitious goal may be achieved by implementing reception conditions that optimize internationally recognized minimal quality standards. For example, standards for the reception of applicants for international protection have been established by Directive 2013/33/EU of the European Parliament. The Directive clearly reports that national authorities should ensure that reception modalities are specifically designed to meet the needs of persons requiring international protection, including legal assistance, document provision, material support, links with local communities, freedom of movement, information about labour

Table 2. Determinants of mental disorder development: logistic regression analyses.

	Unadjusted regressions		Adjusted regressions			
	Odds ratio [95% CI]	<i>p</i> -Value	With HTQ total score		With HTQ factors	
			Odds ratio [95% CI]	<i>p</i> -Value	Odds ratio [95% CI]	<i>p</i> -Value
Age in years	1.001 [0.986; 1.015]	.920	0.996 [0.980; 1.013]	.672	0.998 [0.981; 1.015]	.830
Female gender	1.318 [0.987; 1.759]	.061	0.962 [0.680; 1.360]	.825	0.977 [0.689; 1.385]	.896
Years of education	0.972 [0.941; 1.004]	.088	0.959 [0.923; 0.997]	.034	0.959 [0.922; 0.996]	.032
Unemployed	0.938 [0.702; 1.253]	.664	1.171 [0.832; 1.647]	.365	1.180 [0.836; 1.664]	.347
Country of origin						
Syria	Reference category	<.001	Reference category	.017	Reference category	.006
Iraq	1.180 [0.710; 1.960]		1.592 [0.807; 3.142]		1.672 [0.840; 3.325]	
Nigeria	0.209 [0.101; 0.432]		0.376 [0.142; 0.997]		0.318 [0.118; 0.861]	
Other	0.403 [0.240; 0.674]		0.577 [0.273; 1.222]		0.519 [0.241; 1.114]	
Travel duration (months)						
Less than 1 month	Reference category	<.001	Reference category	.057	Reference category	.057
One to 3 months	0.616 [0.398; 0.952]		0.653 [0.414; 1.028]		0.658 [0.417; 1.041]	
More than 3 months	0.312 [0.198; 0.494]		0.512 [0.267; 0.983]		0.502 [0.260; 0.969]	
Months in host country	1.002 [0.997; 1.007]	.503	1.000 [0.994; 1.006]	.943	1.000 [0.994; 1.006]	.973
Turkish trial	2.058 [1.513; 2.800]	<.001	1.795 [0.997; 3.235]	.051	1.802 [0.998; 3.253]	.051
Number of SH+ sessions	0.888 [0.827; 0.953]	.001	0.853 [0.788; 0.923]	<.001	0.850 [0.785; 0.921]	<.001
GHQ-12	1.188 [1.114; 1.268]	<.001	1.117 [1.036; 1.204]	.004	1.119 [1.038; 1.206]	.003
PHQ-9	1.071 [1.042; 1.100]	<.001	1.061 [1.026; 1.097]	.001	1.061 [1.026; 1.097]	.001
HTQ total score	1.002 [0.970; 1.036]	.883	1.051 [1.008; 1.096]	.020	–	–
HTQ Factor 1 – lack of basic needs	0.999 [0.890; 1.122]	.988	–	–	1.007 [0.863; 1.176]	.928
HTQ Factor 2 – violence and abuse	0.995 [0.929; 1.066]	.894	–	–	1.163 [1.041; 1.299]	.008
HTQ Factor 3 – being close to death	1.018 [0.947; 1.095]	.769	–	–	0.978 [0.888; 1.077]	.650

Note. CI = confidence interval; SH+ = self-help plus; GDP = gross domestic product; GHQ = General Health Questionnaire; PHQ = Patient Health Questionnaire; HTQ = Harvard Trauma Questionnaire. Statistically significant results are reported in bold.

market access, vocational training, social support. Health care, including mental health care, is also mentioned as a key intervention where needed. It would be important to ensure that efforts to support forcibly displaced people are coordinated across the different layers of the social environments in which they are hosted, that is at the level of the individual, their family, the community and the institutions that have governance responsibility for their care and support (R. G. White & Van der Boor, 2021).

In addition to implications in terms of risk reduction policies, the present study has implications in terms of provision of psychological interventions aimed at preventing the development of mental disorders (Miller et al., 2021). As the implementation of preventative psychological interventions to the whole population of asylum seekers and refugees may not be sustainable by host countries, national authorities may consider to offer psychological interventions to at-risk populations. The finding that persons with lower education level, a shorter duration of journey, arriving from countries

with war-related contexts (Iraq, Syria), and with high level of psychological distress and depressive symptoms, are those more at risk of developing mental disorders, appears to identify a target population for focused psychological prevention interventions. The WHO has recently developed a number of low-intensity psychological interventions that may be scaled up as public health strategies to address mental disorders and psychological distress in refugee populations exposed to ongoing adversities (WHO, 2017). In addition to WHO interventions, other psychological treatments have been shown to be effective in alleviating psychological symptoms in asylum seekers and refugees (Turrini et al., 2021; Uphoff et al., 2020), but a preventative effect has been investigated for SH+ only. The present results, by showing an inverse association between number of SH+ sessions and risk of developing a mental disorder, further corroborate the value of SH+ as a prevention intervention. The finding that trauma exposure stands out as a predictor for development of mental disorders would

additionally suggest that interventions may need to include a focus on traumatic memories and experiences, in addition to being focused on low mood.

In parallel with the provision of psychological interventions to at-risk populations, national authorities should ensure regular access to such interventions supporting at risk populations to engage through reducing barriers that might prevent or limit access or use (Fuhr et al., 2019). This can be achieved by optimizing access (e.g. accessible services and supports) to a range of interventions depending on need (e.g. stepped care), and by using different delivery mechanisms (e.g. digital or peer delivered support).

As the number of persons in need of protection is likely to substantially increase globally, driven by long-lasting wars as well as by new conflicts such as the Russo-Ukrainian war which broke out recently (Barbui et al., 2022; Pandi-Perumal et al., 2022), national authorities are urged to develop reception and resettlement programmes meeting the needs of this vulnerable group. These programmes should be designed and implemented attempting to decrease the risk of post-migration stressors that may contribute to worsening the mental health of a population already exposed to potentially traumatic experiences before and during the migration process. Evidence-based focused psychological support should be an important programme component to be delivered to asylum seekers and refugees at-risk of developing a mental disorder.

Contributors

CB, MP and FT conceived the study. FT statistically analysed the data. CB, GO and FT accessed and verified the data. CB wrote the first draft of the manuscript with input from FT and MP. All authors contributed to the conception and design of the study, data interpretation and manuscript revision. All authors read and approved the submitted manuscript and had final responsibility for the decision to submit for publication. CB was the overall guarantor of the content.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the European Commission (grant agreement No. 779255; 'RE-DEFINE: Refugee Emergency: DEFining and Implementing Novel Evidence-based psychosocial interventions').

Disclaimer

The authors alone are responsible for the views expressed in this article, and they do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated.

Ethical approval

This was a secondary analysis of deidentified participant data. The initial study was approved by the WHO Research Ethics Review Committee, and from the Ethics Committees of

all participating sites. Before participation, a written informed consent was obtained from each participant, allowing investigators to use anonymized data for future analysis. Participants gave informed consent to participate in the study before taking part.

ORCID iDs

Corrado Barbui  <https://orcid.org/0000-0003-1073-9282>

Ceren Acarturk  <https://orcid.org/0000-0001-7093-1554>

Lorenzo Tarsitani  <https://orcid.org/0000-0002-1752-966X>

Data availability statement

Data collected in the RE-DEFINE study are stored in the online repository EUDAT B2SHARE (study in Western Europe: <https://b2share.eudat.eu/records/fa7264d624364683830ff37acee01c04>; study in Turkey: <http://doi.org/10.23728/b2share.8ac4f28d2415413e89de7847c05471fc>). The reuse of data will be offered only upon motivated request, which will undergo the scrutiny of the RE-DEFINE General Assembly.

Supplemental material

Supplemental material for this article is available online.

References

- Acarturk, C., McGrath, M., Roberts, B., Ilkkursun, Z., Cuijpers, P., Sijbrandij, M., Sondorp, E., Ventevogel, P., McKee, M., & Fuhr, D. C. (2021) Prevalence and predictors of common mental disorders among Syrian refugees in Istanbul, Turkey: A cross-sectional study. *Social Psychiatry and Psychiatric Epidemiology*, *56*, 475–484.
- Acarturk, C., Uygun, E., Ilkkursun, Z., Carswell, K., Tedeschi, F., Batu, M., Eskici, S., Kurt, G., Anttila, M., Au, T., Baumgartner, J., Churchill, R., Cuijpers, P., Becker, T., Koesters, M., Lantta, T., Nose, M., Ostuzzi, G., Popa, M., Purgato, M., Sijbrandij, M., Turrini, G., Valimaki, M., Walker, L., Wancata, J., Zanini, E., White, R. G., van Ommeren, M., & Barbui, C. (2022). Effectiveness of a WHO self-help psychological intervention for preventing mental disorders among Syrian refugees in Turkey: A randomized controlled trial. *World Psychiatry*, *21*, 88–95.
- Araya, R., Lewis, G., Rojas, G., & Fritsch, R. (2003). Education and income: Which is more important for mental health? *Journal of Epidemiology and Community Health*, *57*, 501–505.
- Barbui, C., Purgato, M., Acarturk, C., Churchill, R., Cuijpers, P., Koesters, M., Sijbrandij, M., Valimaki, M., Wancata, J., & White, R. G. (2022). Preventing the mental health consequences of war in refugee populations. *Epidemiology and Psychiatric Sciences*, *31*, e24.
- Bjelland, I., Krokstad, S., Mykletun, A., Dahl, A. A., Tell, G. S., & Tambs, K. (2008). Does a higher educational level protect against anxiety and depression? The HUNT study. *Social Science and Medicine*, *66*, 1334–145.
- Blackmore, R., Boyle, J. A., Fazel, M., Ranasinha, S., Gray, K. M., Fitzgerald, G., Misso, M., & Gibson-Helm, M. (2020). The prevalence of mental illness in refugees and asylum seekers: A systematic review and meta-analysis. *PLOS Medicine*, *17*, e1003337.

- Blanchard, E. B., Jones-Alexander, J., Buckley, T. C., & Forneris, C. A. (1996). Psychometric properties of the PTSD Checklist (PCL). *Behavioural Research and Therapeutics*, 34, 669–673.
- Charlson, F., van Ommeren, M., Flaxman, A., Cornett, J., Whiteford, H., & Saxena, S. (2019). New WHO prevalence estimates of mental disorders in conflict settings: A systematic review and meta-analysis. *Lancet*, 394, 240–248.
- Crepet, A., Rita, F., Reid, A., Van den Boogaard, W., Deiana, P., Quaranta, G., Barbieri, A., Bongiorno, F., & Di Carlo, S. (2017). Mental health and trauma in asylum seekers landing in Sicily in 2015: A descriptive study of neglected invisible wounds. *Conflict and Health*, 11, 1.
- Dabrowski, C. L., Lating, J. M., Sherman, M. F., Kirkhart, M. W., & Breen, D. (2020). Probable PTSD, symptoms of depression, and an exploratory factor analysis of item responses in Canadian steelworkers after workplace traumas. *Crisis, Stress, and Human Resilience: An International Journal*, 2, 57–71.
- el-Rufaie, O. F., & Daradkeh, T. K. (1996). Validation of the Arabic versions of the thirty- and twelve-item General Health Questionnaires in primary care patients. *British Journal of Psychiatry*, 169, 662–664.
- Epping-Jordan, J. E., Harris, R., Brown, F. L., Carswell, K., Foley, C., Garcia-Moreno, C., Kogan, C., & van Ommeren, M. (2016). Self-help plus (SH+): A new WHO stress management package. *World Psychiatry*, 15, 295–296.
- Fuhr, D. C., Acarturk, C., McGrath, M., Ilkkursun, Z., Sondorp, E., Sijbrandij, M., Ventevogel, P., Cuijpers, P., McKee, M., & Roberts, B. (2019). Treatment gap and mental health service use among Syrian refugees in Sultanbeyli, Istanbul: A cross-sectional survey. *Epidemiology and Psychiatric Sciences*, 29, e70.
- Giacco, D. (2019). Identifying the critical time points for mental health of asylum seekers and refugees in high-income countries. *Epidemiology and Psychiatric Sciences*, 29, e61.
- Gleeson, C., Frost, R., Sherwood, L., Shevlin, M., Hyland, P., Halpin, R., Murphy, J., & Silove, D. (2020). Post-migration factors and mental health outcomes in asylum-seeking and refugee populations: A systematic review. *European Journal of Psychotraumatology*, 11, 1793567.
- Goldberg, D. P., Oldehinkel, T., & Ormel, J. (1998). Why GHQ threshold varies from one place to another. *Psychological Medicine*, 28, 915–921.
- Hajak, V. L., Sardana, S., Verdelli, H., & Grimm, S. (2021). A systematic review of factors affecting mental health and well-being of asylum seekers and refugees in Germany. *Frontiers in Psychiatry*, 12, 643704.
- Henkelmann, J. R., de Best, S., Deckers, C., Jensen, K., Shahab, M., Elzinga, B., & Molendijk, M. (2020). Anxiety, depression and post-traumatic stress disorder in refugees resettling in high-income countries: Systematic review and meta-analysis. *BJPsych Open*, 6, e68.
- Hoell, A., Kourmpeli, E., Salize, H. J., Heinz, A., Padberg, F., Habel, U., Kamp-Becker, I., Hohne, E., Boge, K., & Bajbouj, M. (2021). Prevalence of depressive symptoms and symptoms of post-traumatic stress disorder among newly arrived refugees and asylum seekers in Germany: Systematic review and meta-analysis. *BJPsych Open*, 7, e93.
- Jannesari, S., Hatch, S., & Oram, S. (2020). Seeking sanctuary: Rethinking asylum and mental health. *Epidemiology and Psychiatric Sciences*, 29, e154.
- Jongedijk, R. A., Eising, D. D., van der Aa, N., Kleber, R. J., & Boelen, P. A. (2020). Severity profiles of posttraumatic stress, depression, anxiety, and somatization symptoms in treatment seeking traumatized refugees. *Journal of Affective Disorders*, 266, 71–81.
- Kadri, N., Agoub, M., El Gnaoui, S., Alami Kh, M., Hergueta, T., & Moussaoui, D. (2005). Moroccan colloquial Arabic version of the mini international neuropsychiatric interview (MINI): Qualitative and quantitative validation. *European Psychiatry*, 20, 193–195.
- Kaiser, H. F. (1960). The application of electronic computers to factor analysis. *Educational and Psychological Measurement*, 20, 145–151.
- Kilic, C., Rezaki, M., Rezaki, B., Kaplan, I., Ozgen, G., Sagduyu, A., & Ozturk, M. O. (1997). General health questionnaire (GHQ12 & GHQ28): Psychometric properties and factor structure of the scales in a Turkish primary care sample. *Social Psychiatry and Psychiatric Epidemiology*, 32, 327–331.
- Kirmayer, L. J., Narasiah, L., Munoz, M., Rashid, M., Ryder, A. G., Guzder, J., Hassan, G., Rousseau, C., Pottie, K., Canadian Collaboration for, I., & Refugee, H. (2011). Common mental health problems in immigrants and refugees: General approach in primary care. *CMAJ*, 183, E959–E967.
- Knipscheer, J. W., Sleijpen, M., Mooren, T., Ter Heide, F. J., & van der Aa, N. (2015). Trauma exposure and refugee status as predictors of mental health outcomes in treatment-seeking refugees. *BJPsych Bull*, 39, 178–182.
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16, 606–613.
- Laban, C. J., Gernaat, H. B., Komproe, I. H., Schreuders, B. A., & De Jong, J. T. (2004). Impact of a long asylum procedure on the prevalence of psychiatric disorders in Iraqi asylum seekers in The Netherlands. *Journal of Nervous and Mental Disorders*, 192, 843–851.
- Mesa-Vieira, C., Haas, A. D., Buitrago-Garcia, D., Roa-Diaz, Z. M., Minder, B., Gamba, M., Salvador, D., Jr., Gomez, D., Lewis, M., Gonzalez-Jaramillo, W. C., Pahud de Mortanges, A., Buttia, C., Muka, T., Trujillo, N., & Franco, O. H. (2022). Mental health of migrants with pre-migration exposure to armed conflict: A systematic review and meta-analysis. *Lancet Public Health*, 7, e469–e481.
- Miller, K. E., Jordans, M. J. D., Tol, W. A., & Galappatti, A. (2021). A call for greater conceptual clarity in the field of mental health and psychosocial support in humanitarian settings. *Epidemiology and Psychiatric Sciences*, 30, e5.
- Mollica, R. F., Caspi-Yavin, Y., Bollini, P., Truong, T., Tor, S., & Lavelle, J. (1992). The Harvard Trauma Questionnaire. Validating a cross-cultural instrument for measuring torture, trauma, and posttraumatic stress disorder in Indochinese refugees. *Journal of Nervous and Mental Disorders*, 180, 111–116.
- Nickerson, A., Byrow, Y., Rasmussen, A., O'Donnell, M., Bryant, R., Murphy, S., Mau, V., McMahon, T., Benson, G., & Liddell, B. (2021). Profiles of exposure to potentially traumatic events

- in refugees living in Australia. *Epidemiology and Psychiatric Sciences*, 30, e18.
- Pandi-Perumal, S. R., Kumar, V. M., Pandian, N. G., de Jong, J. T., Andiappan, S., Corlateanu, A., Mahalaksmi, A. M., Chidambaram, S. B., Kumar, R. R., Ramasubramanian, C., Sivasubramaniam, S., Bjorkum, A. A., Cutajar, J., Berk, M., Trakht, I., Vrdoljak, A., Meira, E. C. M., Eyre, H. A., Gronli, J., . . . Seeman, M. V. (2022). Scientists against war: A plea to world leaders for better governance. *Sleep and Vigilance*, 6, 1–6.
- Papola, D., Purgato, M., Gastaldon, C., Bovo, C., van Ommeren, M., Barbui, C., & Tol, W. A. (2020). Psychological and social interventions for the prevention of mental disorders in people living in low- and middle-income countries affected by humanitarian crises. *Cochrane Database of Systematic Reviews*, 9, CD012417.
- Patel, A. R., Newman, E., & Richardson, J. (2022). A pilot study adapting and validating the Harvard Trauma Questionnaire (HTQ) and PTSD checklist-5 (PCL-5) with Indian women from slums reporting gender-based violence. *BMC Womens Health*, 22, 22.
- Plumpton, C. O., Morris, T., Hughes, D. A., & White, I. R. (2016). Multiple imputation of multiple multi-item scales when a full imputation model is infeasible. *BMC Research Notes*, 9, 45.
- Priebe, S., Giacco, D., & El-Nagib, R. (2016). *Public health aspects of mental health among migrants and refugees: A review of the evidence on mental health care for refugees, asylum seekers and irregular migrants in the WHO European region*. WHO Regional Office for Europe.
- Purgato, M., Carswell, K., Acarturk, C., Au, T., Akbair, S., Anttila, M., Baumgartner, J., Bailey, D., Biondi, M., Bird, M., Churchill, R., Eskici, S., Hansen, L. J., Heron, P., Ilkkursun, Z., Kilian, R., Koesters, M., Lantta, T., Nose, M., . . . Van Ommeren, M. (2019). Effectiveness and cost-effectiveness of Self-Help Plus (SH+) for preventing mental disorders in refugees and asylum seekers in Europe and Turkey: Study protocols for two randomised controlled trials. *BMJ Open*, 9, e030259.
- Purgato, M., Carswell, K., Tedeschi, F., Acarturk, C., Anttila, M., Au, T., Bajbouj, M., Baumgartner, J., Biondi, M., Churchill, R., Cuijpers, P., Koesters, M., Gastaldon, C., Ilkkursun, Z., Lantta, T., Nose, M., Ostuzzi, G., Papola, D., Popa, M., . . . Barbui, C. (2021). Effectiveness of self-help plus in preventing mental disorders in refugees and asylum seekers in Western Europe: A multinational randomized controlled trial. *Psychotherapy and Psychosomatics*, 90, 403–414.
- Royston, P. (2005). Multiple imputation of missing values: Update. *Stata Journal*, 5, 188–201.
- Rytwinski, N. K., Scur, M. D., Feeny, N. C., & Youngstrom, E. A. (2013). The co-occurrence of major depressive disorder among individuals with posttraumatic stress disorder: A meta-analysis. *Journal of Traumatic Stress*, 26, 299–309.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59(Suppl 20), 22–33; quiz 34–57.
- StataCorp. (2017). *Stata Statistical Software: Release 15*. StataCorp LLC.
- Sterne, J. A., White, I. R., Carlin, J. B., Spratt, M., Royston, P., Kenward, M. G., Wood, A. M., & Carpenter, J. R. (2009). Multiple imputation for missing data in epidemiological and clinical research: Potential and pitfalls. *BMJ*, 338, b2393.
- Stevens, J. (2002). *Applied multivariate statistics for the social sciences*. Lawrence Erlbaum Associates.
- Tabachnick, B. F., Fidell, L. S., & Ullman, J. B. (2007). *Using multivariate statistics*. Pearson/Allyn & Bacon.
- Turrini, G., Tedeschi, F., Cuijpers, P., Del Giovane, C., Kip, A., Morina, N., Nose, M., Ostuzzi, G., Purgato, M., Ricciardi, C., Sijbrandij, M., Tol, W., & Barbui, C. (2021). A network meta-analysis of psychosocial interventions for refugees and asylum seekers with PTSD. *BMJ Global Health*, 6(6), e005029.
- United Nations High Commissioner for Refugees. (2022). *Refugee data finder*. Author.
- Uphoff, E., Robertson, L., Cabieses, B., Villalon, F. J., Purgato, M., Churchill, R., & Barbui, C. (2020). An overview of systematic reviews on mental health promotion, prevention, and treatment of common mental disorders for refugees, asylum seekers, and internally displaced persons. *Cochrane Database of Systematic Reviews*, 9, CD013458.
- White, I. R., Royston, P., & Wood, A. M. (2011). Multiple imputation using chained equations: Issues and guidance for practice. *Statistics in Medicine*, 30, 377–99.
- White, R. G., & Van der Boor, C. (2021). Enhancing the capabilities of forcibly displaced people: A human development approach to conflict- and displacement-related stressors. *Epidemiology and Psychiatric Sciences*, 30, e34.
- World Health Organization. (2017). *Scalable psychological interventions for people in communities affected by adversity*. Author.
- World Health Organization. (2021a). *Self help plus (SH+): A group-based stress management course for adults. Generic field-trial version 1.0*. Author.
- World Health Organization. (2021b). *Doing what matters in times of stress*. Author.
- Winkler, J. G., Brandl, E. J., Bretz, H. J., Heinz, A., & Schouler-Ocak, M. (2019). The influence of residence status on psychiatric symptom load of asylum seekers in Germany. *Psychiatrische Praxis*, 46, 191–199.
- Wu, Y., Levis, B., Sun, Y., Krishnan, A., He, C., Riehm, K. E., Rice, D. B., Azar, M., Yan, X. W., Neupane, D., Bhandari, P. M., Imran, M., Chiovitti, M. J., Saadat, N., Boruff, J. T., Cuijpers, P., Gilbody, S., McMillan, D., . . . Thombs, B. D. (2020). Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale – Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. *Journal of Psychosomatic Research*, 129, 109892.