

VU Research Portal

The top of the Olduvai subchron in a high-resolution magnetostratigraphy from the West Turkana core WTK13, Hominin Sites and Paleolakes Drilling Project (HSPDP)

Sier, M.J.; Langereis, C.; Dupont-Nivet, G.; Feibel, C.S.; Joordens, J.; van der Lubbe, H.J.L.; Beck, C.; Olago, D.; Cohen, A.

published in

19th EGU General Assembly, EGU2017
2017

document version

Publisher's PDF, also known as Version of record

document license

CC BY

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Sier, M. J., Langereis, C., Dupont-Nivet, G., Feibel, C. S., Joordens, J., van der Lubbe, H. J. L., Beck, C., Olago, D., & Cohen, A. (2017). The top of the Olduvai subchron in a high-resolution magnetostratigraphy from the West Turkana core WTK13, Hominin Sites and Paleolakes Drilling Project (HSPDP). In *19th EGU General Assembly, EGU2017: proceedings from the conference held 23-28 April, 2017 in Vienna, Austria* (EGU General Assembly Conference Abstracts). EGU General Assembly Conference Abstracts.
<https://ui.adsabs.harvard.edu/abs/2017EGUGA..1916960S/abstract>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl



The top of the Olduvai subchron in a high-resolution magnetostratigraphy from the West Turkana core WTK13, Hominin Sites and Paleolakes Drilling Project (HSPDP)

Mark Sier (1), Cor Langereis (2), Guillaume Dupont-Nivet (3,4), Craig Feibel (5), Jose Jordeens (1,6), Jeroen van der Lubbe (6), Catherine Beck (7), Daniel Olago (8), Andrew Cohen (9), and the HSPDP Team members Team (1) Leiden University, Faculty of Archeology, Leiden, The Netherlands, (2) Utrecht University, Paleomagnetic Laboratory, Utrecht, The Netherlands, (3) CNRS - Géosciences Rennes UMR6118, Université de Rennes 1, France, (4) Potsdam University, Institute of Earth and Environmental Science, Potsdam - Golm, Germany, (5) Rutgers University, Department of Anthropology, New Brunswick, NJ, USA, (6) Free University Amsterdam, Sedimentology Department, Amsterdam, The Netherlands, (7) Hamilton College, Geosciences, Clinton NY, USA, (8) University of Nairobi, The Institute for Climate Change and Adaptation, Nairobi, Kenya, (9) University of Arizona, Department of Geosciences, Tucson AZ, USA

One of the major challenges in understanding the evolution of our own species is identifying the role climate change has played in the evolution of earlier hominin species. To clarify the influence of climate, we need long and continuous high-resolution paleoclimate records, preferably obtained from hominin-bearing sediments, that are well-dated by tephro- and magnetostratigraphy and other methods. This is hindered, however, by the fact that fossil-bearing sediments are often discontinuous, and subject to weathering, which may lead to oxidation and remagnetization. To obtain fresh, unweathered sediments, the Hominin Sites and Paleolakes Drilling Project (HSPDP) collected a ~216- meter core (WTK13) in 2013 from deposits of Early Pleistocene paleolake Lorenyang in the western Turkana Basin (Kenya). Here, we present the magnetostratigraphy of the core. Rock magnetic analyses reveal the presence of iron sulphides carrying the remanent magnetizations. To recover polarity orientation from the near-equatorial WTK13 core drilled at 5°N, we developed and successfully applied two independent drill-core reorientation methods taking advantage of (1) the sedimentary fabric as expressed in the Anisotropy of Magnetic Susceptibility (AMS) and (2) the occurrence of a viscous component oriented in the present day field. The reoriented directions reveal a normal to reversed polarity reversal identified as the top of the Olduvai subchron. From this excellent record, we find no evidence for the 'Vrica subchron' previously reported in the area. We suggest that outcrop-based interpretations supporting the presence of the Vrica subchron have been affected by the oxidation of iron sulphides initially present in the sediments as evident in the core record, and by subsequent remagnetization. Based on our new high-resolution magnetostratigraphy and stratigraphic markers, we provide constraints for an initial age model of the WTK13 core. We discuss the implications of the observed geomagnetic record for human evolution studies.