

VU Research Portal

Comorbidity in knee osteoarthritis

de Rooij, M.

2017

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

de Rooij, M. (2017). *Comorbidity in knee osteoarthritis: Development and evaluation of tailored exercise therapy*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Chapter 4

Osteoarthritis of the hip or knee: *which coexisting disorders are disabling?*

Kerstin G. Reeuwijk
Mariëtte de Rooij
Gabriella M. van Dijk
Cindy Veenhof
Martijn P. Steultjens
Joost Dekker

Published in: Clinical Rheumatology 2010; 29:739–747

Abstract

Objective. Exercise therapy is generally recommended in osteoarthritis (OA) of the hip or knee. However, coexisting disorders may bring additional impairments, which may necessitate adaptations to exercise for OA of the hip or knee. For the purpose of developing an adapted protocol for exercise therapy in OA patients with coexisting disorders, information is needed on which specific coexisting disorders in OA are associated with activity limitations and pain.

Methods. To describe the relationship between specific coexisting disorders, activity limitations, and pain in patients with OA of the hip or knee, a cross-sectional cohort study among 288 older adults (50–85 years of age) with OA of hip or knee was conducted. Subjects were recruited from three rehabilitation centers and two hospitals. Demographic data, clinical data, information about coexisting disorders (i.e., comorbidity and other disorders), activity limitations (WOMAC: physical functioning domain), and pain (visual analogue scale (VAS)) were collected by questionnaire. Statistical analysis included descriptive statistics and multivariate regression analysis.

Results. Coexisting disorders associated with activity limitations were chronic back pain or hernia, arthritis of the hand or feet, and other chronic rheumatic diseases (all musculoskeletal disorders); diabetes and chronic cystitis (non-musculoskeletal disorders); hearing impairments in a face-to-face conversation, vision impairments in long distances, and dizziness in combination with falling (all sensory impairments); and overweight and obesity. Coexistent disorders associated with pain were arthritis of the hand or feet, other chronic rheumatic diseases (musculoskeletal disorders), and diabetes (non-musculoskeletal disorder).

Conclusion. Specific disorders coexisting next to OA and associated with additional activity limitations and pain were identified. These coexisting disorders need to be addressed in exercise therapy and rehabilitation for patients with OA of the hip or knee.



Introduction

Osteoarthritis (OA) is one of the diseases with the highest rates of comorbidity¹. Previous studies have reported comorbidity rates of 68% to 85%²⁻⁵. Diseases that frequently occur next to OA are diabetes, hypertension, and cardiovascular disorders; other disorders, including overweight and back pain, occur frequently as well^{1,4,6}. Thus, coexisting disorders—defined as coexisting diseases and coexisting other disorders (e.g., overweight)—are highly prevalent in OA.

Exercise therapy is generally recommended in OA of the hip or knee: Exercise is effective in reducing activity limitations and pain in OA^{7,8}. However, coexisting disorders may bring additional impairments, which necessitate adaptations to the exercise protocol for OA of the hip or knee. For the future purpose of developing such an adapted protocol, information is needed on which specific coexisting disorders in OA are disabling, i.e., which coexisting disorders are associated with activity limitations and pain. Coexisting disorders causing activity limitations and pain are likely to cause restrictions to exercise therapy as well, necessitating adaptations in the exercise protocol.

Previously, we have reported which coexisting disorders are associated with additional activity limitations and pain in OA of hip or knee^{5,9}. In these studies, we used the Cumulative Illness Rating Scale (CIRS)¹⁰⁻¹² to assess coexisting disorders. The CIRS yields information on global categories of coexisting diseases, for example, ear, eye, nose, and throat diseases or endocrine and metabolic diseases. Although quite informative, more detailed information on which specific coexisting disorders are disabling is required in order to be able to develop the exercise therapy protocol with adaptations for coexisting disorders. The objective of the study was to describe the relationship between specific coexisting disorders, activity limitations, and pain in patients with hip or knee OA.

Methods

Design

The present study is a secondary analysis of previously reported data⁵. The design of this cross-sectional study is summarized below. The reader is referred to the original publication for a more detailed description of the design⁵. The study was approved by the Medical Ethical Committee of the VU University Medical Centre, Amsterdam, the Netherlands.

Study population

Participants were recruited from three rehabilitation centers and two hospitals (Departments of Orthopedics, Rheumatology, or Rehabilitation). Inclusion criteria were

(a) diagnosis of OA of the hip or knee by medical specialist according to radiological criteria or clinical criteria of the American College of Rheumatology^{13,14}, (b) 50 years of age or older, (c) referral to hospital or rehabilitation center less than a year before inclusion, (d) at least moderate functional problems (Lequesne algofunctional index score ≥ 5)¹⁵, and (e) informed consent. Exclusion criteria were (a) 85 years of age or older, (b) insufficient understanding of the Dutch language, and (c) expected death due to fatal illness within 1 year after inclusion.

Measurements

Patients were invited to a test location. The data used in the present study were gathered by means of interview (demographic and clinical data) and questionnaires (activity limitations, pain, and coexistent disorders). X-rays were used to evaluate radiological impairment of the hip or knee.

Demographic and clinical data

Demographic and clinical data were collected for each patient including age, gender, height, weight, location of OA, duration of complaints, other joint complaints, level of education, and marital status. Body mass index (BMI) was calculated (weight/height²). If available, X-rays of the hip and knee recorded in the year before inclusion were scored on joint space width and osteophytes, following a standardized procedure^{16,17}. A 0–3 scale was used for rating the radiographs: 0 = normal; 1 = mild or 1–33% abnormal; 2 = moderate or 34–66% abnormal; 3 = severe or 67–100% abnormal. From these scores, Kellgren and Lawrence grades were calculated.

Activity limitations and pain

Activity limitations were measured using the physical functioning domain of the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) (range 0–100)^{18,19}. WOMAC physical functioning score is a standardized score using the formula: $((68 - \text{total score on the physical functioning domain}) \times 100)/68$. A higher score on this standardized WOMAC stands for fewer activity limitations. Pain at the time of assessment was rated on a visual analogue scale (VAS) (range 0–10). A higher score on VAS reflects more pain.

Coexisting disorders

Patients indicated the presence or absence of coexisting disorders in the year prior to the interview, using the list of the Dutch Bureau of Statistics (CBS). This list describes the most common chronic diseases and disorders in the Netherland²⁰. Based on expert consultation, impairments in vision and hearing were added to this list. Overweight was defined as $25 \text{ kg/m}^2 \leq \text{BMI} < 30 \text{ kg/m}^2$; obesity was defined as $\text{BMI} \geq 30 \text{ kg/m}^2$. Coexisting disorders were categorized as musculoskeletal disorders, non-musculoskeletal disorders, sensory impairments, and overweight and obesity.

Statistical analyses

The presence of coexisting disorders, activity limitations, and pain was analyzed using descriptive statistics. Subsequent analyses were limited to those disorders that affected $\geq 5\%$ of the study population. The association between coexisting disorders (present or absent) and activity limitations was evaluated in multivariate regression analyses, correcting for age and gender. Similar analyses were done for the association between coexisting disorders and pain, correcting for age and gender. An association was termed significant if the p value was < 0.05 . For all analyses, the SPSS (version 14.0) was used²¹.

Results

Study population

Initially, 775 patients with osteoarthritis of the hip or knee that visited the department in the year prior to inclusion were contacted by mail and were asked to participate in the study. Of those patients that volunteered ($n = 364$), 288 were included. Seventy-six patients were excluded because they did not meet the inclusion criteria. Reasons of

Table 1. Baseline characteristics (N=288)

Gender: male, n (%)	83 (28.8%)
Age, mean (sd)	66 (8.7)
Body mass index, mean (sd), kg/m ²	27.8 (4.5)
Location OA	
Knee OA, n (%)	139 (48.4%)
Hip OA, n (%)	72 (25.1%)
Both, n (%)	76 (26.5%)
Duration of complains (years), mean (sd)	9.9 (10.7)
Physical functioning (WOMAC), mean (sd)	56.51 (19.85)
Pain (VAS), mean (sd)	4.81 (2.56)
Radiological impairment knee ^a	
K&L grade ≥ 2 ; n (%)	118 (95.2)
Radiological impairment hip ^b	
K&L grade ≥ 2 ; n (%)	83 (97.6)

N, n number, sd standard deviation, OA: osteoarthritis, WOMAC, Western Ontario and McMasters Universities Osteoarthritis Index; VAS, Visual Analogue Scale

^a N = 138 (from only a part of the included patients X-rays were available)

^b N = 90 (from only part of the included patients X-rays were available)



Table 2. Presence of coexisting disorders

	Present n (%)
Musculoskeletal disorders	
Chronic back pain (> 3 month) or hernia	85 (29.5)
Arthritis of the hands or feet	53 (18.4)
Other chronic rheumatic diseases (> 3 month)	29 (10.1)
Non-musculoskeletal disorders	
Asthma or COPD	45 (15.6)
Sinusitis	35 (12.2)
Severe cardiac disorder or coronary disease	23 (8)
Hypertension	92 (31.9)
(Consequences of) a stroke	6 (2.1)
Peptic ulcer or duodenal ulcer disease	10 (3.5)
Severe bowel disorder (> 3 month)	19 (6.6)
Gallstones or inflammation of the gall bladder	6 (2.1)
Liver disorder or cirrhosis of the liver	0 (0)
Kidney stones (calculus renalis)	7 (2.4)
Severe kidney disorder	2 (0.7)
Chronic cystitis	16 (5.6)
Prolapse (only females, N=205)	17 (8.3)
Diabetes	28 (9.7)
Thyroid disorders	25 (8.7)
Epilepsy	2 (0.7)
Migraine	19 (6.6)
Severe skin disease	4 (1.4)
Cancer and malignant diseases	7 (2.4)
Sensory impairments	
Hearing impairments in a group conversation (N=286)	90 (31.5)
Hearing impairments in a face-to-face conversation (N=287)	22 (7.7)
Vision impairments in short distances (N=287)	77 (26.8)
Vision impairments in long distances (N=287)	33 (11.5)
Dizziness in combination with falling	24 (8.3)
Overweight and obesity (N=285)	
Overweight ($25 \leq \text{BMI} < 30$)	149 (51.7)
Obesity ($\text{BMI} \geq 30$)	68 (23.9)

N = 288, unless otherwise stated.

Data in italics are present in >5%

exclusion were age ($n = 2$), language ($n = 4$), less than moderate functional problems ($n = 48$), and referral longer than 1 year before inclusion ($n = 22$). Analyses showed that there were no differences between the group of patients that were initially contacted ($N = 775$) and the patients that were included in the study ($N = 288$) with regard to age and gender. Some differences were found in the location of OA. Compared to our study population, patients that were initially contacted suffered less frequently from both hip and knee OA (6.2%) and more frequently from knee OA (59.5%) and hip OA (34.3%)⁵.

Table 1 shows baseline characteristics of the study population. The majority of the study population was female (71.2%), the mean age was 66 (SD = 8,7) years. The largest part of the subjects (48.4%) suffered from knee OA, 25.1% suffered from hip OA, and 26.5% had both hip and knee OA. The mean physical functioning score on the WOMAC was 56.5 (SD = 19.6), and the mean pain score on the VAS was 4.8 (SD = 2.6). Most patients (80%) were recruited from the departments of orthopedics; the other 20% originated from departments of rheumatology and departments of rehabilitation.

Coexisting disorders

The presence of coexisting disorders in the study population is presented in Table 2. Eighteen coexisting disorders occurred in >5% of the sample (see Table 2). In the category of musculoskeletal disorders, coexisting conditions occurring in > 5% of the patients included chronic back pain or hernia, 29.5%; arthritis of the hand or feet, 18.4%; and other chronic rheumatic diseases, 10.1%. In the category of non-musculoskeletal disorders, coexisting conditions occurring in > 5% of the patients included hypertension, 31.9%; asthma or COPD, 15.6%; sinusitis, 12.2%; diabetes, 9.7%; thyroid disorders, 8.7%; severe cardiac disorder or coronary disease, 8%; severe bowel disorder, 6.6%; migraine, 6.6%; chronic cystitis, 5.6%; and prolapse in 8.3% of the female patients. In the category of sensory impairments, coexisting conditions occurring in > 5% of the patients were hearing impairments in a group conversation, 31.5%; vision impairments in short distances and long distances, 26.8% and 11.5%, respectively; and dizziness in combination with falling, 8.3%. Overweight and obesity occurred in 51.7% and 23.9%, respectively.

Association between coexisting disorders and activity limitations

Table 3 shows the relationship between specific coexisting disorders and activity limitations. Significantly more activity limitations were found in OA patients with chronic back pain or hernia ($p < 0.05$), arthritis of the hand or feet ($p < 0.05$), and other chronic rheumatic diseases ($p < 0.05$) (all in the category of musculoskeletal disorders); with diabetes ($p < 0.05$) and chronic cystitis ($p < 0.05$) (category of non-musculoskeletal disorders); with hearing impairments in a face-to-face conversation ($p < 0.05$), vision impairments in long distances ($p < 0.05$), and dizziness in combination with falling ($p < 0.05$) (category of sensory impairments); and with overweight ($p < 0.01$) and obesity ($p < 0.05$).



Table 3. Association between coexisting disorders, activity limitations, and pain

	Activity limitations (WOMAC)	Pain (VAS)
Musculoskeletal disorders		
Chronic back pain (> 3 month) or hernia		
Present (mean (sd)) ^a	51.7 (18.2)	5.0 (2.5)
Absent (mean (sd)) ^a	58.5 (20.2)	4.7 (2.5)
B (95%CI) ^b	-6.4 (-11.5 ; -1.4)*	0.3 (-0.4 ; 1.0)
R ²	0.033	0.008
Arthritis of hands or feet		
Present (mean (sd)) ^a	50.1 (20.3)	5.8 (2.5)
Absent (mean (sd)) ^a	57.9 (19.5)	4.6 (2.5)
B (95%CI) ^b	-7.4 (-13.4 ; -1.5)*	1.2 (0.4 ; 1.9)**
R ²	0.032	0.037
Other chronic rheumatic diseases (> 3 month)		
Present (mean (sd)) ^a	47.3 (18.3)	5.9 (2.6)
Absent (mean (sd)) ^a	57.5 (19.8)	4.7 (2.5)
B (95%CI) ^b	-10.0 (-17.6 ; -2.4)*	1.1 (0.1 ; 2.1)*
R ²	0.035	0.023
Non-musculoskeletal disorders		
Asthma or COPD		
Present (mean (sd)) ^a	55.2 (19.1)	4.8 (2.6)
Absent (mean (sd)) ^a	56.7 (20.0)	4.8 (2.6)
B (95%CI) ^b	-1.7 (-8.0 ; 4.7)	0.0 (-0.8 ; 0.9)
R ²	0.012	0.005
Sinusitis		
Present (mean (sd)) ^a	54.5 (15.8)	4.9 (2.6)
Absent (mean (sd)) ^a	56.8 (20.4)	4.8 (2.6)
B (95%CI) ^b	-2.0 (-9.1 ; 5.1)	0.1 (-0.8 ; 1.0)
R ²	0.012	0.006
Severe cardiac disorder or coronary disease		
Present (mean (sd)) ^a	51.2 (20.6)	5.1 (2.8)
Absent (mean (sd)) ^a	57.0 (19.8)	4.8 (2.6)
B (95%CI) ^b	-5.5 (-14.1 ; 3.1)	0.4 (-0.7 ; 1.6)
R ²	0.012	0.008
Hypertension		
Present (mean (sd)) ^a	55.8 (21.9)	4.9 (2.5)
Absent (mean (sd)) ^a	56.8 (18.8)	4.8 (2.6)
B (95%CI) ^b	-0.6 (-5.6 ; 4.5)	0.3 (-0.4 ; 0.9)
R ²	0.012	0.007
Severe bowel disorder (> 3 month)		
Present (mean (sd)) ^a	54.3 (20.3)	4.5 (3.2)
Absent (mean (sd)) ^a	56.7 (19.9)	4.8 (2.5)
B (95%CI) ^b	-2.0 (-11.3 ; 7.2)	-0.4 (-1.6 ; 0.8)
R ²	0.012	0.007

Table 3. (cont'd)

	Activity limitations (WOMAC)	Pain (VAS)
Chronic cystitis		
Present (mean (sd)) ^a	45.2 (17.6)	5.4 (2.9)
Absent (mean (sd)) ^a	57.2 (19.8)	4.8 (2.5)
B (95%CI) ^b	-10.8 (-20.9; -0.7) †	0.7 (-0.6; 2.1)
R ²	0.027	0.01
Prolapse (females only)		
Present (mean (sd)) ^a	47.0 (22.0)	5.2 (2.9)
Absent (mean (sd)) ^a	56.4 (19.7)	4.8 (2.6)
B (95%CI) ^b	-9.0 (-18.9; 0.9) ^{***}	0.4 (-0.9; 1.7)
R ²	0.03	0.005
Diabetes		
Present (mean (sd)) ^a	49.5 (18.2)	5.9 (2.2)
Absent (mean (sd)) ^a	57.3 (19.9) [*]	4.7 (2.6)
B (95%CI) ^b	-8.2 (-15.9; -0.4) [*]	1.2 (0.2; 2.2) [*]
R ²	0.026	0.024
Thyroid disorders		
Present (mean (sd)) ^a	51.4 (22.6)	5.6 (2.8)
Absent (mean (sd)) ^a	57.0 (19.6)	4.7 (2.5)
B (95%CI) ^b	-5.8 (-14.2; 2.6)	1.1 (0.0; 2.1) ^{***}
R ²	0.018	0.018
Migraine		
Present (mean (sd)) ^a	56.6 (20.8)	4.5 (2.6)
Absent (mean (sd)) ^a	56.5 (19.8)	4.8 (2.6)
B (95%CI) ^b	0.1 (-9.2; 9.5)	-0.4 (-1.6; 0.8)
R ²	0.012	0.007
Sensory impairments		
Hearing impairments in a group conversation		
Present (mean (sd)) ^a	55.4 (19.1)	4.6 (2.7)
Absent (mean (sd)) ^a	57.0 (20.2)	4.9 (2.5)
B (95%CI) ^{**}	-0.9 (-6.0; 4.2)	-0.3 (-0.9; 0.4)
R ²	0.011	0.004
Hearing impairments in a face-to-face conversation		
Present (mean (sd)) ^a	46.7 (19.9)	5.6 (2.8)
Absent (mean (sd)) ^a	57.2 (19.7)	4.8 (2.5)
B (95%CI) ^b	-9.6 (-18.4; -0.8) [*]	1.0 (-0.1; 2.2) ^{***}
R ²	0.027	0.016
Vision impairments in short distances		
Present (mean (sd)) ^a	54.1 (19.8)	5.0 (2.6)
Absent (mean (sd)) ^a	57.3 (19.8)	4.8 (2.6)
B (95%CI) ^b	-2.9 (-9.1; 2.3)	0.3 (-0.4; 1.0)
R ²	0.016	0.007



Table 3. (cont'd)

	Activity limitations (WOMAC)	Pain (VAS)
Vision impairments in long distances		
Present (mean (sd)) ^a	47.6 (19.1)	5.2 (2.5)
Absent (mean (sd)) ^a	57.6 (19.7)	4.8 (2.6)
B (95%CI) ^b	-9.0 (-16.3 ; -1.6)*	0.6 (-0.4 ; 1.6)
R ²	0.031	0.01
Dizziness in combination with falling		
Present (mean (sd)) ^a	45.4 (23.4)	5.7 (2.9)
Absent (mean (sd)) ^a	57.5 (19.2)	4.7 (2.5)
B (95%CI) ^b	-11.2 (-19.6 ; -2.8)**	1.1 (0.0 ; 2.2)***
R ²	0.035	0.019
Weight		
Overweight (25 ≤ BMI < 30)		
Present (mean (sd)) ^a	54.7 (19.1)	4.9 (2.6)
Absent (mean (sd)) ^a	62.7 (19.0)	4.4 (2.6)
B (95%CI) ^b	-8.0 (-13.7 ; -2.4)**	0.5 (-0.3 ; 1.2)
Obesity (BMI ≥ 30)		
Present (mean (sd)) ^a	54.1 (21.4)	5.0 (2.6)
Absent (mean (sd)) ^a	62.7 (19.0)	4.4 (2.6)
B (95%CI) ^b	-8.6 (-15.3 ; -1.9)*	0.6 (-0.3 ; 1.5)
R ²	0.041	0.014

WOMAC Western Ontario and McMaster Universities Osteoarthritis Index, VAS Visual Analogue Scale, B regression coefficient; 95% CI: 95% Confidence interval

* $p < 0.05$; ** $p < 0.01$; *** p -value < 0.001

^a observed value; ^b adjusted value for age and sex

Association between coexisting disorders and pain

Table 3 also shows the relationship between coexisting disorders and pain. Arthritis of the hand or feet ($p < 0.01$) and other chronic rheumatic diseases ($p < 0.05$) (category of musculoskeletal disorders) and diabetes ($p < 0.05$) (category of non-musculoskeletal disorders) were found to be significantly associated with more pain.

Discussion

The aim of this study was to describe the relationship between specific coexisting disorders, activity limitations, and pain in patients with OA of the hip or knee. We have previously reported that “symptoms of other musculoskeletal disorders” occur frequently in OA⁵: We have now found chronic back pain or hernia, arthritis of the hands or feet,

and other chronic rheumatic diseases to be associated with activity limitations. In our previous study, we found “endocrine and metabolic diseases” to be associated with activity limitations: We now discovered that this applies to diabetes. We also found “other urogenital diseases” to be associated with activity limitations⁵: We now discovered that this applies in particular to chronic cystitis. Previously, we found “ear, eye, nose, and throat diseases” to be associated with activity limitations⁵: We have now observed that this applies in particular to hearing impairments in a face-to-face conversation and vision impairments in long distances. We also found dizziness in combination with falling to be associated with activity limitations. Finally, overweight and obesity were found to be related to activity limitations.

Regarding coexistent disorders and pain, we have now found arthritis of the hands or feet and other chronic rheumatic diseases to be associated with pain. Previously we did not find a relationship between endocrine and metabolic diseases and pain⁵; however, we now found an association between diabetes and pain.

Coexisting disorders may bring additional impairments, which necessitate adaptations to exercise for patients with OA of the hip or knee. For the future purpose of developing a protocol for exercise in OA patients with coexisting disorders, information is needed on which specific coexisting disorders in OA are associated with activity limitations and pain. The present study brings highly relevant new information in this respect. To our knowledge, this is the first study suggesting that hearing impairments, vision impairments, dizziness, and chronic cystitis need to be addressed in order to optimize functioning of patients with OA. Only Peters et al. have previously reported on the (near significant) relationship between eye diseases and future disability in patients with knee OA²². The present results shows that coexisting sensory impairments in hearing, vision, and balance, as well as motor impairments in bladder control (in the case of cystitis), have impact on functioning of patients with OA of the hip or knee: Apparently, these impairments need to be addressed in the rehabilitation of OA patients.

It is well known that OA is associated with other musculoskeletal disorders²³. In contrast, very little information is available on the impact of coexisting musculoskeletal disorders on activity limitations and pain. Cimmino et al. describe greater pain in patients with generalized OA than in patients with hip or knee OA alone²⁴. The present study found an association between arthritis of the hand or feet, “other” chronic rheumatic diseases, chronic back pain or hernia, and activity limitations. We also found a relationship between arthritis of the hand or feet, other chronic rheumatic diseases, and pain.

The present study confirms earlier observations on the impact of diabetes, overweight, and obesity on activity limitations in OA. Diabetes causes physiological restrictions to exercise therapy²⁵, whereas overweight and obesity introduce behavioral restrictions to exercise²⁶. Combining exercise therapy with a weight loss program has been shown to result in moderately improved outcome, compared to exercise therapy alone²⁷. Further work on adaptations to exercise therapy for OA patients with diabetes, overweight, and obesity is urgently required, because of the high prevalence and the functional impact of these coexisting disorders.



In our previous study, we found 54% of the study population to have cardiac diseases, and a significant relationship with activity limitations was found⁵. Caporali et al. found cardiac diseases (myocardial infarction and/or angina pectoris) to influence pain, quality of life, and joint function in OA patients⁴. Ettinger et al. also describe a higher likelihood of disabilities in OA patients with heart diseases²⁸. In the present study, no significant relationship was found between cardiac diseases and activity limitations or pain. A likely explanation is that subjects rated the presence of “severe heart diseases or coronary diseases.” This item might have implicated a too serious stage of disease, causing us to miss patients with mild or moderate heart disease. Despite our present results, adaptations for cardiac disease in exercise therapy for OA patients seem to be required.

The number of coexisting disorders associated with activity limitations ($n = 10$) was larger than the number of coexisting disorders associated with pain ($n = 3$). It seems that coexisting disorders easily impact on activity limitations, while the impact on pain is less strong. Future research is needed to elucidate the mechanisms of how coexisting disorders have an impact on activity limitations and pain. This research may also clarify the differential impact of coexisting disorders on activity limitations and pain.

Some limitations of the present study need to be discussed. Firstly, we relied on self-report to assess coexisting disorders, instead of medical record review. However, self-reported coexisting disorders tend to correspond rather well with coexisting disorders derived from medical record review (29): The percentage of agreement exceeded 90% for all coexisting disorders except for tumors; the kappa statistic ranged from 0.35 to 0.85, reflecting fair to substantial agreement. Fair agreement ($\text{kappa} < 0.40$) was found for ulcer disease, end organ damage resulting from diabetes, and connective tissue disease (not assessed in the present study); substantial agreement ($\text{kappa} > 0.60$) was found for myocardial infarction, stroke, and renal disease. Thus, self-report on coexisting disorders seems to be valid, with some exceptions noted above. Secondly, in assessing coexistent disorders, we used the list of the Dutch Bureau of Statistics (CBS). Unfortunately, this list did not include psychiatric disorders. From previous studies, it is known that depression plays an important role in developing activity limitations^{30,31}. Thirdly, patients included in this study were recruited from hospitals and rehabilitation centers; these patients may have received some form of rehabilitation. Thus, the results of the present study cannot be generalized to the general population suffering from OA. Nevertheless, the results are applicable to the group of patients consulting in hospitals and rehabilitation centers: This is a highly relevant group. Fourthly, the study focused on patients with OA of the hip or knee. We did not have information on a reference group of subjects without OA or subjects with another index disease. In subjects without OA and subjects with another index disease, comorbidity is likely to be associated with activity limitations and pain as well. In future studies, it would be interesting to compare impact of comorbidity in OA patients and other subjects. Finally, we did not adjust the analysis for multiple comparisons: This could easily result in false-negative errors (type II errors) because of the limited number of subjects with a specific coexisting disorder. It is acknowledged that this may have introduced some risk of false-positive errors (type I errors).

In conclusion, specific disorders coexisting next to OA and associated with additional activity limitations and pain were identified. These coexisting disorders need to be addressed in exercise therapy and rehabilitation for patients with OA of the hip or knee.

Acknowledgments

This study was in part supported by a grant from Netherlands Organization for Health Research and Development (ZON-MW Revalidatieonderzoek) and a grant from Royal Dutch Society for Physiotherapy (KNGF).



References

1. Schellevis FG, van der Velde J, van der Lesdonk E, van Eijk JTM, van Weel C. Comorbidity of chronic diseases in general practice. *J Clin Epidemiol* 1993; 5:469-473.
2. Juhakoski R, Tenhonen S, Anttonen T, Kauppinen T, Arokose JP. Factors affecting self-reported pain and physical functioning in patients with hip osteoarthritis. *Arch Phys Med Rehabil* 2008; 89:1066-1073.
3. Caporali R, Cimmino MA, Sarzi Puttini P, Scarpa R, Parazzini F, Zaninelli A, et al. Comorbid conditions in the AMCIA study patients: effects on the quality of life and drug prescriptions by general practitioners and specialists. *Semin Arthritis Rheum* 2005; 35(suppl 1):31-37.
4. Tuominen U, Blom M, Hirvonen J, Seitsalo S, Lehto M, Paavolainen P, et al. The effect of comorbidities on health related quality of life in patients placed on the waiting list for total joint replacement. *Health Qual Life Outcomes* 2007; 5:16.
5. van Dijk GM, Veenhof C, Schellevis F, Hulsmans H, Bakker JJP, Arwert H, et al. Comorbidity, limitations in activities and pain in patients with osteoarthritis of the hip or knee. *BMC Musculoskel Disord* 2008; 9:95.
6. Tukker A, Visscher TL, Picavet HS. Overweight and health problems of the lower extremities: osteoarthritis, pain and disability. *Public Health Nutr* 2009; 12(3):359-368.
7. Pendleton A, Arden N, Dougados M, Doherty M, Bannwarth B, Bijlsma JWJ, et al. EULAR recommendations for the management of knee osteoarthritis: report of a task force of the Standing Committee for International Clinical Studies Including Therapeutic Trials (ESCISIT). *Ann Rheum Dis* 2000; 59:936-944.
8. Zhang W, Moskowitz RW, Nuld G, Abramson S, Altman D, Arden N, et al. OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartil* 2008; 16:137-162 .
9. van Dijk GM, Veenhof C, Spreeuwenberg P, Coene N, Burger BJ, van Schaardenburg D, et al. Prognosis of limitations in activities in osteoarthritis of the hip or knee: a 3-year cohort study. *Arch Phys Med Rehabil* 2010; 91:58-66 .
10. Linn BS, Linn MW, Gurel L. Cumulative Illness Rating Scale. *J Am Geriatr Soc* 1968; 16(5):622-626.
11. Conwell Y, Forbes N, Cox C, Caine E. Validation of a measure of physical illness burden at autopsy the Cumulative Illness Rating Scale. *J Am Geriatr Soc* 1993; 41:38-41 .
12. Fortin M, Hudon C, Dubois M, Almirall J, Lapointe L, Soubhi H. Comparative assessment of three different indices of multi-morbidity for studies on health-related quality of life. *Health Qual Life Outcomes*. 2005; 3:74 .
13. Altman R, Alarcon G, Appelrouth D, Bloch D, Borenstein D, Brandt K, et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hip. *Arthritis Rheum* 1991; 34:505-514 .
14. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the knee. *Arthritis Rheum* 1986; 29:1039-1049 .
15. Lequesne M, Mery C, Samson M, Gerard D. Indices of severity for osteoarthritis of the hip and knee. *Scand J Rheumatol* 1987; 18(Suppl 65):85-89 .
16. Altman R, Hochberg M, Murphy W, Wolfe F, Lequesne M. Atlas of individual radiographic features in osteoarthritis. *Osteoarthritis Cartil* 1995; 3(suppl A):3-70 .
17. Dieppe P. Recommended methodology for assessing the progression of osteoarthritis of



- the hip and knee joints. *Osteoarthr Cartil* 1995; 3:73-77 .
18. Bellamy N, Buchanan W, Goldsmith C, Campbell J, Stitt L. Validation study of WOMAC: a health status instrument for measuring clinically-important patient-relevant outcomes following total hip or knee arthroplasty in osteoarthritis. *J Orthop Rheumatol* 1988; 1:95-108 .
 19. Roorda LD, Jones CA, Waltz M, Lankhorst GJ, Bouter LM, van der Eijken JW. Satisfactory cross cultural equivalence of the Dutch WOMAC in patients with hip osteoarthritis waiting for arthroplasty. *Ann Rheum Dis* 2004; 63:36-42 20.
 20. CBS Zelfgerapporteerde medische consumptie, gezondheid en leefstijl. Den Haag/Heerlen: CBS. 2008; <http://statline.cbs.nl/StatWeb/publication/?DM=SLNL&PA=03799&D1=94-122&D2=0-17&D3=0&D4=a&VW=T> (consulted at 09.12.2008).
 21. SPSS (2005) Windows, Rel. 14.0.0. SPSS Inc, Chicago.
 22. Peters TJ, Sanders C, Dieppe P, Donovan J. Factors associated with change in pain and disability over time: a community-based prospective observational study of hip and knee osteoarthritis. *Br J Gen Pract* 2005; 55:205-211 23.
 23. Kadam UT, Jordan K, Croft PR. Clinical comorbidity in patients with osteoarthritis: a case-control study of general practice consultants in England and Wales. *Ann Rheum Dis* 2004; 63:408-414.
 24. Cimmino MA, Sarzi-Puttini P, Scarpa R, Caporali R, Parazzini R, Zaninelli A, et al. Clinical presentation of osteoarthritis in general practice: determinants of pain in Italian patients in the AMICA study. *Arthritis Rheum* 2005; 35(suppl 1):17-23.
 25. Cryer PE, Davis SN, Shamooh H. Hypoglycemia in diabetes. *Diabetes Care* 2003; 26(6):1902-1912.
 26. Fuijoka K. Management of obesity as a chronic disease: nonpharmacologic, pharmacologic and surgical options. *Obes Res* 2002;10 (suppl2):116S-123S.
 27. Messier SP, Loeser RF, Miller GD, Morgan TM, Rejeski WJ, Sevick M, et al. Exercise and dietary weight loss in overweight and obese older adults with knee osteoarthritis. *Arthritis Rheum* 2004; 50:1501-1510.
 28. Ettinger WH, Davis MA, Neuhaus JM, Mallon KP. Long- term physical functioning in persons with knee osteoarthritis from NHANES I: effects of comorbid medical conditions. *J Clin Epidemiol* 1994; 47:809-815.
 29. Katz JN, Chan LC, Sangha O, Fossel AH, Bates DW. Can comorbidity be measured by questionnaires rather than medical record review? *Med Care* 1996; 34:73-84.
 30. Dunlop DD, Luons JS, Manheim LM, Song J, Chang RW. Arthritis and heart disease as risk factors for major depression: the role of functional limitations. *Med Care* 2004; 42:502-511.
 31. Graney MJ. The reciprocal relationship between disability and depression. *J Am Geriatr Soc* 2000; 48:452-453.