

VU Research Portal

Celiac disease

Rostami-Nejad, M.

2012

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Rostami-Nejad, M. (2012). *Celiac disease: Epidemiology, Genetic and Clinical Behavior in Iran*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Dissertation title: celiac disease, epidemiology, genetic and clinical behavior in Iran

Summary English

Celiac associated disorders and the questions around these associations seem to be very important for health professionals who deal with this condition. This is because this common systemic disorder has a multifactorial etiology with a multitude of symptoms and complications inside and outside small bowel. Almost all those reported conditions with an association with coeliac disease are or must be HLA-DQ2/DQ8 positive. That means that they all have a common characteristic sharing a similar genetic background. The environmental factors associated with a complex immune-genetics are leading to enteropathy that will be resulting in malabsorption syndrome. The co-morbidity between celiac disease and other disorders has been clearly established and there have been many reports of numerous intestinal and extra intestinal coexistent disorders with CD. In this short systematic review the nature of these associations has been explored and classified according to their pathogenesis.

Study on epidemiology, associations with other conditions like *H.pylori*, Rotaviruses, *Toxoplasma gondii* clinical behavior in Eastern countries and challenges in diagnosis and treatment of celiac disease are the outline of my research and training over the past 4-5 years. The study we performed so far includes 12 Original Article, 3 Review Articles and 2 Letters. According to these studies coeliac disease (CD) is as common in Iran as in Western countries with similar association pattern with a prevalence of around 1% in Iranian population. Our preliminary study on genetic showed that the proportion of HLA-DQ2 in coeliac patients was higher than control and this finding is in agreement with other studies in Western populations.