

VU Research Portal

Fat Distribution and Arterial Stiffness

Schouten, F.

2012

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Schouten, F. (2012). *Fat Distribution and Arterial Stiffness: The Amsterdam Growth and Health Longitudinal study*. Gildeprint Drukkerijen.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

1

General Introduction

INTRODUCTION

Excessive body fat and, in particular, a central pattern of fat distribution, is an independent predictor of cardiovascular risk [1-3]. Furthermore, it has been demonstrated that individuals with obesity are likely to have an increase in arterial stiffness [4], which is an independent predictor of cardiovascular disease [5,6]. Obesity-related increases in arterial stiffness have been proposed as one of the potential pathways through which (central) obesity could lead to cardiovascular disease (CVD) [4,7].

Arterial stiffness, what is it?

The arteries have two important functions. The first one is to act as a *conduit* from heart to arterioles, in order to deliver blood to tissues and organs according to their needs. The second function is to act as a *cushion*. By smoothing pressure fluctuations, flow variability imposed by the intermittent contracting heart is reduced, so that the blood is directed to organs and tissues in an almost steady stream [8].

Structurally, the arterial tree can be divided in two compartments: a (proximal) elastic and a (distal) muscular part. Although all arteries of the vascular tree participate in the two aforementioned functions, proximal arteries (i.e. the aorta and its main branches) have a dominant role in cushioning, whereas distal muscular arteries also contribute to flow regulation. Due to their anatomical structure, the proximal compartment is characterized by optimal elasticity/compliance, whereas the distal compartment is characterized by higher stiffness [9]. In accordance, ageing, drugs and risk factors have differential effects along the arterial tree [10-12]. Arterial stiffening impairs the ability of the arterial system to handle the pressure wave that occurs during cardiac election. Stiffening of the arterial tree leads to increased systolic blood pressure, decreased diastolic blood pressure, increased left ventricular mass, and decreased diastolic perfusion [13,14]. The principal clinical complications of arterial stiffness, such as myocardial infarction, stroke and heart failure, usually occur in middle-aged or older people. However, arterial stiffening already starts in childhood [15].

Body fatness and body composition, how do they contribute to CVD?

Central (or abdominal) fat can be distinguished into subcutaneous abdominal fat and intra-abdominal fat. It has been postulated that especially intra-abdominal fat (also referred to as 'visceral fat') is detrimental because it directly releases free fatty acids into the portal vein, causing reduced hepatic insulin sensitivity, increased gluconeogenesis and dyslipidemia [16,17]. In contrast, peripheral fat (i.e. limbs) has been suggested to have a protective effect through uptake and storage of free fatty acids [18].

The secretory functions of adipose tissue are probably also subject to regional variations [19,20]. Abdominal fat and peripheral fat may differ in secretion of adipose derived proteins that influence the metabolism, vascular function, vascular structure and this way arterial stiffness. The pathobiological mechanisms explaining why especially central body fatness is associated with arterial stiffness remains largely unknown, however. One mechanism may be an altered profile of adipokine secretion, such as hyperleptinaemia [21-24] and/or hypoadiponectinaemia [25-33]. As an alternative to the adipokine hypothesis, other adiposity-related factors may in part explain the relationship between central body fatness and arterial stiffness. These include low-grade systemic inflammation [14,27,34] and endothelial dysfunction [14,35].

These potential mechanisms might (partly) explain the relation between (abdominal) obesity and arterial stiffness, and therefore we investigated these mechanisms in this thesis.

Amsterdam Growth and Health Longitudinal Study

The study design of the present thesis has been developed within the ongoing *Amsterdam Growth and Health Longitudinal Study* (AGAHLS), an observational longitudinal study that started in 1976 with a total inclusion of 698 boys and girls [36-38]. Its initial goal was to describe the natural development of growth, health and lifestyle of adolescents, and to investigate longitudinal relationships between biological and lifestyle variables. The mean age of the subjects at the beginning of the study was 13.1 (± 0.8) years. Since then, subjects have been measured 3 to 9 times during a 30-year follow-up period. At each measurement, anthropometrical,

biological and lifestyle variables were assessed, as detailed elsewhere [36-38]. In 2000, when the subjects' mean age was 36.6 (± 0.6) years, the following measurements were added to the study for the first time: body composition (including body fat distribution) by means of dual-energy x-ray absorptiometry (DXA) and properties of three large arteries by non-invasive ultrasound imaging (Figure 1).




Figure 1. The Amsterdam Growth and Health Study design.

Objectives and outline of this thesis

Previously, it was unknown whether obesity-related vascular pathology also manifests itself in vessel wall alterations, and at what stage of life these changes could become apparent. Over the last years, studies that were mostly cross-sectional in design have shown (central) obesity to be independently related to increased arterial stiffness [39-41]. In addition, previous data from the AGAHL cohort, which consists of apparently healthy participants, demonstrated cross-sectionally that such effects can be detected relatively early in life (at a mean age of 36 years) [42,43]. At this age, 1.1% of the participants were underweight ($BMI < 18.5 \text{ kg/m}^2$), 65.3% normal weight ($BMI 18.5-25.0$), 28.5% overweight ($BMI 25-30$) and 5.1% obese ($BMI > 30$). This implies that obesity-related effects in arterial stiffness are not confined to individuals with overweight or obesity, but are seen across the entire range of levels of body mass/fatness. However, prospective studies on the effects of body fat

(distribution) on vessel wall properties as well as potential mechanistic insights into these associations in young, apparently healthy adults are limited.

Therefore, in the present thesis, we investigated the (longitudinal) association between body fatness, fat distribution and arterial stiffness, and in addition, we investigated a few candidate pathophysiological mechanisms of this relationship. Specifically, we formulated the following main research questions (as illustrated in Figure 2):

1. What are the associations of body fatness and body fat distribution, and changes herein during a 6-year follow-up, with (changes in) arterial stiffness?
2. Are these associations mediated by biomarkers of low-grade inflammation (i.e., C-reactive protein, serum amyloid A, interleukin 6, interleukin 8, tumor necrosis factor α and soluble intercellular adhesion molecule 1), and/or biomarkers of endothelial dysfunction (i.e., soluble endothelial selectin, thrombomodulin and both vascular- and intercellular adhesion molecules 1 and von Willebrand factor)?
3. Are these associations (from research question 1) mediated by adipokines (i.e., leptin, adiponectin)?




Figure 2. Illustration of the hypothesized associations investigated in this thesis.

In order to answer the above research questions, the outline of this thesis is as follows:

Chapter 2 addresses the associations between changes in central (i.e. trunk) fat versus peripheral (i.e. limbs) fat and lean masses with changes in arterial stiffness over a 6-year follow-up period.

In **Chapter 3**, we investigated the associations between, on the one hand, biomarkers of endothelial dysfunction (i.e., soluble endothelial selectin, thrombomodulin and both vascular- and intercellular adhesion molecules 1 and von Willebrand factor) and of low-grade inflammation (i.e., C-reactive protein, serum amyloid A, interleukin 6, interleukin 8, tumor necrosis factor α and soluble intercellular adhesion molecule 1) and, on the other hand, arterial stiffness, over a 6-year period.

Chapter 4 addresses the associations between leptin, adiponectin, and the leptin-to-adiponectin ratio (LAR) with arterial stiffness, over a 6-yr period.

In **Chapter 5** we investigated to what extent levels of adipokines (adiponectin, leptin, the leptin-to-adiponectin ratio (LAR)), biomarkers of low-grade inflammation and endothelial dysfunction explain the relationship between central fatness and arterial stiffness over a 6-year period.

In **Chapter 6** we studied to what extent the longitudinal development of fatness parameters (BMI, sum of skinfolds and skinfold ratio) from the age of 13 to 36 years precede favourable or unfavourable levels of adipokines at the age of 36 years. In addition, we investigated if we could identify a certain critical age/period in which the groups with favourable and unfavourable levels of adipokines began to differ substantially.

Finally, in the general discussion in **Chapter 7**, the main findings of the studies included in this thesis are discussed, including some methodological issues. Furthermore, the practical implications and suggestions for future research are addressed.

REFERENCES

1. Seidell JC, Verschuren WM, van Leer EM, Kromhout D. Overweight, underweight, and mortality. A prospective study of 48,287 men and women. *Arch Intern Med* 1996; 156:958-63.
2. Poirier P, Eckel RH. Obesity and cardiovascular disease. *Curr Atheroscler Rep* 2002; 4:448-53.
3. Grundy SM. Obesity, metabolic syndrome, and cardiovascular disease. *J Clin Endocrinol Metab* 2004; 89:2595-600.
4. Safar ME, Czernichow S, Blacher J. Obesity, arterial stiffness, and cardiovascular risk. *J Am Soc Nephrol* 2006; 17:S109-11.
5. Laurent S, Cockcroft J, Van Bortel L, Boutouyrie P, Giannattasio C, Hayoz D , et al. Expert consensus document on arterial stiffness: methodological issues and clinical applications. *Eur Heart J* 2006; 27:2588-605.
6. Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol* 2010; 55:1318-27.
7. Seals DR, Gates PE. Stiffening our resolve against adult weight gain. *Hypertension* 2005; 45:175-7.
8. Safar ME, O'Rourke MF. Arterial stiffness in hypertension.: Elsevier B.V. the Netherlands; 2006.
9. Safar ME, Levy BI, Struijker-Boudier H. Current perspectives on arterial stiffness and pulse pressure in hypertension and cardiovascular diseases. *Circulation* 2003; 107:2864-9.
10. Safar ME, editor. Arteries in clinical hypertension. . Philadelphia (PA) Lippincott-Raven; 1996.
11. Bortolotto LA, Hanon O, Franconi G, Boutouyrie P, Legrain S, Girerd X. The aging process modifies the distensibility of elastic but not muscular arteries. *Hypertension* 1999; 34:889-92.
12. Benetos A, Laurent S, Hoeks AP, Boutouyrie PH, Safar ME. Arterial alterations with aging and high blood pressure. A noninvasive study of carotid and femoral arteries. *Arterioscler Thromb* 1993; 13:90-7.
13. O'Rourke M. Mechanical principles in arterial disease. *Hypertension* 1995; 26:2-9.
14. Stehouwer CD, Henry RM, Ferreira I. Arterial stiffness in diabetes and the metabolic syndrome: a pathway to cardiovascular disease. *Diabetologia* 2008; 51:527-39.
15. Tounian P, Aggoun Y, Dubern B, Varille V, Guy-Grand B, Sidi D , et al. Presence of increased stiffness of the common carotid artery and endothelial dysfunction in severely obese children: a prospective study. *Lancet* 2001; 358:1400-4.
16. Despres JP, Lemieux S, Lamarche B, Prud'homme D, Moorjani S, Brun LD , et al. The insulin resistance-dyslipidemic syndrome: contribution of visceral obesity and therapeutic implications. *Int J Obes Relat Metab Disord* 1995; 19 Suppl 1:S76-86.
17. Bjorntorp P. "Portal" adipose tissue as a generator of risk factors for cardiovascular disease and diabetes. *Arteriosclerosis* 1990; 10:493-6.
18. Rebuffe-Scrive M, Enk L, Crona N, Lonnroth P, Abrahamsson L, Smith U, Bjorntorp P. Fat cell metabolism in different regions in women. Effect of menstrual cycle, pregnancy, and lactation. *J Clin Invest* 1985; 75:1973-6.
19. Arner P. Regional differences in protein production by human adipose tissue. *Biochem Soc Trans* 2001; 29:72-5.
20. Bays H, Mandarino L, DeFronzo RA. Role of the adipocyte, free fatty acids, and ectopic fat in pathogenesis of type 2 diabetes mellitus: peroxisomal proliferator-activated receptor agonists provide a rational therapeutic approach. *J Clin Endocrinol Metab* 2004; 89:463-78.
21. Singhal A, Farooqi IS, Cole TJ, O'Rahilly S, Fewtrell M, Kattenhorn M , et al. Influence of leptin on arterial distensibility: a novel link between obesity and cardiovascular disease? *Circulation* 2002; 106:1919-24.
22. Windham BG, Griswold ME, Farasat SM, Ling SM, Carlson O, Egan JM , et al. Influence of leptin, adiponectin, and resistin on the association between abdominal adiposity and arterial stiffness. *Am J Hypertens* 2010; 23:501-7.

23. Schutte R, Huisman HW, Schutte AE, Malan NT. Leptin is independently associated with systolic blood pressure, pulse pressure and arterial compliance in hypertensive African women with increased adiposity: the POWIRS study. *J Hum Hypertens* 2005; 19:535-41.
24. Scuteri A, Orru M, Morrell C, Piras MG, Taub D, Schlessinger D , et al. Independent and additive effects of cytokine patterns and the metabolic syndrome on arterial aging in the SardinIA Study. *Atherosclerosis* 2011; 215:459-64.
25. Snijder MB, Flyvbjerg A, Stehouwer CD, Frystyk J, Henry RM, Seidell JC , et al. Relationship of adiposity with arterial stiffness as mediated by adiponectin in older men and women: the Hoorn Study. *Eur J Endocrinol* 2009; 160:387-95.
26. Ohashi N, Ito C, Fujikawa R, Yamamoto H, Kihara Y, Kohno N. The impact of visceral adipose tissue and high-molecular weight adiponectin on cardio-ankle vascular index in asymptomatic Japanese subjects. *Metabolism* 2009; 58:1023-9.
27. Tsioufis C, Dimitriadis K, Selima M, Thomopoulos C, Mihas C, Skiadas I , et al. Low-grade inflammation and hypoadiponectinaemia have an additive detrimental effect on aortic stiffness in essential hypertensive patients. *Eur Heart J* 2007; 28:1162-9.
28. Nguyen QM, Srinivasan SR, Xu JH, Chen W, Berenson GS. Racial (black-white) divergence in the association between adiponectin and arterial stiffness in asymptomatic young adults: the Bogalusa heart study. *Am J Hypertens* 2008; 21:553-7.
29. Mahmud A, Feely J. Adiponectin and arterial stiffness. *Am J Hypertens* 2005; 18:1543-8.
30. Araki T, Emoto M, Yokoyama H, Maeno T, Hatsuda S, Mori K , et al. The association of plasma adiponectin level with carotid arterial stiffness. *Metabolism* 2006; 55:587-92.
31. Sung SH, Chuang SY, Sheu WH, Lee WJ, Chou P, Chen CH. Relation of adiponectin and high-sensitivity C-reactive protein to pulse-wave velocity and N-terminal pro-B-type natriuretic peptide in the general population. *Am J Cardiol* 2009; 103:1411-6.
32. Urbina EM, Khoury P, Martin LJ, D'Alessio D, Dolan LM. Gender differences in the relationships among obesity, adiponectin and brachial artery distensibility in adolescents and young adults. *Int J Obes (Lond)* 2009; 33:1118-25.
33. Higashiura K, Ura N, Ohata J, Togashi N, Takagi S, Saitoh S , et al. Correlations of adiponectin level with insulin resistance and atherosclerosis in Japanese male populations. *Clin Endocrinol (Oxf)* 2004; 61:753-9.
34. Diamant M, Lamb HJ, van de Ree MA, Endert EL, Groeneveld Y, Bots ML , et al. The association between abdominal visceral fat and carotid stiffness is mediated by circulating inflammatory markers in uncomplicated type 2 diabetes. *J Clin Endocrinol Metab* 2005; 90:1495-501.
35. Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW. C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol* 1999; 19:972-8.
36. Kemper H, editor. *Growth, Health and Fitness of Teenagers. Longitudinal research in international perspective.* . Basel (Switzerland): Karger; 1985. (Medicine and Sport Science).
37. Kemper H, editor. *Amsterdam Growth and Health Longitudinal Study (AGAHLs) A 23-Year Follow-Up from Teenager to Adult about Lifestyle and Health.* Basel (Switzerland): Karger; 2004. (Medicine and Sport Science).
38. Kemper HC, van Mechelen W, Post GB, Snel J, Twisk JW, van Lenthe FJ, Welten DC. The Amsterdam Growth and Health Longitudinal Study. The past (1976-1996) and future (1997-?). *Int J Sports Med* 1997; 18 Suppl 3:S140-50.
39. Sutton-Tyrrell K, Newman A, Simonsick EM, Havlik R, Pahor M, Lakatta E , et al. Aortic stiffness is associated with visceral adiposity in older adults enrolled in the study of health, aging, and body composition. *Hypertension* 2001; 38:429-33.
40. Wildman RP, Mackey RH, Bostom A, Thompson T, Sutton-Tyrrell K. Measures of obesity are associated with vascular stiffness in young and older adults. *Hypertension* 2003; 42:468-73.
41. Snijder MB, Henry RM, Visser M, Dekker JM, Seidell JC, Ferreira I , et al. Regional body composition as a determinant of arterial stiffness in the elderly: The Hoorn Study. *J Hypertens* 2004; 22:2339-47.

42. Ferreira I, Snijder MB, Twisk JW, van Mechelen W, Kemper HC, Seidell JC, Stehouwer CD. Central fat mass versus peripheral fat and lean mass: opposite (adverse versus favorable) associations with arterial stiffness? The Amsterdam Growth and Health Longitudinal Study. *J Clin Endocrinol Metab* 2004; 89:2632-9.
43. Ferreira I, Twisk JW, van Mechelen W, Kemper HC, Seidell JC, Stehouwer CD. Current and adolescent body fatness and fat distribution: relationships with carotid intima-media thickness and large artery stiffness at the age of 36 years. *J Hypertens* 2004; 22:145-55.