

## VU Research Portal

### **Genetic & Environmental risk factors for Obsessive-Compulsive Symptoms: Do they affect the same brain?**

den Braber, A.

2012

#### **document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

#### **citation for published version (APA)**

den Braber, A. (2012). *Genetic & Environmental risk factors for Obsessive-Compulsive Symptoms: Do they affect the same brain?* [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

#### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# References

- Abel KM, Drake R, Goldstein JM. Sex differences in schizophrenia. *Int Rev Psychiatry* 2010; 22: 417-28.
- Adler CM, McDonough-Ryan P, Sax KW, Holland SK, Arndt S, Strakowski SM. fMRI of neuronal activation with symptom provocation in unmedicated patients with obsessive compulsive disorder. *J Psychiatr Res* 2000; 34: 317-24.
- Albert U, Maina G, Bogetto F, Ravizza L. The role of recent life events in the onset of obsessive-compulsive disorder. *CNS Spectr* 2000; 5: 44-50.
- Alexander GE, DeLong MR, Strick PL. Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu Rev Neurosci* 1986; 9: 357-81.
- Allen JS, Damasio H, Grabowski TJ, Bruss J, Zhang W. Sexual dimorphism and asymmetries in the gray-white composition of the human cerebrum. *Neuroimage* 2003; 18: 880-94.
- Alonso P, Menchon JM, Mataix-Cols D, Pifarre J, Urretavizcaya M, Crespo JM, et al. Perceived parental rearing style in obsessive-compulsive disorder: relation to symptom dimensions. *Psychiatry res.* 2004; 127: 267-78.
- Alptekin K, Degirmenci B, Kivircik B, Durak H, Yemez B, Derebek E, et al. Tc-99m HMPAO brain perfusion SPECT in drug-free obsessive-compulsive patients without depression. *Psychiatry res.* 2001; 107: 51-6.
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorder: DSM-IV (4th ed.)*. Washington, DC: American Psychiatric Association 1994.
- Aouizerate B, Guehl D, Cuny E, Rougier A, Bioulac B, Tignol J, et al. Pathophysiology of obsessive-compulsive disorder: a necessary link between phenomenology, neuropsychology, imagery and physiology. *Prog Neurobiol* 2004; 72: 195-221.
- Arnold PD, Sicard T, Burroughs E, Richter MA, Kennedy JL. Glutamate transporter gene SLC1A1 associated with obsessive-compulsive disorder. *Arch Gen Psychiatry* 2006; 63: 769-76.
- Ashburner J, Friston KJ. Voxel-based morphometry--the methods. *Neuroimage* 2000; 11: 805-21.
- Ashburner J, Friston KJ. Why voxel-based morphometry should be used. *Neuroimage* 2001; 14: 1238-43.
- Atmaca M, Yildirim H, Ozdemir H, Tezcan E, Poyraz AK. Volumetric MRI study of key brain regions implicated in obsessive-compulsive disorder. *Prog Neuro-psychopharmacol Biol Psychiatry* 2007; 31: 46-52.
- Baker SC, Rogers RD, Owen AM, Frith CD, Dolan RJ, Frackowiak RSJ, et al. Neural systems engaged by planning: A PET study of the Tower of London task. *Neuropsychologia* 1996; 34: 515-26.
- Bannon S, Gonsalvez CJ, Croft RJ, Boyce PM. Executive functions in obsessive-compulsive disorder: state or trait deficits? *Aust N Z J Psychiatry* 2006; 40: 1031-8.

- Bao AM, Swaab DF. Sex differences in the brain, behavior, and neuropsychiatric disorders. *Neuroscientist* 2010; 16: 550-65.
- Basser PJ, Pierpaoli C. Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor MRI. *J Magn Reson B* 1996; 111: 209-19.
- Beaulieu C. The basis of anisotropic water diffusion in the nervous system - a technical review. *NMR Biomed* 2002; 15: 435-55.
- Bechara A, Damasio H, Damasio AR. Emotion, decision making and the orbitofrontal cortex. *Cereb Cortex* 2000; 10: 295-307.
- Beck AT, Rial WY, Rickels K. Short form of depression inventory: cross-validation. *Psychol Rep* 1974; 34: 1184-6.
- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961; 4: 561-71.
- Bekker MH, van Mens-Verhulst J. Anxiety disorders: sex differences in prevalence, degree, and background, but gender-neutral treatment. *Gend Med* 2007; 4 Suppl B: S178-S193.
- Bengel D, Greenberg BD, Cora-Locatelli G, Altemus M, Heils A, Li Q, et al. Association of the serotonin transporter promoter regulatory region polymorphism and obsessive-compulsive disorder. *Mol Psychiatry* 1999; 4: 463-6.
- Berntson GG, Bigger JT, Jr., Eckberg DL, Grossman P, Kaufmann PG, Malik M, et al. Heart rate variability: origins, methods, and interpretive caveats. *Psychophysiology* 1997; 34: 623-48.
- Bhattacharyya PK, Lowe MJ. Cardiac-induced physiologic noise in tissue is a direct observation of cardiac-induced fluctuations. *Magn Reson Imaging* 2004; 22: 9-13.
- Billett EA, Richter MA, Sam F, Swinson RP, Dai XY, King N, et al. Investigation of dopamine system genes in obsessive-compulsive disorder. *Psychiatr Genet* 1998; 8: 163-9.
- Binder JR, Rao SM, Hammeke TA, Yetkin FZ, Jesmanowicz A, Bandettini PA, et al. Functional magnetic resonance imaging of human auditory cortex. *Ann Neurol* 1994; 35: 662-72.
- Birn RM, Diamond JB, Smith MA, Bandettini PA. Separating respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in fMRI. *Neuroimage* 2006; 31: 1536-48.
- Bishop KM, Wahlsten D. Sex differences in the human corpus callosum: myth or reality? *Neurosci Biobehav Rev* 1997; 21: 581-601.
- Bogetto F, Venturello S, Albert U, Maina G, Ravizza L. Gender-related clinical differences in obsessive-compulsive disorder. *Eur Psychiatry* 1999; 14: 434-41.
- Boly M, Faymonville ME, Peigneux P, Lambermont B, Damas P, Del FG, et al. Auditory processing in severely brain injured patients: differences between the minimally conscious state and the persistent vegetative state. *Arch Neurol* 2004; 61: 233-8.

- Boomsma DI, de Geus EJ, Vink JM, Stubbe JH, Distel MA, Hottenga JJ, et al. Netherlands Twin Register: from twins to twin families. *Twin Res Hum Genet* 2006; 9: 849-57.
- Boomsma DI, Vink JM, van Beijsterveldt TC, de Geus EJ, Beem AL, Mulder EJ, et al. Netherlands Twin Register: a focus on longitudinal research. *Twin Res* 2002; 5: 401-6.
- Breiter HC, Rauch SL, Kwong KK, Baker JR, Weisskoff RM, Kennedy DN, et al. Functional magnetic resonance imaging of symptom provocation in obsessive-compulsive disorder. *Arch Gen Psychiatry* 1996; 53: 595-606.
- Brown MA, Semelka RC. *MRI: Basic Principles and Applications*, 4 ed. Hoboken, NJ: John Wiley & Sons Inc 2010.
- Brunetti M, Babiloni C, Ferretti A, Del GC, Merla A, Olivetti BM, et al. Hypothalamus, sexual arousal and psychosexual identity in human males: a functional magnetic resonance imaging study. *Eur J Neurosci* 2008; 27: 2922-7.
- Buchsbaum MS, Buchsbaum BR, Chokron S, Tang C, Wei TC, Byne W. Thalamocortical circuits: fMRI assessment of the pulvinar and medial dorsal nucleus in normal volunteers. *Neuroscience letters* 2006; 404: 282-7.
- Burgaleta M, Head K, Alvarez-Linera J, Martinez K, Escorial S, Haier R, et al. Sex differences in brain volume are related to specific skills, not to general intelligence. *Intelligence* 2011. In press.
- Burgess PW, Gilbert SJ, Dumontheil I. Function and localization within rostral prefrontal cortex (area 10). *Philos Trans R Soc Lond B Biol Sci* 2007; 362: 887-99.
- Busatto GF, Buchpiguel CA, Zamignani DR, Garrido GE, Glabus MF, Rosario-Campos MC, et al. Regional cerebral blood flow abnormalities in early-onset obsessive-compulsive disorder: an exploratory SPECT study. *J Am Acad Child Adolesc Psychiatry* 2001; 40: 347-54.
- Buss DM. Desires in human mating. *Ann N Y Acad Sci* 2000; 907: 39-49.
- Buxton RB. *Introduction to functional magnetic resonance imaging. Principles and Techniques*, 2 ed. Cambridge, NY: Cambridge University Press 2009.
- Cabeza R, Nyberg L. *Imaging cognition II: An empirical review of 275 PET and fMRI studies*. *J Cog Neuroscience* 2000; 12: 1-47.
- Cannistraro PA, Makris N, Howard JD, Wedig MM, Hodge SM, Wilhelm S, et al. A diffusion tensor imaging study of white matter in obsessive-compulsive disorder. *Depress Anxiety* 2006.
- Carlsson ML. On the role of prefrontal cortex glutamate for the antithetical phenomenology of obsessive compulsive disorder and attention deficit hyperactivity disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2001; 25: 5-26.
- Castle DJ, Deale A, Marks IM. Gender differences in obsessive compulsive disorder. *Aust N Z J Psychiatry* 1995; 29: 114-7.
- Cath DC, Spinhoven P, Hoogduin CA, Landman AD, van Woerkom TC, van de

- Wetering BJ, et al. Repetitive behaviors in Tourette's syndrome and OCD with and without tics: what are the differences? *Psychiatry res.* 2001; 101: 171-85.
- Cath DC, van Grootheest DS, Willemsen G, van Oppen P, Boomsma DI. Environmental factors in obsessive-compulsive behavior: evidence from discordant and concordant monozygotic twins. *Behav Genet* 2008; 38: 108-20.
- Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and behavioural correlates. *Brain* 2006; 129: 564-83.
- Chamberlain SR, Blackwell AD, Fineberg NA, Robbins TW, Sahakian BJ. The neuropsychology of obsessive compulsive disorder: the importance of failures in cognitive and behavioural inhibition as candidate endophenotypic markers. *Neurosci Biobehav Rev* 2005; 29: 399-419.
- Chamberlain SR, Fineberg NA, Blackwell AD, Clark L, Robbins TW, Sahakian BJ. A neuropsychological comparison of obsessive-compulsive disorder and trichotillomania. *Neuropsychologia* 2007; 45: 654-62.
- Chamberlain SR, Menzies L. Endophenotypes of obsessive-compulsive disorder: rationale, evidence and future potential. *Expert Rev Neurother* 2009; 9: 1133-46.
- Chang C, Cunningham JP, Glover GH. Influence of heart rate on the BOLD signal: the cardiac response function. *Neuroimage* 2009; 44: 857-69.
- Chen XL, Xie JX, Han HB, Cui YH, Zhang BQ. MR perfusion-weighted imaging and quantitative analysis of cerebral hemodynamics with symptom provocation in unmedicated patients with obsessive-compulsive disorder. *Neuroscience letters* 2004; 370: 206-11.
- Chou KH, Cheng Y, Chen IY, Lin CP, Chu WC. Sex-linked white matter microstructure of the social and analytic brain. *Neuroimage* 2011; 54: 725-33.
- Chouinard PA, Paus T. The primary motor and premotor areas of the human cerebral cortex. *Neuroscientist* 2006; 12: 143-52.
- Ciesielski KT, Hamalainen MS, Lesnik PG, Geller DA, Ahlfors SP. Increased MEG activation in OCD reflects a compensatory mechanism specific to the phase of a visual working memory task. *Neuroimage* 2005; 24: 1180-91.
- Clifford CA, Murray RM, Fulker DW. Genetic and environmental influences on obsessional traits and symptoms. *Psychol Med* 1984; 14: 791-800.
- Conners CK, Erhardt D, Sparrow E. *Conners' Adult ADHD Rating Scales (CAARS)*. North Tonawanda, NY: Multi-Health systems Inc 1999; 144.
- Cosgrove KP, Mazure CM, Staley JK. Evolving knowledge of sex differences in brain structure, function, and chemistry. *Biol Psychiatry* 2007; 62: 847-55.
- Crespo-Facorro B, Roiz-Santianez R, Perez-Iglesias R, Mata I, Rodriguez-Sanchez JM, Tordesillas-Gutierrez D, et al. Sex-specific variation of MRI-based cortical morphometry in adult healthy volunteers: the effect on cognitive functioning. *Prog Neuropsychopharmacol Biol Psychiatry* 2011; 35: 616-23.
- Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ. Neural systems supporting

- interoceptive awareness. *Nat Neurosci* 2004; 7: 189-95.
- Cummings JL. Frontal-subcortical circuits and human behavior. *Arch Neurol* 1993; 50: 873-80.
- Cummings JL. Anatomic and behavioral aspects of frontal-subcortical circuits. *Ann N Y Acad Sci* 1995; 769: 1-13.
- Dagher A, Owen AM, Boecker H, Brooks DJ. Mapping the network for planning: a correlational PET activation study with the Tower of London task. *Brain* 1999; 122 ( Pt 10): 1973-87.
- de Courten-Myers GM. The human cerebral cortex: gender differences in structure and function. *J Neuropathol Exp Neurol* 1999; 58: 217-26.
- de Geus EJ, van't Ent D, Wolfensberger SP, Heutink P, Hoogendijk WJ, Boomsma DI, et al. Intrapair differences in hippocampal volume in monozygotic twins discordant for the risk for anxiety and depression. *Biol Psychiatry* 2007; 61: 1062-71.
- de Mathis MA, de AP, Funaro G, Torresan RC, Moraes I, Torres AR, et al. Gender differences in obsessive-compulsive disorder: a literature review. *Rev Bras Psiquiatr* 2011; 33: 390-9.
- de Munck JC, Goncalves SI, Faes TJ, Kuijter JP, Pouwels PJ, Heethaar RM, et al. A study of the brain's resting state based on alpha band power, heart rate and fMRI. *Neuroimage* 2008; 42: 112-21.
- Delorme R, Gousse V, Roy I, Trandafir A, Mathieu F, Mouren-Simeoni MC, et al. Shared executive dysfunctions in unaffected relatives of patients with autism and obsessive-compulsive disorder. *Eur Psychiatry* 2007; 22: 32-8.
- den Braber A, van 't Ent D, Boomsma DI, Cath DC, Veltman DJ, Thompson PM, et al. White Matter Differences in Monozygotic Twins Discordant or Concordant for Obsessive-Compulsive Symptoms: A Combined Diffusion Tensor Imaging/Voxel-Based Morphometry Study. *Biol Psychiatry* 2011.
- den Braber A, van 't Ent D, Cath DC, Wagner J, Boomsma DI, de Geus EJ. Brain activation during cognitive planning in twins discordant or concordant for obsessive-compulsive symptoms. *Brain* 2010; 133: 3123-40.
- den Braber A, van't Ent D, Blokland GA, van Grootheest DS, Cath DC, Veltman DJ, et al. An fMRI study in monozygotic twins discordant for obsessive-compulsive symptoms. *Biol Psychol* 2008; 79: 91-102.
- Denys D, de GF, van Megen HJ, Westenberg HG. Symptom dimensions in obsessive-compulsive disorder: factor analysis on a clinician-rated scale and a self-report measure. *Psychopathology* 2004; 37: 181-9.
- Dickel DE, Veenstra-Vanderweele J, Cox NJ, Wu X, Fischer DJ, Van Etten-Lee M, et al. Association testing of the positional and functional candidate gene SLC1A1/EAAC1 in early-onset obsessive-compulsive disorder. *Arch Gen Psychiatry* 2006; 63: 778-85.
- Dillon DG, Pizzagalli DA. Inhibition of Action, Thought, and Emotion: A Selective Neurobiological Review. *Appl Prev Psychol* 2007; 12: 99-114.

- Diwadkar VA, Carpenter PA, Just MA. Collaborative activity between parietal and dorso-lateral prefrontal cortex in dynamic spatial working memory revealed by fMRI. *Neuroimage* 2000; 12: 85-99.
- Eagly AH, Wood W. The origins of sex differences in human behavior - Evolved dispositions versus social roles. *American Psychologist* 1999; 54: 408-23.
- Eley TC, Bolton D, O'Connor TG, Perrin S, Smith P, Plomin R. A twin study of anxiety-related behaviours in pre-school children. *J Child Psychol Psychiatry* 2003; 44: 945-60.
- Enoch MA, Greenberg BD, Murphy DL, Goldman D. Sexually dimorphic relationship of a 5-HT2A promoter polymorphism with obsessive-compulsive disorder. *Biol Psychiatry* 2001; 49: 385-8.
- Faw B. Pre-frontal executive committee for perception, working memory, attention, long-term memory, motor control, and thinking: a tutorial review. *Consciousness and cognition* 2003; 12: 83-139.
- Filipek PA, Richelme C, Kennedy DN, Caviness VS, Jr. The young adult human brain: an MRI-based morphometric analysis. *Cereb Cortex* 1994; 4: 344-60.
- Fineberg NA, Potenza MN, Chamberlain SR, Berlin HA, Menzies L, Bechara A, et al. Probing compulsive and impulsive behaviors, from animal models to endophenotypes: a narrative review. *Neuropsychopharmacology* 2010; 35: 591-604.
- Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A* 2000; 97: 11050-5.
- Fitzgerald KD, Welsh RC, Gehring WJ, Abelson JL, Himle JA, Liberzon I, et al. Error-related hyperactivity of the anterior cingulate cortex in obsessive-compulsive disorder. *Biol Psychiatry* 2005; 57: 287-94.
- Friedlander L, Desrocher M. Neuroimaging studies of obsessive-compulsive disorder in adults and children. *Clinical Psychology Review* 2006; 26: 32-49.
- Frost RO, Steketee G. Perfectionism in obsessive-compulsive disorder patients. *Behav Res Ther* 1997; 35: 291-6.
- Fullana MA, Mataix-Cols D, Caspi A, Harrington H, Grisham JR, Moffitt TE, et al. Obsessions and compulsions in the community: prevalence, interference, help-seeking, developmental stability, and co-occurring psychiatric conditions. *Am J Psychiatry* 2009; 166: 329-36.
- Garibotto V, Scifo P, Gorini A, Alonso CR, Brambati S, Bellodi L, et al. Disorganization of anatomical connectivity in obsessive compulsive disorder: a multi-parameter diffusion tensor imaging study in a subpopulation of patients. *Neurobiol Dis* 2010; 37: 468-76.
- Geisser S, Greenhouse SW. An Extension of Box's Results on the Use of the F Distribution in Multivariate-Analysis. *Annals of Mathematical Statistics* 1958; 29: 885-91.
- Geller DA, Wieland N, Carey K, Vivas F, Petty CR, Johnson J, et al. Perinatal factors affecting expression of obsessive compulsive disorder in children and



- adolescents. *J Child Adolesc Psychopharmacol* 2008; 18: 373-9.
- Giedd JN, Castellanos FX, Rajapakse JC, Vaituzis AC, Rapoport JL. Sexual dimorphism of the developing human brain. *Prog Neuropsychopharmacol Biol Psychiatry* 1997; 21: 1185-201.
- Giedd JN, Rapoport JL, Garvey MA, Perlmutter S, Swedo SE. MRI assessment of children with obsessive-compulsive disorder or tics associated with streptococcal infection. *Am J Psychiatry* 2000; 157: 281-3.
- Gilbert AR, Moore GJ, Keshavan MS, Paulson LA, Narula V, Mac Master FP, et al. Decrease in thalamic volumes of pediatric patients with obsessive-compulsive disorder who are taking paroxetine. *Arch Gen Psychiatry* 2000; 57: 449-56.
- Glover GH, Li TQ, Ress D. Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. *Magn Reson Med* 2000; 44: 162-7.
- Goldman-Rakic PS. Topography of cognition: parallel distributed networks in primate association cortex. *Annu Rev Neurosci* 1988; 11: 137-56.
- Goldstein JM, Seidman LJ, Horton NJ, Makris N, Kennedy DN, Caviness VS, Jr., et al. Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic resonance imaging. *Cereb Cortex* 2001; 11: 490-7.
- Good CD, Johnsrude I, Ashburner J, Henson RN, Friston KJ, Frackowiak RS. Cerebral asymmetry and the effects of sex and handedness on brain structure: a voxel-based morphometric analysis of 465 normal adult human brains. *Neuroimage* 2001; 14: 685-700.
- Goodale MA, Milner AD. Separate Visual Pathways for Perception and Action. *Trends in neurosciences* 1992; 15: 20-5.
- Goodman WK, Price LH, Rasmussen SA, Mazure C, Delgado P, Heninger GR, et al. The Yale-Brown Obsessive Compulsive Scale. II. Validity. *Arch Gen Psychiatry* 1989a; 46: 1012-6.
- Goodman WK, Price LH, Rasmussen SA, Mazure C, Fleischmann RL, Hill CL, et al. The Yale-Brown Obsessive Compulsive Scale. I. Development, use, and reliability. *Arch Gen Psychiatry* 1989b; 46: 1006-11.
- Grabe HJ, Meyer C, Hapke U, Rumpf HJ, Freyberger HJ, Dilling H, et al. Prevalence, quality of life and psychosocial function in obsessive-compulsive disorder and subclinical obsessive-compulsive disorder in northern Germany. *Eur Arch Psychiatry Clin Neurosci* 2000; 250: 262-8.
- Grasby PM, Frith CD, Friston KJ, Bench C, Frackowiak RS, Dolan RJ. Functional mapping of brain areas implicated in auditory-verbal memory function. *Brain* 1993; 116 ( Pt 1): 1-20.
- Graybiel AM, Rauch SL. Toward a neurobiology of obsessive-compulsive disorder. *Neuron* 2000; 28: 343-7.
- Grisham JR, Anderson TM, Sachdev PS. Genetic and environmental influences on obsessive-compulsive disorder. *Eur Arch Psychiatry Clin Neurosci* 2008; 258: 107-16.

- Gu BM, Park JY, Kang DH, Lee SJ, Yoo SY, Jo HJ, et al. Neural correlates of cognitive inflexibility during task-switching in obsessive-compulsive disorder. *Brain* 2008; 131: 155-64.
- Gur RC, Gunning-Dixon F, Bilker WB, Gur RE. Sex differences in temporo-limbic and frontal brain volumes of healthy adults. *Cereb Cortex* 2002; 12: 998-1003.
- Gur RC, Turetsky BI, Matsui M, Yan M, Bilker W, Hughett P, et al. Sex differences in brain gray and white matter in healthy young adults: correlations with cognitive performance. *J Neurosci* 1999; 19: 4065-72.
- Ha TH, Kang DH, Park JS, Jang JH, Jung WH, Choi JS, et al. White matter alterations in male patients with obsessive-compulsive disorder. *Neuroreport* 2009; 20: 735-9.
- Haber SN. The primate basal ganglia: parallel and integrative networks. *J Chem Neuroanat* 2003; 26: 317-30.
- Halpern DF. Sex differences in intelligence. Implications for education. *Am Psychol* 1997; 52: 1091-102.
- Heijmans BT, Tobi EW, Lumey LH, Slagboom PE. The epigenome: Archive of the prenatal environment. *Epigenetics* 2009; 4.
- Herrero MT, Barcia C, Navarro JM. Functional anatomy of thalamus and basal ganglia. *Childs Nerv Syst* 2002; 18: 386-404.
- Hesse S, Muller U, Lincke T, Barthel H, Villmann T, Angermeyer MC, et al. Serotonin and dopamine transporter imaging in patients with obsessive-compulsive disorder. *Psychiatry res.* 2005; 140: 63-72.
- Hettema JM, Neale MC, Kendler KS. A review and meta-analysis of the genetic epidemiology of anxiety disorders. *AM J Psychiatry* 2001; 158: 1568-78.
- Hoshi E, Tanji J. Integration of target and body-part information in the premotor cortex when planning action. *Nature* 2000; 408: 466-70.
- Hudziak JJ, VAN Beijsterveldt CE, Althoff RR, Stanger C, Rettew DC, Nelson EC, et al. Genetic and environmental contributions to the Child Behavior Checklist Obsessive-Compulsive Scale: a cross-cultural twin study. *Arch Gen Psychiatry* 2004; 61: 608-16.
- Husted DS, Shapira NA, Goodman WK. The neurocircuitry of obsessive-compulsive disorder and disgust. *Prog Neuro-psychopharmacol Biol Psychiatry* 2006; 30: 389-99.
- Im K, Lee JM, Lee J, Shin YW, Kim IY, Kwon JS, et al. Gender difference analysis of cortical thickness in healthy young adults with surface-based methods. *Neuroimage* 2006; 31: 31-8.
- Insel TR, Winslow JT. Neurobiology of obsessive compulsive disorder. *Psychiatr Clin North Am* 1992; 15: 813-24.
- Iwamura Y. Hierarchical somatosensory processing. *Curr Opin Neurobiol.* 1998; 8: 522-8.
- Jonnal AH, Gardner CO, Prescott CA, Kendler KS. Obsessive and compulsive symptoms in a general population sample of female twins. *Am J Med Genet*

- 2000; 96: 791-6.
- Kang X, Herron TJ, Woods DL. Regional variation, hemispheric asymmetries and gender differences in pericortical white matter. *Neuroimage* 2011; 56: 2011-23.
- Karno M, Golding JM, Sorenson SB, Burnam MA. The epidemiology of obsessive-compulsive disorder in five US communities. *Arch Gen Psychiatry* 1988; 45: 1094-9.
- Kastner S, Pinsk MA. Visual attention as a multilevel selection process. *Cogn Affect Behav Neurosci* 2004; 4: 483-500.
- Katura T, Tanaka N, Obata A, Sato H, Maki A. Quantitative evaluation of interrelations between spontaneous low-frequency oscillations in cerebral hemodynamics and systemic cardiovascular dynamics. *Neuroimage* 2006; 31: 1592-600.
- Kerns JG, Cohen JD, MacDonald AW, III, Cho RY, Stenger VA, Carter CS. Anterior cingulate conflict monitoring and adjustments in control. *Science* 2004; 303: 1023-6.
- Kim JJ, Lee MC, Kim J, Kim IY, Kim SI, Han MH, et al. Grey matter abnormalities in obsessive-compulsive disorder: statistical parametric mapping of segmented magnetic resonance images. *Br J Psychiatry* 2001; 179: 330-4.
- Kitamura H, Shioiri T, Kimura T, Ohkubo M, Nakada T, Someya T. Parietal white matter abnormalities in obsessive-compulsive disorder: a magnetic resonance spectroscopy study at 3-Tesla. *Acta Psychiatr Scand* 2006; 114: 101-8.
- Koechlin E, Hyafil A. Anterior prefrontal function and the limits of human decision-making. *Science* 2007; 318: 594-8.
- Kuelz AK, Hohagen F, Voderholzer U. Neuropsychological performance in obsessive-compulsive disorder: a critical review. *Biol Psychol* 2004; 65: 185-236.
- Kwon JS, Kim JJ, Lee DW, Lee JS, Lee DS, Kim MS, et al. Neural correlates of clinical symptoms and cognitive dysfunctions in obsessive-compulsive disorder. *Psychiatry res.* 2003; 122: 37-47.
- Labad J, Menchon JM, Alonso P, Segalas C, Jimenez S, Jaurrieta N, et al. Gender differences in obsessive-compulsive symptom dimensions. *Depress Anxiety* 2008; 25: 832-8.
- Laberge D, Buchsbaum MS. Positron Emission Tomographic Measurements of Pulvinar Activity During An Attention Task. *J Neurosci* 1990; 10: 613-9.
- Lacerda AL, Dalgalarondo P, Caetano D, Camargo EE, Etchebehere EC, Soares JC. Elevated thalamic and prefrontal regional cerebral blood flow in obsessive-compulsive disorder: a SPECT study. *Psychiatry res.* 2003; 123: 125-34.
- Lacoste-Utamsing C, Holloway RL. Sexual dimorphism in the human corpus callosum. *Science* 1982; 216: 1431-2.
- Laplane D, Levasseur M, Pillon B, Dubois B, Baulac M, Mazoyer B, et al. Obsessive-compulsive and other behavioural changes with bilateral basal ganglia

- lesions. A neuropsychological, magnetic resonance imaging and positron tomography study. *Brain* 1989; 112 ( Pt 3): 699-725.
- Lazeron RH, Rombouts SA, Machielsen WC, Scheltens P, Witter MP, Uylings HB, et al. Visualizing brain activation during planning: the tower of London test adapted for functional MR imaging. *AJNR Am J Neuroradiol* 2000; 21: 1407-14.
- Le Bihan D, Mangin JF, Poupon C, Clark CA, Pappata S, Molko N, et al. Diffusion tensor imaging: concepts and applications. *J Magn Reson Imaging* 2001; 13: 534-46.
- Leichnetz GR. Connections of the medial posterior parietal cortex (area 7m) in the monkey. *Anat Rec* 2001; 263: 215-36.
- Lenroot RK, Giedd JN. Sex differences in the adolescent brain. *Brain Cogn* 2010; 72: 46-55.
- Lensi P, Cassano GB, Correddu G, Ravagli S, Kunovac JL, Akiskal HS. Obsessive-compulsive disorder. Familial-developmental history, symptomatology, comorbidity and course with special reference to gender-related differences. *Br J Psychiatry* 1996; 169: 101-7.
- Lin H, Katsovich L, Ghebremichael M, Findley DB, Grantz H, Lombroso PJ, et al. Psychosocial stress predicts future symptom severities in children and adolescents with Tourette syndrome and/or obsessive-compulsive disorder. *J Child Psychol Psychiatry* 2007; 48: 157-66.
- Locke S, Angevine JB, Jr., Yakovlev PI. Limbic nuclei of thalamus and connections of limbic cortex. II. Thalamocortical projection of the lateral dorsal nucleus in man. *Arch Neurol* 1961; 4: 355-64.
- Loring-Meier S, Halpern DF. Sex differences in visuospatial working memory: components of cognitive processing. *Psychon Bull Rev* 1999; 6: 464-71.
- Lucey JV, Costa DC, Blanes T, Busatto GF, Pilowsky LS, Takei N, et al. Regional cerebral blood flow in obsessive-compulsive disordered patients at rest. Differential correlates with obsessive-compulsive and anxious-avoidant dimensions. *Br J Psychiatry* 1995; 167: 629-34.
- Luders E, Narr KL, Thompson PM, Rex DE, Jancke L, Steinmetz H, et al. Gender differences in cortical complexity. *Nat Neurosci* 2004; 7: 799-800.
- Luders E, Narr KL, Thompson PM, Rex DE, Woods RP, Deluca H, et al. Gender effects on cortical thickness and the influence of scaling. *Hum Brain Mapp* 2006; 27: 314-24.
- Lv B, Li J, He H, Li M, Zhao M, Ai L, et al. Gender consistency and difference in healthy adults revealed by cortical thickness. *Neuroimage* 2010; 53: 373-82.
- MacDonald AW III, Cohen JD, Stenger VA, Carter CS. Dissociating the role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. *Science* 2000; 288: 1835-8.
- Magno E, Foxe JJ, Molholm S, Robertson IH, Garavan H. The anterior cingulate and error avoidance. *J Neurosci* 2006; 26: 4769-73.
- Maltby N, Tolin DF, Worhunsky P, O'Keefe TM, Kiehl KA. Dysfunctional action

- monitoring hyperactivates frontal-striatal circuits in obsessive-compulsive disorder: an event-related fMRI study. *Neuroimage* 2005; 24: 495-503.
- Mann VA, Sasanuma S, Sakuma N, Masaki S. Sex differences in cognitive abilities: a cross-cultural perspective. *Neuropsychologia* 1990; 28: 1063-77.
- Mars RB, Bestmann S, Rothwell JC, Haggard P. Effects of motor preparation and spatial attention on corticospinal excitability in a delayed-response paradigm. *Exp Brain Res*. 2007; 182: 125-9.
- Martin N, Boomsma D, Machin G. A twin-pronged attack on complex traits. *Nat Genet* 1997; 17: 387-92.
- Mataix-Cols D, van den Heuvel OA. Common and distinct neural correlates of obsessive-compulsive and related disorders. *Psychiatr Clin North Am* 2006; 29: 391-410, viii.
- Mataix-Cols D, Wooderson S, Lawrence N, Brammer MJ, Speckens A, Phillips ML. Distinct neural correlates of washing, checking, and hoarding symptom dimensions in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2004; 61: 564-76.
- McCarthy MM, Arnold AP. Reframing sexual differentiation of the brain. *Nat Neurosci* 2011; 14: 677-83.
- Melcher T, Falkai P, Gruber O. Functional brain abnormalities in psychiatric disorders: neural mechanisms to detect and resolve cognitive conflict and interference. *Brain Res Rev* 2008; 59: 96-124.
- Menzies L, Achard S, Chamberlain SR, Fineberg N, Chen CH, Del Campo N, et al. Neurocognitive endophenotypes of obsessive-compulsive disorder. *Brain* 2007.
- Menzies L, Chamberlain SR, Laird AR, Thelen SM, Sahakian BJ, Bullmore ET. Integrating evidence from neuroimaging and neuropsychological studies of obsessive-compulsive disorder: the orbitofronto-striatal model revisited. *Neurosci Biobehav Rev* 2008a; 32: 525-49.
- Menzies L, Williams GB, Chamberlain SR, Ooi C, Fineberg N, Suckling J, et al. White matter abnormalities in patients with obsessive-compulsive disorder and their first-degree relatives. *Am J Psychiatry* 2008b; 165: 1308-15.
- Menzler K, Belke M, Wehrmann E, Krakow K, Lengler U, Jansen A, et al. Men and women are different: diffusion tensor imaging reveals sexual dimorphism in the microstructure of the thalamus, corpus callosum and cingulum. *Neuroimage* 2011; 54: 2557-62.
- Michael GA, Buron V. The human pulvinar and stimulus-driven attentional control. *Behav Neurosci* 2005; 119: 1353-67.
- Miguel EC, Leckman JF, Rauch S, do Rosario-Campos MC, Hounie AG, Mercadante MT, et al. Obsessive-compulsive disorder phenotypes: implications for genetic studies. *Mol Psychiatry* 2005; 10: 258-75.
- Mori S, Zhang J. Principles of diffusion tensor imaging and its applications to basic neuroscience research. *Neuron* 2006; 51: 527-39.

- Nakamae T, Narumoto J, Sakai Y, Nishida S, Yamada K, Nishimura T, et al. Diffusion tensor imaging and tract-based spatial statistics in obsessive-compulsive disorder. *J Psychiatr Res* 2011; 45: 687-90.
- Nakamae T, Narumoto J, Shibata K, Matsumoto R, Kitabayashi Y, Yoshida T, et al. Alteration of fractional anisotropy and apparent diffusion coefficient in obsessive-compulsive disorder: a diffusion tensor imaging study. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32: 1221-6.
- Nakao T, Nakagawa A, Yoshiura T, Nakatani E, Nabeyama M, Sanematsu H, et al. Duration effect of obsessive-compulsive disorder on cognitive function: a functional MRI study. *Depress Anxiety* 2009; 26: 814-23.
- Nakao T, Nakagawa A, Yoshiura T, Nakatani E, Nabeyama M, Yoshizato C, et al. A functional MRI comparison of patients with obsessive-compulsive disorder and normal controls during a Chinese character Stroop task. *Psychiatry res.* 2005; 139: 101-14.
- Nestadt G, Samuels J, Riddle M, Bienvenu OJ, III, Liang KY, LaBuda M, et al. A family study of obsessive-compulsive disorder. *Arch Gen Psychiatry* 2000; 57: 358-63.
- Newman SD, Carpenter PA, Varma S, Just MA. Frontal and parietal participation in problem solving in the Tower of London: fMRI and computational modeling of planning and high-level perception. *Neuropsychologia* 2003; 41: 1668-82.
- Nicolini H, Arnold P, Nestadt G, Lanzagorta N, Kennedy JL. Overview of genetics and obsessive-compulsive disorder. *Psychiatry Res* 2009; 170: 7-14.
- Nobre AC, Allison T, McCarthy G. Word Recognition in the Human Inferior Temporal-Lobe. *Nature* 1994; 372: 260-3.
- Nopoulos P, Flaum M, O'Leary D, Andreasen NC. Sexual dimorphism in the human brain: evaluation of tissue volume, tissue composition and surface anatomy using magnetic resonance imaging. *Psychiatry Res* 2000; 98: 1-13.
- Noshirvani HF, Kasvikis Y, Marks IM, Tsakiris F, Monteiro WO. Gender-divergent aetiological factors in obsessive-compulsive disorder. *Br J Psychiatry* 1991; 158: 260-3.
- O'Donnell S, Noseworthy MD, Levine B, Dennis M. Cortical thickness of the frontopolar area in typically developing children and adolescents. *Neuroimage* 2005; 24: 948-54.
- Ogawa S, Tank DW, Menon R, Ellermann JM, Kim SG, Merkle H, et al. Intrinsic signal changes accompanying sensory stimulation: functional brain mapping with magnetic resonance imaging. *Proc Natl Acad Sci U S A* 1992; 89: 5951-5.
- Page LA, Rubia K, Deeley Q, Daly E, Toal F, Mataix-Cols D, et al. A functional magnetic resonance imaging study of inhibitory control in obsessive-compulsive disorder. *Psychiatry Res* 2009; 174: 202-9.
- Parker G, Brotchie H. Gender differences in depression. *Int Rev Psychiatry* 2010; 22: 429-36.
- Pauls DL. The genetics of obsessive-compulsive disorder: a review. *Dialogues Clin*

- Neurosci 2010; 12: 149-63.
- Pauls DL, Alsobrook JP, Goodman W, Rasmussen S, Leckman JF. A family study of obsessive-compulsive disorder. *Am J Psychiatry* 1995; 152: 76-84.
- Pauls DL, Towbin KE, Leckman JF, Zahner GE, Cohen DJ. Gilles de la Tourette's syndrome and obsessive-compulsive disorder. Evidence supporting a genetic relationship. *Arch Gen Psychiatry* 1986; 43: 1180-2.
- Peper JS, Brouwer RM, Schnack HG, van Baal GC, van LM, van den Berg SM, et al. Sex steroids and brain structure in pubertal boys and girls. *Psychoneuroendocrinology* 2009a; 34: 332-42.
- Peper JS, Brouwer RM, van Baal GC, Schnack HG, van LM, Boomsma DI, et al. Does having a twin brother make for a bigger brain? *Eur J Endocrinol* 2009b; 160: 739-46.
- Petrides M, Pandya DN. Projections to the frontal cortex from the posterior parietal region in the rhesus monkey. *J Comp Neurol* 1984; 228: 105-16.
- Pollatos O, Gramann K, Schandry R. Neural systems connecting interoceptive awareness and feelings. *Hum Brain Mapp* 2007; 28: 9-18.
- Pujol J, Soriano-Mas C, Alonso P, Cardoner N, Menchon JM, Deus J, et al. Mapping structural brain alterations in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2004; 61: 720-30.
- Purcell R, Maruff P, Kyrios M, Pantelis C. Cognitive deficits in obsessive-compulsive disorder on tests of frontal-striatal function. *Biol Psychiatry* 1998a; 43: 348-57.
- Purcell R, Maruff P, Kyrios M, Pantelis C. Neuropsychological deficits in obsessive-compulsive disorder: a comparison with unipolar depression, panic disorder, and normal controls. *Arch Gen Psychiatry* 1998b; 55: 415-23.
- Rachman S, de Silva P. Abnormal and normal obsessions. *Behaviour research and therapy* 1978; 16: 233-48.
- Radua J, Mataix-Cols D. Voxel-wise meta-analysis of grey matter changes in obsessive-compulsive disorder. *Br J Psychiatry* 2009; 195: 393-402.
- Radua J, van den Heuvel OA, Surguladze S, Mataix-Cols D. Meta-analytical comparison of voxel-based morphometry studies in obsessive-compulsive disorder vs other anxiety disorders. *Arch Gen Psychiatry* 2010; 67: 701-11.
- Ramnani N, Owen AM. Anterior prefrontal cortex: insights into function from anatomy and neuroimaging. *Nat Rev Neurosci* 2004; 5: 184-94.
- Rao NP, Reddy YC, Kumar KJ, Kandavel T, Chandrashekar CR. Are neuropsychological deficits trait markers in OCD? *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32: 1574-9.
- Rauch SL, Wedig MM, Wright CI, Martis B, McMullin KG, Shin LM, et al. Functional magnetic resonance imaging study of regional brain activation during implicit sequence learning in obsessive-compulsive disorder. *Biol Psychiatry* 2007; 61: 330-6.
- Remijnse PL, Nielen MM, van Balkom AJ, Cath DC, van Oppen P, Uylings HB, et al. Reduced orbitofrontal-striatal activity on a reversal learning task in obsessive-

- compulsive disorder. *Arch Gen Psychiatry* 2006; 63: 1225-36.
- Reynolds CA, Turner A, Gatz M, Pedersen NL. Comparative rating measures of health and environmental exposures: how well do twins agree? *Twin Res Hum Genet* 2005; 8: 113-9.
- Ridderinkhof KR, van den Wildenberg WP, Segalowitz SJ, Carter CS. Neurocognitive mechanisms of cognitive control: the role of prefrontal cortex in action selection, response inhibition, performance monitoring, and reward-based learning. *Brain Cogn* 2004; 56: 129-40.
- Rosenberg DR, Keshavan MS. A.E. Bennett Research Award. Toward a neurodevelopmental model of obsessive-compulsive disorder. *Biol Psychiatry* 1998; 43: 623-40.
- Rosenberg DR, Keshavan MS, Dick EL, Bagwell WW, Macmaster FP, Birmaher B. Corpus callosal morphology in treatment-naive pediatric obsessive compulsive disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 1997; 21: 1269-83.
- Rotge JY, Guehl D, Dilharreguy B, Tignol J, Bioulac B, Allard M, et al. Meta-analysis of brain volume changes in obsessive-compulsive disorder. *Biol Psychiatry* 2009; 65: 75-83.
- Rowe JB, Owen AM, Johnsrude IS, Passingham RE. Imaging the mental components of a planning task. *Neuropsychologia* 2001; 39: 315-27.
- Rucklidge JJ. Gender differences in attention-deficit/hyperactivity disorder. *Psychiatr Clin North Am* 2010; 33: 357-73.
- Saito Y, Nobuhara K, Okugawa G, Takase K, Sugimoto T, Horiuchi M, et al. Corpus callosum in patients with obsessive-compulsive disorder: diffusion-tensor imaging study. *Radiology* 2008; 246: 536-42.
- Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RS, Busa E, et al. Thinning of the cerebral cortex in aging. *Cereb Cortex* 2004; 14: 721-30.
- Salkovskis PM, Harrison J. Abnormal and normal obsessions--a replication. *Behav Res Ther* 1984; 22: 549-52.
- Sanavio E. Obsessions and compulsions: the Padua Inventory. *Behav Res Ther* 1988; 26: 169-77.
- Savage CR, Baer L, Keuthen NJ, Brown HD, Rauch SL, Jenike MA. Organizational strategies mediate nonverbal memory impairment in obsessive-compulsive disorder. *Biol Psychiatry* 1999a; 45: 905-16.
- Savage CR, Deckersbach T, Wilhelm S, Rauch SL, Baer L, Jenike MA. Learning strategies disrupt verbal and nonverbal memory in OCD. *Biol Psychiatry* 1999b; 45: 185.
- Saxena S, Brody AL, Ho ML, Alborzian S, Ho MK, Maidment KM, et al. Cerebral metabolism in major depression and obsessive-compulsive disorder occurring separately and concurrently. *Biol Psychiatry* 2001; 50: 159-70.
- Saxena S, Brody AL, Maidment KM, Smith EC, Zohrabi N, Katz E, et al. Cerebral glucose metabolism in obsessive-compulsive hoarding. *AM J Psychiatry*



- 2004; 161: 1038-48.
- Saxena S, Rauch SL. Functional neuroimaging and the neuroanatomy of obsessive-compulsive disorder. *Psychiatr Clin North Am* 2000; 23: 563-86.
- Schienze A, Schafer A, Stark R, Walter B, Vaitl D. Neural responses of OCD patients towards disorder-relevant, generally disgust-inducing and fear-inducing pictures. *Int J Psychophysiol* 2005; 57: 69-77.
- Schlosser RG, Wagner G, Schachtzabel C, Peikert G, Koch K, Reichenbach JR, et al. Fronto-cingulate effective connectivity in obsessive compulsive disorder: A study with fMRI and dynamic causal modeling. *Hum Brain Mapp* 2010; 31: 1834-50.
- Schmithorst VJ, Holland SK, Dardzinski BJ. Developmental differences in white matter architecture between boys and girls. *Hum Brain Mapp* 2008; 29: 696-710.
- Schultz RT, Evans DW, Wolff M. Neuropsychological models of childhood obsessive-compulsive disorder. *Child Adolesc Psychiatr Clin N Am* 1999; 8: 513-31, viii.
- Schultz W, Dayan P, Montague PR. A neural substrate of prediction and reward. *Science* 1997; 275: 1593-9.
- Shallice T. Specific impairments of planning. *Philos Trans R Soc Lond B Biol Sci* 1982; 298: 199-209.
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 1998; 59 Suppl 20: 22-33.
- Shenton ME, Dickey CC, Frumin M, McCarley RW. A review of MRI findings in schizophrenia. *Schizophr Res* 2001; 49: 1-52.
- Sherman SM, Guillery RW. The role of the thalamus in the flow of information to the cortex. *Philos Trans R Soc Lond B Biol Sci*. 2002; 357: 1695-708.
- Shin YW, Kim DJ, Ha TH, Park HJ, Moon WJ, Chung EC, et al. Sex differences in the human corpus callosum: diffusion tensor imaging study. *Neuroreport* 2005; 16: 795-8.
- Shmueli K, van GP, de Zwart JA, Horovitz SG, Fukunaga M, Jansma JM, et al. Low-frequency fluctuations in the cardiac rate as a source of variance in the resting-state fMRI BOLD signal. *Neuroimage* 2007; 38: 306-20.
- Singer HS, Minzer K. Neurobiology of Tourette's syndrome: concepts of neuroanatomic localization and neurochemical abnormalities. *Brain Dev* 2003; 25 Suppl 1: S70-S84.
- Sisk CL, Zehr JL. Pubertal hormones organize the adolescent brain and behavior. *Front Neuroendocrinol* 2005; 26: 163-74.
- Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE, et al. Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data. *Neuroimage* 2006; 31: 1487-505.
- Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TE, Johansen-Berg

- H, et al. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 2004; 23 Suppl 1: S208-S219.
- Soriano-Mas C, Pujol J, Alonso P, Cardoner N, Menchon JM, Harrison BJ, et al. Identifying patients with obsessive-compulsive disorder using whole-brain anatomy. *Neuroimage* 2007; 35: 1028-37.
- Sowell ER, Peterson BS, Kan E, Woods RP, Yoshii J, Bansal R, et al. Sex differences in cortical thickness mapped in 176 healthy individuals between 7 and 87 years of age. *Cereb Cortex* 2007; 17: 1550-60.
- Spielberger CD, Gorsuch RL, Lusehe RE. Manual for the State-Trait Anxiety Inventory (STAI). Palo Alto, CA: Consulting Psychologists Press; 1970.
- Spielberger CD, Jacobs GA, Russell S, Crane RS. Butcher J, Spielberger CD, editors. Assessment of anger: The State Trait Anger Scale. Hillsdale NJ: Lawrence Erlbaum Associates; 1983.
- Stewart SE, Fagerness JA, Platko J, Smoller JW, Scharf JM, Illmann C, et al. Association of the SLC1A1 glutamate transporter gene and obsessive-compulsive disorder. *Am J Med Genet B Neuropsychiatr Genet* 2007a; 144B: 1027-33.
- Stewart SE, Platko J, Fagerness J, Birns J, Jenike E, Smoller JW, et al. A genetic family-based association study of OLIG2 in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2007b; 64: 209-14.
- Sullivan EV, Rosenbloom MJ, Desmond JE, Pfefferbaum A. Sex differences in corpus callosum size: relationship to age and intracranial size. *Neurobiol Aging* 2001; 22: 603-11.
- Swaab DF, Chung WC, Kruijver FP, Hofman MA, Ishunina TA. Sexual differentiation of the human hypothalamus. *Adv Exp Med Biol* 2002; 511: 75-100.
- Szeszko PR, Ardekani BA, Ashtari M, Malhotra AK, Robinson DG, Bilder RM, et al. White matter abnormalities in obsessive-compulsive disorder: a diffusion tensor imaging study. *Arch Gen Psychiatry* 2005; 62: 782-90.
- Szeszko PR, MacMillan S, McMeniman M, Chen S, Baribault K, Lim KO, et al. Brain structural abnormalities in psychotropic drug-naïve pediatric patients with obsessive-compulsive disorder. *AM J Psychiatry* 2004; 161: 1049-56.
- Szeszko PR, Vogel J, Ashtari M, Malhotra AK, Bates J, Kane JM, et al. Sex differences in frontal lobe white matter microstructure: a DTI study. *Neuroreport* 2003; 14: 2469-73.
- Thompson PM, Cannon TD, Narr KL, van ET, Poutanen VP, Huttunen M, et al. Genetic influences on brain structure. *Nat Neurosci* 2001; 4: 1253-8.
- Toga AW, Thompson PM. Genetics of brain structure and intelligence. *Annu Rev Neurosci* 2005; 28: 1-23.
- Togao O, Yoshiura T, Nakao T, Nabeyama M, Sanematsu H, Nakagawa A, et al. Regional gray and white matter volume abnormalities in obsessive-compulsive disorder: a voxel-based morphometry study. *Psychiatry Res* 2010; 184: 29-37.
- Tong Y, Lindsey KP, deB FB. Partitioning of physiological noise signals in the brain

- with concurrent near-infrared spectroscopy and fMRI. *J Cereb Blood Flow Metab* 2011; 31: 2352-62.
- Tukel R, Polat A, Genc A, Bozkurt O, Atli H. Gender-related differences among Turkish patients with obsessive-compulsive disorder. *Compr Psychiatry* 2004; 45: 362-6.
- Ursu S, Stenger VA, Shear MK, Jones MR, Carter CS. Overactive action monitoring in obsessive-compulsive disorder: evidence from functional magnetic resonance imaging. *Psychol Sci* 2003; 14: 347-53.
- Valente AA Jr., Miguel EC, Castro CC, Amaro E, Jr., Duran FL, Buchpiguel CA, et al. Regional gray matter abnormalities in obsessive-compulsive disorder: a voxel-based morphometry study. *Biol Psychiatry* 2005; 58: 479-87.
- van 't Ent D, van Beijsterveldt TC, Derks EM, Hudziak JJ, Veltman DJ, Todd RD, et al. Neuroimaging of response interference in twins concordant or discordant for inattention and hyperactivity symptoms. *Neuroscience* 2009.
- van den Heuvel OA, Groenewegen HJ, Barkhof F, Lazeron RH, van Dyck R, Veltman DJ. Frontostriatal system in planning complexity: a parametric functional magnetic resonance version of Tower of London task. *Neuroimage* 2003; 18: 367-74.
- van den Heuvel OA, Remijnse PL, Mataix-Cols D, Vrenken H, Groenewegen HJ, Uylings HB, et al. The major symptom dimensions of obsessive-compulsive disorder are mediated by partially distinct neural systems. *Brain* 2009; 132: 853-68.
- van den Heuvel OA, Veltman DJ, Groenewegen HJ, Cath DC, van Balkom AJ, van Hartkamp J, et al. Frontal-striatal dysfunction during planning in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2005a; 62: 301-9.
- van den Heuvel OA, Veltman DJ, Groenewegen HJ, Witter MP, Merkelbach J, Cath DC, et al. Disorder-specific neuroanatomical correlates of attentional bias in obsessive-compulsive disorder, panic disorder, and hypochondriasis. *Arch Gen Psychiatry* 2005b; 62: 922-33.
- van Grootheest DS, Cath DC, Beekman AT, Boomsma DI. Twin studies on obsessive-compulsive disorder: a review. *Twin Res Hum Genet* 2005; 8: 450-8.
- van Grootheest DS, Cath DC, Beekman AT, Boomsma DI. Genetic and environmental influences on obsessive-compulsive symptoms in adults: a population-based twin-family study. *Psychol Med* 2007; 37: 1635-44.
- van Houdt PJ, Ossenblok PP, Boon PA, Leijten FS, Velis DN, Stam CJ, et al. Correction for pulse height variability reduces physiological noise in functional MRI when studying spontaneous brain activity. *Hum Brain Mapp* 2010; 31: 311-25.
- van Oppen P. Obsessions and compulsions: dimensional structure, reliability, convergent and divergent validity of the Padua Inventory. *Behav Res Ther* 1992; 30: 631-7.
- van Oppen P, Arntz A. Cognitive therapy for obsessive-compulsive disorder. *Behav Res Ther* 1994; 32: 79-87.

- van Oppen P, Hoekstra RJ, Emmelkamp PM. The structure of obsessive-compulsive symptoms. *Behav Res Ther* 1995; 33: 15-23.
- Veale DM, Sahakian BJ, Owen AM, Marks IM. Specific cognitive deficits in tests sensitive to frontal lobe dysfunction in obsessive-compulsive disorder. *Psychol Med* 1996; 26: 1261-9.
- Viard A, Flament MF, Artiges E, Dehaene S, Naccache L, Cohen D, et al. Cognitive control in childhood-onset obsessive-compulsive disorder: a functional MRI study. *Psychol Med* 2005; 35: 1007-17.
- Wendland JR, Moya PR, Timpano KR, Anavitarte AP, Kruse MR, Wheaton MG, et al. A haplotype containing quantitative trait loci for SLC1A1 gene expression and its association with obsessive-compulsive disorder. *Arch Gen Psychiatry* 2009; 66: 408-16.
- Wicker B, Keysers C, Plailly J, Royet JP, Gallese V, Rizzolatti G. Both of us disgusted in My insula: the common neural basis of seeing and feeling disgust. *Neuron* 2003; 40: 655-64.
- Wilcox HC, Grados M, Samuels J, Riddle MA, Bienvenu OJ, III, Pinto A, et al. The association between parental bonding and obsessive compulsive disorder in offspring at high familial risk. *J Affect Disord* 2008; 111: 31-9.
- Windischberger C, Langenberger H, Sycha T, Tschernko EM, Fuchsjaeger-Mayerl G, Schmetterer L, et al. On the origin of respiratory artifacts in BOLD-EPI of the human brain. *Magn Reson Imaging* 2002; 20: 575-82.
- Wolfensberger SP, Veltman DJ, Hoogendijk WJ, Boomsma DI, de Geus EJ. Amygdala responses to emotional faces in twins discordant or concordant for the risk for anxiety and depression. *Neuroimage* 2008; 41: 544-52.
- Yakovlev PI, Locke S. Limbic nuclei of thalamus and connections of limbic cortex. III. Corticocortical connections of the anterior cingulate gyrus, the cingulum, and the subcallosal bundle in monkey. *Arch Neurol* 1961; 5: 364-400.
- Yeterian EH, Pandya DN. Striatal connections of the parietal association cortices in rhesus monkeys. *J Comp Neurol* 1993; 332: 175-97.
- Yoo SY, Jang JH, Shin YW, Kim DJ, Park HJ, Moon WJ, et al. White matter abnormalities in drug-naive patients with obsessive-compulsive disorder: a diffusion tensor study before and after citalopram treatment. *Acta Psychiatr Scand* 2007; 116: 211-9.
- Yucel M, Harrison BJ, Wood SJ, Fornito A, Wellard RM, Pujol J, et al. Functional and biochemical alterations of the medial frontal cortex in obsessive-compulsive disorder. *Arch Gen Psychiatry* 2007; 64: 946-55.