

## VU Research Portal

### **Cardiovascular magnetic resonance imaging and computed tomography in patients with suspected coronary artery disease**

Groothuis, J.G.J.

2013

#### **document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

#### **citation for published version (APA)**

Groothuis, J. G. J. (2013). *Cardiovascular magnetic resonance imaging and computed tomography in patients with suspected coronary artery disease*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

#### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# Addendum 3

## Traumatic myocardial infarction visualized by computed tomography angiography


Jan G.J. Groothuis, Martijn R. Meijerink, Albert C. van Rossum

*Netherlands Heart Journal 2012 Dec;20(12):516-7.*

A 38-year old woman was shot in her chest with a handgun. Upon arrival in the emergency department computed tomography angiography (CTA) was performed.

The entry location of the bullet was clearly visible (A, arrow) with bilateral hemato-pneumothorax with atelectasis and consolidation of both lungs. Image B showed the pathway of the bullet, rupturing the left anterior descending coronary artery (arrow). On the arterial and late phase images there was no enhancement of the anteroseptal and anterior myocardial wall visible (A and C, arrowheads), suggestive of myocardial infarction. Electrocardiography (D) showed ST-segment elevation in leads V2-5. Trans-esophageal echocardiography revealed severe pericardial effusion and akinesia of the left ventricular anterior and anteroseptal myocardial wall. During emergency thoracotomy, traumatic rupture of the mid left anterior descending coronary artery was confirmed. Coronary bypass surgery was performed by a venous graft on aorta and distal left anterior descending coronary artery.

Previously, it was shown that CTA can detect healed myocardial infarcts. (1) This case shows how CTA can visualize an acute traumatic myocardial infarction.



**Figure 1.** Computed tomography angiography images in transverse orientation (A-C) showing the entry location (A, arrow) and pathway of the bullet rupturing the left anterior descending coronary artery (B, arrow). On the arterial and late phase images there was no enhancement of the anteroseptal and anterior myocardial wall visible (A and C, arrowheads), suggestive of myocardial infarction. Electrocardiography (D) showed ST-segment elevation in leads V2-5. \* Indicates left ventricle.

## REFERENCES

1. Henneman MM, Schuijff JD, Dibbets-Schneider P, Stokkel MP, van der Geest RJ, van der Wall EE, Bax JJ. Comparison of multislice computed tomography to gated single-photon emission computed tomography for imaging of healed myocardial infarcts. *Am J Cardiol.* 2008;101:144-148