

VU Research Portal

Neuroendocrine regulation in PCOS

Hendriks, M.L.

2014

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Hendriks, M. L. (2014). *Neuroendocrine regulation in PCOS*. [, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Chapter 2

Why does ovarian surgery in PCOS help? Insight into the endocrine implications of ovarian surgery for ovulation induction in polycystic ovary syndrome

Marja-Liisa Hendriks
Hans J.C.F. Ket
Peter G.A. Hompes
Roy Homburg
Cornelis B. Lambalk

Hum Reprod Update 2007;13(3):249-64

ABSTRACT

Polycystic ovary syndrome (PCOS) is a complex disorder with heterogeneity of clinical and endocrine features. Ovarian surgery for ovulation induction has been used in the management of clomifene citrate-resistant anovulatory women with PCOS. Various types of ovarian surgery have been employed (wedge resection, electrocautery, laser vaporization, multiple ovarian biopsies and others) and all operations result in an altered endocrine profile after surgery. The mechanism behind the reversal of endocrinological dysfunction in PCOS after ovarian surgery remains incompletely understood. This review scans the literature systematically to identify the endocrine changes after ovarian surgery in PCOS, in order to glean some knowledge of the mechanism involved. After ovarian surgery in PCOS, a rapid reduction of all ovarian hormones is seen, in combination with increased serum levels of pituitary hormones. Folliculogenesis is then initiated and ovarian hormone production increases, synchronically with a reduction of pituitary hormones. Continuation of follicle growth in subsequent cycles after ovarian surgery occurs in an environment with less androgens and lower LH and FSH levels compared with pretreatment levels. The endocrine changes found after ovarian surgery in PCOS women seem to be governed by the ovaries themselves. Rapid reduction of all ovarian hormones restores feedback to the hypothalamus and pituitary, resulting in appropriate gonadotropin secretion. Initiation of follicular development seems to be induced by increasing FSH levels following a reduction of the follicle excess and (intra-ovarian) androgen levels. Additionally, anti-Müllerian hormone and gonadotropin surge attenuating factor probably have a role in the endocrine changes.

INTRODUCTION

Polycystic ovary syndrome (PCOS) is the most common cause of anovulatory infertility (Adams et al., 1986) and is classically characterized by clinical symptoms of hyperandrogenism (hirsutism and acne) and oligo- or amenorrhoea. The presentation of PCOS symptoms is very heterogeneous, varies from mild to severe and has a wide spectrum. The confusion caused by this heterogeneity of clinical and endocrine features regarding the definition of PCOS will hopefully be settled by the adoption of the recent Rotterdam consensus (2004). Common endocrine abnormalities in PCOS include chronic high luteinising hormone (LH) levels (Berger et al., 1975; Panidis et al., 2005), hyperandrogenism

(Wild et al., 1985; Kumar et al., 2005), hyperinsulinaemia, insulin resistance (Dunaif et al., 1989) and dyslipidoproteinaemia (Wild et al., 1985; Wild, 2002). These endocrine disturbances interfere with folliculogenesis, are (partly) responsible for oligo- or anovulation and are likely to constitute an increased risk for cardiovascular diseases and diabetes (Legro et al., 1999; Elting et al., 2001). In spite of years of research, PCOS remains incompletely understood.

Ovarian surgery for ovulation induction restores the menstrual cycle in the majority of patients suffering from PCOS and favourably changes their endocrine profile (Cohen, 1996). Seventy years after the first report of successful surgical ovarian intervention in PCOS (Stein and Leventhal, 1935), the mechanism behind the reversal of endocrinological dysfunction after ovarian surgery remains unclear. Surgical ovarian intervention for PCOS is sparingly used at the present time, as acceptable ovulation and pregnancy rates are achieved using clomifene citrate, metformin and gonadotropins. However, some women remain anovulatory or cannot be successfully treated medically, and ovarian surgery becomes an option.

Types of ovarian surgery

Several surgical approaches for restoring ovulation in women with PCOS have been studied over the years, for example classical wedge resection, multiple ovarian biopsies, laser vaporization and electrocautery. All types of ovarian surgery share a common goal of creating ovarian damage and from an endocrine point of view can be seen as equivalent procedures (Cohen, 1996). A clear distinction has to be made, however, between ovarian drilling to induce ovulation and more extensive ovarian drilling to prevent ovarian stimulation syndrome. The latter method requires more profound ovarian destruction, done by using a lot of power to 'pepper pot' the ovaries, whereas the methods for ovulation induction attempt to use the lowest possible dose to achieve the desired effect. Within this review, only the studies with ovarian surgery for ovulation induction are discussed.

Mechanisms involved

The cause for reversal of the endocrinological dysfunction after ovarian surgery in PCOS is unclear. Many theories have been proposed for the cause of the re-establishment of menstrual cycles after ovarian surgery. Originally, the removal of a mechanical barrier has been postulated (Katz et al., 1978; Ben Shlomo et al., 1998) and reducing the size of the ovary was thought to allow gonadotropins to act more effectively after the ovarian surgery (Katz et al., 1978). Others have suggested that surgery may cause increased blood flow to the ovaries, resulting in increased delivery of gonadotropins (Cohen, 1996; Takeuchi et al., 2002). Reduction of androgens after ovarian surgery causes lower peripheral aromatization to estrogens and could theoretically result in restoring feedback to the hypothalamus and pituitary (Felemban et al., 2000; Takeuchi et al., 2002). More recent theories include gonadotropin surge attenuating factor (GnSAF, also called gonadotropin surge inhibiting factor: GnSIF). GnSAF is a hormone produced by the ovaries (Messinis et al., 1991) and its function lies in regulating and suppressing LH secretion by reducing pituitary sensitivity (Fowler et al., 2003). A deficiency of GnSAF in PCOS patients has been hypothesized as the cause of elevated LH levels (Balen and Jacobs, 1991; de Koning et al., 2001). Ovarian surgery could cause increased insulin-like growth factor-I (IGF-I) production, which interacts with follicle stimulating hormone (FSH) and results in follicular development. Follicle growth causes increased GnSAF production, and GnSAF subsequently suppresses LH secretion (Balen and Jacobs, 1994). Another suggested mechanism is reduced inhibin levels after ovarian surgery leading to increased FSH levels (Al Ojaimi, 2004). Less is known about the importance of Anti-Müllerian hormone (AMH) regarding ovarian surgery in PCOS. AMH is an ovarian product and a local inhibitor of FSH action (Durlinger et al., 2001, 2002) and may play a role in restoration of ovulation after ovarian surgery.

No extended review has yet been published to augment the endocrine implications of ovarian surgery and this review combines the existing literature (systematically searched) to identify essential endocrine changes after ovarian surgery in PCOS in order to glean some knowledge of the mechanism involved and further insight into the pathophysiology of PCOS.

MATERIALS AND METHODS

A systematic literature search was conducted in PubMed (January 2006), Embase.com and the Cochrane Library-Wiley (December 2005) databases. Three groups of search terms were used in every possible spelling, as synonyms, acronyms, key- or text words and these groups were combined with an 'AND'-operator. (i) PCOS, (ii) surgical interventions: (electro)cautery, (electro)coagulation, diathermy, drilling, laparoscopic ovarian drilling (LOD), laser, biopsy, excision, wedge, endoscopy, laparoscopy, surgery and (iii) hormones: LH, FSH, testosterone, gonadotropins, gonadotropin releasing hormone (GnRH), prolactin, androgens, androstenedione, dehydroepiandrosterone

(sulphate) (DHEA(S)), dihydrotestosterone (DHT), sex hormone-binding globulin (SHBG), estrogen, progesterone, anti-Müllerian hormone (AMH), gonadotropin surge-inhibiting factor/gonadotrophin surge-attenuating factor (GnSIF/GnSAF), hormones, endocrinology. Case reports and animal-studies were excluded. The full search strategy can be requested from the corresponding author.

A trial was eligible for inclusion if it was a publication in English and the treatment consisted of an ovarian surgery procedure to induce or facilitate ovulation induction in subfertile women with established PCOS, in combination with endocrine results before and after surgery. Study description of characteristics of PCOS subject had to be in accordance with the Rotterdam consensus (2004) criteria. Two reviewers selected the eligible studies, when discordance existed, a uniform decision was made by re-evaluating the study. A total of 63 papers were eligible for this review. Figures shown in this review represent weighted average hormonal values on indicated days, constructed by weighing the mean value of an individual study with their group size (Hazewinkel, 2002) and were constructed when multiple studies were available at different time points to summarize the effect of the surgery. The figures represent an estimate of endocrine changes after ovarian surgery and are shown without units, due to limitations of combining data from different studies with different hormonal assay methodology.

2

RESULTS

Ovarian hormones

Testosterone and androstenedione

Androgens are frequently and classically elevated in PCOS, with or without clinical symptoms (DeVane et al., 1975; Kumar et al., 2005). The potent androgens testosterone and androstenedione are both secreted by the ovary and the adrenal gland (Burger, 2002) and testosterone is additionally derived from peripheral production (Burger, 2002). Considerable amounts of testosterone are bound to SHBG, only the unbound proportion is biologically active (free) testosterone (Burger, 2002).

Testosterone

Serum testosterone concentrations in PCOS patients decreased after ovarian surgery from post-operative day 1 (Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987; Greenblatt and Casper, 1987; van der Weiden and Alberda, 1987; van der Weiden et al., 1989; Tasaka et al., 1990; Utsunomiya et al., 1990; Kovacs et al., 1991; Gadir et al., 1992; Campo et al., 1993; Naether et al., 1993; Liguori et al., 1996; Parsanezhad et al., 2005). One study showed a non-significant decrease one day after surgery (Kojima et al., 1989). One study showed increased testosterone values three hours after surgery preceding a rapid decrease on the first postoperative day (Judd et al., 1976). Testosterone concentrations reached a nadir around

3 days after surgery (Greenblatt and Casper, 1987; Campo et al., 1993; Liguori et al., 1996), followed by a small increase, but never reaching pretreatment values.

In the first days, weeks and years following ovarian surgery, testosterone levels remained low in the majority of the studies (Vejlsted and Albrechtsen, 1976; Aakvaag and Gjonnaess, 1985; Greenblatt and Casper, 1987; van der Weiden and Alberda, 1987; Sumioki et al., 1988; van der Weiden et al., 1989; Armar et al., 1990; Gadir et al., 1990; Keckstein et al., 1990; Tasaka et al., 1990; Utsunomiya et al., 1990; Kovacs et al., 1991; Rossmanith et al., 1991; Abdel et al., 1993; Campo et al., 1993; Naether et al., 1993; Tiitinen et al., 1993; Verhelst et al., 1993; Alborzi, 1994; Liguori et al., 1996; Taskin et al., 1996; Anttila et al., 1998; Gjonnaess, 1998, 1999; Kaaijk et al., 1999; Soliman et al., 2000; Wu et al., 2000, 2004; Zullo et al., 2000; Alborzi et al., 2001; Amer et al., 2002a; Asada et al., 2002; Takeuchi et al., 2002; Amin et al., 2003; Duleba et al., 2003; Malkawi et al., 2003; Parsanezhad et al., 2003; Al Ojaimi, 2004; Api et al., 2005; Kucuk and Kilic-Okman, 2005; Malkawi and Qublan, 2005; Parsanezhad et al., 2005). A few studies found no change in testosterone concentrations after ovarian surgery (Campo et al., 1983; Kojima et al., 1989; Szilagyi et al., 1990; Farhi et al., 1995; Fukaya et al., 1995; Farquhar et al., 2002). Although most studies did not find differences in baseline and/or post-treatment testosterone values between post-treatment ovulatory and non-ovulatory women, between pregnant and nonpregnant women or between obese versus lean patients (Abdel et al., 1993; Campo et al., 1993; Duleba et al., 2003; Al Ojaimi, 2004; Wu et al., 2004; Parsanezhad et al., 2005), some studies did find lower (Aakvaag and Gjonnaess, 1985; Amer et al., 2003b; Api et al., 2005; Hayashi et al., 2005; Kucuk and Kilic-Okman, 2005) or higher (Farhi et al., 1995) testosterone concentrations pre- and/or post-operatively in post-treatment ovulatory PCOS women.

A dose/'puncture' response relationship was found in one study, showing lower testosterone levels after surgery when using more punctures. This relationship was seen using low numbers of punctures per ovary (Amer et al., 2003a), but was not found in a study using more punctures (Malkawi and Qublan, 2005).

Regularly ovulating controls undergoing diagnostic laparoscopy or laparoscopic tubal ligation showed no testosterone change in the first days to weeks after surgery in two studies (Greenblatt and Casper, 1987; Liguori et al., 1996). However, other studies found a testosterone decrease after surgery in controls, although less marked than in PCOS women (Gjonnaess and Norman, 1987; Utsunomiya et al., 1990; Parsanezhad et al., 2005).

Overall, profound reduction of testosterone was seen from the first day after ovarian surgery. Testosterone levels reached a nadir around the third post-operative day and thereafter increased gradually, without reaching pre-operative values (Figure 1A). Responders to ovarian surgery

seemed to have comparable pre- and post-treatment testosterone concentrations compared with non-responders (Figure 1B), although some studies gave conflicting results. The reduction of testosterone seemed to be dose dependent (the more tissue destruction, the lower the testosterone levels after surgery), although this relationship was only found using low numbers of punctures per ovary and a maximal testosterone reduction was quickly reached.

Regularly ovulating controls showed a slight decline of androgens after laparoscopic surgery, but this was less marked than that in PCOS women (Figure 1A).




Figure 1. (A) Mean testosterone serum levels (nmol/L) in polycystic ovary syndrome (PCOS) women and regularly ovulating controls prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase (EFP). PCOS values are derived from 35 papers, a maximum of 1123 patients and control values from 6 papers with a maximum of 121 patients.

(B) Mean testosterone serum levels (nmol/L) in post-treatment PCOS responders and non-responders. Shown are data from prior to ovarian surgery (days=0) and in the subsequent EFP after ovarian surgery. Values are derived from 9 papers and a maximum of 307 responders and 109 non-responders.

Androstenedione

Serum androstenedione levels decreased significantly in the first days after treatment in most studies (Judd et al., 1976; Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987; Greenblatt and Casper, 1987; Sumioki et al., 1988; Kojima et al., 1989; van der Weiden et al., 1989; Armar et al., 1990; Sakata et al., 1990; Szilagyi et al., 1990; Campo et al., 1993; Liguori et al., 1996; Zullo et al., 2000), although two studies found a non-significant reduction (van der Weiden and Alberda, 1987; Utsunomiya et al., 1990). Androstenedione levels during the surgical procedure increased in both PCOS women (Judd et al., 1976; Armar et al., 1990) and regularly ovulating controls (Judd et al., 1976). A nadir was reached 2–4 days after treatment (Judd et al., 1976; Greenblatt and Casper, 1987; Armar et al., 1990; Sakata et al., 1990; Campo et al., 1993; Liguori et al., 1996).

Week(s) to years after ovarian surgery, androstenedione levels in PCOS remained low, although there was a tendency to increase slightly over time (Judd et al., 1976; Aakvaag and Gjonnaess, 1985; Greenblatt and Casper, 1987; van der Weiden and Alberda, 1987; Sumioki et al., 1988; Kojima et al., 1989; van der Weiden et al., 1989; Armar et al., 1990; Keckstein et al., 1990; Sakata et al., 1990; Rossmann et al., 1991; Campo et al., 1993; Szilagyi et al., 1993; Tiitinen et al., 1993; Verhelst et al., 1993; Taskin et al., 1996; Anttila et al., 1998; Gjonnaess, 1998, 1999; Wu et al., 2000; Amer et al., 2002a, 2003a; Malkawi et al., 2003; Malkawi and Qublan, 2005). A few studies showed no androstenedione change following ovarian surgery (Utsunomiya et al., 1990; Kovacs et al., 1991; Cibula et al., 2000). Post-treatment ovulatory women showed lower androstenedione levels prior to and after surgery compared with non-responders (Aakvaag and Gjonnaess, 1985). A dose/‘puncture’ response relationship was found in one study, showing lower androstenedione levels after surgery when using more punctures. Comparable to testosterone, the dose response relationship for androstenedione was seen using low numbers of punctures per ovary (Amer et al., 2003a), but was not found in a study using more punctures (Malkawi and Qublan, 2005).

Regularly ovulating controls showed no change of androstenedione levels after surgery (Judd et al., 1976; Utsunomiya et al., 1990; Liguori et al., 1996; Amer et al., 2002a), although one study found decreased levels after treatment (Gjonnaess and Norman, 1987).

To summarize, reduction of androstenedione in PCOS was seen from the first day after ovarian surgery, but during the ovarian surgery elevated androstenedione levels were found. Androstenedione levels reached a nadir around the fourth post-operative day, and thereafter androgen values increased gradually (Figure 2). PCOS patients responding to ovarian surgery seem to have lower pre- and post-treatment androstenedione values compared with non-responders (no figure due to limited numbers of studies). Reduction of androstenedione could be dose dependent, although this relationship was only found using low numbers of punctures per ovary. Regularly ovulating controls showed no change of androstenedione after laparoscopic surgery (Figure 2).




Figure 2. Mean androstenedione serum levels (nmol/L) in PCOS women and regularly ovulating controls prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. PCOS values are derived from 25 papers, a maximum of 454 patients and control values from 5 papers with a maximum of 91 patients.

Estradiol

Ovarian granulosa cells produce estrogens in response to FSH stimulation (Havelock et al., 2004). Women with PCOS frequently have chronic stable estrogen levels due to insufficient follicle growth.

Estradiol concentrations remained stable in the first days following ovarian surgery in PCOS women in most studies (Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987; Sumioki et al., 1988; Kojima et al., 1989; van der Weiden et al., 1989; Sakata et al., 1990; Szilagyi et al., 1990; Utsunomiya et al., 1990; Abdel et al., 1993), although decreased levels were also found (Judd et al., 1976; Tanaka et al., 1978; Greenblatt and Casper, 1987). Early and mid follicular estradiol concentrations weeks after surgery were comparable to pretreatment levels (van der Weiden et al., 1989; Gadir et al., 1990; Sakata et al., 1990; Tasaka et al., 1990; Verhelst et al., 1993; Wu et al., 2004; Kucuk and Kilic-Okman, 2005; Kandil and Selim, 2005) or were decreased (Rossmannith et al., 1991; Szilagyi et al., 1993).

Late follicular, peri-ovulatory and luteal estrogen levels were higher than before ovarian surgery, especially in responders (Aakvaag and Gjonnaess, 1985; Kojima et al., 1989; Verhelst et al., 1993; Gjonnaess, 1998). Many studies compared pre- and post-treatment estradiol measurements without mentioning the cycle days. These studies will not be discussed here, because results cannot be interpreted.

Regularly ovulating controls showed no estradiol change following surgery (Gjonnaess and Norman, 1987).

Overall, estradiol decreased slightly in the first days after surgery, followed by an increase approximately three weeks after treatment (Figure 3A). The early follicular estradiol levels in subsequent cycles were slightly decreased compared with pretreatment values. Peri-ovulatory and luteal estradiol concentrations were increased, especially in responders (Figure 3B).




Figure 3. (A) Mean estradiol serum levels (pmol/L) in PCOS women prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. Values are derived from 13 papers and a maximum of 139 patients.

(B) Mean estradiol serum levels (pmol/L) in posttreatment PCOS responders and non-responders. Shown are data from prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. Values are derived from 3 papers and a maximum of 73 responders and 23 non-responders.

Inhibins

Inhibins are produced by granulosa cells and production is stimulated by FSH, and inhibins, in turn, inhibit FSH secretion (Laven and Fauser, 2004). Inhibin B is mainly produced by pre- and small antral follicles and concentrations are highest in the midfollicular phase (Laven and Fauser, 2004). Women with PCOS have inhibin B levels within the normal range (Magoffin and Jakimiuk, 1998; Norman et al., 2001; Cortet-Rudelli et al., 2002; Welt et al., 2002), in spite of many small follicles. It has been speculated that inhibin levels in PCOS are relatively suppressed by LH and insulin (Welt et al., 2002).

Three studies reported inhibin levels before and after ovarian surgery. Pre-operatively inhibin B levels on cycle day 5 were higher than in regularly ovulating controls and no pattern of regular pulsatility was seen (Lockwood et al., 1998). A rapid decrease of inhibin post-operatively followed by a subsequent rise (maximum at 3 weeks) was found by Kovacs et al. (1991) (although this study did not define its PCOS population used), and initiation of inhibin B pulsatility was seen after ovarian surgery (Lockwood et al., 1998).

Two studies showed lower inhibin B levels in the early follicular phase, 1–3 months after bilateral drilling or diathermy (Lockwood et al., 1998; Kandil and Selim, 2005) whereas no significant change was seen after unilateral drilling (Kandil and Selim, 2005).

Inhibin levels seem to decrease rapidly after ovarian surgery, followed by a subsequent rise and restoration of inhibin pulsatility. Inhibin levels rose 3–5 weeks after ovarian surgery. Early follicular phase levels in subsequent cycles after ovarian surgery showed lower inhibin levels. Due to limited studies measuring inhibin after surgery, it was not possible to compose a figure.

Progesterone

Progesterone is produced by luteinized granulosa cells under LH stimulation (Havelock et al., 2004).

Progesterone concentrations remained stable for up to 2 weeks after ovarian surgery and in subsequent follicular phases (Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987; Verhelst et al., 1993; Wu et al., 2004). One study showed a decline of progesterone 8 days after surgery in ovulatory women (Armar et al., 1990). Approximately 3 weeks after ovarian surgery and in following luteal phases, progesterone levels rose and were higher than pretreatment values (Aakvaag and Gjonnaess, 1985; Verhelst et al., 1993; Gjonnaess, 1998, 1999).

Progesterone levels in regularly ovulating controls did not show changes after surgery (Gjonnaess and Norman, 1987).

Ovarian surgery does not seem to influence progesterone levels directly after the procedure (Figure 4A and 4B). Weeks after the treatment, progesterone levels rose temporarily in patients responding to surgery and returned to pretreatment levels in subsequent follicular phases.




Figure 4. (A) Mean progesterone serum levels (nmol/L) in PCOS women prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. Values are derived from 4 papers and a maximum of 109 patients.

(B) Mean progesterone serum levels (nmol/L) in post-treatment PCOS responders and non-responders. Shown are data from prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. Values are derived from 2 papers and a maximum of 51 responders and 16 non-responders.

Anti-Müllerian hormone

AMH is an ovarian product, a marker of ovarian reserve (van Rooij et al., 2002, 2005) and is a local inhibitor of FSH action (Durlinger et al., 2001, 2002). Ovarian production of AMH is higher in PCOS patients compared with women with regular menstrual cycles (Cook et al., 2002; Pigny et al., 2006), probably due to an excess of antral follicles (Jonard and Dewailly, 2004).

So far, no studies have been published about the possible AMH changes after ovarian surgery in PCOS.

Pituitary hormones*Luteinising hormone*

LH stimulates androstenedione production by ovarian theca cells (Havelock et al., 2004) and is responsible for ovulation and luteinization. Chronically elevated LH concentrations are common and characteristic of women with PCOS (van Santbrink et al., 1997) and are (partly) responsible for the problems associated with this syndrome.

The day after ovarian surgery, LH concentrations in PCOS women increased in most studies (Gjonnaess and Norman, 1987; Greenblatt and Casper, 1987; Sakata et al., 1990; Tasaka et al., 1990; Naether et al., 1993; Liguori et al., 1996), although no (significant) change (Judd et al., 1976; Kojima et al., 1989; van der Weiden et al., 1989; Utsunomiya et al., 1990; Abdel et al., 1993; Campo et al., 1993) and an LH decrease were also found (Parsanezhad et al., 2005). Subsequently, LH levels decreased and remained low (with the exception of peri-ovulatory peaks) for years after surgery (Tanaka et al., 1978; Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987; Greenblatt and Casper, 1987; Sumioki et al., 1988; van der Weiden et al., 1989; Armar et al., 1990; Gadir et al., 1990, 1992; Sakata et al., 1990; Tasaka et al., 1990; Rossmanith et al., 1991; Szilagyi et al., 1993; Tiitinen et al., 1993; Alborzi, 1994; Farhi et al., 1995; Fukaya et al., 1995; Liguori et al., 1996; Taskin et al., 1996; Anttila et al., 1998; Gjonnaess, 1998, 1999; Soliman et al., 2000; Wu et al., 2000, 2004; Zullo et al., 2000; Alborzi et al., 2001; Amer et al., 2002a, b, 2003a, b; Takeuchi et al., 2002; Amin et al., 2003; Duleba et al., 2003; Malkawi et al., 2003; Parsanezhad et al., 2003; Al Ojaimi, 2004; Kamel et al., 2004; Api et al., 2005; Kucuk and Kilic-Okman, 2005; Malkawi and Qublan, 2005; Parsanezhad et al., 2005). A few studies found unchanged LH levels weeks to months after surgery or increased LH values (mostly peri-ovulatory measurements) (Judd et al., 1976; Kojima et al., 1989; Szilagyi et al., 1990; Utsunomiya et al., 1990; Kovacs et al., 1991; Campo et al., 1993; Naether et al., 1993; Verhelst et al., 1993; Cibula et al., 2000; Asada et al., 2002).

Baseline LH values were higher in women who responded to surgery (Al Ojaimi, 2004; Hayashi et al., 2005) and in lean women (Wu et al., 2004), whereas others found no differences (Duleba

et al., 2003) or lower LH values in responders (Api et al., 2005). The magnitude of the LH-decrease after surgery was higher in PCOS responders (ovulatory or pregnant) compared with non-responders (Aakvaag and Gjonnaess, 1985; Abdel et al., 1993; Jamal, 2000; Amer et al., 2003b; Parsanezhad et al., 2003; Al Ojaimi, 2004; Api et al., 2005; Hayashi et al., 2005).

A dose/‘puncture’ dependency was found, showing a more pronounced increase in LH levels after surgery when using more punctures. This relationship was seen using a small number of punctures per ovary (Amer et al., 2002b, 2003a), but this was not shown with more punctures (Malkawi and Qublan, 2005). After surgery, the LH amplitude decreased while the frequency remained stable (Sumioki et al., 1988; Rossmanith et al., 1991). Regularly ovulating controls did not show LH change after surgery (Judd et al., 1976; Gjonnaess and Norman, 1987; Utsunomiya et al., 1990; Amer et al., 2002a; Parsanezhad et al., 2005).

Thus LH concentrations showed a transient increase the day after ovarian surgery in PCOS, followed by a gradual decrease (Figure 5A). Weeks to years after ovarian surgery, LH levels remained low, with the exception of peri-ovulatory peaks. Lower LH pulse amplitude seems to cause the lower LH levels, while the LH pulse frequency remained stable. Minimal ovarian damage was needed to lower the LH concentrations and a possible dose response relationship was seen. Both responders to ovarian surgery and non-responders showed an LH increase the day after surgery. PCOS women responding to ovarian surgery showed a more pronounced decline of LH levels in the early follicular phase after the procedure compared with non-responders (Figure 5B).

Follicle stimulating hormone

FSH stimulates granulosa cell production of estradiol, inhibin B and GnSAF (Messinis et al., 1991; Havelock et al., 2004; Laven and Fauser, 2004). Increasing pituitary FSH production during the early follicular phase allows the growth of a follicle cohort. Anovulation in PCOS is thought to be partly due to a relative intrinsic inhibition of FSH action. The follicular arrest can be reversed by increasing the amount of FSH by FSH supplementation or stimulation of endogenous FSH production by clomifene citrate administration. Pituitary FSH secretion is selectively suppressed by inhibin A and B (Laven and Fauser, 2004).

FSH levels rose in the first days after ovarian surgery in women with PCOS (Greenblatt and Casper, 1987; Sakata et al., 1990; Tasaka et al., 1990; Naether et al., 1993; Liguori et al., 1996; Parsanezhad et al., 2003), although some studies found no (significant) change (Judd et al., 1976; Gjonnaess and Norman, 1987; Kojima et al., 1989; van der Weiden et al., 1989; Utsunomiya et al., 1990; Campo et al., 1993; Parsanezhad et al., 2005).




Figure 5. (A) Mean luteinising hormone (LH) serum levels (IU/L) in PCOS women and regularly ovulating controls prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. PCOS values are derived from 38 papers, a maximum of 1127 patients and control values from 4 papers with a maximum of 65 patients.

(B) Mean LH serum levels (IU/L) in post-treatment PCOS responders and non-responders. Shown are data from prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. Values are derived from 10 papers and a maximum of 311 responders and 106 non-responders.

In the first weeks after surgery, FSH concentrations declined to levels comparable to baseline values (Judd et al., 1976; Tanaka et al., 1978; Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987; Greenblatt and Casper, 1987; Sumioki et al., 1988; Kojima et al., 1989; Armar et al., 1990; Sakata et al., 1990; Szilagyi et al., 1990, 1993; Tasaka et al., 1990; Kovacs et al., 1991; Campo et al., 1993; Naether et al., 1993; Tiitinen et al., 1993; Verhelst et al., 1993; Alborzi,

1994; Farhi et al., 1995; Fukaya et al., 1995; Liguori et al., 1996; Anttila et al., 1998; Soliman et al., 2000; Wu et al., 2000, 2004; Zullo et al., 2000; Amer et al., 2002a, 2003a, b; Takeuchi et al., 2002; Duleba et al., 2003; Malkawi et al., 2003; Kandil and Selim, 2005; Kucuk and Kilic-Okman, 2005; Malkawi and Qublan, 2005; Parsanezhad et al., 2005), whereas others found increased (van der Weiden et al., 1989; Gadir et al., 1990; Utsunomiya et al., 1990; Rossmannith et al., 1991; Abdel et al., 1993; Taskin et al., 1996; Amer et al., 2002b; Amin et al., 2003; Parsanezhad et al., 2003; Al Ojaimi, 2004; Kamel et al., 2004; Api et al., 2005) or decreased (Szilagyi et al., 1990; Gjonnaess, 1998, 1999; Alborzi et al., 2001) FSH values after ovarian surgery.

A dose/'puncture' dependency was not found (Amer et al., 2003a; Malkawi and Qublan, 2005), although one study showed a possible trend towards higher FSH levels when using more punctures (Amer et al., 2002b). FSH pulse frequency and amplitude did not change after the treatment (Sumioki et al., 1988; Rossmannith et al., 1991).

PCOS responders showed a more pronounced increase in FSH in the first days after surgery compared with non-responders (Aakvaag and Gjonnaess, 1985; Armar et al., 1990; Abdel et al., 1993; Campo et al., 1993). In most studies, FSH values in the weeks after surgery were comparable between responders and non-responders and the initial difference seen after surgery was lost (Jamal, 2000; Amer et al., 2003b; Al Ojaimi, 2004; Api et al., 2005), although one study found elevated FSH in responders (Parsanezhad et al., 2003) and another in non-responders (Hayashi et al., 2005).

Regularly ovulating controls showed no FSH change after surgery (Judd et al., 1976; Gjonnaess and Norman, 1987; Utsunomiya et al., 1990; Amer et al., 2002a; Parsanezhad et al., 2005).

Overall, FSH concentrations increased in the first days after ovarian surgery, with a more pronounced elevation in responders (Figures 6A and 6B). Following this FSH elevation, the levels returned gradually to pre-operative values and the initial difference shortly after surgery between responders and non-responders was lost (Figure 6B). FSH pulsatility did not change after surgery.

Prolactin

Prolactin is mainly secreted by the anterior pituitary and secretion is under the tonic inhibition of dopamine. Prolactin has a diurnal variation and various factors like estrogens and (surgical) stress stimulate prolactin secretion (Tyson et al., 1972; Sassin et al., 1973; Brandt et al., 1976). Studies with large PCOS populations showed average prolactin levels within the (high) normal range (Balen et al., 1995; Hernandez et al., 2000). In spite of normal prolactin levels in PCOS, the reaction to a dopamine receptor antagonist (metoclopramide) is less than in ovulatory controls, suggesting a low dopamine hypothalamic tone (Minakami et al., 1988; Hernandez et al., 2000).




Figure 6. (A) Mean follicle stimulating hormone (FSH) serum levels (IU/L) in PCOS women and regularly ovulating controls prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. PCOS values are derived from 34 papers, a maximum of 1092 patients and control values from 4 papers with a maximum of 60 patients.

(B) Mean FSH serum levels (IU/L) in post-treatment PCOS responders and non-responders. Shown are data from prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. Values are derived from 8 papers and a maximum of 302 responders and 103 non-responders.

After surgery, prolactin levels rose on the first day in both PCOS patients and in ovulatory controls (Gjonnaess and Norman, 1987; van der Weiden et al., 1989; Parsanezhad et al., 2005). Prolactin concentrations were not altered weeks to years after the procedure in the majority of the studies (van der Weiden et al., 1989; Gadir et al., 1990; Szilagyi et al., 1990; Verhelst et al., 1993; Gjonnaess, 1998; Alborzi et al., 2001; Duleba et al., 2003; Al Ojaimi, 2004; Wu et al.,

2004; Kucuk and Kilic-Okman, 2005). One study showed a decline in prolactin after the ovarian procedure (Alborzi, 1994), another study an increase in prolactin up to 10 weeks after ovarian surgery (Parsanezhad et al., 2005). Parsanezhad et al. (2005) showed that the majority of the women remaining anovulatory after ovarian surgery were hyperprolactinaemic 6–10 weeks after surgery. Prolactin response to dopaminergic blockade by metoclopramide was increased after ovarian surgery (Szilagyi et al., 1993).

A transient prolactin elevation after surgery was seen in both PCOS patients and in regularly ovulatory controls. Ovarian surgery does not seem to influence basal prolactin levels weeks to months after the procedure, although dopaminergic inhibition of prolactin seems to increase as indicated by the increased response to metoclopramide.

Gonadotropin releasing hormone challenge test

The sensitivity and priming of the pituitary to gonadotropin releasing hormone (GnRH or LHRH) can be measured by the magnitude of FSH and LH response to GnRH administration. Patients with PCOS have a higher LH response to GnRH compared with ovulatory controls (Heineman et al., 1984). This suggests that in women with PCOS the pituitary is already primed. The cause of this increased sensitivity of the pituitary in PCOS is unclear. It has been suggested that the low concentrations of the ovarian hormone GnSAF may be responsible for this priming. GnSAF normally antagonizes the priming (Fowler et al., 1990; Dafopoulos et al., 2004).

Ovarian surgery diminished pituitary LH response to GnRH administration in the first weeks after ovarian surgery in PCOS (Tanaka et al., 1978; Sumioki et al., 1988; Keckstein et al., 1990; Mio et al., 1991; Rossmannith et al., 1991; Campo et al., 1993; Fukaya et al., 1995). Only on the first day after surgery was the LH response to GnRH found to be increased (Gjonnaess and Norman, 1987).

Adrenal hormones

Dehydroepiandrosterone

Dehydroepiandrosterone (DHEA) and DHEA-sulphate (DHEAS) act as inactive precursor steroids for peripheral conversion into more potent androgens (Burger, 2002; Labrie et al., 2005). They are predominantly of adrenocortical origin and are frequently elevated in PCOS (DeVane et al., 1975; Kumar et al., 2005). Due to its adrenocortical origin, DHEAS is used as a marker for adrenal androgen production (Kumar et al., 2005).

In the first days after ovarian surgery, DHEA(S) levels decreased in half of the studies (Gjonnaess and Norman, 1987; Szilagyi et al., 1990; Utsunomiya et al., 1990; Liguori et al., 1996), whereas the other studies showed no significant change (Aakvaag and Gjonnaess, 1985; Naether et al., 1993; Cibula et al., 2000; Parsanezhad et al., 2005).

Weeks to months after ovarian surgery, DHEAS and/or DHEA levels were comparable with pre-operative values in the majority of the studies (Vejlsted and Albrechtsen, 1976; Campo et al., 1983; Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987; Sumioki et al., 1988; Armar et al., 1990; Szilagyi et al., 1990; Utsunomiya et al., 1990; Kovacs et al., 1991; Rossmannith et al., 1991; Naether et al., 1993; Tiitinen et al., 1993; Verhelst et al., 1993; Gjonnaess, 1998; Cibula et al., 2000; Wu et al., 2000, 2004; Takeuchi et al., 2002; Duleba et al., 2003; Api et al., 2005; Kucuk and Kilic-Okman, 2005; Parsanezhad et al., 2005), although some studies showed decreased DHEA(S) levels (Naether et al., 1993; Liguori et al., 1996; Gjonnaess, 1999; Alborzi et al., 2001; Malkawi et al., 2003; Malkawi and Qublan, 2005).

No differences in DHEAS levels before and/or after surgery were found between the post-treatment ovulatory group compared with the non-ovulatory group (Aakvaag and Gjonnaess, 1985; Parsanezhad et al., 2005), between the patients who became pregnant compared with the non-pregnant women (Duleba et al., 2003; Cleemann et al., 2004; Api et al., 2005) and the obese versus lean patients (Wu et al., 2004).

DHEAS levels in control groups with regularly ovulatory women undergoing laparoscopy varied from a decrease the first days after surgery (Gjonnaess and Norman, 1987; Utsunomiya et al., 1990) to no change for up to months after laparoscopy (Parsanezhad et al., 2005).

Thus, the influence of ovarian surgery on DHEA(S) production seems unclear. Overall, the DHEA(S) levels seems to decrease slightly in the first days following surgery (Figure 7). This was seen in both PCOS women and in controls. Weeks to months after the treatment, DHEA(S) levels were comparable or slightly reduced in PCOS patients. DHEAS levels before and after surgery did not differ between responders and non-responders and between lean and obese patients.

Cortisol and ACTH testing

Cortisol levels remained stable in three studies measuring cortisol levels for up to months after ovarian surgery (Sumioki et al., 1988; Gadir et al., 1990; Wu et al., 2000).

The adrenal steroid response (17-hydroxy (OH) progesterone, androstenedione, DHEA(S) and cortisol) to adrenocorticotrophic hormone (ACTH) was not influenced by ovarian surgery in two studies (Cibula et al., 2000; Saleh et al., 2001). Another study showed lower 17 ketosteroid, 17-OH progesterone and androstenedione secretions as a response to ACTH stimulation after ovarian surgery (Wu et al., 2000).




Figure 7. Mean dehydroepiandrosterone (DHEAS) serum levels (nmol/L) in PCOS women and in regularly ovulating controls prior to ovarian surgery (days=0), the first 28 days following ovarian surgery and in the subsequent early follicular phase. PCOS values are derived from 18 papers, a maximum of 806 patients and control values from 3 papers with a maximum of 60 patients.

17-hydroxy progesterone

17-OH progesterone decreased on the first day after ovarian surgery in women with PCOS and in regularly ovulating controls (Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987). Subsequently 17-OH progesterone levels returned in PCOS women to pretreatment values (Aakvaag and Gjonnaess, 1985; Verhelst et al., 1993; Malkawi et al., 2003; Malkawi and Qublan, 2005), although one study showed an increase in 17-OH progesterone (Gjonnaess, 1998).

Other hormones

Dihydrotestosterone

DHT is a potent androgenic metabolite and is primarily derived from peripheral conversion (through enzyme 5-alpha-reductase) of testosterone, androstenedione and DHEA and is partly secreted by the adrenal gland (Silva et al., 1987; Burger, 2002). Increased 5-alpha-reductase activity is found in PCOS and could (partly) be responsible for excess DHT production in PCOS (Fassnacht et al., 2003; Jakimiuk et al., 1999). Besides substantial ovarian and adrenal production of androgens, increased peripheral conversion contributes to higher total androgen levels in PCOS (Fassnacht et al., 2003).

On the first day after ovarian surgery, DHT levels decreased in both PCOS patients and in ovulatory controls (in the follicular phase) (Aakvaag and Gjonnaess, 1985; Gjonnaess and Norman, 1987). Weeks to years after ovarian surgery, DHT levels remained stable (Aakvaag and Gjonnaess, 1985; Gjonnaess, 1999) or alternated within the same study between normal and

lower values (Verhelst et al., 1993; Gjonnaess, 1998). Post-treatment ovulatory women showed lower pretreatment DHT values than non-ovulatory women (Aakvaag and Gjonnaess, 1985).

Overall DHT concentration decreased the day after surgery in both PCOS patients and ovulatory controls. After the initial reduction, DHT levels returned to pre-operative values or remained decreased.

Sex hormone binding globulin

SHBG is a major transport protein for testosterone and its production is inhibited by insulin and prolactin and stimulated by testosterone and estradiol (Plymate et al., 1988). Women with PCOS show lower serum levels of SHBG than ovulatory controls (Kumar et al., 2005).

Most studies showed no significant change in SHBG concentrations after ovarian surgery in PCOS women (Judd et al., 1976; Gjonnaess and Norman, 1987; van der Weiden et al., 1989; Szilagyi et al., 1990; Kovacs et al., 1991; Rossmanith et al., 1991; Tiitinen et al., 1993; Verhelst et al., 1993; Taskin et al., 1996; Anttila et al., 1998; Wu et al., 2000; Amer et al., 2003a; Duleba et al., 2003), although increased SHBG levels were sometimes found after surgery (Aakvaag and Gjonnaess, 1985; Gjonnaess, 1998, 1999; Wu et al., 2004; Kucuk and Kilic-Okman, 2005). Overall SHBG levels seem to increase modestly after ovarian surgery.

Insulin, glucose and insulin resistance

PCOS women often have some degree of insulin resistance and compensatory hyperinsulinaemia (Dunaif et al., 1989; Legro et al., 2004). Insulin increases the GnRH stimulated LH release, potentiates LH-induced ovarian androgen synthesis (Adashi et al., 1981; Bergh et al., 1993) and has a role in the pathogenesis of PCOS. The gold standard for whole-body insulin sensitivity assessment is the euglycemic-hyperinsulinaemic clamp (Legro et al., 2004). Unfortunately, this test is time consuming and expensive and the oral glucose tolerance tests (OGTT) is the best practical alternative (Legro et al., 2004).

Fasting insulin levels in PCOS patients remained unaltered in the first weeks to months after ovarian surgery in the majority of the studies (Tiitinen et al., 1993; Tulandi et al., 2000; Saleh et al., 2001; Duleba et al., 2003; Malkawi et al., 2003; Wu et al., 2004; Malkawi and Qublan, 2005). Two studies showed a small, but significant decline of fasting insulin levels one and six months after ovarian surgery (Wu et al., 2000; Api et al., 2005).

One study showed a rise in fasting insulin levels and insulin pulse amplitude months after the procedure (Gadir et al., 1990). Ovarian surgery did not alter the level of fasting glucose (Tulandi et al., 2000; Wu et al., 2000, 2004; Saleh et al., 2001; Duleba et al., 2003; Malkawi et al., 2003; Api

et al., 2005; Malkawi and Qublan, 2005). Insulin and glucose response to OGTT were modestly but significantly decreased after ovarian surgery in hyperinsulinaemic PCOS women only (Saleh et al., 2001). Combining the results of the lean and obese PCOS patients, the average OGTT did not change (Tulandi et al., 2000; Duleba et al., 2003). The insulin sensitivity index worsened in one study after the ovarian surgery (Duleba et al., 2003) and insulin/glucose decreased in another (Wu et al., 2000). Two studies showed unaltered glucose/insulin ratios (Wu et al., 2004; Api et al., 2005). The euglycemic–hyperinsulinaemic clamp was performed in one study and showed no difference in eight patients before and after ovarian surgery (although this study did not define its PCOS population) (Lemieux et al., 1999). Baseline parameters of women who achieved pregnancy after the surgery showed significant lower insulin levels and higher insulin sensitivity indexes prior to surgery compared with women who did not get pregnant (Duleba et al., 2003; Dale et al., 2004; Api et al., 2005).

Overall, insulin sensitivity does not seem to change after ovarian surgery, although more research is needed for more clarity. Hyperinsulinaemic women may become less (but remain) insulin resistant after surgery. The limitation of some of these studies is the exclusion of women who became pregnant before endocrine follow up, potentially resulting in selection bias. These pregnant women may have had more favourable insulin changes. Analysing the data from the normoinsulinaemic, hyperinsulinaemic, lean or obese women separately may reveal different results.

Leptin

Leptin is secreted by adipose tissue, modulates energy balance and influences reproductive function (Lonnqvist et al., 1995; Broberger, 2005). Leptin serum levels correlate positively with body mass index (BMI) (Lonnqvist et al., 1995; Telli et al., 2002) and obesity is associated with a resistance to leptin (Chan and Mantzoros, 2005). Insulin has been shown to stimulate leptin production (Kolaczynski et al., 1996) and there is evidence that androgens suppress leptin production (Jockenhovel et al., 1997; Remsberg et al., 2002). Many studies measured leptin levels in PCOS compared to weight matched controls. Most of these studies showed comparable leptin levels between PCOS and normal ovulatory women (Telli et al., 2002; Remsberg et al., 2002; Carmina et al., 2005), while others found higher leptin levels in PCOS (El et al., 1999; Baranowska et al., 1999). A relative leptin deficiency in overweight PCOS women is speculated, given the (visceral) fat mass and hyperinsulinaemia in these women (Remsberg et al., 2002). This relative leptin deficiency might (partly) be due to insulin resistance limiting insulin stimulated leptin secretion (Jacobs and Conway, 1999).

So far, the effect of surgery on leptin has been evaluated in one study, showing mean leptin levels comparable with pretreatment values 3 months after ovarian surgery. In obese PCOS

patients, leptin significantly decreased contrary to stable levels in lean PCOS women after ovarian surgery (Wu et al., 2004).

Renin

The renin-angiotensin system is classically known for its effects on blood pressure. Besides this effect on the circulation, it may also have an important role in reproduction (Vinson et al., 1997). In (normotensive) women with PCOS, elevated serum renin levels have been found compared with normal ovulatory women (Jaatinen et al., 1995; Hacıhanefioglu et al., 2000).

Ovarian surgery did not influence renin levels significantly after the first menstrual cycle, measured in 11 PCOS patients (Anttila et al., 1998). It must be noted that total renin was measured. Changes in active renin concentrations could not be ruled out and local ovarian activity could have changed (Anttila et al., 1998).

Insulin-like growth factor and binding protein

Insulin-like growth factors (IGF) I and II have important roles in modulation of ovarian function (Fowler et al., 2000); IGF-I regulates a variety of processes in ovarian cells and stimulates androgen production (Carmina et al., 1995; Giudice, 1995). IGF binding globulins (IGFBP) influence the activity of IGFs in a cell- and tissue-specific manner (Beattie et al., 2006) and five of the six IGFBPs (IGFBP 1, 2, 3, 4 and 5) have been identified in the human ovary (Giudice et al., 1995). IGFBP-I alters serum IGF bioavailability and influences IGF-I action on the ovary (Fowler et al., 2000). Insulin is thought to act (partially) through ovarian IGF-I receptors, augmenting hyperandrogenism in PCOS (Gdansky et al., 1997). PCOS women have lower IGFBP-I concentrations compared with normal ovulatory women (Carmina et al., 1995).

IGF-I and IGFBP-I remained unchanged when measured 1–3 months after ovarian surgery in PCOS patients (Tiitinen et al., 1993; Amin et al., 2003; Wu et al., 2004). There are no publications on IGF-II and other IGFBP's levels before and after ovarian surgery in PCOS.

Vascular endothelial growth factor

Vascular endothelial growth factor (VEGF) is secreted by the ovaries, has angiogenic properties and plays an important role in cyclic ovarian angiogenesis (Leung et al., 1989; Geva and Jaffe, 2000). VEGF levels are positively correlated with ovarian blood flow (Agrawal et al., 2002; bd El Aal et al., 2005). Elevated VEGF values have been found in women with PCO(S) (Agrawal et al., 1998, 2002; bd El Aal et al., 2005) and VEGF secretion is stimulated by LH, FSH and insulin in combination with hCG (Christenson and Stouffer, 1997; Agrawal et al., 2002).

Baseline VEGF levels were higher in PCOS women compared with normal ovulatory controls (Tulandi et al., 2000; Amin et al., 2003). Ovarian surgery lowered VEGF levels in combination with reduced ovarian stromal blood flow in one study of 25 PCOS patients (Amin et al., 2003). Another study showed no changes in VEGF concentrations after surgery in 27 PCOS patients (Tulandi et al., 2000). Results of the latter study may have been biased by the exclusion of six women who became pregnant before endocrine follow up.

Homocysteine

Homocysteine is a risk factor for atherosclerosis and has been found to be elevated in women with PCOS (Yarali et al., 2001; Loverro et al., 2002; Vrbikova et al., 2003), especially in insulin resistant individuals (Schachter et al., 2003; Wijeyaratne et al., 2004), although some studies have not found any difference (Sills et al., 2001; Morgante et al., 2002). It has been suggested that high homocysteine levels are linked to androgen levels (Vrbikova et al., 2003).

Homocysteine levels decreased 3 months after ovarian surgery, although this was measured in only one study of 22 PCOS patients (Kucuk and Kilic-Okman, 2005).

Lipid profile

Dyslipidaemia is common in patients with PCOS (Talbot et al., 1995; Orio, Jr. et al., 2004). PCOS women have higher total cholesterol, triglycerides and very low density lipoprotein and lower high-density lipoprotein (HDL) levels compared with ovulatory controls (Wild et al., 1985; Wild, 2002; Orio, Jr. et al., 2004; Vural et al., 2005).

Lipid levels did not seem affected by ovarian surgery. Ovarian surgery did not change total cholesterol, HDL and LDL concentrations in two studies (Duleba et al., 2003; Kucuk and Kilic-Okman, 2005).

DISCUSSION

When ovarian surgery causes damage to the PCOS ovaries, independent of the type of the procedure, the ovarian–pituitary-axis regulation changes. For optimal identification of essential endocrine changes after ovarian surgery, the endocrine events will be discussed in chronological order.

The day after surgery in PCOS patients, serum levels of ovarian produced steroids, and non-steroidal substances declined in combination with increased serum levels of pituitary hormones and increased pituitary sensitivity (Figure 8). Invariably lower serum levels of the ovarian hormones (testosterone, androstenedione, estradiol, inhibin) may (partially) be a direct effect

of the ovarian surgery. Apparently, granulosa and theca cell damage caused by the surgical procedure is responsible, possibly in combination with a temporary disruption of ovarian function. A consequence is altered ovarian feedback to the hypothalamus and pituitary, subsequently causing changes in pituitary sensitivity and secretion. This results in an increase of serum levels of pituitary hormones (LH, FSH, prolactin) shortly after the surgery. Both LH and FSH are normally suppressed by estrogens (Messinis, 2006), and the lower estrogen concentrations after ovarian surgery could augment the FSH and LH increase. In addition, FSH secretion is likely to be stimulated due to the fall of inhibin which normally restrains GnRH independent pituitary FSH secretion (Lambalk et al., 1998). However, a temporal change in secretion and sensitivity of the hypothalamus and pituitary induced by the anaesthesia or operation itself cannot be excluded. Temporally increased pituitary secretion has been found after surgery in regularly ovulating women (Messinis et al., 1999), although most studies found stable or even decreased pituitary hormone secretion after surgery in controls and PCOS patients (Adashi et al., 1980; Hagen et al., 1980).




Figure 8. Mean levels of LH, FSH, testosterone (TT) and estradiol prior to ovarian surgery (days=0) and the first days following surgery in PCOS women. The figure is a compilation of Figures 1A, 3A, 5A and 6A.

A sufficient increase of FSH secretion can subsequently induce the growth of a follicle cohort, explaining the more distinct incline of FSH in responders compared with non-responders after surgery. Unfortunately, there are no studies reporting inhibin levels in the first hours after surgery, and therefore it is unclear if inhibin drops prior to the rise of FSH.

The LH levels that increase shortly after ovarian surgery are associated with increased pituitary sensitivity. The higher pituitary sensitivity occurs in combination with lower estrogen levels.

Estrogens normally exert a negative feedback on LH predominantly by blocking hypothalamic GnRH secretion, whereas pituitary sensitivity increases (Messinis, 2006). This more sensitive pituitary however does not secrete large amounts of LH because of the relative absence of endogenous GnRH. Lower estrogen levels after surgery are expected to diminish pituitary sensitivity, but possibly the remaining endogenous GnRH will temporarily provide enough substrate for increased LH production. It is also possible that sensitivity of the pituitary is increased by decreased secretion of GnSAF. GnSAF normally reduces pituitary sensitivity and suppresses LH secretion (Fowler et al., 2003). Ovarian cell damage by the surgery may cause a drop in the already-low GnSAF production immediately after the procedure, resulting in higher pituitary sensitivity thus explaining both the rise in LH levels and the increased pituitary response to GnRH. So far, GnSAF has not been purified, and although plausible, this GnSAF theory remains unproven.

With regard to prolactin, its elevation seen after surgery is probably related to stress of operation and/or anaesthetics, since elevation was seen in both PCOS patients and in regularly ovulatory controls, and a transitory prolactin elevation caused by surgical stress is a known phenomenon (Brandt et al., 1976; Adashi et al., 1980; Soules et al., 1980).

Besides changes in ovarian and pituitary hormone production, the adrenal production of DHEA(S) and 17-OH progesterone decreased in the first days after surgery. Because reduction of adrenal hormones is seen in PCOS women as well in regularly ovulating controls, a more general effect of surgery is suggested as a cause of this decrease.

Both PCOS patients and ovulatory controls show decreased DHT concentration the day after surgery. Lower circulating testosterone and androstenedione concentrations after ovarian surgery probably contribute to the lower DHT levels, as DHT is a peripheral product of these androgenic substrates.

Five days after the ovarian surgery, the ovarian production of estradiol and inhibin starts to rise after the initial reduction. These increments probably represent ovarian functional recovery and folliculogenesis. Synchronic with the estradiol elevation, FSH levels decrease (Figure 8). A plausible explanation for this suppression of FSH secretion are the higher estradiol and inhibin levels produced by growing follicles, a process similar to the normal ovulatory cycle (Jonard and Dewailly, 2004; Messinis, 2006). The elevated progesterone levels three weeks after ovarian surgery suggest that a proportion of the women were ovulating.

Reduced ovarian hormone production (testosterone, androstenedione, estradiol, inhibin) is seen in subsequent follicular phases after the ovarian surgery in PCOS. Previous studies have

shown that partial destruction of ovarian tissue due to the ovarian surgery results in a decreased ovarian reserve (Kandil and Selim, 2005). A decreased amount of ovarian tissue could result in less endocrine capacity and cause the reduction in ovarian hormone production.

Lower androgen levels after ovarian surgery are a known phenomenon. Destruction of androgen producing tissue in combination with lower LH levels after ovarian surgery is probably the main cause for this reduction. Besides lower androgen levels, SHBG increases modestly, possibly due to higher estradiol levels in ovulatory cycles (Eden et al., 1989). Increased SHBG levels result in lower levels of biologically active (free) androgens after surgery. Intra-ovarian hyperandrogenism is seen as the main culprit for follicle excess and arrest (Jonard and Dewailly, 2004), and lower androgen levels after ovarian surgery may reduce the follicle excess and facilitate appropriate folliculogenesis.

Lower ovarian production of inhibin B after ovarian surgery could be a reflection of the decreased ovarian reserve after the treatment, since inhibin B levels are a marker of follicular reserve (Tinkanen et al., 2001). So far, only limited data on inhibin are available.

Altered ovarian feedback through the reduced ovarian hormone production could be the cause of the restoration of pituitary function over time. Serum LH levels diminish, as its pulse amplitude decreases, in combination with a stable frequency. Weeks after the ovarian surgery, the pituitary sensitivity decreases, expressed by lower pituitary LH response to GnRH. Theoretically, restoration of GnSAF production by healthy growing follicles after surgery (Balen and Jacobs, 1994) in combination with decreased estradiol levels in the early follicular phases could result in diminished pituitary priming, consequently explaining the lower LH response to GnRH and the lower amplitude of pituitary LH secretion. Increased GnSAF production by growing follicles could also elucidate the lower LH levels found in responders compared with non-responders. Non-responders show no (adequate) follicle growth, resulting in less GnSAF production and higher LH levels.

Some time after surgery FSH levels become comparable to baseline values, in spite of lower inhibin B levels in the early follicular phase. Remarkably, such FSH levels were not able to initiate follicular growth before treatment, but after surgery these FSH levels are sufficient for folliculogenesis. This strongly suggests an increased effect of FSH after surgery. A more favourable endocrine environment may increase the ovarian sensitivity to FSH and in combination with an intercycle FSH rise, this may cause follicle growth to be induced. The endocrine environment may also be altered by changes in ovarian AMH tone. AMH levels are higher in PCOS compared with regularly ovulating women (Cook et al., 2002; Pigny et al., 2006) and reduction of the follicle excess and reserve after surgery could hypothetically reduce ovarian AMH production, resulting

in less AMH inhibition of FSH action. Ovarian sensitivity could therefore be increased by lower AMH production. FSH serum levels after surgery would become locally more effective, resulting in reversion of the follicular arrest.

The influence of ovarian surgery on adrenal hormones seems limited, as could be expected. Weeks to months after surgery cortisol, 17-OH progesterone levels and ACTH tests are comparable to pretreatment results and DHEA(S) is comparable or only slightly reduced. Resumption of ovulation after ovarian surgery seems independent of adrenal functional changes.

After the initial reduction, DHT levels return to pre-operative values or remain slightly lower. High peripheral conversion of testosterone, androstenedione and DHEA due to increased 5-alpha-reductase activity in PCOS could provide substantial DHT production in spite of reduced androgens. DHT normalization does not seem compulsory for induction of ovulation.

Insulin sensitivity does not seem to change after ovarian surgery, although hyperinsulinaemic women showed modest improvement. Assuming that an intrinsic defect in insulin action, the insulin androgenic pathway and *b*-cell dysfunction in PCOS are major contributors to insulin resistance (Dunaif et al., 1989; Ehrmann et al., 1995; Dunaif, 1997; Corbould et al., 2005), surgery is not expected to influence the hyperinsulinaemia and insulin resistance. Earlier studies showed that suppression of hyperandrogenism did not result in significant changes in insulin levels in hyperinsulinaemic, obese or lean PCOS women (Dunaif et al., 1990; Diamanti-Kandarakis et al., 1995) or lead to only modest improvements (Elkind-Hirsch et al., 1993; Moghetti et al., 1996). Reduction of androgen producing ovarian tissue and lower LH levels after surgery are apparently sufficient for reducing androgen levels, in spite of continuous stimulation of androgen production by insulin. However, a distinction has to be made between lean and obese PCOS patients: obese PCOS patients will have hyperinsulinaemia as a principal driver of ovarian hyperandrogenism and would probably benefit more from insulin lowering therapy, whereas lean PCOS patients would benefit from the lower LH levels found after ovarian surgery. Nevertheless, resolving the follicular arrest can occur without concomitant changes of the insulin metabolism.

Leptin, VEGF and homocysteine levels seem to decrease after ovarian surgery in PCOS, as reported by a few studies. Leptin decreased in obese subjects in spite of constant insulin levels, stable BMI and lower androgen levels. It is unknown how leptin secretion is attenuated in these obese PCOS women after the surgery. Lower VEGF levels may have been caused by lower LH concentrations, resulting in less VEGF stimulatory effects, in combination with a reduction of ovarian VEGF production due to decreased ovarian reserve after surgery (Kandil and Selim, 2005). Homocysteine levels decrease after ovarian surgery. The cause of this improvement

is unclear. It may be due to lower androgen levels, which are suggested to be linked with homocysteine levels (Vrbikova et al., 2003).

It has been suggested that IGF-I plays a role in the re-initiation of folliculogenesis (Balen and Jacobs, 1994). Local injury of the ovary could be followed by a cascade of inflammatory factors, among IGF-I (Balen and Jacobs, 1994), and IGF-I in turn would increase FSH action (Adashi et al., 1988) and follicular development could ensue. Looking at this hypothesis with the selected studies in mind, the studies measuring IGF-I or IGF-I BP concentrations after ovarian surgery all reported no changes (Tiitinen et al., 1993; Amin et al., 2003; Wu et al., 2004), but the IGF-I was measured month(s) after surgery. Assuming that the cascade of inflammatory factors will be diminished by that time, the results of the studies neither support nor reject this hypothesis. Furthermore, only IGF-II is expressed in the human ovary and it seems to play a more important role than IGF-I in normal follicular development (Giudice et al., 1995). Furthermore at this point in time, there is no convincing evidence that IGF system abnormalities have a central place in the follicular arrest in PCOS patients (Jonard and Dewailly, 2004).

The general limitation of using serum values remains the absence of information about the local ovarian changes, autocrine and interfollicular influences.

Conclusion

Endocrine changes after ovarian surgery for ovulation induction in PCOS seem to be governed by the ovaries. The rapid changes occurring after surgery strongly argue for inappropriate gonadotropin secretion secondary to inadequate ovarian feedback.

Reduction of all ovarian produced hormones immediately after surgery restores feedback to the hypothalamus and pituitary resulting in increased pituitary sensitivity and temporarily higher LH and particularly FSH levels. Initiation of folliculogenesis is induced by the increased FSH levels, in combination with a reduction of the follicle excess and reduced (intra-ovarian) androgen levels.

In addition, upon ovarian surgery the endocrine environment changes over time. After the initial reduction, ovarian production of estradiol and inhibin increases, proving folliculogenesis. Subsequently through ovarian feedback, the sensitivity of the pituitary decreases and reduced FSH and LH levels are found. In addition, increased GnSAF production by growing follicles could explain the reduced pituitary sensitivity to GnRH, the lower LH levels and could antagonize the stimulatory effects of estradiol on GnRH-induced LH secretion (Fowler et al., 2003).

Continuous follicle growth in subsequent cycles after ovarian surgery occurs in a more favourable internal environment with less (free) androgens, lower LH levels, reduced follicle excess and

possibly lower local AMH production. This leads to a prolonged altered endocrine situation that seems to concord with increased ovarian sensitivity to FSH, since FSH levels comparable to pretreatment levels become adequate for inducing folliculogenesis after surgery.

REFERENCES

- Aakvaag A and Gjonnaess H (1985) Hormonal response to electrocautery of the ovary in patients with polycystic ovarian disease. *Br J Obstet Gynaecol* 92,1258–1264.
- Abdel GA, Khatim MS, Alnaser HM, Mowafi RS and Shaw RW (1993) Ovarian electrocautery: responders versus non-responders. *Gynecol Endocrinol* 7,43–48.
- Adams J, Polson DW and Franks S (1986) Prevalence of polycystic ovaries in women with anovulation and idiopathic hirsutism. *Br Med J (Clin Res Ed)* 293,355–359.
- Adashi EY, Hsueh AJ and Yen SS (1981) Insulin enhancement of luteinizing hormone and follicle-stimulating hormone release by cultured pituitary cells. *Endocrinology* 108,1441–1449.
- Adashi EY, Rebar RW, Ehara Y, Naftolin F and Yen SS (1980) Impact of acute surgical stress on anterior pituitary function in female subjects. *Am J Obstet Gynecol* 138,609–614.
- Adashi EY, Resnick CE, Hernandez ER, May JV, Knecht M, Svoboda ME and Van Wyk JJ (1988) Insulin-like growth factor-I as an amplifier of follicle-stimulating hormone action: studies on mechanism(s) and site(s) of action in cultured rat granulosa cells. *Endocrinology* 122,1583–1591.
- Agrawal R, Sladkevicius P, Engmann L, Conway GS, Payne NN, Bekis J, Tan SL, Campbell S and Jacobs HS (1998) Serum vascular endothelial growth factor concentrations and ovarian stromal blood flow are increased in women with polycystic ovaries. *Hum Reprod* 13,651–655.
- Agrawal R, Jacobs H, Payne N and Conway G (2002) Concentration of vascular endothelial growth factor released by cultured human luteinized granulosa cells is higher in women with polycystic ovaries than in women with normal ovaries. *Fertil Steril* 78,1164–1169.
- Al Ojaimi EH (2004) Endocrine changes after laparoscopic ovarian drilling in clomifene citrate-resistant women with polycystic ovarian syndrome. *Saudi Med J* 25,1032–1039.
- Alborzi S (1994) Laparoscopic ovarian cauterization in patients with polycystic ovaries; clinical outcome and endocrine changes. *Iran J Med Sci* 19,120–124.
- Alborzi S, Khodaei R and Parsanejad ME (2001) Ovarian size and response to laparoscopic ovarian electro-cauterization in polycystic ovarian disease. *Int J Gynaecol Obstet* 74,269–274.
- Amer SA, Banu Z, Li TC and Cooke ID (2002a) Long-term follow-up of patients with polycystic ovary syndrome after laparoscopic ovarian drilling: endocrine and ultrasonographic outcomes. *Hum Reprod* 17,2851–2857.
- Amer SAK, Li TC and Cooke ID (2002b) Laparoscopic ovarian diathermy in women with polycystic ovarian syndrome: a retrospective study on the influence of the amount of energy used on the outcome. *Hum Reprod* 17,1046–1051.
- Amer SA, Li TC and Cooke ID (2003a) A prospective dose-finding study of the amount of thermal energy required for laparoscopic ovarian diathermy. *Hum Reprod* 18,1693–1698.
- Amer SA, Li TC and Cooke ID (2003b) Repeated laparoscopic ovarian diathermy is effective in women with anovulatory infertility due to polycystic ovary syndrome. *Fertil Steril* 79,1211–1215.
- Amin AF, Abd el-Aal DE, Darwish AM and Meki AR (2003) Evaluation of the impact of laparoscopic ovarian drilling on Doppler indices of ovarian stromal blood flow, serum vascular endothelial growth factor, and insulin-like growth factor-1 in women with polycystic ovary syndrome. *Fertil Steril* 79,938–941.
- Anttila L, Penttila TA, Matinlauri I, Koskinen P and Irjala K (1998) Serum total renin levels after ovarian electrocautery in women with polycystic ovary syndrome. *Gynecol Endocrinol* 12,327–331.
- Api M, Gorgen H and Cetin A (2005) Laparoscopic ovarian drilling in polycystic ovary syndrome. *Eur J Obstet Gynecol Reprod Biol* 119,76–81.

- Armar NA, McGarrigle HH, Honour J, Holownia P, Jacobs HS and Lachelin GC (1990) Laparoscopic ovarian diathermy in the management of anovulatory infertility in women with polycystic ovaries: endocrine changes and clinical outcome. *Fertil Steril* 53,45–49.
- Asada H, Kishi I, Kaseda S, Nakagawa H, Maruyama T and Yoshimura Y (2002) Laparoscopic treatment of polycystic ovaries with the holmium: YAG laser. *Fertil Steril* 77,852–853.
- Balen AH and Jacobs HS (1991) Gonadotropin surge attenuating factor: a missing link in the control of LH secretion? *Clin Endocrinol (Oxf)* 35,399–402.
- Balen AH and Jacobs HS (1994) A prospective study comparing unilateral and bilateral laparoscopic ovarian diathermy in women with the polycystic ovary syndrome. *Fertil Steril* 62,921–925.
- Balen AH, Conway GS, Kaltsas G, Techatrasak K, Manning PJ, West C and Jacobs HS (1995) Polycystic ovary syndrome: the spectrum of the disorder in 1741 patients. *Hum Reprod* 10,2107–2111.
- Baranowska B, Radzikowska M, Wasilewska-Dziubinska E, Kaplinski A, Roguski K and Plonowski A (1999) Neuropeptide Y, leptin, galanin and insulin in women with polycystic ovary syndrome. *Gynecol Endocrinol* 13,344–351.
- Beattie J, Allan GJ, Lochrie JD and Flint DJ (2006) Insulin-like growth factor-binding protein-5 (IGFBP-5): a critical member of the IGF axis. *Biochem J* 395,1–19.
- Ben Shlomo I, Homburg R and Shalev E (1998) Hyperandrogenic anovulation (the polycystic ovary syndrome)—back to the ovary? *Hum Reprod Update* 4,296–300.
- Berger MJ, Taymor ML and Patton WC (1975) Gonadotropin levels and secretory patterns in patients with typical and atypical polycystic ovarian disease. *Fertil Steril* 26,619–626.
- Bergh C, Carlsson B, Olsson JH, Selleskog U and Hillensjo T (1993) Regulation of androgen production in cultured human thecal cells by insulin-like growth factor I and insulin. *Fertil Steril* 59,323–331.
- Brandt M, Kehlet H, Binder C, Hagen C and McNeilly AS (1976) Effect of epidural analgesia on the glycoregulatory endocrine response to surgery. *Clin Endocrinol (Oxf)* 5,107–114.
- Broberger C (2005) Brain regulation of food intake and appetite: molecules and networks. *J Intern Med* 258,301–327.
- Burger HG (2002) Androgen production in women. *Fertil Steril* 77(Suppl 4), S3–S5.
- Campo S, Garcea N, Caruso A and Siccardi P (1983) Effect of celioscopic ovarian resection in patients with polycystic ovaries. *Gynecol Obstet Invest* 15,213–222.
- Campo S, Felli A, Lamanna MA, Barini A and Garcea N (1993) Endocrine changes and clinical outcome after laparoscopic ovarian resection in women with polycystic ovaries. *Hum Reprod* 8,359–363.
- Carmina E, Stanczyk FZ, Morris RS, Lee PD, Savjani G and Lobo RA (1995) Altered regulation of insulin-like growth factor binding protein-1 in patients with polycystic ovary syndrome. *J Soc Gynecol Invest* 2,743–747.
- Carmina E, Orio F, Palomba S, Cascella T, Longo RA, Colao AM, Lombardi G and Lobo RA (2005) Evidence for altered adipocyte function in polycystic ovary syndrome. *Eur J Endocrinol* 152,389–394.
- Chan JL and Mantzoros CS (2005) Role of leptin in energy-deprivation states: normal human physiology and clinical implications for hypothalamic amenorrhoea and anorexia nervosa. *Lancet* 366,74–85.
- Christenson LK and Stouffer RL (1997) Follicle-stimulating hormone and luteinizing hormone/chorionic gonadotropin stimulation of vascular endothelial growth factor production by macaque granulosa cells from pre- and periovulatory follicles. *J Clin Endocrinol Metab* 82,2135–2142.

- Cibula D, Kuzel D, Rezacsek K, Vrbikova J, Zivny J and Starka PL (2000) Multiple ovarian biopsy in the treatment of women with polycystic ovary syndrome (PCOS). *Clin Exp Obstet Gynecol* 27,194–196.
- Cleemann L, Lauszus FF and Trolle B (2004) Laparoscopic ovarian drilling as first line of treatment in infertile women with polycystic ovary syndrome. *Gynecol Endocrinol* 18,138–143.
- Cohen J (1996) Laparoscopic procedures for treatment of infertility related to polycystic ovarian syndrome. *Hum Reprod Update* 2,337–344.
- Cook CL, Siow Y, Brenner AG and Fallat ME (2002) Relationship between serum mullerian-inhibiting substance and other reproductive hormones in untreated women with polycystic ovary syndrome and normal women. *Fertil Steril* 77,141–146.
- Corbould A, Kim YB, Youngren JF, Pender C, Kahn BB, Lee A and Dunaif A (2005) Insulin resistance in the skeletal muscle of women with PCOS involves intrinsic and acquired defects in insulin signaling. *Am J Physiol Endocrinol Metab* 288,E1047–E1054.
- Cortet-Rudelli C, Pigny P, Decanter C, Leroy M, Maunoury-Lefebvre C, Thomas-Desrousseaux P and Dewailly D (2002) Obesity and serum luteinizing hormone level have an independent and opposite effect on the serum inhibin B level in patients with polycystic ovary syndrome. *Fertil Steril* 77,281–287.
- Dafopoulos K, Kotsovassilis CG, Milingos S, Kallitsaris A, Galazios G, Zintzaras E, Sotiropoulos P and Messinis IE (2004) Changes in pituitary sensitivity to GnRH in estrogen-treated postmenopausal women: evidence that gonadotropin surge attenuating factor plays a physiological role. *Hum Reprod* 19,1985–1992.
- Dale PO, Tanbo T, Ertzeid G, Bjerkce S, Oldereid N, Fedorcsak P and Abyholm T (2004) The impact of insulin resistance on the outcome of laparoscopic ovarian electrocautery in infertile women with the polycystic ovary syndrome. *Gynecol Endocrinol* 19,182–189.
- DeVane GW, Czekala NM, Judd HL and Yen SS (1975) Circulating gonadotropins, estrogens and androgens in polycystic ovarian disease. *Am J Obstet Gynecol* 121,496–500.
- Diamanti-Kandarakis E, Mitrakou A, Hennes MM, Platanissiotis D, Kaklas N, Spina J, Georgiadou E, Hoffmann RG, Kissebah AH and Raptis S (1995) Insulin sensitivity and antiandrogenic therapy in women with polycystic ovary syndrome. *Metabolism* 44,525–531.
- Duleba AJ, Banaszewska B, Spaczynski RZ and Pawelczyk L (2003) Success of laparoscopic ovarian wedge resection is related to obesity, lipid profile, and insulin levels. *Fertil Steril* 79,1008–1014.
- Dunaif A (1997) Insulin resistance and the polycystic ovary syndrome: mechanism and implications for pathogenesis. *Endocr Rev* 18,774–800.
- Dunaif A, Segal KR, Futterweit W and Dobrjansky A (1989) Profound peripheral insulin resistance, independent of obesity, in polycystic ovary syndrome. *Diabetes* 38,1165–1174.
- Dunaif A, Green G, Futterweit W and Dobrjansky A (1990) Suppression of hyperandrogenism does not improve peripheral or hepatic insulin resistance in the polycystic ovary syndrome. *J Clin Endocrinol Metab* 70,699–704.
- Durlinger AL, Visser JA and Themmen AP (2002) Regulation of ovarian function: the role of anti-Mullerian hormone. *Reproduction* 124,601–609.
- Durlinger AL, Gruijters MJ, Kramer P, Karels B, Kumar TR, Matzuk MM, Rose UM, De Jong FH, Uilenbroek JT and Grootegoed JA et al. (2001) Anti-Mullerian hormone attenuates the effects of FSH on follicle development in the mouse ovary. *Endocrinology* 142,4891–4899.
- Eden JA, Place J, Carter GD, Alagband-Zadeh J and Pawson ME (1989) The role of chronic anovulation in the polycystic ovary syndrome: normalization of sex-hormone-binding globulin levels after clomifene-induced ovulation. *Clin Endocrinol (Oxf)* 30,323–332.

- Ehrmann DA, Sturis J, Byrne MM, Karrison T, Rosenfield RL and Polonsky KS (1995) Insulin secretory defects in polycystic ovary syndrome. Relationship to insulin sensitivity and family history of non-insulindependent diabetes mellitus. *J Clin Invest* 96,520–527.
- El OH, Ghalia AA, Khalifa A, Mahfouz H, El SA and Shoieb N (1999) Serum leptin as an additional possible pathogenic factor in polycystic ovary syndrome. *Clin Biochem* 32,71–75.
- bd El Aal DE, Mohamed SA, Amine AF and Meki AR (2005) Vascular endothelial growth factor and insulin-like growth factor-1 in polycystic ovary syndrome and their relation to ovarian blood flow. *Eur J Obstet Gynecol Reprod Biol* 118,219–224.
- Elkind-Hirsch KE, Valdes CT and Malinak LR (1993) Insulin resistance improves in hyperandrogenic women treated with Lupron. *Fertil Steril* 60,634–641.
- Elting MW, Korsen TJ, Bezemer PD and Schoemaker J (2001) Prevalence of diabetes mellitus, hypertension and cardiac complaints in a follow-up study of a Dutch PCOS population. *Hum Reprod* 16,556–560.
- Farhi J, Soule S and Jacobs HS (1995) Effect of laparoscopic ovarian electrocautery on ovarian response and outcome of treatment with gonadotropins in clomifene citrate-resistant patients with polycystic ovary syndrome. *Fertil Steril* 64,930–935.
- Farquhar CM, Williamson K, Gudex G, Johnson NP, Garland J and Sadler L (2002) A randomized controlled trial of laparoscopic ovarian diathermy versus gonadotropin therapy for women with clomifene citrate-resistant polycystic ovary syndrome. *Fertil Steril* 78,404–411.
- Fassnacht M, Schlenz N, Schneider SB, Wudy SA, Alolio B and Arlt W (2003) Beyond adrenal and ovarian androgen generation: increased peripheral 5 alpha-reductase activity in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 88,2760–2766.
- Felemban A, Tan SL and Tulandi T (2000) Laparoscopic treatment of polycystic ovaries with insulated needle cautery: a reappraisal. *Fertil Steril* 73,266–269.
- Fowler PA, Messinis IE and Templeton AA (1990) Inhibition of LHRH-induced LH and FSH release by gonadotropin surge-attenuating factor (GnSAF) from human follicular fluid. *J Reprod Fertil* 90,587–594.
- Fowler DJ, Nicolaidis KH and Miell JP (2000) Insulin-like growth factor binding protein-1 (IGFBP-1): a multifunctional role in the human female reproductive tract. *Hum Reprod Update* 6,495–504.
- Fowler PA, Sorsa-Leslie T, Harris W and Mason HD (2003) Ovarian gonadotropin surge-attenuating factor (GnSAF): where are we after 20 years of research? *Reproduction* 126,689–699.
- Fukaya T, Murakami T, Tamura M, Watanabe T, Terada Y and Yajima A (1995) Laser vaporization of the ovarian surface in polycystic ovary disease results in reduced ovarian hyperstimulation and improved pregnancy rates. *Am J Obstet Gynecol* 173,119–125.
- Gadir AA, Khatim MS, Mowafi RS, Alnaser HM, Alzaid HG and Shaw RW (1990) Hormonal changes in patients with polycystic ovarian disease after ovarian electrocautery or pituitary desensitization. *Clin Endocrinol (Oxf)* 32,749–754.
- Gadir AA, Alnaser HM, Mowafi RS and Shaw RW (1992) The response of patients with polycystic ovarian disease to human menopausal gonadotropin therapy after ovarian electrocautery or a luteinizing hormone-releasing hormone agonist. *Fertil Steril* 57,309–313.
- Gdanský E, Diamant YZ, Laron Z, Silbergeld A, Kaplan B and Eshet R (1997) Increased number of IGF-I receptors on erythrocytes of women with polycystic ovarian syndrome. *Clin Endocrinol (Oxf)* 47,185–190.
- Geva E and Jaffe RB (2000) Role of vascular endothelial growth factor in ovarian physiology and pathology. *Fertil Steril* 74,429–438.

- Giudice LC (1995) The insulin-like growth factor system in normal and abnormal human ovarian follicle development. *Am J Med* 98,48S–54S.
- Giudice LC, van Dessel HJ, Cataldo NA, Chandrasekher YA, Yap OW and Fauser BC (1995) Circulating and ovarian IGF binding proteins: potential roles in normo-ovulatory cycles and in polycystic ovarian syndrome. *Prog Growth Factor Res* 6,397–408.
- Gjonnaess H (1998) Late endocrine effects of ovarian electrocautery in women with polycystic ovary syndrome. *Fertil Steril* 69,697–701.
- Gjonnaess H (1999) Comparison of ovarian electrocautery and oral contraceptives in the treatment of hyperandrogenism in women with polycystic ovary syndrome. *Acta Obstet Gynecol Scand* 78,530–533.
- Gjonnaess H and Norman N (1987) Endocrine effects of ovarian electrocautery in patients with polycystic ovarian disease. *Br J Obstet Gynaecol* 94, 779–783.
- Greenblatt E and Casper RF (1987) Endocrine changes after laparoscopic ovarian cautery in polycystic ovarian syndrome. *Am J Obstet Gynecol* 156,279–285.
- Hacihanefioglu B, Seyisoglu H, Karsidag K, Elter K, Aksu F, Yilmaz T and Gurol AO (2000) Influence of insulin resistance on total renin level in normotensive women with polycystic ovary syndrome. *Fertil Steril* 73,261–265.
- Hagen C, Brandt MR and Kehlet H (1980) Prolactin, LH, FSH, GH, and cortisol response to surgery and the effect of epidural analgesia. *Acta Endocrinol (Copenh)* 94,151–154.
- Havelock JC, Rainey WE and Carr BR (2004) Ovarian granulosa cell lines. *Mol Cell Endocrinol* 228,67–78.
- Hayashi H, Ezaki K, Endo H and Urashima M (2005) Preoperative luteinizing hormone levels predict the ovulatory response to laparoscopic ovarian drilling in patients with clomifene citrate-resistant polycystic ovary syndrome. *Gynecol Endocrinol* 21,307–311.
- Hazewinkel M (2002) *Encyclopaedia of Mathematics*. Springer-Verlag, Berlin, Heidelberg, New York.
- Heineman MJ, Thomas CM, Doesburg WH and Rolland R (1984) Pituitary and ovarian responses to LHRH stimulation in women with clinical features of the polycystic ovary syndrome. *Eur J Obstet Gynecol Reprod Biol* 17,273–284.
- Hernandez I, Parra A, Mendez I, Cabrera V, Cravioto MC, Mercado M, Diaz-Sanchez V and Larrea F (2000) Hypothalamic dopaminergic tone and prolactin bioactivity in women with polycystic ovary syndrome. *Arch Med Res* 31,216–222.
- Jaatinen TA, Matinlauri I, Anttila L, Koskinen P, Erkkola R and Irjala K (1995) Serum total renin is elevated in women with polycystic ovarian syndrome. *Fertil Steril* 63,1000–1004.
- Jacobs HS and Conway GS (1999) Leptin, polycystic ovaries and polycystic ovary syndrome. *Hum Reprod Update* 5,166–171.
- Jakimiuk AJ, Weitsman SR and Magoffin DA (1999) 5 α -reductase activity in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 84,2414–2418.
- Jamal HS (2000) Bilateral or unilateral KTP laser ovarian drilling in polycystic ovarian disease. *Ann Saudi Med* 20,165–167.
- Jockenhovel F, Blum WF, Vogel E, Englaro P, Muller-Wieland D, Reinwein D, Rascher W and Krone W (1997) Testosterone substitution normalizes elevated serum leptin levels in hypogonadal men. *J Clin Endocrinol Metab* 82,2510–2513.
- Jonard S and Dewailly D (2004) The follicular excess in polycystic ovaries, due to intra-ovarian hyperandrogenism, may be the main culprit for the follicular arrest. *Hum Reprod Update* 10,107–117.
- Judd HL, Rigg LA, Anderson DC and Yen SS (1976) The effects of ovarian wedge resection on circulating gonadotropin and ovarian steroid levels in patients with polycystic ovary syndrome. *J Clin Endocrinol Metab* 43,347–355.

- Kaaijk EM, Hamerlynck JVTH, Beek JF and van der Veen F (1999) Clinical outcome after unilateral oophorectomy in patients with polycystic ovary syndrome. *Hum Reprod* 14,889–892.
- Kamel MA, Abdel Hamid AED, Abdel-Rahim M and Mostafa SA (2004) Laparoscopic ovarian re-electro cautery versus ovulation induction with FSH for persistent anovulation after laparoscopic PCOS treatment. *Middle East Fertil Soc J* 9,70–78.
- Kandil M and Selim M (2005) Hormonal and sonographic assessment of ovarian reserve before and after laparoscopic ovarian drilling in polycystic ovary syndrome. *BJOG* 112,1427–1430.
- Katz M, Carr PJ, Cohen BM and Millar RP (1978) Hormonal effects of wedge resection of polycystic ovaries. *Obstet Gynecol* 51,437–444.
- Keckstein G, Rossmannith W, Spatzier K, Schneider V, Borchers K and Steiner R (1990) The effect of laparoscopic treatment of polycystic ovarian disease by CO₂-laser or Nd:YAG laser. *Surg Endosc* 4,103–107.
- Kojima E, Yanagibori A, Otaka K and Hirakawa S (1989) Ovarian wedge resection with contact Nd:YAG laser irradiation used laparoscopically. *J Reprod Med* 34,444–446.
- Kolaczynski JW, Nyce MR, Considine RV, Boden G, Nolan JJ, Henry R, Mudaliar SR, Olefsky J and Caro JF (1996) Acute and chronic effects of insulin on leptin production in humans: Studies in vivo and in vitro. *Diabetes* 45,699–701.
- de Koning J, Lambalk CB, Helmerhorst FM and Helder MN (2001) Is GnRH self-priming an obligatory feature of the reproductive cycle? *Hum Reprod* 16,209–214.
- Kovacs G, Buckler H, Bangah M, Outch K, Burger H, Healy D, Baker G and Phillips S (1991) Treatment of anovulation due to polycystic ovarian syndrome by laparoscopic ovarian electrocautery. *Br J Obstet Gynaecol* 98,30–35.
- Kucuk M and Kilic-Okman T (2005) Hormone profiles and clinical outcome after laparoscopic ovarian drilling in women with polycystic ovary syndrome. *Med Sci Monit* 11,CR29–CR34.
- Kumar A, Woods KS, Bartolucci AA and Azziz R (2005) Prevalence of adrenal androgen excess in patients with the polycystic ovary syndrome (PCOS). *Clin Endocrinol (Oxf)* 62,644–649.
- Labrie F, Luu-The V, Belanger A, Lin SX, Simard J, Pelletier G and Labrie C (2005) Is dehydroepiandrosterone a hormone? *J Endocrinol* 187,169–196.
- Lambalk CB, De Koning CH and Braat DD (1998) The endocrinology of dizygotic twinning in the human. *Mol Cell Endocrinol* 145,97–102.
- Laven JS and Fauser BC (2004) Inhibins and adult ovarian function. *Mol Cell Endocrinol* 225,37–44.
- Legro RS, Castracane VD and Kauffman RP (2004) Detecting insulin resistance in polycystic ovary syndrome: purposes and pitfalls. *Obstet Gynecol Surv* 59,141–154.
- Legro RS, Kunselman AR, Dodson WC and Dunaif A (1999) Prevalence and predictors of risk for type 2 diabetes mellitus and impaired glucose tolerance in polycystic ovary syndrome: a prospective, controlled study in 254 affected women. *J Clin Endocrinol Metab* 84,165–169.
- Lemieux S, Lewis GF, Ben Chetrit A, Steiner G and Greenblatt EM (1999) Correction of hyperandrogenemia by laparoscopic ovarian cautery in women with polycystic ovarian syndrome is not accompanied by improved insulin sensitivity or lipid-lipoprotein levels. *J Clin Endocrinol Metab* 84,4278–4282.
- Leung DW, Cachianes G, Kuang WJ, Goeddel DV and Ferrara N (1989) Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 246,1306–1309.
- Liguori G, Tolino A, Moccia G, Scognamiglio G and Nappi C (1996) Laparoscopic ovarian treatment in infertile patients with polycystic ovarian syndrome (PCOS): endocrine changes and clinical outcome. *Gynecol Endocrinol* 10,257–264.

- Lockwood GM, Muttukrishna S, Groome NP, Matthews DR and Ledger WL (1998) Mid-follicular phase pulses of inhibin B are absent in polycystic ovarian syndrome and are initiated by successful laparoscopic ovarian diathermy: a possible mechanism regulating emergence of the dominant follicle. *J Clin Endocrinol Metab* 83,1730–1735.
- Lonnqvist F, Arner P, Nordfors L and Schalling M (1995) Overexpression of the obese (ob) gene in adipose tissue of human obese subjects. *Nat Med* 1,950–953.
- Loverro G, Lorusso F, Mei L, Depalo R, Cormio G and Selvaggi L (2002) The plasma homocysteine levels are increased in polycystic ovary syndrome. *Gynecol Obstet Invest* 53,157–162.
- Magoffin DA and Jakimiuk AJ (1998) Inhibin A, inhibin B and activin A concentrations in follicular fluid from women with polycystic ovary syndrome. *Hum Reprod* 13,2693–2698.
- Malkawi HY and Qublan HS (2005) Laparoscopic ovarian drilling in the treatment of polycystic ovary syndrome: how many punctures per ovary are needed to improve the reproductive outcome? *J Obstet Gynaecol Res* 31,115–119.
- Malkawi HY, Qublan HS and Hamaideh AH (2003) Medical vs. surgical treatment for clomifene citrate-resistant women with polycystic ovary syndrome. *J Obstet Gynaecol* 23,289–293.
- Messinis IE (2006) Ovarian feedback, mechanism of action and possible clinical implications. *Hum Reprod* 12(5),557–571.
- Messinis IE, Hirsch P and Templeton AA (1991) Follicle stimulating hormone stimulates the production of gonadotropin surge attenuating factor (GnSAF) in vivo. *Clin Endocrinol (Oxf)* 35,403–407.
- Messinis IE, Milingos SD, Alexandris E, Kariotis I, Kollios G and Seferiadis K (1999) Leptin concentrations in normal women following bilateral ovariectomy. *Hum Reprod* 14,913–918.
- Minakami H, Abe N, Oka N, Kimura K, Tamura T and Tamada T (1988) Prolactin release in polycystic ovarian syndrome. *Endocrinol Jpn* 35,303–310.
- Mio Y, Toda T, Tanikawa M, Terado H, Harada T and Terakawa N (1991) Transvaginal ultrasound-guided follicular aspiration in the management of anovulatory infertility associated with polycystic ovaries. *Fertil Steril* 56,1060–1065.
- Moggetti P, Tosi F, Castello R, Magnani CM, Negri C, Brun E, Furlani L, Caputo M and Muggeo M (1996) The insulin resistance in women with hyperandrogenism is partially reversed by antiandrogen treatment: evidence that androgens impair insulin action in women. *J Clin Endocrinol Metab* 81,952–960.
- Morgante G, La MA, Setacci F, Setacci C, Petraglia F and De LV (2002) The cardiovascular risk factor homocysteine is not elevated in young women with hyperandrogenism or hypoestrogenism. *Gynecol Obstet Invest* 53,200–203.
- Naether OG, Fischer R, Weise HC, Geiger-Kotzler L, Delfs T and Rudolf K (1993) Laparoscopic electrocoagulation of the ovarian surface in infertile patients with polycystic ovarian disease. *Fertil Steril* 60,88–94.
- Norman RJ, Milner CR, Groome NP and Robertson DM (2001) Circulating follistatin concentrations are higher and activin concentrations are lower in polycystic ovarian syndrome. *Hum Reprod* 16,668–672.
- Orio F, Jr, Palomba S, Jr, Spinelli L, Jr, Cascella T, Jr, Tauchmanova L, Jr, Zullo F, Jr, Lombardi G Jr and Colao A, Jr (2004) The cardiovascular risk of young women with polycystic ovary syndrome: an observational, analytical, prospective case-control study. *J Clin Endocrinol Metab* 89,3696–3701.
- Panidis D, Farmakiotis D, Rousso D, Katsikis I, Kourtis A and Diamanti-Kandarakis E (2005) Serum luteinizing hormone levels are markedly increased and significantly correlated with Delta4-androstenedione levels in lean women with polycystic ovary syndrome. *Fertil Steril* 84,538–540.

- Parsanezhad ME, Bagheri MH, Alborzi S and Schmidt EH (2003) Ovarian stromal blood flow changes after laparoscopic ovarian cauterization in women with polycystic ovary syndrome. *Hum Reprod* 18,1432–1437.
- Parsanezhad ME, Alborzi S, Zolghadril J, Parsanezhad M, Keshavarzi G, Omrani GR and Schmidt EH (2005) Hyperprolactinemia after laparoscopic ovarian drilling: an unknown phenomenon. *Reprod Biol Endocrinol* 3,6.
- Pigny P, Jonard S, Robert Y and Dewailly D (2006) Serum anti-Mullerian hormone as a surrogate for antral follicle count for definition of the polycystic ovary syndrome. *J Clin Endocrinol Metab* 91,941–945.
- Plymate SR, Matej LA, Jones RE and Friedl KE (1988) Inhibition of sex hormone-binding globulin production in the human hepatoma (HepG2) cell line by insulin and prolactin. *J Clin Endocrinol Metab* 67,460–464.
- Remsberg KE, Talbott EO, Zborowski JV, Evans RW and McHugh-Pemu K (2002) Evidence for competing effects of body mass, hyperinsulinemia, insulin resistance, and androgens on leptin levels among lean, overweight, and obese women with polycystic ovary syndrome. *Fertil Steril* 78,479–486.
- van Rooij I, Broekmans FJ, te Velde ER, Fauser BC, Bancsi LF, de Jong FH and Themmen AP (2002) Serum anti-Mullerian hormone levels: a novel measure of ovarian reserve. *Hum Reprod* 17,3065–3071.
- van Rooij I, Broekmans FJ, Scheffer GJ, Looman CW, Habbema JD, de Jong FH, Fauser BJ, Themmen AP and te Velde ER (2005) Serum antimullerian hormone levels best reflect the reproductive decline with age in normal women with proven fertility: a longitudinal study. *Fertil Steril* 83,979–987.
- Rossmannith WG, Keckstein J, Spatzier K and Lauritzen C (1991) The impact of ovarian laser surgery on the gonadotropin secretion in women with polycystic ovarian disease. *Clin Endocrinol (Oxf)* 34,223–230.
- Rotterdam Consensus (2004) Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Hum Reprod* 19,41–47.
- Sakata M, Tasaka K, Kurachi H, Terakawa N, Miyake A and Tanizawa O (1990) Changes of bioactive luteinizing hormone after laparoscopic ovarian cauterization in patients with polycystic ovarian syndrome. *Fertil Steril* 53,610–613.
- Saleh A, Morris D, Tan SL and Tulandi T (2001) Effects of laparoscopic ovarian drilling on adrenal steroids in polycystic ovary syndrome patients with and without hyperinsulinemia. *Fertil Steril* 75,501–504.
- van Santbrink EJ, Hop WC and Fauser BC (1997) Classification of normogonadotropic infertility: polycystic ovaries diagnosed by ultrasound versus endocrine characteristics of polycystic ovary syndrome. *Fertil Steril* 67,452–458.
- Sassin JF, Frantz AG and Kapen S Weitzman ED (1973) The nocturnal rise of human prolactin is dependent on sleep. *J Clin Endocrinol Metab* 37,436–440.
- Schachter M, Raziell A, Friedler S, Strassburger D, Bern O and Ron-El R (2003) Insulin resistance in patients with polycystic ovary syndrome is associated with elevated plasma homocysteine. *Hum Reprod* 18,721–727.
- Sills ES, Genton MG, Perloe M, Schattman GL, Bralley JA and Tucker MJ (2001) Plasma homocysteine, fasting insulin, and androgen patterns among women with polycystic ovaries and infertility. *J Obstet Gynaecol Res* 27,163–168.
- Silva PD, Gentzschlein EE and Lobo RA (1987) Androstenedione may be a more important precursor of tissue dihydrotestosterone than testosterone in women. *Fertil Steril* 48,419–422.
- Soliman EM, Attia AM, Elebrashi AN, Younis AS and Salit ME (2000) Laparoscopic ovarian electrocauterization improves ovarian response to gonadotropins in clomifene citrate resistant patients with polycystic ovary syndrome. *Middle East Fertil Soc J* 5,120–125.

- Soules MR, Sutton GP, Hammond CB and Haney AF (1980) Endocrine changes at operation under general anesthesia: reproductive hormone fluctuations in young women. *Fertil Steril* 33,364–371.
- Stein I and Leventhal M (1935) Amenorrhea associated with bilateral polycystic ovaries. *Am J Obstet Gynecol* 29,181–91.
- Sumioki H, Utsunomiya T, Matsuoka K, Korenaga M and Kadota T (1988) The effect of laparoscopic multiple punch resection of the ovary on hypothalamo-pituitary axis in polycystic ovary syndrome. *Fertil Steril* 50,567–572.
- Szilagyi A, Rossmanith W, Csermely T and Csaba I (1990) Changes in circulating hormone levels after ovarian wedge resection in patients with polycystic ovary syndrome. *Arch Gynecol Obstet* 248,31–35.
- Szilagyi A, Hole R, Keckstein J and Rossmanith WG (1993) Effects of ovarian surgery on the dopaminergic and opioidergic control of gonadotropin and prolactin secretion in women with polycystic ovarian disease. *Gynecol Endocrinol* 7,159–166.
- Takeuchi S, Futamura N, Takubo S, Noda N, Minoura H and Toyoda N (2002) Polycystic ovary syndrome treated with laparoscopic ovarian drilling with a harmonic scalpel. A prospective, randomized study. *J Reprod Med* 47,816–820.
- Talbott E, Guzick D, Clerici A, Berga S, Detre K, Weimer K and Kuller L (1995) Coronary heart disease risk factors in women with polycystic ovary syndrome. *Arterioscler Thromb Vasc Biol* 15,821–826.
- Tanaka T, Fujimoto S and Kutsuzawa T (1978) The effect of ovarian wedge resection and incision on circulating gonadotropin in patients with polycystic ovarian disease. *Int J Fertil* 23,93–99.
- Tasaka K, Sakata M, Kurachi H, Komura H, Miyake A and Tanizawa O (1990) Electrocautery in polycystic ovary syndrome. *Horm Res* 33(Suppl 2), 40–42.
- Taskin O, Yalcinoglu AI, Kafkasli A, Burak F and Ozekici U (1996) Comparison of the effects of ovarian cauterization and gonadotropin-releasing hormone agonist and oral contraceptive therapy combination on endocrine changes in women with polycystic ovary disease. *Fertil Steril* 65,1115–1118.
- Telli MH, Yildirim M and Noyan V (2002) Serum leptin levels in patients with polycystic ovary syndrome. *Fertil Steril* 77,932–935.
- Tiitinen A, Tenhunen A and Seppala M (1993) Ovarian electrocauterization causes LH-regulated but not insulin-regulated endocrine changes. *Clin Endocrinol (Oxf)* 39,181–184.
- Tinkanen H, Blauer M, Laippala P, Tuohimaa P and Kujansuu E (2001) Correlation between serum inhibin B and other indicators of the ovarian function. *Eur J Obstet Gynecol Reprod Biol* 94,109–113.
- Tulandi T, Saleh A, Morris D, Jacobs HS, Payne NN and Tan SL (2000) Effects of laparoscopic ovarian drilling on serum vascular endothelial growth factor and on insulin responses to the oral glucose tolerance test in women with polycystic ovary syndrome. *Fertil Steril* 74,585–588.
- Tyson JE, Hwang P, Guyda H and Friesen HG (1972) Studies of prolactin secretion in human pregnancy. *Am J Obstet Gynecol* 113,14–20.
- Utsunomiya T, Sumioki H and Taniguchi I (1990) Hormonal and clinical effects of multifollicular puncture and resection on the ovaries of polycystic ovary syndrome. *Horm Res* 33(Suppl 2),35–39.
- Vejlsted H and Albrechtsen R (1976) Biochemical and clinical effect of ovarian wedge resection in the polycystic ovary syndrome. *Obstet Gynecol* 47,575–580.
- Verhelst J, Gerris J, Joostens M, van der Meer S, van Royen E and Mahler C (1993) Clinical and endocrine effects of laser vaporization in patients with polycystic ovarian disease. *Gynecol Endocrinol* 7,49–55.

- Vinson GP, Saridogan E, Puddefoot JR and Djahanbakhch O (1997) Tissue renin-angiotensin systems and reproduction. *Hum Reprod* 12,651–662.
- Urbikova J, Tallova J, Bicikova M, Dvorakova K, Hill M and Starka L (2003) Plasma thiols and androgen levels in polycystic ovary syndrome. *Clin Chem Lab Med* 41,216–221.
- Vural B, Caliskan E, Turkoz E, Kilic T and Demirci A (2005) Evaluation of metabolic syndrome frequency and premature carotid atherosclerosis in young women with polycystic ovary syndrome. *Hum Reprod* 20,2409–2413.
- van der Weiden RM and Alberda AT (1987) Laparoscopic ovarian electrocautery in patients with polycystic ovarian disease resistant to clomifene citrate. *Surg Endosc* 1,217–219.
- van der Weiden RM, Alberda AT, de Jong FH and Brandenburg H (1989) Endocrine effects of laparoscopic ovarian electrocautery in patients with polycystic ovarian disease, resistant to clomifene citrate. *Eur J Obstet Gynecol Reprod Biol* 32,157–162.
- Welt CK, Taylor AE, Martin KA and Hall JE (2002) Serum inhibin B in polycystic ovary syndrome: regulation by insulin and luteinizing hormone. *J Clin Endocrinol Metab* 87,5559–5565.
- Wijeyaratne CN, Nirantharakumar K, Balen AH, Barth JH, Sheriff R and Belchetz PE (2004) Plasma homocysteine in polycystic ovary syndrome: does it correlate with insulin resistance and ethnicity? *Clin Endocrinol (Oxf)* 60,560–567.
- Wild RA (2002) Long-term health consequences of PCOS. *Hum Reprod Update* 8,231–241.
- Wild RA, Painter PC, Coulson PB, Carruth KB and Ranney GB (1985) Lipoprotein lipid concentrations and cardiovascular risk in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 61,946–951.
- Wu MH, Huang MF, Tsai SJ, Pan HA, Cheng YC and Lin YS (2004) Effects of laparoscopic ovarian drilling on young adult women with polycystic ovarian syndrome. *J Am Assoc Gynecol Laparosc* 11,184–190.
- Wu XK, Zhou SY, Sallinen K, Pollanen P and Erkkola R (2000) Ovarian-adrenal cross-talk in polycystic ovary syndrome: evidence from wedge resection. *Eur J Endocrinol* 143,383–388.
- Yarali H, Yildirim A, Aybar F, Kabakci G, Bukulmez O, Akgul E and Oto A (2001) Diastolic dysfunction and increased serum homocysteine concentrations may contribute to increased cardiovascular risk in patients with polycystic ovary syndrome. *Fertil Steril* 76,511–516.
- Zullo F, Pellicano M, Zupi E, Guida M, Mastrantonio P and Nappi C (2000) Minilaparoscopic ovarian drilling under local anesthesia in patients with polycystic ovary syndrome. *Fertil Steril* 74,376–379.