

VU Research Portal

Retinoblastoma

Dommering, C.J.

2015

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Dommering, C. J. (2015). *Retinoblastoma: molecular genetics and clinical consequences*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl



CHAPTER

RB1 MUTATIONS AND SECOND PRIMARY MALIGNANCIES AFTER HEREDITARY RETINOBLASTOMA

3

Charlotte J. Dommering*, Tamara Marees*, Annemarie H. van der Hout,
Saskia M. Imhof, Hanne Meijers-Heijboer, Peter J. Ringens, Flora E. van Leeuwen,
Annette C. Moll

* Both authors contributed equally to the manuscript

Fam Cancer. 2012;11(2):225-33

ABSTRACT

Survivors of hereditary retinoblastoma have a high risk of second primary malignancies, but it has not been investigated whether specific *RB1* germline mutations are associated with greater risk of second primary malignancies in a large cohort. We conducted a retrospective cohort study of 199 survivors of hereditary retinoblastoma with a documented *RB1* germline mutation diagnosed between 1905-2005.

In total, 44 hereditary retinoblastoma survivors developed a second primary malignancy after a median follow-up of 30.2 years (range, 1.33-76.0). A significantly increased risk of second primary malignancy was observed among carriers of one of the 11 recurrent CGA>TGA nonsense *RB1* mutations (hazard ratio (HR)=3.53; [95% confidence interval (CI)=1.82-6.84]; P=.000), and there was a significantly lower risk for subjects with a low penetrance mutation (HR=.19; [95% CI=.05-.81]; P=.025).

Our findings suggest a genotype-phenotype correlation for second primary cancers of retinoblastoma survivors and may impact on long-term surveillance protocols of patients with hereditary retinoblastoma, if confirmed by future studies.

INTRODUCTION

Retinoblastoma is the most common primary intraocular malignancy of childhood.¹ Mutational inactivation of both alleles of the *RB1* tumor suppressor gene in the developing retina initiates the formation of retinoblastoma.^{2,3} The *RB1* gene consists of 27 exons and is located on chromosome 13q14 (GenBank accession number L11910, MIM#180200). The gene encodes a ubiquitously expressed nuclear protein, which is involved in cell cycle regulation, cellular differentiation and survival.⁴ About 40% of retinoblastoma patients have a hereditary predisposition, caused by a heterozygous germline mutation in the *RB1* gene and are usually bilaterally affected.⁵ Over 600 different pathogenic mutations have been described. Patients with nonhereditary retinoblastoma only have one eye affected, no germline mutation in the *RB1* gene and two somatic retinal *RB1* mutations.

As is known from long-term follow-up studies⁶⁻¹⁰, hereditary retinoblastoma subjects have a strongly increased risk for second primary malignancies, (including osteosarcoma, soft tissue sarcoma, melanoma and epithelial cancers) which is associated with excess mortality.¹¹⁻¹³ So far, it has not been examined in a large cohort of retinoblastoma patients whether specific *RB1* mutations might be associated with greater risk of second malignancy.

The objective of the present study was to investigate the *RB1* genotype in relation to second malignancy risk in hereditary retinoblastoma subjects.

MATERIALS AND METHODS

Patients

In the Netherlands we have data available of Dutch retinoblastoma subjects diagnosed from 1862 onwards. Detailed information on data collection and follow-up has been described previously.⁹ Relevant data collected for the present study were family history of retinoblastoma, tumor laterality, treatment for retinoblastoma, reports on invasive cancers, and date and (underlying) cause of death. Only the first cancer after retinoblastoma was included in this study. Time at risk for a second primary cancer began at diagnosis of retinoblastoma and ended on the date of second malignancy diagnosis, emigration, the date last known to be alive, the date of death, or the closing date of the study, whichever came first.

Patients with bilateral disease, a positive family history of retinoblastoma, or a germline mutation in the *RB1* gene detected by chromosomal or DNA analysis were classified as hereditary. The remaining patients, those with unilateral retinoblastoma, no family history of retinoblastoma, and no germline mutation detected in the *RB1* gene, were classified as having non-hereditary retinoblastoma.

Eligible subjects for the current study included all hereditary retinoblastoma patients from the Dutch retinoblastoma cohort (1862-2005), in whom a germline

RB1 mutation was documented. If a retinoblastoma patient had died before DNA-testing could be performed, but a *RB1* mutation was determined in the family, the patient was considered to be a carrier of the familial mutation (n = 26). Every affected family member was handled as a single case in the analysis. Of the 1028 retinoblastoma patients in the Dutch cohort, we identified a total of 410 (39.89) hereditary cases. Two-hundred eleven patients were excluded because no DNA analysis could be performed (n = 180) or DNA analysis did not detect a *RB1* mutation (n = 31). The remaining 199 patients were included in this study (see **Figure 1**), of whom 168 were alive at the time of inclusion.

This study was approved by the Medical Ethics Committees of all participating hospitals, and was conducted in accordance with the principles of the Helsinki declaration.

Mutation screening


Since the beginning of the 1990s all newly diagnosed retinoblastoma patients in the Netherlands undergo germline *RB1* mutation analysis. Many patients who were diagnosed before that time underwent DNA-testing between 1990 and 2005, when they were referred to the clinical genetics department (n = 90). Rb patients diagnosed prior to 1990, in whom mutation testing had not yet been performed at the time of the study and who wanted to participate in the study, were invited to undergo DNA-testing and were offered genetic counseling (n = 23).

DNA analysis included direct sequencing of exons 1 and 15, and the *RB1* promoter and Denaturing Gradient Gel Electrophoresis (DGGE) analysis of the other exons and flanking intronic sequences. To detect large deletions and duplications Multiplex Ligation-dependent Probe Amplification (MLPA) analysis was performed. If warranted, e.g. when dysmorphic features or mental retardation was noted, karyotyping was performed to detect chromosomal rearrangements. With these techniques we have been able to detect 90% of mutations in familial and/or bilateral cases.

Type of *RB1* mutation

In the *RB1* gene are several methylated CGA codons known to lead to 11 recurrent nonsense mutations by C>T transitions.¹⁴⁻¹⁶ An important factor in the high recurrence of mutations at these sites was shown to be deamination of 5-methylcytosine.¹⁷

For this study mutations in the promoter, exon 1, missense mutations and deletions of the complete *RB1* gene were regarded as low penetrance mutations, based on previous studies.¹⁸⁻²¹ Four out of six familial splice mutations were also regarded as low penetrance mutations, based on a diseased eye ratio (DER) of ≤ 1.5 , defined as the total number of affected eyes per family divided by the number of mutation carriers in the family.^{22,23} To exclude possible mosaicism as a cause of milder expression the first mutation carrier in these families was excluded from the analysis.



Rb = retinoblastoma, SPT = second primary tumor

Figure 1. Flow chart showing reasons for inclusion and exclusion of retinoblastoma patients with hereditary retinoblastoma from our cohort. In the total group of 410 hereditary retinoblastoma patients from our cohort, 99 second primary tumors (SPT) have been diagnosed. In the flow chart is also depicted in which in- or excluded group these SPT's have occurred. Percentage is calculated from the total of 99 SPT's.

Statistical Methods

We compared the frequency of second primary cancers among hereditary retinoblastoma survivors with specific documented *RB1* mutations, and tested for differences using chi-square tests.

Multivariate Cox regression analysis was performed to quantify the effects of specific *RB1* mutations on the risk of second primary malignancies. Therapy, age, laterality, sex, and familial or sporadic occurrence were taken into account as possible confounders (SPSS, Chicago, IL).

RESULTS

Median age of all patients included in the study cohort was 30.0 years (range 1.0-75.0). Of the total of 199 participants, 111 were familial cases and 88 concerned sporadic patients. After a median follow-up time of 30.2 years (range=1.33-76.0), 44 carriers of a *RB1* mutation from 31 different families developed a second primary malignancy. **Table 1** shows the number of germline *RB1* mutations according to

type of mutation and lists the number of second primary tumors according to type of mutation. The mutations found in patients who developed a second malignancy were distributed throughout most of the *RB1* gene and did not appear to cluster in one region (**Figure 2**). There was no correlation between the different types of second malignancies diagnosed in these patients and the type of mutation or the region of the gene where the mutation was located. In the group of retinoblastoma survivors who developed a second malignancy, only nonsense and frameshift mutations, certain splice mutations and large rearrangements were observed.




Figure 2. Graphical representation of *RB1* and mutations found among hereditary retinoblastoma subjects diagnosed with a second primary malignancy (n = 44). Exons 1 through 27 are not drawn to scale. Every symbol represents a retinoblastoma subject diagnosed with a second primary malignancy. Black symbols represent sporadic hereditary retinoblastoma subjects. Greyscale coloured symbols represent subjects with familial retinoblastoma. Downward-pointing symbols represent mutations in exons, and upward-pointing symbols represent mutations in introns. The horizontal lines below depict large rearrangements.

Table 1. Number and type of second primary tumor (SPT) by mutation type

Type of <i>RB1</i> mutation ^a	Type of SPT					
	Number of carriers n (%) ^b	Number of cases with SPT n (%) ^b	Sarcoma ^c	Melanoma	Epithelial cancer	Other ^d
Nonsense/Frame-shift mutation	117 (58.8)	31 (26.5)	11	8	10	2
Recurrent nonsense mutation	49 (41.9)	17 (34.7)	7	7	2	1
Low penetrance mutation=exon 1	7 (6)	1 (14.3)			1	
Splice mutation	34 (17.1)	7 (20.6)	2	1	4	0
Low penetrance mutation	11 (32.4)	0				
Large rearrangements	35 (17.6)	6 (17.1)	2	2	2	0
Low penetrance mutation	21 (60)	1 (4.8)	1			
Missense mutation	11 (5.5)	0	0	0	0	0
Low penetrance mutation	11 (100)	0				
Promoter mutation	2 (1)	0	0	0	0	0
Low penetrance mutation	2 (100)	0				
Total	199	44 (22.1)	15	11	16	2

^aSubclassification of the mutation type is shown in italics, percentage as compared to total number of cases with this mutation type

^bPercentage of cases with an SPT as compared to total number of cases with this mutation type

^cIncluding soft tissue sarcoma, osteosarcoma, and chondrosarcoma

^dMalignant tumor not otherwise specified and brain tumor

Table 2 displays the mutations of all patients who developed a second primary malignancy, along with clinical details, listed according to subcategories of germline *RB1* mutation type.

We assessed the risk of second malignancy in relation to type of mutation by multivariable Cox model analysis, adjusted for age and therapy. This showed that subjects carrying one of the recurrent nonsense mutations had a significantly elevated risk of developing second malignancies, compared to subjects carrying other mutations (hazard ratio [HR], 3.53; [95% confidence interval (CI), 1.82-6.84]; $P=.000$). Since 5 members of family 8 developed a second malignancy, we did the same analysis while excluding this family. This did not significantly change the outcome (HR, 3.17; [95% CI, 1.50-6.69]; $P=.002$). Leaving both family 8 and all low penetrance mutations out of the analysis, showed a lower but still statistically significantly increased risk for recurrent nonsense mutations as compared to other mutations (HR, 2.46; [95% CI, 1.14-5.28]; $P=.02$).

Table 3 shows the recurrent nonsense mutations known in the *RB1* gene, and displays which of these mutations are found in our cohort in relation to the number of patients and the number of second primary cancers in these patients. A statistically significantly decreased risk for a second primary malignancy was found in the 52 patients with a low penetrance mutation when compared to other mutations (HR, .19; [95% CI, .05-.81]; $P=.025$). Of all 34 patients carrying a splice mutation, eleven were carrier of a low penetrance mutation according to our definition, i.e. a DER ≤ 1.5 . None of the carriers of a low penetrance splice mutation developed a second primary cancer. Out of 21 carriers of a deletion involving the whole *RB1* gene, just one developed a sarcoma and one out of seven carriers of a mutation in exon 1 developed breast cancer at the age of 59. No second malignancies were observed in carriers of a *RB1* promoter or missense mutation ($n=13$).

Table 2. Descriptions of retinoblastoma patients with a second primary malignancy according to subcategories of *RB1* mutation type

Subject no.* ¹	Family ²	Number of RB patients in the family without second primary malignancy (age at inclusion)	Site	Mutation description ³	Protein	Laterality	First in family	Therapy	Type of second primary malignancy ⁴	Second primary malignancy (age at diagnosis)
Nonsense or frameshift mutation										
1	F22	NA	Exon 1	<i>g.2121dupC</i>	<i>p.Ala22GlyfsX9</i>	BiI	Yes	RT	Epithelial	B (59)
2	F11	1 (37)	Exon 2	<i>g.5446G>T</i>	<i>p.Glu54X</i>	BiI	Yes	Enucl	Epithelial	BI (52)
3	F12	1 (12)	Exon 2	<i>g.5505_5506dup</i>	<i>p.Ala74GlufsX4</i>	BiI	Yes	RT + CH	Other	(44)
4	F16	4 (59,40,14,6)	Exon 6	<i>g.45855delIT</i>	<i>p.Leu199TyrfsX2</i>	BiI	No	RT	Osteosarcoma	(5)
5*	F31	NA	Exon 8	<i>g.59695C>T</i>	<i>p.Arg255X</i>	BiI	Yes	RT	Osteosarcoma	(12)
6	F17	2 (41,29)	Exon 8	<i>g.59702_59703dup</i>	<i>p.Asn258ArgfsX7</i>	UI	Yes	Enucl	Epithelial	Pa (56)
7	F17		Exon8	<i>g.59702_59703dup</i>	<i>p.Asn258ArgfsX7</i>	BiI	No	RT	Epithelial	BI (51)
8	F17		Exon8	<i>g.59702_59703dup</i>	<i>p.Asn258ArgfsX7</i>	BiI	No	RT + CH	Epithelial	L (39)
9*	F4	0	Exon 10	<i>g.64348C>T</i>	<i>p.Arg320X</i>	BiI	Yes	Enucl	Epithelial	C (49)
10*	F4		Exon 10	<i>g.64348C>T</i>	<i>p.Arg320X</i>	BiI	No	RT	Melanoma	(25)
11*	F5	2 (26,11)	Exon 11	<i>g.65386C>T</i>	<i>p.Arg358X</i>	BiI	Yes	RT + CH	Sarcoma	LMS (41)
12	F6	2 (47,41)	Exon 12	<i>g.70261C>T</i>	<i>p.Gln383X</i>	UI	Yes	Enucl	Epithelial	BI (62)
13*	F8	3 (26,11,49)	Exon 14	<i>g.76430C>T</i>	<i>p.Arg445X</i>	BiI	Yes	Enucl	Epithelial	L (65)
14*	F8		Exon 14	<i>g.76430C>T</i>	<i>p.Arg445X</i>	BiI	No	Enucl	Melanoma	(23)

Table 2. Continued

Subject no.* ₁	Family ²	Number of RB patients in the family without second primary malignancy (age at inclusion)	Site	Mutation description ³	Protein	Laterality	First in family	Therapy	Type of second primary malignancy ⁴	Second primary malignancy (age at diagnosis)
15*	F8		Exon 14	<i>g.76430C>T</i>	<i>p.Arg445X</i>	Bil	No	RT	Melanoma	(37)
16*	F8		Exon 14	<i>g.76430C>T</i>	<i>p.Arg445X</i>	Bil	No	RT	Melanoma	(45)
17*	F8		Exon 14	<i>g.76430C>T</i>	<i>p.Arg445X</i>	Bil	No	RT	Melanoma	(21)
18*	F23	NA	Exon 15	<i>g.76898C>T</i>	<i>p.Arg467X</i>	Bil	No	RT + CH	Melanoma	(28)
19*	F9	1 (54)	Exon 17	<i>g.78238C>T</i>	<i>p.Arg552X</i>	Bil	Yes	RT	Sarcoma	LMS (33)
20*	F9		Exon 17	<i>g.78238C>T</i>	<i>p.Arg552X</i>	Bil	No	RT + CH	Osteosarcoma	(12)
21*	F10	0	Exon 18	<i>g.150037C>T</i>	<i>p.Arg579X</i>	Ul	Yes	Missing	Other	(59)
22*	F10		Exon 18	<i>g.150037C>T</i>	<i>p.Arg579X</i>	Bil	No	RT	Melanoma	(35)
23*	F24	NA	Exon 18	<i>g.150037C>T</i>	<i>p.Arg579X</i>	Bil	No	RT	Osteosarcoma	(10)
24	F25	NA	Exon 19	<i>g.153211T>A</i>	<i>p.Tyr606X</i>	Bil	No	RT	Melanoma	(35)
25	F13	3 (42,70,51)	Exon 20	<i>g.156787_156788 delGC</i>	<i>p.Thr687ProfsX4</i>	Bil	No	Enucl	Epithelial	O (72)
26	F13		Exon 20	<i>g.156787_156788 delGC</i>	<i>p.Thr687ProfsX4</i>	Ul	No	Enucl	Sarcoma	Unknown location (50)
27	F14	2 (53,50)	Exon 21	<i>g.160832G>T</i>	<i>p.Glu737X</i>	Ul	Yes	Enucl	Epithelial	O (58)
28	F14		Exon 21	<i>g.160832G>T</i>	<i>p.Glu737X</i>	Bil	No	RT	Osteosarcoma	(20)
29*	F27	NA	Exon 23	<i>g.162237C>T</i>	<i>p.Arg787X</i>	Bil	No	RT + CH	Sarcoma	LMS (32)
30*	F28	NA	Exon 23	<i>g.162237C>T</i>	<i>p.Arg787X</i>	Bil	No	RT	Sarcoma	Lipo (12)
31	F26	NA	Exon 23	<i>g.162266delC</i>	<i>p.Leu797TyrfsX13</i>	Bil	No	RT + CH	Osteosarcoma	(26)

Table 2. Continued

Subject no. ^{*,1}	Family ²	Number of RB patients in the family without second primary malignancy (age at inclusion)	Site	Mutation description ³	Protein	Laterality	First in family	Therapy	Type of second primary malignancy ⁴	Second primary malignancy (age at diagnosis)
Splice mutation										
32	F7	1 (29)	Exon 13	g.73869G>A	p.Gln444Gln exon 13 skipped	Bil	Yes	RT	Epithelial	L (48)
33	F20	NA	Intron 6	g.45867G>A/ IVS6+1G>A	n.i.	Bil	No	RT	Epithelial	B (57)
34	F21	NA	Intron 14	g.76491G>T/ IVS14+5G>T	n.i.	Bil	No	RT	Melanoma	(51)
35	F19	NA	Intron 17	g.149997G>A/ IVS17-1G>A	n.i.	Bil	No	RT	Sarcoma	Rhab (8)
36	F1	0	Intron 20	g.160729G>C/ IVS20-1G>C	n.i.	Bil	Yes	RT	Epithelial	B1 (62)
37	F1		Intron 20	g.160729G>C/ IVS20-1G>C	n.i.	Bil	No	RT	Sarcoma	Hfs (31)
38	F1		Intron 20	g.160729G>C/ IVS20-1G>C	n.i.	Bil	No	RT	Epithelial	B (43)
Large rearrangements										
39	F15	1 (5)	Duplication exon 3	g.39446- ?_39561+?dup	n.i.	Ul	Yes	Enucl	Melanoma	(31)
40	F3	2 (45,9)	Deletion RB1	c.-138-?_27841+?del	-	Bil	No	RT	Sarcoma	Rhab (11)

Table 2. Continued

Subject no.*,1	Family ²	Number of RB patients in the family without second primary malignancy (age at inclusion)	Site	Mutation description ³	Protein	Laterality	First in family	Therapy	Type of second primary malignancy ⁴	Second primary malignancy (age at diagnosis)
41	F2	2 (17,15)	Deletion exon 10/11	Del exon 10 or 10 and 11	n.i.	Bil	Yes	RT	Melanoma	(31)
42	F29	NA	Deletion exon 3-17	g.39446-?_78279+7del	n.i.	Bil	No	RT	Sarcoma	LMS (32)
43	F30	NA	Deletion exon 6-17	g.45799-?_78279+7del	n.i.	Bil	No	RT	Epithelial	Seb (38)
44	F18	2 (68,42)	Deletion exon 9-27	g.61730-?_g177078+7del	n.i.	UI	Yes	Enucl	Epithelial	C (57)

Abbreviations: RB, retinoblastoma; NA, not applicable, i.e. sporadic retinoblastoma patient; n.i., not investigated; Bil, bilateral disease; UI, unilateral disease; RT, radiation therapy; CH, chemotherapy; Enucl, enucleation.

B, breast cancer; Bl, bladder cancer; Pa, pancreatic cancer; L, lung cancer; C, colon cancer; LMS, leiomyosarcoma; O, ovarian cancer; Lipo, liposarcoma; Rhab, rhabdomyosarcoma; His, histiocytoma; Seb, sebaceous adenocarcinoma.

* Subject numbers marked with an asterisk (*) are carriers of a recurrent stop mutation.

† Died of other cause than second primary cancer.

¹ Italicized cases represent family members also affected with a second primary malignancy.

² Family numbers 1-18: have more affected family members with RB; family numbers 19-31: sporadic retinoblastoma patients.

³ Reference sequence GenBank #L11910.⁴ Underlined types of second primary malignancies represent tumours diagnosed inside the field of radiation; defined as tumor in the eye lids, orbits, periorcular sinuses, temporal bones, or skin overlying the temporal bone region.

DISCUSSION

Our study is the first to examine the association between specific *RB1* germline mutations and the risk of second primary malignancies in a nationwide well-documented cohort. Adjusting for age and therapy, we found a higher second primary malignancy risk for retinoblastoma subjects carrying a recurrent nonsense mutation, and a lower risk for carriers of a low penetrance mutation. We first compared the risk of second primary malignancies for carriers of recurrent nonsense mutations to all other mutations in the *RB1* gene. Because one family (F8) with many family members affected by a second malignancy may have influenced the outcome too much, we excluded this family from the analysis. This still showed a statistically significantly increased risk. It is remarkable that 4 members of this family developed a melanoma. This could be due to common genetic background, though this phenomenon does not hold true for other families. As far as we are aware, the family does not display any signs of dysplastic nevus syndrome. We further hypothesized that the increased risk of second malignancies for recurrent mutations compared to all other mutations may have been caused by a substantial contribution of the lower risk of second malignancies for low penetrance mutations, included in the comparison. Therefore we also left the low penetrance mutations out of the analysis. This still showed a significantly increased risk of second primary cancers for recurrent nonsense mutations. Three recurrent nonsense mutations did not demonstrate any second malignancies in our study cohort (**Table 3**). Whether these mutations do not lead to a higher second malignancy risk, needs to be clarified in future studies. What could be the cause of the higher risk for second malignancies in carriers of recurrent nonsense mutations? Nonsense and frameshift mutations are associated with bilateral retinoblastoma and high (>90%) penetrance¹⁹, irrespective of the location of the premature stop mutation. This is attributed to nonsense mediated mRNA decay (NMD): a mechanism of mRNA surveillance that prevents the expression of truncated proteins, by breaking down mutant mRNA containing a premature termination codon.²⁴ NMD can be beneficial, eliminating truncated transcripts that could lead to proteins with possible dominant negative or gain-of function effects, but may also be harmful when preventing translation of truncated protein that would otherwise still be partly functional.^{25,26} Studies have also shown that NMD efficacy may vary between tissues.^{27,28} An explanation for our findings could be that specific nonsense mutations escape NMD in certain tissues, which will result in the expression of a truncated protein. This truncated protein may either have residual activity resulting in a milder effect or may have a dominant negative effect, as has been described for other tumor-suppressor genes.^{25,29} The higher risk in recurrent nonsense mutation carriers may then be explained by a differential effect of NMD between these mutations and other truncating mutations. How such a differential effect would specifically exist between the two different types of truncating mutations remains to be

determined. Alternatively, while the recurrent nonsense mutations result in the loss of a 5-methylcytosine within the gene, this may affect the chromatin structure and/or expression of the gene and thereby increasing the chance of transformation. However, in spite of a clearly elevated risk for recurrent nonsense mutations in our cohort, we cannot rule out that the elevated risk is a chance finding.

Table 3. List of the 11 recurrent *RB1* nonsense mutations, the number of patients in our cohort carrying the mutation and the number of patients with this mutation who developed a second primary tumor (SPT)

Exon	Mutation	Protein	Number of patients in our cohort	Number of patients who developed a SPT
8	g.59683C>T	Arg.251X	2	0
8	g.59695C>T	Arg.255X	1	1
10	g.64348C>T	Arg.320X	5	2
11	g.65386C>T	Arg.358X	3	1
14	g.76430C>T	Arg.445X	11	5
14	g.76460C>T	Arg.455X	8	0
15	g.76898C>T	Arg.467X	1	1
17	g.78238C>T	Arg.552X	3	2
17	g.78250C>T	Arg.556X	5	0
18	g.150037C>T	Arg.579X	7	3
23	g.162237C>T	Arg.787X	3	2
Total			49	17

Genotype-phenotype correlations of *RB1* mutations have been described for specific types of mutations: certain splice mutations, promoter, exon 1 and missense mutations lead to reduced expressivity (unilateral retinoblastoma) and incomplete penetrance (unaffected carriers) of retinoblastoma.^{18,19,21,23,30} This is attributed to a reduction in the amount of normal protein that is produced or to residual activity of mutant protein.^{18,19} Reduced expressivity and incomplete penetrance have also been described for deletions of the complete *RB1* gene. This is thought to be caused by co-deletion of adjacent unknown genes leading to a greater chance of apoptosis, when the wildtype allele is lost as the second hit in the tumor.²⁰ In line with reduced expressivity in the retina, we demonstrated a lower risk of second primary cancers for carriers of these low penetrance mutations. In our study group none of the carriers (n=24) of a missense, promoter or low penetrance splice *RB1* mutation developed a second malignancy, and just one out of 7 carriers with a mutation in exon 1 developed another cancer, i.e.

breast cancer at age 59. The latter may also be attributed to the high population risk of breast cancer. A lower risk for second primary malignancies was noted for carriers of a complete deletion of the *RB1* gene as well: only one out of 21 carriers developed a second malignancy (i.e. rhabdomyosarcoma at age 11).

Strength of our study is the unique large data set of genotyped survivors of retinoblastoma from a population based cohort. Very few studies on genotype-phenotype relations of *RB1* mutations mention second primary cancers. Two studies on mutations in de *RB1* gene stated that they could not detect an association between the mutation and manifestation of a second primary cancer or tumor type.^{21,31} These studies included only a few patients with second primary cancers, however. Some limitations of our study should be considered. First, our mutation detection rate was 90% for all familial and bilateral cases at the time of the study. Leaving 10% of hereditary patients out of the analysis may have influenced the outcome. Some mutations may have been missed, because they are present in a mosaic state; others because they might be located deep in an intron. Second, some types of mutations (e.g. promoter mutations) are relatively rare and therefore some mutation type subgroups were small, making it difficult to draw firm conclusions. A third limitation is that *RB1* mutation detection has become available just 20 years ago. Although many patients from the Dutch retinoblastoma cohort have been genotyped in the past years, this study still comprises a relatively young group of retinoblastoma subjects; almost 50% of all known patients with hereditary retinoblastoma could be included. Quite a few patients diagnosed with (osteo)sarcoma had already died before it was possible to perform DNA analysis. The exclusion of these (osteo)sarcomas may have limited our ability to detect possible associations between specific mutations and sarcoma risk.

In conclusion, our results suggest a genotype-phenotype correlation for second primary malignancies of retinoblastoma survivors and may impact on long-term surveillance protocols of patients with hereditary retinoblastoma, if confirmed by future studies.

ACKNOWLEDGEMENTS

The authors would like to thank all patients who participated and Q. Waisfisiz, PhD, Department of Clinical Genetics, VU University Medical Center, professor dr. H.P.J. te Riele, Netherlands Cancer Institute and professor dr. D.R. Lohmann, Institut für Humangenetik, Universitätsklinikum Essen, Germany for their valuable contributions to the manuscript.

FUNDING

The study was supported by grand VU 2004-3046 from the Dutch Cancer Society.

REFERENCES

- 1 Mahoney MC, Burnett WS, Majerovics A, Tanenbaum H: The epidemiology of ophthalmic malignancies in New York State. *Ophthalmology* 1990; 97: 1143-1147.
- 2 Friend SH, Bernards R, Rogelj S et al.: A human DNA segment with properties of the gene that predisposes to retinoblastoma and osteosarcoma. *Nature* 1986; 323: 643-646.
- 3 Knudson AG, Jr.: Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci U S A* 1971; 68: 820-823.
- 4 Burkhart DL, Sage J: Cellular mechanisms of tumour suppression by the retinoblastoma gene. *Nat Rev Cancer* 2008; 8: 671-682.
- 5 Leiderman YI, Kiss S, Mukai S: Molecular genetics of *RB1*--the retinoblastoma gene. *Semin Ophthalmol* 2007; 22: 247-254.
- 6 Moll AC, Imhof SM, Bouter LM et al.: Second primary tumors in patients with hereditary retinoblastoma: a register-based follow-up study, 1945-1994. *Int J Cancer* 1996; 67: 515-519.
- 7 Fletcher O, Easton D, Anderson K, Gilham C, Jay M, Peto J: Lifetime risks of common cancers among retinoblastoma survivors. *J Natl Cancer Inst* 2004; 96: 357-363.
- 8 Kleinerman RA, Tucker MA, Tarone RE et al.: Risk of New Cancers After Radiotherapy in Long-Term Survivors of Retinoblastoma: An Extended Follow-Up. *J Clin Oncol* 2005; 23: 2272-2279.
- 9 Marees T, Moll AC, Imhof SM, de Boer MR, Ringens PJ, van Leeuwen FE: Risk of second malignancies in survivors of retinoblastoma: more than 40 years of follow-up. *J Natl Cancer Inst* 2008; 100: 1771-1779.
- 10 Woo KI, Harbour JW: Review of 676 second primary tumors in patients with retinoblastoma: association between age at onset and tumor type. *Arch Ophthalmol* 2010; 128: 865-870.
- 11 Eng C, Li FP, Abramson DH et al.: Mortality from second tumors among long-term survivors of retinoblastoma. *J Natl Cancer Inst* 1993; 85: 1121-1128.
- 12 Yu CL, Tucker MA, Abramson DH et al.: Cause-specific mortality in long-term survivors of retinoblastoma. *J Natl Cancer Inst* 2009; 101: 581-591.
- 13 Marees T, van Leeuwen FE, de Boer MR, Imhof SM, Ringens PJ, Moll AC: Cancer mortality in long-term survivors of retinoblastoma. *Eur J Cancer* 2009; 45: 3245-3253.
- 14 Mancini D, Singh S, Ainsworth P, Rodenhiser D: Constitutively methylated CpG dinucleotides as mutation hot spots in the retinoblastoma gene (*RB1*). *Am J Hum Genet* 1997; 61: 80-87.
- 15 Lohmann DR: *RB1* gene mutations in retinoblastoma. *Hum Mutat* 1999; 14: 283-288.
- 16 Valverde JR, Alonso J, Palacios I, Pestana A: *RB1* gene mutation up-date, a meta-analysis based on 932 reported mutations available in a searchable database. *BMC Genet* 2005; 6: 53
- 17 Duncan BK, Miller JH: Mutagenic deamination of cytosine residues in DNA. *Nature* 1980; 287: 560-561.
- 18 Harbour JW: Molecular basis of low-penetrance retinoblastoma. *Arch Ophthalmol* 2001; 119: 1699-1704.
- 19 Lohmann DR, Gallie BL: Retinoblastoma: revisiting the model prototype of inherited cancer. *Am J Med Genet C Semin Med Genet* 2004; 129C: 23-28.
- 20 Albrecht P, Ansperger-Rescher B, Schuler A, Zeschnigk M, Gallie B, Lohmann DR: Spectrum of gross deletions and insertions in the *RB1* gene in patients with retinoblastoma and association with phenotypic expression. *Hum Mutat* 2005; 26: 437-445.

- 21 Taylor M, Dehainault C, Desjardins L et al.: Genotype-phenotype correlations in hereditary familial retinoblastoma. *Hum Mutat* 2007; 28: 284-293.
- 22 Lohmann DR, Brandt B, Hopping W, Passarge E, Horsthemke B: Distinct *RB1* gene mutations with low penetrance in hereditary retinoblastoma. *Hum Genet* 1994; 94: 349-354.
- 23 Scheffer H, van der Vlies P, Burton M et al.: Two novel germline mutations of the retinoblastoma gene (*RB1*) that show incomplete penetrance, one splice site and one missense. *J Med Genet* 2000; 37: E6
- 24 Chang YF, Imam JS, Wilkinson MF: The nonsense-mediated decay RNA surveillance pathway. *Annu Rev Biochem* 2007; 76: 51-74.
- 25 Holbrook JA, Neu-Yilik G, Hentze MW, Kulozik AE: Nonsense-mediated decay approaches the clinic. *Nat Genet* 2004; 36: 801-808.
- 26 Bhuvanagiri M, Schlitter AM, Hentze MW, Kulozik AE: NMD: RNA biology meets human genetic medicine. *Biochem J* 2010; 430: 365-377.
- 27 Resta N, Susca FC, Di Giacomo MC et al.: A homozygous frameshift mutation in the *ESCO2* gene: evidence of intertissue and interindividual variation in Nmd efficiency. *J Cell Physiol* 2006; 209: 67-73.
- 28 Linde L, Boelz S, Neu-Yilik G, Kulozik AE, Kerem B: The efficiency of nonsense-mediated mRNA decay is an inherent character and varies among different cells. *Eur J Hum Genet* 2007; 15: 1156-1162.
- 29 Karam R, Carvalho J, Bruno I et al.: The NMD mRNA surveillance pathway downregulates aberrant E-cadherin transcripts in gastric cancer cells and in *CDH1* mutation carriers. *Oncogene* 2008; 27: 4255-4260.
- 30 Zhang K, Nowak I, Rushlow D, Gallie BL, Lohmann DR: Patterns of missplicing caused by *RB1* gene mutations in patients with retinoblastoma and association with phenotypic expression. *Hum Mutat* 2008; 29: 475-484.
- 31 Lohmann DR, Brandt B, Hopping W, Passarge E, Horsthemke B: The spectrum of *RB1* germ-line mutations in hereditary retinoblastoma. *Am J Hum Genet* 1996; 58: 940-949.

