

VU Research Portal

Patients' expectations:

Haanstra, T.M.

2015

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Haanstra, T. M. (2015). *Patients' expectations: determinants, mechanisms and impact on clinical outcomes*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

CHAPTER 3

Can optimism, pessimism, hope, treatment credibility and treatment expectancy be distinguished in patients undergoing Total Hip and Total Knee Arthroplasty?

T.M. Haanstra, C. Tilbury, S.J. Kamper, R.L. Tordoir,
T.P.M. Vliet Vlieland, R.G.H.H. Nelissen, P. Cuijpers,
H.C.W. de Vet, J. Dekker, D.L. Knol, R.W.J.G. Ostelo

Abstract

Objectives

The constructs optimism, pessimism, hope, treatment credibility and treatment expectancy are associated with outcomes of medical treatment. While these constructs are grounded in different theoretical models, they nonetheless show some conceptual overlap. The purpose of this study was to examine whether currently available measurement instruments for these constructs capture the conceptual differences between these constructs within a treatment setting.

Methods

Patients undergoing Total Hip and Total Knee Arthroplasty (THA and TKA) (Total N= 361; 182 THA; 179 TKA), completed the Life Orientation Test-Revised for optimism and pessimism, the Hope Scale, the Credibility Expectancy Questionnaire for treatment credibility and treatment expectancy. Confirmatory factor analysis was used to examine whether the instruments measure distinct constructs. Four theory-driven models with one, two, four and five latent factors were evaluated using multiple fit indices and $\Delta\chi^2$ tests, followed by some posthoc models.

Results

The results of the theory driven confirmatory factor analysis showed that a five factor model in which all constructs loaded on separate factors yielded the most optimal and satisfactory fit. Posthoc, a bifactor model in which (besides the 5 separate factors) a general factor is hypothesized accounting for the commonality of the items showed a significantly better fit than the five factor model. All specific factors, except for the hope factor, showed to explain a substantial amount of variance beyond the general factor.

Conclusion

Based on our primary analyses we conclude that optimism, pessimism, hope, treatment credibility and treatment expectancy are distinguishable in THA and TKA patients. Postdoc, we determined that all constructs, except hope, showed substantial specific variance, while also sharing some general variance.

Introduction

Growing evidence supports the importance of psychological constructs in predicting outcomes of medical treatment including surgery¹⁻⁷. Usually epidemiological studies investigating the relationship between psychological factors and outcome of treatment restrict their assessment to one or two psychological questionnaires. However, in order to disentangle their unique contribution to outcome, the roles of separate constructs need to be explored simultaneously¹⁸. It is therefore necessary that the instruments that aim to measure these constructs are able to discriminate between them.

Much attention has been given to the future oriented constructs 'optimism', and 'hope'^{1,9}. Both these constructs reflect expectancies about one's future. More specifically, optimism has been defined as "generalized positive outcome expectancies"¹⁰ and hope as "a cognitive set that is based on a reciprocally derived sense of successful agency (goal directed determination) and pathways (planning of ways to meet goals)"¹¹. Theory suggests that a hopeful person is more explicitly concerned with self-initiated actions that will enable him to achieve a favourable future while an optimistic person believes that somehow (through either internal or external factors) his future will be successful¹². Substantial empirical work investigating optimism and hope has been done within mental health settings. But evidence suggests these constructs may also be related to outcomes in medical treatments like surgery. For example: optimism explains close to 10% of the variance in post-surgical pain after total hip arthroplasty (THA) and total knee arthroplasty (TKA)¹³. Besides these general future oriented constructs interest in treatment specific psychological constructs like 'treatment expectancy' and 'treatment credibility' has also grown¹⁴. Treatment expectancy is defined as "improvements that clients believe will be achieved" and treatment credibility as "how believable, convincing and logical the treatment is"¹⁵. Conceptually, expectations for a given treatment may develop (at least partly) from how credible the treatment seems. Both these constructs may be related to treatment outcomes. For example it was found that expectancies about treatment outcome help predict return to work outcomes¹⁶.

While the abovementioned psychological constructs are grounded in different theoretical models, some studies have hypothesized that there is some conceptual overlap between them^{1,10-12,17-21}. Optimism, hope, treatment credibility and treatment expectancy have for instance all been conceptualized as an anticipatory state and beliefs about the future^{11,12,22,23}. Others have emphasized the conceptual differences between the constructs. Some suggest hope is an emotional state, while optimism is a cognitive state²⁴. Treatment credibility also has been defined as a cognitive concept, whereas treatment expectancy as a more affective or emotional concept, similar to hope¹⁵. Furthermore, treatment credibility and treatment expectancy are conceptualized to be situational (i.e. treatment specific), in contrast to hope and optimism which are dispositional^{15,25}. Multiple studies have empirically explored the distinction between the constructs optimism (and pessimism) and hope. A recent meta-analysis concluded that these

constructs are positively associated but not redundant ($\rho < 0.8$) and that hope and optimism have differential relationships with outcomes like well-being or personality⁹. Treatment expectancy and treatment credibility however have not been included in studies examining the distinctiveness of future oriented constructs, yet. It may be that in medical situations like upcoming elective surgery patients answer items belonging to the hope and optimism questionnaires more in a situational way, referring to their treatment or illness. Consequently in medical treatments, and more specifically in the invasive treatments like surgery, optimism and hope could possibly show similarities to treatment credibility and treatment expectancy.

This study aims to examine whether the instruments for optimism, pessimism, hope treatment credibility and treatment expectancy measure distinct constructs in a population of patients scheduled for THA or TKA.

Materials and Methods

Participants and procedures

This study was part of a larger prospective cohort study on the outcomes of THA and TKA. It included consecutive patients undergoing a primary THA or TKA because of osteoarthritis in the Rijnland Hospital in Leiderdorp, the Netherlands between October 2010 and September 2012. Assessments were done pre-operatively and 12 months after surgery. Between July 2011 and September 2012 a subgroup of participants received additional questionnaires concerning optimism, hope and expectancies pre-operatively. For the present analysis pre-operative data of this subgroup were used. The larger study, as well as the extension for the subgroup was approved by the Medical Ethical Committee of the Rijnland General Hospital, Leiderdorp, the Netherlands (registration number 10/07). All participants gave written informed consent.

Measurement

One day prior to surgery all participants completed a questionnaire including sociodemographic, disease characteristics, Quality of Life and the Life Orientation Test-Revised (LOT-R), the Hope Scale (HS) and the Credibility Expectancy Questionnaire (CEQ). Demographic characteristics included: age (years), sex and education level. Disease characteristics pain and functioning were measured using the Pain and ADL subscales of the HOOS (for THA patients)²⁶ and KOOS (for TKA patients)²⁷ questionnaires. Quality of Life was measured using the SF-36 questionnaire²⁸ from which mental component scores (MCS) and physical component scores (PCS) were derived.

Optimism (and Pessimism)

The Life Orientation Test- Revised²⁰ is a 10 item self-reported questionnaire that aims to measure optimism. The questionnaire consists of 3 positively formulated items (e.g. I'm always optimistic about my future), 3 negatively formulated items (e.g. I rarely count on

good things happening to me) and 4 filler items (e.g. It's easy for me to relax), all items are answered on a 5 point Likert-type scale. The LOT-R was originally developed by Scheier and Carver in 1994²⁰ who called the LOT-R a unidimensional questionnaire in which the observed variables represent one latent factor called trait optimism. However others have argued that the items in the LOT-R represent two latent factors namely optimism and pessimism^{29,30}. Hence, sumscores range from 3-15 when two subscale scores are calculated or from 3-30 when one total score is calculated. The factor structure of the Dutch version of the LOT-R was tested recently³¹. Results showed that the two factor model had the best fit.

Hope

The Hope Scale consists of 12 items of which 4 items measure 'pathways' (e.g. There are lots of ways around the problem), 4 items measure 'agency' (e.g. I meet the goals that I have set for myself) and 4 are filler items (e.g. I worry about my health)¹¹. All items are answered on an 8 point scale with two anchors (1=totally disagree and 8= totally agree). The hope scale is considered to be a unidimensional scale in which agency and pathways together represent the construct 'trait hope'. Analysis of the Dutch version of the HS has shown good model fits for a one factor structure³². Hence, a sumscore which ranges from 8-64 points is derived by summing the 8 items of the HS.

Treatment Credibility and Treatment Expectancy

The Credibility Expectancy Questionnaire is a self-reported six item questionnaire that aims to measure treatment credibility and expectancy for improvement. Originally it was developed by Devilly et al in 2000¹⁵, and validated in several groups. The Dutch translation was done by Smeets et al in 2008¹⁴. In both the original and the Dutch version three items (e.g. at this point, how successfully do you think the surgery will be in reducing your complaints) were found to load on the credibility factor and three items (e.g. at this point, how much do you really feel that the surgery will help to reduce your complaints) on the expectancy factor. Introductory instructions tell the patient that beliefs about how well the therapy might help contain both thoughts and feelings about the therapy and that these may be the same or different¹⁴. Items 1 to 3 and 5 are answered on a scale ranging from 1 (not at all) to 9 (very much), Items 4 and 6 are answered on a 0 (not at all) to 100% (very much). In accordance with Smeets et al scores on item 4 and 6 were transformed with a minimum of 1 and a maximum of 9, and a sum score was formed for each factor ranging from 3 to 27.

Statistical analysis

Confirmatory factor analysis (CFA) for ordered categorical items was used to examine whether the constructs optimism (LOT-R), pessimism (LOT-R), hope (HS), treatment credibility (CEQ credibility subscale) and treatment expectancy (CEQ expectancy subscale), are distinguishable. Because observed variables were all answered on ordinal scales, a matrix based on polychoric correlations was used for CFA. Negatively formulated items of the LOT-R were reverse scored prior to entry into the CFA models. Analyses were conducted using the weighted least squares mean and variance adjusted estimator

(WLSMV) in Mplus 6.12. For the total group of THA and TKA patients four theory-driven models with five, four, two and one latent factors, in which the factors were allowed to correlate within the CFA models, were evaluated using multiple fit indices and compared using $\Delta\chi^2$ test³³. The following fit indices and thresholds were used to denote a satisfactory model: Tucker-Lewis index (TLI) >0.95; comparative fit index (CFI) >0.95 and the root mean square error of approximation (RMSEA) <0.06³⁴. A significant $\Delta\chi^2$ test indicates that the model with the smallest χ^2 (in this case the least stringent model) has a significantly better fit.

Model 1 hypothesised a full differentiation between the five constructs treatment credibility, treatment expectancy, hope, optimism and pessimism. Thus items of each construct was forced to load on a separate factor. Model 2 hypothesised a differentiation between four constructs; the treatment credibility, treatment expectancy and hope items were still forced to load on separate factors, but in this model the optimism and reverse-scored pessimism items were forced to load on one factor as it is controversial whether LOT-R has a uni- or bidimensional structure^{29;30}. Model 3 hypothesised a two factor structure in which the optimism, pessimism and hope (LOT-R and HS) items were forced to load on one factor representing 'generalized positive beliefs about the future' and the treatment credibility and treatment expectancy (CEQ) items were forced to load on one factor representing 'treatment specific beliefs about the future'. This model was tested because of the theoretical plausibility that patients may have general and situational, in this case treatment specific, beliefs about the future. Model 4 hypothesised that treatment credibility, treatment expectancy, hope, optimism and pessimism items load on a single underlying latent factor. This model was tested because when it is assumed that optimism, pessimism, hope, treatment credibility and treatment expectancy are not distinguishable at all, the data should fit this one factor model. If necessary (eg because of ambiguities or high correlations between factors) post-hoc models were tested. Guttman's lambda 2 was used to determine internal consistency reliability of each subscale. A value > 0.7 was considered indicative of good internal consistency reliability^{35;36}. All the analyses above were done using the total sample of THA and TKA patients.

When using the same questionnaire in different groups Factorial Invariance (FI) should be established to show that the items of the questionnaire measure the particular latent construct similarly across groups. In our study both TKA and THA patients were included, and as patients with scheduled for knee arthroplasty may face different difficulties to patients scheduled for hip arthroplasty, the constructs measured in this study may also have different meanings for these groups.

Assessing factorial invariance involves a process of comparing the fit indices for a series of models with increasingly stringent constraints on the relationships between the model parameters. The best-fitting model for the total sample (TKA and THA) identified in the previous analysis was assessed in multigroup CFA's to test for factorial invariance across the TKA and THA group³⁷: Four multigroup CFA models with increasingly stringent model constraints were tested (Table 1).

- A baseline model (configural invariance): in which only the factor structure (number of factors and the pattern of the free and fixed loadings) was constrained to be equal across groups. In this model no equality constraints were imposed on the intercepts and factor loadings.
- A weak FI model: in which the factor structure and factor loadings were constrained to be equal across groups, intercepts were allowed to vary among groups and factor variances were fixed to one in both groups.
- A strong FI model: in which factor structure and loadings and intercepts (thresholds) were constrained to be equal across groups.
- A strict FI model: in which factor structure, factor loadings, intercepts and residual variances were constrained to be equal across groups.

To evaluate the degree of measurement invariance, the recommendations by Cheung and Rensvold were followed, which state that the null hypothesis (invariance) is kept if the incremental change in comparative fit index (CFI) is equal to or smaller than 0.01³⁸. Acceptance of the strong or the strict invariance model was sufficient to assume that the measurement instruments used measure the same constructs in all participants (both THA and TKA).

Missing data were incorporated by using the default option available in Mplus. For WLSMV estimation, Mplus computes polychoric correlations based on pairwise present data between two variables, treating missing data as missing completely at random (MCAR). Under MCAR, the missingness is assumed to occur entirely at random and not depend on observed covariates or on the response itself.

Table 1: levels of factorial invariance

FI Models	Model parameters constrained to be equal across groups
No FI	None
Weak FI	Factor loadings
Strong FI	Factor loadings and item intercepts (thresholds)
Strict FI	Factor loadings and item intercepts (thresholds) and residual item variances/covariances

Results

Characteristics of the sample and internal consistency reliability of the subscales

A total of 745 patients were admitted for THA and 614 patients were admitted for TKA from October 2010 to September 2012. Of these, 420 THA (63.2%) and 395 TKA (65.9%) patients consented to participate and completed the surveys. A subgroup of 184 THA and 191 TKA patients, the ones enrolled in the study between July 2011 and September 2012, received additional questionnaires including the LOT-R, the HS and the CEQ. Of these, 14 had missing responses on all items and were therefore excluded, leaving in total 361 patients for analysis (182 THA, 179 TKA). Characteristics of the subgroup of participants that completed the additional questionnaires and mean scores (sd) on the subscales of these questionnaires are presented in table 2 for THA and TKA groups separately. In both TKA and THA groups the majority of patients were females, the mean age was 67 years for both groups. The mean pain score was 41.9 for THA patients and 39.7 for TKA patients. The mean functioning score (HOOS/KOOS ADL) was 43.8 for THA patients and 45.6 for TKA patients. THA patients on average scored 23.7 on the credibility and 22.5 on the expectancy subscale of the CEQ, TKA patients scored 23.5 and 22.1 on these subscales respectively. HS scores were 43.2 for THA and 41.3 for TKA patients. Optimism was scored 9.9 for THA patients and 10.0 for TKA patients, Pessimism scores were 10.8 for THA and 10.4 for TKA patients. Internal consistency reliability (lambda 2) of each of the subscales was acceptable (Table 2). For 14 patients responses on all items were missing and therefore they were excluded from analysis. All the questionnaire items had missing responses, though in most items < 7% responses were missing. An exception was one of the HS items which had 42% missing responses (item 6) due to a printing error in the questionnaire. The amount of data in the pairwise coefficients ranged between 0.57 and 0.98.

Confirmatory factor analysis


Table 3 shows the model fit indices for the five, four, two and one factor models, as well as $\Delta\chi^2$ tests comparing the five factor model with the four factor model, the four factor model with the two factor model, and the two factor model with the one factor model. The five factor model showed fit indices that satisfied the cut-off criteria determined by Hu and Bentler³⁴, whilst the models with four, two and one latent factor did not satisfy these criteria. Further, $\Delta\chi^2$ tests also indicated that the four factor model fit significantly worse than the five factor model, the two factor model fit significantly worse than the four factor model and the one factor model fit significantly worse than the two factor model. Thus, of the four models tested the five factor model is to be preferred based on all fit indices. The five factor model including the standardized factor loadings and correlations between factors is presented in Figure 1. In this five factor model a very strong correlation was seen between the treatment credibility and treatment expectancy factors, and a strong correlation between the optimism and hope factors.

Table 2: characteristics of the sample included in this study and lambda 2 values for the subscales included in the Confirmatory Factor Analysis.

	Total hip arthroplasty N=182 mean (SD)/ %	Total knee arthroplasty N=179 mean (SD)/%	Total sample N=361 Mean (SD)/%	Guttman's lambda 2 for the total sample
Gender % female	58.1%	71.6%	64.8%	-
Age	67.1 (9.9)	67.6 (9.3)	67.4 (9.6)	-
Education level				-
Low	33.7%	50.3%	42.0%	
Medium	32.0%	31.2%	31.6%	
High	34.3%	18.5%	26.4%	
BMI	27.5 (4.7)	29.5 (4.8)	28.5 (4.9)	-
SF-36 Physical summary scale (range 0-100)	38.8 (7.1)	39.1 (7.7)	38.9 (7.4)	-
Sf-36 Mental summary scale (range 0-100)	51.8 (10.7)	52.2 (11.2)	52.0 (10.9)	-
HOOS/KOOS* pain (range 0 -100)	41.9 (17.8)	39.7 (16.7)	40.6 (17.2)	-
HOOS/KOOS* Activities of Daily Living (range 0-100)	43.8 (17.6)	45.6 (17.6)	44.7 (17.5)	-
CEQ Credibility (range 3-27)	23.7 (3.0)	23.5 (3.1)	23.6 (3.1)	0.714
CEQ Expectancy (range 3-27)	22.5 (3.0)	22.1 (3.0)	22.3 (3.0)	0.779
HS Hope (range 8-64)	43.2 (11.6)	41.3 (11.8)	42.2 (11.7)	0.941
LOT-R Optimism (range 3-15)	9.9 (2.9)	10.0 (2.8)	9.9 (2.8)	0.834
LOT-R Pessimism, reverse scored (range 3-15)	10.8 (2.8)	10.4 (2.7)	10.6 (2.8)	0.709

*The THA patients completed the HOOS questionnaire and the TKA patients completed the KOOS questionnaire HOOS =the Hip injury and Osteoarthritis Outcome Score, KOOS=the Knee injury and Osteoarthritis Outcome Score, CEQ= Credibility Expectancy Questionnaire, HS= Hope Scale, LOT-R= Life Orientation Test Revised.

Figure 1. Path diagram and standardized factor loadings and correlations between factors for the 5 factor model



Factorial Invariance testing (see Table 4) showed that the baseline model was well-fitting and thereby supported configural invariance. For the increasingly stringent models none of the subsequent null-hypotheses of measurement invariance were rejected using the recommendations of Cheung and Rensvold³⁸ which state that the null hypothesis (invariance) is not rejected if the incremental change in CFI is equal to or smaller than 0.01. Thus, strict invariance could be supported.

Table 3: χ^2 difference tests and model fit indices for the models tested for the total group (THA and TKA)

	χ^2 (df)	P -value	$\Delta\chi^2$ (df)	P -value	TLI	CFI	RMSEA
Five factor model	400.9 (160)	<0.01	237.1 (4) §	<0.01	0.981	0.984	0.065
Four factor model	1121.8(164)	<0.01	86.7 (5) #	<0.01	0.927	0.937	0.127
Two factor model	1220.4(169)	<0.01	271.1 (1) §	<0.01	0.922	0.930	0.131
One factor model	3081.9 (170)	<0.01			0.785	0.807	0.218

§ five factor model compared to four factor model # four factor model compared to two factor model § two factor model compared to one factor TLI= Tucker-Lewis Index, CFI= comparative fit index, RMSEA=root mean square error of approximation.

Table 4: model fit indices of the multigroup models for factorial invariance testing across THA and TKA

Factorial Invariance models	χ^2 (df)	P value	TLI	CFI	RMSEA
Baseline model (configural invariance)	640.1 (415)	<0.01	0.986	0.985	0.055
Weak Invariance	653.6 (430)	<0.01	0.987	0.985	0.054
Strong Invariance	672.4 (405)	<0.01	0.983	0.982	0.060
Strict Invariance	718.6 (425)	<0.01	0.983	0.981	0.062

Because of the very strong correlation between treatment expectancy and treatment credibility, and between hope and optimism, three post-hoc analyses were performed. A four factor model with separate factors for hope, optimism and pessimism but in which expectancy and credibility items were forced to load on one factor (Additional figure 1) showed fit indices equal to the five factor model (TLI = 0.98, CFI =0.98, RMSEA = 0.06, χ^2 (df) = 412.5 (164)) The $\Delta\chi^2$ test indicated that the four factor model fit significantly worse than the five factor model ($\Delta\chi^2$ (df) = 16.1 (4) $p<0.01$). A four factor model with separate factors for treatment expectancy, treatment credibility and pessimism but in which

optimism and hope were forced to load on one factor (Additional figure 2), had a slightly worse fit compared to the five factor model (TLI = 0.97, CFI = 0.97, RMSEA = 0.08, χ^2 (df) = 568.7 (164)). The $\Delta\chi^2$ test indicated that the four factor model fit significantly worse than the five factor model ($\Delta\chi^2$ (df) = 102.7 (4) $p < 0.01$).

Further, a bifactor model in which (besides the 5 separate factors) there is a general factor (gf) that is hypothesized to account for the commonality of the items of the 5 separate constructs was tested (Additional figure 3). This bifactor model showed better fit indices than the 5 factor model and the four factor model in which the expectancy and credibility items were forced to load on one factor (TLI = 0.99, CFI = 0.99, RMSEA = 0.06, χ^2 (df) = 304.6 (140)). The $\Delta\chi^2$ test indicated that the bifactor model fit statistically significantly better than the five factor model ($\Delta\chi^2$ (df) = 86.8 (20) $p < 0.01$). For the bifactor model we calculated the proportion of variance accounted for by all factors (ω_k), the proportion of variance accounted for by the general factor (ω_H). For each of the 5 separate factors we calculated the proportion of variance unique from the general factor (ω_{Nk}). For these unique proportions of variance a value of $\omega_{Nk} \geq 0.30$ was regarded as substantial, a value of $0.20 \leq \omega_{Nk} < 0.30$ was regarded as moderate, and a value of $\omega_{Nk} < 0.20$ was regarded low. Table 5 shows that the total amount of variance accounted for by all factors is large (0.94). Also, a substantial amount of variance of all factors (ω_k) is accounted for by variation in the general factor (0.79). This suggests that all items indeed measure a common construct. However the specific factors differ in how much variance they account for unique from the general factor. Treatment expectancy, treatment credibility, optimism and pessimism explain a substantial amount of variance unique from the general factor, however hope does not explain a substantial amount of variance unique from the general factor (Table 5).

Table 5: the proportion of variance explained by all factors (ω_k), the proportion of variance of the total scale explained by the general factor (ω_H) and the proportion of variance of the separate constructs explained by the specific factors (ω_{Nk})

Scale	ω_k	ω_H	ω_{Nk}
Total model (general factor)	0,942	0,787	
Separate constructs			
Treatment credibility	0,838		0,789
Treatment expectancy	0,809		0,782
Hope	0,956		0,015
Optimism	0,857		0,329
Pessimism	0,769		0,769

Discussion

This study examined whether the existing instruments for optimism, pessimism, hope, treatment credibility and treatment expectancy measure distinct psychological constructs in patients undergoing TKA or THA. Because it was not our purpose to develop new instruments or to revise the existing ones, we chose a confirmatory approach (CFA) in all our analyses instead of an exploratory approach (EFA). Moreover, we aimed to use all instruments in the same way as they are currently utilized in research and practice and therefore did not delete items with low factor loadings.

The results of the theory driven CFA showed that a five factor model in which optimism (LOT-R subscale optimism), pessimism (LOT-R subscale pessimism), hope (HS), treatment credibility (CEQ subscale credibility) and treatment expectancy (CEQ subscale expectancy) had the most optimal fit. However, there were two interesting observations. First, a strong correlation ($r = 0.82$) was observed between expectancy and credibility. Therefore a post-hoc analysis was performed in which a four factor model in which expectancy and credibility were forced to load on one factor was tested. Although fit indices were very similar as the five factor model, the $\Delta\chi^2$ test indicated the five factor model was the preferred model. Earlier studies found moderate to very high correlations between expectancy and credibility ($r = 0.56^{14}$, $r = 0.68^{15}$ and $r = 0.83^{15}$), though exploratory as well as confirmatory factor analyses suggest expectancy and credibility are two separate factors^{14,15}. Although the current study cannot provide the definite answer regarding the distinctiveness of the constructs treatment expectancy and treatment credibility in patients undergoing THA or TKA, it seems reasonable that the a-priori, theory driven 5 factor model is preferred. Additionally the five factor model is also supported by the results of those earlier studies on the constructs expectancy and credibility. Nevertheless, future studies should investigate the factorial structure of the CEQ and the distinctiveness of the constructs treatment credibility and treatment expectancy to determine if our findings are replicable or unique to our study sample.

Secondly, a strong correlation was also observed between optimism and hope ($r = 0.79$). A priori we hypothesized that these two factors would be correlated but still distinct because both are defined as general future oriented constructs but do have considerable theoretical differences, though we did not expect such a strong correlation. We therefore also performed a post-hoc analysis to investigate the influence of the strong correlation between hope and optimism on model fit. Results showed that the five factor model had a significantly better fit than the four factor model in which optimism and hope were forced to load on one factor.

In a third post-hoc analysis we investigated the possibility that a model in which, besides the five separate factors, a general factor is included that accounts for shared variance in all items would fit the data. Results showed that this model fit the data better than any other model tested and that there is a strong general factor that accounts for a large

amount of the variance of the total bifactor model. Thus, we suggest that there is a general 'outlook on future' factor that underlies each of the items. Separately, there are four more specific factors namely treatment credibility, treatment expectancy, optimism and pessimism that each account for unique variance above this general factor. Hope however did not account for a substantial amount unique variance above the general factor.

Our findings are consistent with previous factor analyses that have shown hope and optimism to be related but distinct constructs^{9,12}. Our study has extended these findings by demonstrating this in patients undergoing THA and TKA as well as by additionally considering treatment specific future oriented psychological constructs. Our results however slightly differ from Magaletta and Olivers¹² study because we found that the five factor model which included pessimism as a separate factor showed better model fit compared a four factor model in which all items of the LOT-R loaded on one factor. This could be a result of the use of the Dutch translation of the LOT-R which has shown to have a two-dimensional structure³¹ Similar to previous studies we found that optimism and hope are positively related and that both of these are negatively related to pessimism^{9,22}.

Studying the conceptual overlap of psychological constructs seems to gain more importance. A reason for this is that many psychological measures have been developed in the last decades and all of them have individually shown to measure important constructs in medical care but considerable overlap may exist between these constructs (and measures), causing lack of conceptual clarity and confusion among researchers and care providers about which psychological measures to use in studies and daily practice. Therefore, studies investigating conceptual overlap or distinctiveness of these constructs within a medical care setting are important. Recently, two of such studies have been published. De Rooij et al³⁹ investigated the conceptual overlap between cognitive concepts in patients with chronic widespread pain and found that 16 different cognitive subscales could be reduced to three factors namely 1. negative emotional cognitions, 2. active cognitive coping and 3. control belief and expectations of chronicity. Campbell et al studied the conceptual overlap of psychological constructs in low back pain patients and found that 20 subscales of psychological questionnaires could be reduced into four factors namely 1. pain-related distress, 2. cognitive coping, 3. causal beliefs and 4. perceptions of the future. Our study also addresses this issue; however we had a slightly different approach. De Rooij et al and Campbell et al performed factor analyses on a subscale level thereby aiming to identify the most complete though comprehensive set of cognitive (de Rooij) or psychological (Campbell) constructs. We however assessed whether individual item of questionnaires measuring the constructs of interest indeed load on the factors as intended by the developers of the subscale. Our approach therefore, may be seen as the first in a two-step approach in examining overlap between constructs. Once distinctive measurement has been established on an item level, a next step could then be assessing overlap between subscales as de Rooij and Campbell did.

Strengths and Limitations

A strength of this study is that the most widely accepted measurement instruments that aim to measure the included constructs were used, ensuring comparability to future CFA's in other patient groups. A limitation of this study is the limited sample size for CFA's, which made us decide to test our primary hypotheses on the complete sample of THA and TKA patients. A multigroup analysis was only done to test for factorial invariance between the THA and TKA group. Although the FI models converged well and results suggest that strict invariance holds for our data, we do recognize that these analyses may be slightly underpowered and therefore these results should be interpreted with caution. Factorial invariance testing showed similar factorial structures in both groups implying that the constructs measured in this study have the same meaning in both patient groups, thereby suggesting generalizability of our results. Another limitation is the high percentage of missing responses on one of the items of the HS, which was caused by a printing error in the questionnaire. The WLSMV estimator in Mplus statistical software incorporates missing data by pairwise presence, though this is under the assumption that missing data are missing completely at random (MCAR). Because of the reason of the missing data and the fact that that participants with and without missing data (on item 6 of the HS) did not significantly differ on baseline characteristics we believe that the MCAR assumption may hold for our data.

Furthermore, we used $\Delta\chi^2$ -test to compare models, which is controversial as the χ^2 is influenced by sample size; therefore we also included other model fit statistics and based our conclusions on a combination of a-priori defined cut-points. CFA is a test of acceptance of a-priori defined models that are not data driven. A limitation of this method is however, that besides the theoretically plausible models tested in this study, there might be other models that show an even better fit to the data. Our results have furthermore not been validated in an external dataset we therefore encourage future research in TKA and THA and also in other patient groups.


Conclusions and Implications

Based on the results of the current study and previous work we suggest that the constructs treatment expectancy, treatment credibility, hope, optimism and pessimism are distinguishable in THA and TKA patients. Posthoc, a bifactor model in which (besides the 5 separate factors) a general factor is hypothesized accounting for the commonality of the items showed a significantly better fit than the five factor model. All specific factors, except for the hope factor, showed to explain a substantial amount of variance beyond the general factor. Future studies should investigate the factorial structure of the CEQ. Our results may be valuable for the design of clinical studies aiming to measure one or more of these constructs as well as for the evaluation of interventions focussed on altering treatment expectancy which have been initiated lately by several groups⁴¹⁻⁴³. As optimism and hope have been hypothesized to be relatively stable traits, it is necessary for researchers evaluating interventions aimed at altering treatment expectancy, to measure the possibly alterable treatment expectancy distinct from optimism and hope. A next step in making these constructs of benefit for the patient undergoing THA and TKA


is to investigate the relationships between these factors (e.g. Does optimism influence treatment specific expectancies?) and to find out which one or which combination of constructs predicts with more accuracy treatment outcomes after THA and TKA like pain, quality of life and physical well-being the best. In the future clinicians may use these constructs in addition to other tools, in order to identify patients with a high-risk for poor outcome in their decision for the type of intervention, either surgical or conservative.

Can optimism, pessimism, hope, treatment credibility and treatment expectancy be distinguished in patients undergoing Total Hip and Total Knee Arthroplasty?


additional figure 1: Post-hoc model 1, four factor model in which treatment expectancy and treatment credibility load on one factor



Additional figure 2: Post-hoc model 2, four factor model in which the optimism and hope items load on one factor



Additional figure 3: Post-hoc model 3, the bifactor model.



Reference List

- 1 Achat H, Kawachi I, Spiro A, III, DeMolles DA, Sparrow D. Optimism and depression as predictors of physical and mental health functioning: the Normative Aging Study. *Ann Behav Med* 2000;22(2):127-130.
- 2 Dawn AG, Lee PP. Patient expectations for medical and surgical care: a review of the literature and applications to ophthalmology. *Surv Ophthalmol* 2004 Sep;49(5):513-524.
- 3 Flood AB, Lorence DP, Ding J, McPherson K, Black NA. The role of expectations in patients' reports of post-operative outcomes and improvement following therapy. *Med Care* 1993 Nov;31(11):1043-1056.
- 4 Fontaine KR, Cheskin LJ. Optimism and obesity treatment outcomes. *J Clin Psychol* 1999 Jan;55(1):141-143.
- 5 Main CJ, Foster N, Buchbinder R. How important are back pain beliefs and expectations for satisfactory recovery from back pain? *Best Pract Res Clin Rheumatol* 2010 Apr;24(2):205-217.
- 6 Sears SF, Serber ER, Lewis TS, et al. Do positive health expectations and optimism relate to quality-of-life outcomes for the patient with an implantable cardioverter defibrillator? *J Cardiopulm Rehabil* 2004 Sep;24(5):324-331.
- 7 Vissers MM, Bussmann JB, Verhaar JA, Busschbach JJ, Bierma-Zeinstra SM, Reijman M. Psychological factors affecting the outcome of total hip and knee arthroplasty: a systematic review. *Semin Arthritis Rheum* 2012 Feb;41(4):576-588.
- 8 Smith TW, MacKenzie J. Personality and risk of physical illness. *Annu Rev Clin Psychol* 2006;2:435-467.
- 9 Alarcon GM, Bowling NA, Khazon S. Great expectations: A meta-analytic examination of optimism and hope. *Personality and Individual Differences* 2013 Jan.
- 10 Scheier MF, Carver CS. Optimism, coping, and health: assessment and implications of generalized outcome expectancies. *Health Psychol* 1985;4(3):219-247.
- 11 Snyder CR, Harris C, Anderson JR, et al. The will and the ways: Development and validation of an individual-differences measure of hope. *J Pers Soc Psychol* 1991 Apr;60(4):570-585.
- 12 Magaletta PR, Oliver JM. The hope construct, will, and ways: their relations with self-efficacy, optimism, and general well-being. *J Clin Psychol* 1999 May;55(5):539-551.
- 13 Pinto PR, McIntyre T, Ferrero R, Almeida A, raujo-Soares V. Predictors of acute postsurgical pain and anxiety following primary total hip and knee arthroplasty. *J Pain* 2013 May;14(5):502-515.
- 14 Smeets RJ, Beelen S, Goossens ME, Schouten EG, Knottnerus JA, Vlaeyen JW. Treatment expectancy and credibility are associated with the outcome of both physical and cognitive-behavioral treatment in chronic low back pain. *Clin J Pain* 2008 May;24(4):305-315.
- 15 Devilly GJ, Borkovec TD. Psychometric properties of the credibility/expectancy questionnaire. *J Behav Ther Exp Psychiatry* 2000 Jun;31(2):73-86.
- 16 Iles RA, Davidson M, Taylor NF, O'Halloran P. Systematic review of the ability of recovery expectations to predict outcomes in non-chronic non-specific low back pain. *J Occup Rehabil* 2009 Mar;19(1):25-40.
- 17 Snyder CR, Simpson SC, Michael ST, Cheavens J. Optimism and hope constructs: Variants on a positive expectancy theme. In: Chang EC (ed.) *Optimism & pessimism: Implications for theory, research, and practice*. Washington, DC US: American Psychological Association, 2001. 101-125.
- 18 Snyder CR, Michael ST, Cheavens JS. Hope as a psychotherapeutic foundation of common factors, placebos, and expectancies. In: Hubble MA, Duncan BL, Miller SD (eds.) *The heart and soul of change: What works in therapy*. Washington, DC US: American Psychological Association, 1999. 179-200.
- 19 Carver CS, Scheier MF, Segerstrom SC. Optimism. *Clin Psychol Rev* 2010 Nov;30(7):879-889.
- 20 Scheier MF, Carver CS, Bridges MW. Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): a reevaluation of the Life Orientation Test. *J Pers Soc Psychol* 1994 Dec;67(6):1063-1078.
- 21 Chemers MM, Watson CB, May ST. Dispositional affect and leadership effectiveness: A comparison of self-esteem, optimism, and efficacy. *Personality and Social Psychology Bulletin* 2000 Mar;26(3):267-277.

- 22 Bryant FB, Cvengros JA. Distinguishing hope and optimism: Two sides of a coin, or two separate coins? *Journal of Social and Clinical Psychology* 2004 Apr;23(2):273-302.
- 23 Gallagher MW, Lopez SJ. Positive expectancies and mental health: Identifying the unique contributions of hope and optimism. *The Journal of Positive Psychology* 2009 Nov;4(6):548-556.
- 24 Bruininks P, Malle BF. Distinguishing Hope from Optimism and Related Affective States. *Motivation and Emotion* 2005 Dec;29(4):327-355.
- 25 Snyder CR. *Handbook of hope: Theory, measures, and applications*. San Diego, CA US: Academic Press, 2000.
- 26 de Groot I, Reijman M, Terwee CB, et al. Validation of the Dutch version of the Hip disability and Osteoarthritis Outcome Score. *Osteoarthritis Cartilage* 2007 Jan;15(1):104-109.
- 27 de Groot I, Favejee MM, Reijman M, Verhaar JA, Terwee CB. The Dutch version of the Knee Injury and Osteoarthritis Outcome Score: a validation study. *Health Qual Life Outcomes* 2008;6:16.
- 28 Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992 Jun;30(6):473-483.
- 29 Herzberg PY, Glaesmer H, Hoyer J. Separating optimism and pessimism: a robust psychometric analysis of the revised Life Orientation Test (LOT-R). *Psychol Assess* 2006 Dec;18(4):433-438.
- 30 Glaesmer H, Rief W, Martin A, et al. Psychometric properties and population-based norms of the Life Orientation Test Revised (LOT-R). *Br J Health Psychol* 2012 May;17(2):432-445.
- 31 ten Klooster PM, Weekers AM, Eggelmeijer F, et al. Optimisme en/of pessimisme: Factorstructuur van de Nederlandse Life Orientation Test-Revised. *Psychologie & Gezondheid* 2010 May;38(2):89-100.
- 32 Brouwer D, Meijer RR, Weekers AM, Baneke JJ. On the dimensionality of the Dispositional Hope Scale. *Psychol Assess* 2008 Sep;20(3):310-315.
- 33 Muthén L.K., Muthén B.O. *Mplus User's Guide*. Sixth edition ed. Los Angeles, CA: Muthén&Muthén, 2010.
- 34 Hu Lt, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling* 1999;6(1):1-55.
- 35 Sijtsma K. On the Use, the Misuse, and the Very Limited Usefulness of Cronbach's Alpha. *Psychometrika* 2009 Mar;74(1):107-120.
- 36 Guttman L. A basis for analyzing test-retest reliability. *Psychometrika* 1945;10:255-282.
- 37 Millsap RE, Yun-Tein J. Assessing Factorial Invariance in Ordered-Categorical Measures. *Multivariate Behavioral Research* 2004;39(3):479-515.
- 38 Cheung GW, Rensvold RB. Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling* 2002;9(2):233-255.
- 39 de Rooij A, Steultjens MP, Siemonsma PC, et al. Overlap of cognitive concepts in chronic widespread pain: an exploratory study. *BMC Musculoskelet Disord* 2011;12:218.
- 40 Campbell P, Bishop A, Dunn KM, Main CJ, Thomas E, Foster NE. Conceptual overlap of psychological constructs in low back pain. *Pain* 2013 Sep;154(9):1783-1791.
- 41 Mancuso CA, Graziano S, Briskie LM, et al. Randomized trials to modify patients' preoperative expectations of hip and knee arthroplasties. *Clin Orthop Relat Res* 2008 Feb;466(2):424-431.
- 42 Iles RA, Taylor NF, Davidson M, O'Halloran P. An effective coaching intervention for people with low recovery expectations and low back pain: A content analysis. *J Back Musculoskelet Rehabil* 2013 Jul 25.
- 43 Laferton JA, Shedden MM, Auer CJ, Moosdorf R, Rief W. Enhancing the efficacy of heart surgery by optimizing patients' preoperative expectations: study protocol of a randomized controlled trial. *Am Heart J* 2013 Jan;165(1):1-7.