

VU Research Portal

Being On-line Together

Batenburg, A.E.

2015

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Batenburg, A. E. (2015). *Being On-line Together: The Effects of Peer-led Online Support Community Participation on Breast Cancer Patients' Psychological Well-being*. Vrije Universiteit.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

CHAPTER 3

The Effects of Online Peer-led Support Community Participation on Breast Cancer Patients' Psychological Well-being: The Influence of Emotional Approach Coping Differences

This chapter is published as: Batenburg, A., & Das, E. (2014). Emotional approach coping and the effects of online peer-led support group participation among patients with breast cancer: a longitudinal study. *Journal of Medical Internet Research*, 16(11), e256. doi:10.2196/jmir.3517

Abstract

There is a lack of longitudinal studies explaining if and when online peer-led support groups are beneficial. How patients cope with emotions that come along with the cancer diagnosis might influence effectiveness of online participation. In this longitudinal study, we hypothesize that mixed findings in previous research are partly caused by individual differences in coping with emotions, which may moderate the effects of online support group participation on patients' well-being. A total of 133 Dutch patients with breast cancer filled out a baseline (T0) and a follow-up (T1, 6 months later) questionnaire assessing intensity of online participation within the online support community, emotional approach coping, and psychological well-being. A number of 109 patients visited an online support community at both points in time. Findings showed no negative effects of intense support group participation. Participating frequently online was especially helpful for patients who approach their emotions less actively; their emotional well-being increased over time. In contrast, frequent online users who actively approach their emotions experienced no change in well-being, reporting highest levels of well-being overall. For patients who participate less intensively within the support community, coping style seems to outweigh effects of online participation; over time, patients who actively approached emotions experienced an increase in psychological well-being, whereas patients with a low score on emotional approach coping reported no change in depression and emotional well-being.

Introduction

An increasing amount of individuals are diagnosed with breast cancer, and this number is expected to grow the coming years due to early diagnosis, average increase in human life expectancy, and more survivors due to improved treatment¹. A growing number of patients turn to the Internet to search for illness-related information and seek support among peers. Patients with breast cancer are among the most active online seekers compared to other patient groups². Therefore, the question if such online communities are beneficial for patients becomes increasingly important.

The lack of longitudinal studies testing causal effects of online peer-led support group participation is most likely due to its uncontrolled setting. Testing effects is difficult because patients participate anonymously and autonomously—they determine when and how much they want to participate. Participants become members at different time points, frequency and length of visits vary among patients, and some patients are active posters whereas others only read messages from others (i.e., lurkers)³. Perhaps as a result, studies covering peer-led support communities are often descriptive in nature (i.e., interview studies, content analyses, cross-sectional surveys). These studies point to the presence of empowering and therapeutic processes, such as emotional and informational support, recognition, and understanding, but also disempowering processes such as being confronted with negative sides of the disease and complainers⁴⁻⁸. However, effects are rarely tested although these peer-led platforms are easily accessible and common online.

Current outcome studies on online peer support mostly concern online interventions set up by health professionals showing positive effects, such as decreased depression, posttraumatic stress, and perceived stress⁹⁻¹². However, it has been discussed that these online interventions often include several other therapeutic aspects besides peer support (e.g., decision-making tools, skill training, or professional moderation). It is unclear if improvements among patients are specifically caused by the sheer support from peers or by other aspects of the intervention¹³. Only a few studies empirically tested the effects of online peer-led support communities. For example, Lieberman and Goldstein¹⁴ showed a positive change in emotional well-being, depression, and posttraumatic growth. However, other studies reported no significant major effects^{15,16} or negative effects¹⁷. If we assume that null findings may not always get published due to publication bias¹³, the meager evidence for a direct relationship between online peer support and well-being may even be an overrepresentation. This requires more studies testing when and why online peer support is beneficial. Therefore,

the first goal of the present research was to provide a robust test of changes in patients' psychological well-being over time due to patients' intensity of online peer-led support group participation.

We put forward that patients' intensity of online participation is important to assess when we aim to test the effectiveness of online peer support group participation because differences in patients' online behavior might affect health outcomes. For example, a cross-sectional study showed that posters felt that they received more benefits (i.e., emotional support, helping others, and emotional expression) from online communities than lurkers, and only for posters higher levels of emotional support/helper therapy and advice were related to lower levels of anxiety³. In the current longitudinal study, we aim to reveal if patients who are more active within a peer-led support community (in terms of frequency and length of visits and amount of posts) benefit more in terms of health outcomes than patients who are less active online.

Apart from intensity of online participation, patients also differ on aspects outside the online environment, which may affect online behavior and effectiveness of online support group participation as well. For example, recent studies suggest that individual differences in health self-efficacy¹⁸, emotional communication competence¹⁹, and differences in coping with emotions²⁰ might be moderating factors. In the current study, we focus on patients' level of emotional approach coping. Evidence from related fields such as clinical psychology shows substantial differences between patients in coping with cancer-related emotions, which significantly impacts ones' well-being. Because online support platforms are merely used to share illness-related experiences (i.e., patients write about their experiences and related thoughts and emotions), variations in patients' emotional coping styles might be one of the reasons why a direct relation between online participation and well-being is not always found. Studies showed that actively coping with emotions (i.e., recognizing and feeling the meaning of losses)²¹ is often related to better well-being. For example, actively approaching emotions is related to decreased depressive symptoms, distress, increased vigor, improved perceived health status, and fewer medical appointments²²⁻²⁵. Hence, how patients cope with emotions that come along with the cancer diagnosis might influence effectiveness of online participation as well.

Our presumption that the relationship between intensity of online participation and well-being might be moderated by patients' emotional coping style is substantiated by two studies on writing style within online peer-led support communities and a cross-sectional study on emotional approach coping. The first two studies showed that the use of words related to expression of emotions and to learning and understanding was related to changes in well-being^{26,27}. However, whether word use caused changes in well-being or was rather a reflection of well-being remains unclear. Nevertheless, these findings show that patients differ in how they deal with illness and emotions in online environments. Another recent cross-sectional study showed that emotional approach coping (i.e., actively processing and expressing emotions) was especially relevant for patients' who participated frequently within an online peer-led support

group (i.e., for patients who visit the online community and post messages relatively often). Patients who actively dealt with their emotions and frequently participated online reported higher psychological well-being than patients who were frequent users but approached their emotions less actively. No difference in well-being was found for patients who participated less often within the online support group²⁰.

Although these findings underscore the importance of individual differences in emotional approach coping when assessing effects of online support group participation, they provide no conclusive evidence regarding causal patterns. Therefore, to further assess individual emotional coping differences we aimed to extend these cross-sectional findings by tracking patients over an extended period of time. In-line with previous cross-sectional findings, we expected that patients who actively deal with emotions may benefit especially from online support groups. Because online support communities often confront patients with emotionally distressing content from peers²⁸, patients who participate intensely, but cope with emotions less actively, might experience additional stress ending up in a downward spiral. However, one might also argue that over time online support and the recognition patients find in stories from others might be especially helpful for patients with more repressive coping styles because they need it most.

In this research project, we conducted a 2-wave longitudinal study among Dutch patients with breast cancer participating in online peer-led support communities, in which we assessed individual differences in emotional coping style. Specifically, we assessed patients' intensity of online support group participation, emotional approach coping²⁵, and three measures of well-being that are generally associated with breast cancer diagnosis at two points in time: emotional well-being²⁹, depression³⁰, and breast cancer-related concerns³¹. We included potential covariates (i.e., factors often associated with the psychological well-being of patients with breast cancer), such as social support from family and friends^{32,33}, disease status, and received professional psychological help. Based on previous findings regarding emotional approach coping, we propose an interaction effect of emotional coping style and the intensity of patients' online participation. With our longitudinal approach, we aim to reveal the long-term effects of this interaction on patients' psychological well-being.

Method

Participants and procedure

We searched the Internet with Google to identify all online support communities for patients with breast cancer in the Netherlands. Criteria for inclusion were that (1) the website was in the Dutch language, (2) the website was purely de-

signed as 24-hour available message boards or part of the website was designed as a 24-hour available message board, and (3) the discussion board was still active (new messages were posted within the past month). With approval of the website owners, we posted a request to participate in an online survey about Internet use of patients with breast cancer on seven support websites. Participants filled out a baseline (T0) survey in June 2011, including demographics, disease status, the intensity of online support group participation, and psychological well-being (the specific measurements used in the current study are described subsequently). After 6 months, we sent this group of patients a follow-up (T1) questionnaire (December 2011) to reassess their psychological well-being and intensity of online support group participation.

This survey was part of a more extensive research project on online peer support among Dutch patients with breast cancer. The research was carried out in accordance with the American Psychological Association's ethics guidelines³⁴ and complied with European Union legislation³⁵ and Dutch legislation³⁶ on data protection. All procedures were approved by the Department of Communication Science of the VU University Amsterdam.

The introduction page of the survey included the length and purpose of the survey, ensured anonymity, and contact information of the investigator. A sample of 134 Dutch breast cancer survivors filled out both questionnaires. Because this sample included 133 females and only one male, we decided to exclude this male from the data analyses to keep a homogeneous group. Response rates are unknown because we had no access to page views of the participating websites. The online survey tool tracked Internet protocol (IP) addresses to prevent users from retaking the survey. Responses to questions were obligatory, but participants were provided with an "I don't know" or "not applicable" option.

Measurements

Demographics, illness characteristics, and control variables. The T0 baseline questionnaire included questions on patients' age, gender, education level, and working status (i.e., if patients were currently working). We measured current disease status (i.e., if cancer cells were currently detected in the patient's body or not), the number of medical appointments in the past three months regarding breast cancer, and if patients were under treatment at the moment. Because social support from resources other than online peers may affect psychological well-being, we asked if patients received any psychological help from a professional and assessed the social support they received from their friends and family based on the six social well-being items from the Functional Assessment of Cancer Therapy-Breast (FACT-B)²⁹. Items referring to support from friends were adjusted into items that clearly referred to offline friends. Respondents rated on a 5-point scale if the statements applied to them, ranging from "not at all" to "totally" (Cronbach alpha=.753).

Intensity of online support group participation. At both T0 and T1, patients' intensity of online support group participation was assessed by four different questions regarding frequency of visits, average length of visits, contribution (i.e., reading, responding, starting new topics or questions), and frequency of posts in the last four weeks^{20,37}. Frequency of visits was assessed on a 7-point scale; the other items were assessed on a 4-point scale. To merge these different scales into one index, all items were transformed into z scores. The scale was internally consistent at T0 (Cronbach alpha=.799) and T1 (Cronbach alpha=.796). See Appendix 1 for the specific items.

Emotional approach coping. At T0 the emotional approach coping scale²⁵ was used to measure participants' coping style concerning emotions, including four items referring to emotional processing (eg, "I realize that my feelings are justified and important") and four items regarding emotional expression (eg, "I take the time to express my emotions"). Participants rated on a 4-point scale if the statements applied to them. The mean scores (emotional processing: M=2.85, SD=0.63; emotional expression: M=2.85, SD=0.57) were comparable to those of a sample of patients with breast cancer from a previous study²³ (emotional processing: mean 3.00, SD 0.72; emotional expression: M=2.95, SD=0.84) and a group of healthy women²⁵ (emotional processing: M=2.85, SD=0.63; emotional expression: M=2.79, SD=0.73). Because factor analyses showed that all factors loaded between 0.58 and 0.86 on one component and explained 44% of the variance, we created one index for emotional approach coping with all eight items. Ratings were summed and averaged across items (Cronbach alpha=.864).

Psychological well-being. At both T0 and T1, we assessed psychological well-being with 3 different concepts: depression, breast cancer-related concerns, and emotional well-being. Depression was measured with the CES-D10³⁰. The scale consisted of 10 items (e.g., "I felt that everything I did took me quite some effort"). Participants rated on a 4-point scale if the statements applied to them the past week from "less than one day" to "5 to 7 days." The scale was internally consistent in both questionnaires (T0: Cronbach alpha=.740; T1: Cronbach alpha=.815), but was positively skewed at T1. A log transformation was performed for depression T0 and T1 to meet the assumptions of multiple regression analysis³⁸. Breast cancer-related concerns (Profile of Concerns about Breast cancer³¹) were measured with an index of 28 items assessed on a 5-point scale (e.g., "As you think about your illness, how much are you concerned that chemotherapy or radiation therapy will damage your body in some way?"). The index showed consistency (T0: Cronbach alpha=.909; T1: Cronbach alpha=.918). Emotional well-being was measured according to six items from the FACT-B on a 5-point scale (eg, "I'm proud of how I am coping with my illness"²⁹ and showed scale consistency at T0 (Cronbach alpha=.821) and T1 (Cronbach alpha=.876).

Analyses

We conducted repeated measures ANOVAs on the three measures of psychological well-being with intensity of online participation at T0 (-1 SD *vs.* $+1$ SD) and emotional approach coping (-1 SD *vs.* $+1$ SD) at T0 as between-subjects factors and time (T0 *vs.* T1) as a within-subjects factor (see ³⁹ for this specific regression analysis). This estimation procedure allows tests of differences between participants with low *vs.* high levels of online support group participation and participants with low *vs.* high levels of emotional approach coping without conducting a median split, thus retaining all observations in the analysis⁴⁰. We added the intensity of online participation at T1 as covariate in our model to control for changes in online participation over time. In addition, we also included disease status at T1 as control variable into the model. Finally, all other variables that correlated significantly with the independent and dependent variables (i.e., intensity of online participation T0, emotional approach coping T0 or psychological well-being T0 and T1) were entered into the model as covariates (see Results).

Results

Sample characteristics

Table 1 presents the patients' characteristics. Our sample of 133 patients with breast cancer included women with a mean age of 48,44 years (SD 8.60). Most patients had an average to high level of education ($n=86$, 65,15%), and more than half ($n=79$, 59,40%) were (still) actively performing their job. 67,5% reported that no cancer cells were detected at the moment. Slightly more than half of the participants ($n=68$, 51,1%) were under treatment, the other half ($n=65$, 48,9%) were not under treatment or only monitored by a physician at the moment. The average number of breast cancer-related medical appointments in the previous 3 months before participating in the current study was 3.6 appointments. Less than half of the participants received psychological guidance during the period of illness ($n=57$, 43,2%).

Correlations

We first ran a correlation matrix (Tables 2 and 3) to assess associations between independent and dependent variables, and to detect potential covariates. No direct correlations between the intensity of online participation at T0, emotional approach coping at T0, breast cancer-related concerns (at both T0 and T1), emotional well-being (at both T0 and T1), and depression (at both T0 and T1) were found. Because of significant correlations with the independent or dependent

measures, we added age, education level, disease status, offline social support, and psychological help from a professional into our models as covariates.

Table 1. Demographics and Health Characteristics of Participants, n=133

	n	%
Age		
Mean (SD)	48.44 (8.60)	
Minimum	23	
Maximum	67	
Highest education level^a (1 unknown)		
Elementary school	4	3.0
Secondary education		
Low	20	15.2
Middle	12	9.1
High	1	0.8
Tertiary education		
Low ^b	9	6.8
Middle	36	27.3
High	41	31.1
Scientific degree	9	6.8
Working status		
Not working	53	40.2
Working	79	59.8
Disease status (10 unknown)		
No cancer cells	83	67.5
Cancer cells	40	32.5
Under treatment		
Yes	68	51.1
No	65	48.9
Amount of cancer-related medical appointments in the last 3 months		
Mean (SD)	3.61 (6.34)	
Minimum	0	
Maximum	40	
Psychological help during period of illness		
Yes	57	43.2
No	75	56.8

Notes. ^a Levels within the Dutch education system: education is divided over 3 schools for different age groups, which are divided in streams for different educational levels. ^b LBO/LTS (i.e., lowest level of vocational school) existed until 1992.

Table 2. Means, Standard Deviations, and Intercorrelations of Independent variables, Covariates, and Dependent Variables (part 1)

	N	M	SD	1	2	3	4	5	6	7
1 Intensity of OSG participation T0 ^a	125	.00	0.78	-						
2 Intensity of OSG participation T1 ^a	113	-.01	0.79	.70 ^c						
3 Emotional approach coping	132	2.86	0.50	.04	-.03					
4 Age	133	48.44	8.60	-.23 ^c	-.21 ^d	-.14				
5 Education	132	6.38	1.95	-.06	-.07	.06	-.21 ^d			
6 Working status ^b	132	0.60	0.49	-.06	-.02	-.05	-.21 ^d	.37 ^c		
7 Offline social support	133	3.80	0.62	.04	.01	.04	.04	.24 ^c	.14	
8 Psychological help ^b	132	0.43	0.50	-.01	-.01	.12	-.45 ^c	.08	.15	-.23 ^c
9 Disease status T1 ^b	123	0.33	0.47	.03	.11	.03	-.07	-.09	-.13	-.06
10 Under treatment ^b	133	0.51	0.50	.13	.17	.03	-.10	-.02	-.07	.09
11 Medical appointments	132	3.61	6.34	.15	.22 ^d	.12	.02	-.11	-.20 ^d	.09
12 Depression T0 (log)	132	0.25	0.11	-.04	-.04	.01	-.13	-.20 ^d	-.13	-.41 ^c
13 Depression T1 (log)	133	0.22	0.13	.05	.09	-.16	-.12	-.23 ^c	-.16	-.37 ^c
14 Emotional well-being T0	133	3.53	0.80	-.04	-.04	-.08	.12	.18 ^d	.20 ^d	.43 ^c
15 Emotional well-being T1	133	3.70	0.90	-.08	-.12	.11	.13	.18 ^d	.12	.37 ^c
16 BC related concerns T0	133	2.61	0.60	.09	.13	.09	-.18 ^d	-.23 ^c	-.24 ^c	-.46 ^c
17 BC related concerns T1	133	2.52	0.63	-.03	-.01	-.03	-.10	-.26 ^c	-.26 ^c	-.41 ^c

Notes. ^a Standardized into Z scores. ^b Coded 0=no, 1=yes. ^c Correlations significant at the .01 level. ^dCorrelations significant at the .05 level.

Table 3. Means, Standard Deviations, and Intercorrelations of Independent variables, Covariates, and Dependent Variables (part 2)

	8	9	10	11	12	13	14	15	16
9 Disease status T1 ^b	.00								
10 Under treatment ^b	-.12	.28 ^c							
11 Medical appointments	.05	.17	.10						
12 Depression T0 (log)	.34 ^c	-.08	-.12	.04					
13 Depression T1 (log)	.30 ^c	.14	-.01	.09	.54 ^c				
14 Emotional well-being T0	-.29 ^c	-.10	.13	-.11	-.57 ^c	-.48 ^c			
15 Emotional well-being T1	-.27 ^c	-.10	.07	-.05	-.39 ^c	-.62 ^c	.65 ^c		
16 BC related concerns T0	.32 ^c	.01	-.09	.001	.51 ^c	.50 ^c	-.60 ^c	-.42 ^c	
17 BC related concerns T1	.29 ^c	.02	-.01	-.003	.50 ^c	.57 ^c	-.52 ^c	-.55 ^c	.65 ^c

Notes. ^b Coded 0=no, 1=yes. ^c Correlations significant at the .01 level.

Effect testing

From the 133 patients, 125 patients reported they were visiting an online breast cancer support community at T0; eight patients were not. The study participants only had access to the questionnaire by visiting the online support community; therefore, they might have misunderstood our question. Furthermore, because these eight patients claimed they did not visit an online support group, they also did not fill out our questions on intensity of participation and were excluded from data analyses.

At T1, 113 patients claimed they were visiting an online support group. From T0 to T1, 16 participants stopped visiting the online support community and four participants started visiting an online community after the first survey. To measure the change in psychological well-being over time caused by the intensity of online support group participation, we only included the patients that were visiting an online support community at both time points ($n=109$)ⁱ.

Of the 109 participants who visited an online community at T0 and T1, nine participants were not aware of their disease status, one participant did not report her level of education (see Table 1), and the level of depression at T0 was missing for one participant. Because we included these variables as covariates, SPSS excluded these participants from the ANOVA analyses. Table 4 shows the within-subjects ANOVAs, indicating a 3-way interaction of time, intensity of online participation at T0, and emotional coping T0 on depression and emotional well-being. No 3-way interaction effect on breast cancer-related concerns was found.

i To prevent our study from unnecessary data loss, one of the reviewers recommended adding an additional lowest category to the items measuring intensity of online participation T1 to include the forum users who stopped visiting the forum from T0 to T1. However, this new variable violated the normal distribution norms for regression analyses. Therefore, we could not include these support group dropouts for data analyses. Nevertheless, when including this new variable in the analyses, the 3-way interactions were –as the results in table 4– still significant (depression: $F(1, 104)=9.825, p=.002$; emotional well-being: $F(1, 105)=4.232, p=.04$).

Table 4. Repeated Measures ANOVA's on Well-being Measures at T0 and T1

	Depression			Emotional well-being			BC concerns		
	(n=98)			(n=99)			(n=99)		
Between subjects effects^a	F(1,88)	p	η^2_p	F(1,89)	p	η^2_p	F(1,89)	p	η^2_p
Intensity of OSG participation T0	0,429	.514	.005	1,197	.277	.013	0,343	.559	.004
Emotional approach coping	4,62	.034	.050	1,285	.260	.014	0,017	.897	.000
Education	6,032	.016	.064	3,278	.074	.036	4,295	.041	.046
Offline support	7,358	.008	.077	10,684	.002	.107	17,834	.000	.167
Psych help	14,095	.000	.138	8,174	.005	.084	7,033	.009	.073
Age	0,181	.671	.002	0,005	.945	.000	0,814	.370	.009
Online activity T1	0,564	.454	.006	3,428	.067	.037	1,295	.258	.014
Disease status	0,103	.749	.001	0,288	.593	.003	0,47	.495	.005
Intensity of OSG participation T0 x Emotional approach coping	1,738	.191	.019	1,568	.214	.017	0,049	.825	.001
Within subjects effects									
Time	0,773	.382	.009	1,041	.310	.012	2,355	.128	.026
Time x Intensity of OSG participation T0	0,104	.747	.001	0,102	.750	.001	0,272	.603	.003
Time x Emotional approach coping	1,185	.279	.013	1,166	.283	.013	0,612	.436	.007
Time x education	0,343	.560	.004	0,104	.748	.001	0,031	.861	.000
Time x offline support	0,107	.107	.745	0,664	.417	.007	2,552	.114	.028
Time x psych help	0,282	.282	.597	0,068	.795	.001	0,334	.565	.004
Time x age	0,663	.418	.007	0,027	.871	.000	0,409	.524	.005
Time x Intensity of OSG participation T1	1,231	.270	.014	0,711	.401	.008	2,352	.129	.026
Time x disease status	9,21	.003	.095	0,084	.772	.001	0,489	.486	.005
Time x Intensity of OSG participation T0 x Emotional approach coping	8,167	.005	.085	4,232	.043	.045	2,45	.121	.027


Note. ^a for between-subjects effects, the dependent variable is the mean of the score of T0 and T1

Depression

No main effect of time on depression was found; overall depression did not change significantly from T0 to T1 (see Table 4). Concerning covariates, only disease status caused a change in depression. However, education, offline social support, and professional psychological help were related to the average score of depression at T0 and T1. No 2-way interactions of time and intensity on online participation, or time and emotional approach coping on depression were found. However, a 3-way interaction effect of time, the intensity of online participation, and emotional approach coping was found. Pairwise comparisons

revealed that patients who scored low on online participation and high on emotional approach coping showed a significant decrease in depression over time ($M_{diff}=0.060$, $SE=0.027$, $p=.03$ (see Figure 1). In addition, online frequent users who scored high on emotional approach coping did not show a change in depression over time ($p=.80$), but showed lowest overall levels of depression at both points in time (Figure 1).


Figure 1. Depression over Time (T0, T1) as a Function of Intensity of Online Participation at T0 (low vs .high) and Emotional Approach Coping (low vs. high)



Emotional well-being

Table 4 shows no main effect of time on emotional well-being; overall emotional well-being did not change significantly from T0 to T1. No covariates were related to changes in well-being, although offline social support and receiving professional psychological help were related to the average score of emotional well-being. No 2-way interaction of time and intensity of online participation, or time and emotional approach coping were found. However, a significant 3-way interaction of time, online intensity of participation, and emotional approach coping on emotional well-being was found. Pairwise comparisons showed that emotional well-being increased over time for patients who scored low on intensity of participation but high on emotional approach coping ($M_{diff}=0.479$, $SE=0.169$ $p=.006$; see Figure 2). No significant change in well-being was observed among patients who scored low on online participation and low on emotional approach coping ($p=.86$). In addition, for patients who scored high on online participation and low on emotional approach coping, emotional well-being increased over time ($M_{diff}=0.391$, $SE=0.181$ $p=.03$; see Figure 2). For frequent online users who scored high on emotional approach coping, no significant change in well-being was found, but showed highest levels of well-being at both points in time ($p=.22$).

Figure 2. Emotional Well-being over Time (T0, T1) as a Function of Intensity of Online Participation at T0 (low vs. high) and Emotional Approach Coping (low vs. high)



Breast cancer-related concerns

Only education, psychological help from a professional, and offline social support were related to the average score of breast cancer-related concerns. No other main or interaction effects were found.

Discussion

Principal findings

This 2-wave longitudinal study extends previous findings on the effects of online peer support in two ways: by testing changes in well-being as a function of intensity of online support group participation and by examining the role of individual differences in emotional coping styles. Findings showed that participating frequently online was especially helpful for patients who approach their emotions less actively; their emotional well-being increased over time. In contrast, frequent online users who actively approach their emotions experienced no change in well-being, reporting highest levels of well-being overall. For patients who participate less intensively within the support community, coping style seems to outweigh effects of online participation; over time, patients who actively approached emotions experienced an increase in psychological well-being, whereas patients with a low score on emotional approach coping reported no change in depression and emotional well-being.

This study was stimulated by findings from a cross-sectional study that showed that approaching one's emotions was especially relevant for patients who were highly active within an online support community²⁰. Specifically, frequent users with a low score on emotional processing and expression reported significantly lower psychological well-being than equally active patients but with a high score on emotional processing and expression. In contrast, current longitudinal findings showed that especially the frequent users who approach emotions less actively benefit from intense online participation; they experienced an increase in emotional well-being over time. An explanation might be that frequent online participation compensates for the negative effects of not approaching emotions. Patients may benefit from several therapeutic processes within the online support group, such as the support from peers, empowerment by the provision of relevant illness-related information, and recognition in stories from others. These helpful processes might compensate for negative effects of not actively approaching emotions and cause a positive change in patients' well-being.

The current results reconcile previous null findings of support group participation on well-being. We found no main effect of the level of online participation on changes in well-being, we only found interactions of online participation with patients' emotional coping style. This underscores the importance of considering individual differences in dealing with illness when examining health outcomes of online support communities. Although we found no negative effects for highly frequent users, researchers should be careful not to treat an online support community as a "one-cure-fits-all" solution because, apparently, they are not.

The present findings also showed the importance of other factors outside the online environment. We found, for example, correlations between patients' well-being and offline social support and professional psychological help. Patients who felt supported by family and friends reported a higher well-being than patients' feeling less supported, and patients receiving professional help reported a lower well-being. Although such covariates showed no influence on well-being change (except for disease status), we should be careful not to overestimate the effects of online peer support. Since no direct relation between online intensity of participation and well-being was found, such 'offline' factors seem to have a stronger influence on patients' well-being than the level of online participation within an online peer support group. Online peer support might contribute to patients' well-being, but it probably does not compensate for the potential negative effects of other important factors, such as a lack of support from relatives and deteriorated health status.

Regarding practical implications, with caution we could state that patients might be encouraged to participate actively in online peer communities. Although we cannot compare the current participating group with patients who decide not to visit online support communities, for the present sample frequent participation was beneficial to patients' well-being in the long run, regardless of their coping style, and did not cause harm to any patient group. Especially

patients who occasionally visit an online community and approach their emotions less actively should be stimulated to participate more frequently because they seem to benefit most from frequent participation.

Limitations and future research

No interaction effects of our key independent variables on breast cancer-related concerns were found. One explanation might be that breast cancer-related concerns are no accurate measure for well-being. Although the other psychological well-being measures were related to breast cancer-related concerns, concerns might be less appropriate to use as an outcome measure because concerns might intensify online participation. To be more specific, concerns may induce the need for online information and support, and for some patients online peer support, in turn, increases well-being over time. Another explanation is that concerns depend mostly on disease factors rather than online peer support. For example, current treatment, physical well-being, or recurrence might induce concerns. Such factors might intensify concerns at different (shorter) time points; therefore, concerns may fluctuate more intensively over time than more "stable" measures of well-being, such as depression. Future studies should include a more detailed assessment of the disease process patients are in and the number of months they have been participating in an online forum to test this assumption.

In the current study, we measured emotional approach coping only in the baseline questionnaire because coping style is merely viewed as an aspect of ones' personality that is rather stable over time and, therefore, often measured just once in longitudinal studies (e.g.,²³). However, studies with a cognitive-behavioral approach suggest that specific treatment has the potential to teach patients certain coping skills⁴¹. Although peer-led support communities do not include professional guidance, it might still be possible that online conversations stimulate approaching emotions, which has a positive influence on their well-being. From a modeling or skills perspective⁴², frequent participation in online support communities may help patients with more repressive coping styles to learn over time how to approach illness-related emotions. In future studies, we recommend to reassess coping style to test if patients learn to cope with emotions due to online participation.

Examining peer-led online communities is difficult because of its' anonymity and "fluidness" (i.e., patients come and go whenever they like). Because participating patients were anonymous, there is no record of what participants did before T0, no information on dropouts, and the findings rely on self-report (i.e., there is no actual information on what they precisely did online). We have no information on participants' starting date of online participation. Possibly, advantages of online participation might particularly occur for patients who just started to participate online, for example because they are recently diagnosed and especially in need of support. It is possible that changes in well-being might be less pronounced in patients who are part of the community for a longer period. Future research should examine this possibility.

Furthermore, we may have encountered selection bias. Patients who were willing to participate in the current study might also be more involved in online support and, therefore, already behave differently than patients less active online. With respect to the current limitations, in future research it would be interesting to track patients' actual online behavior. Although it is rather difficult when it comes to anonymous peer-led online communities, investigating patients' online writings and connecting these findings to measures of well-being and individual characteristics would provide a better understanding of online processes and its effects. For instance, it would be interesting to see if certain coping skills reflect online writing style.

Although this is one of the rare attempts to investigate longitudinal effects of peer-led support communities, tracking patients for an even longer period with more points of measurement might answer more questions on this topic. For example, online participation may not only influence well-being, patients' level of well-being might also incite online participation. For instance, patients' negative experiences lower well-being, which may stimulate online support seeking and, in turn, online interactions with peers helping patients to cope with the current situation, which in turn increases well-being.

Moreover, in the current study we investigated emotional approach coping. However, other individual differences might be of interest as well. For instance, are patients satisfied with the support they receive online and how does this affect well-being? Or how do patients compare their own situation to the condition of peers and how does this affect them? Future research should address other underlying psychological mechanism that could influence the effects of online support group participation.

Conclusion

To our knowledge, this study is one of the first studies to examine the effects of online peer-led support group participation, taking into account individual differences in emotional coping style. Over time, frequent participation in an online peer support community increased emotional well-being, in particular for patients low on emotional approach coping. Findings suggest that intense support group participation may be especially helpful for patients who approach emotions less actively.

Acknowledgments

This study was financially supported by the Network Institute of the VU University Amsterdam (formerly named CAMeRA Centre of Advanced Media Research Amsterdam). We would like to thank the website owners and patients for their support and participation.

References

1. IKNL: Integraal kankercentrum Nederland. Cijfers over kanker - Prevalentie. 2013. Available at: http://www.cijfersoverkanker.nl/selecties/Dataset_2/img52849a55363ad. Accessed November 14, 2013.
2. Nagler RH, Gray SW, Romantan A, et al. Differences in information seeking among breast, prostate, and colorectal cancer patients: results from a population-based survey. *Patient Educ Couns*. 2010;81 Suppl:S54–62. doi:10.1016/j.pec.2010.09.010.
3. Setoyama Y, Yamazaki Y, Namayama K. Benefits of peer support in online Japanese breast cancer communities: differences between lurkers and posters. *J Med Internet Res*. 2011;13(4):e122. doi:10.2196/jmir.1696.
4. Fernsler JI, Manchester LJ. Evaluation of a computer-based cancer support network. *Cancer Pract*. 1997;5(1):46–51.
5. Mo PKH, Coulson NS. Exploring the communication of social support within virtual communities: a content analysis of messages posted to an online HIV/AIDS support group. *Cyberpsychol Behav*. 2008;11(3):371–4. doi:10.1089/cpb.2007.0118.
6. Rodgers S, Chen Q. Internet community group participation: psychosocial benefits for women with breast cancer. *J Comput Commun*. 2005;10(4):00–00. doi:10.1111/j.1083-6101.2005.tb00268.x.
7. Setoyama Y, Nakayama K, Yamazaki Y. Peer support from online community on the internet among patients with breast cancer in Japan. *Stud Health Technol Inform*. 2009;146:886.
8. Van Uden-Kraan CF, Drossaert CHC, Taal E, Shaw BR, Seydel ER, van de Laar MAJF. Empowering processes and outcomes of participation in online support groups for patients with breast cancer, arthritis, or fibromyalgia. *Qual Health Res*. 2008;18(3):405–17. doi:10.1177/1049732307313429.
9. Hong Y, Peña-Purcell NC, Ory MG. Outcomes of online support and resources for cancer survivors: a systematic literature review. *Patient Educ Couns*. 2012;86(3):288–96. doi:10.1016/j.pec.2011.06.014.

10. Klemm P, Bunnell D, Cullen M, Soneji R, Gibbons P, Holecek A. Online cancer support groups: a review of the research literature. *Comput Inform Nurs*. 2003;21(3):136–42.
11. Lieberman MA, Golant M, Giese-Davis J, et al. Electronic support groups for breast carcinoma: a clinical trial of effectiveness. *Cancer*. 2003;97(4):920–5. doi:10.1002/cncr.11145.
12. Winzelberg AJ, Classen C, Alpers GW, et al. Evaluation of an internet support group for women with primary breast cancer. *Cancer*. 2003;97(5):1164–73. doi:10.1002/cncr.11174.
13. Eysenbach G, Powell J, Englesakis M, Rizo C, Stern A. Health related virtual communities and electronic support groups: systematic review of the effects of online peer to peer interactions. *BMJ*. 2004;328(May):1–6. doi:10.1136/bmj.328.7449.1166.
14. Lieberman MA, Goldstein B a. Self-help on-line: an outcome evaluation of breast cancer bulletin boards. *J Health Psychol*. 2005;10(6):855–62. doi:10.1177/1359105305057319.
15. Høybye MT, Dalton SO, Deltour I, Bidstrup PE, Frederiksen K, Johansen C. Effect of Internet peer-support groups on psychosocial adjustment to cancer: a randomised study. *Br J Cancer*. 2010;102(9):1348–54. doi:10.1038/sj.bjc.6605646.
16. Owen JE, Klapow JC, Roth DL, et al. Randomized pilot of a self-guided internet coping group for women with early-stage breast cancer. *Ann Behav Med*. 2005;30(1):54–64. doi:10.1207/s15324796abm3001_7.
17. Salzer MS, Palmer SC, Kaplan K, et al. A randomized, controlled study of Internet peer-to-peer interactions among women newly diagnosed with breast cancer. *Psychooncology*. 2010;19(4):441–6. doi:10.1002/pon.1586.
18. Namkoong K, Shah D V, Han JY, et al. Expression and reception of treatment information in breast cancer support groups: How health self-efficacy moderates effects on emotional well-being. *Patient Educ Couns*. 2010;81 Suppl:S41–7. doi:10.1016/j.pec.2010.09.009.
19. Yoo W, Namkoong K, Choi M, et al. Giving and receiving emotional support online: communication competence as a moderator of psychosocial benefits for women with breast cancer. *Comput Human Behav*. 2014;30:13–22. doi:10.1016/j.chb.2013.07.024.

20. Batenburg A, Das E. Emotional coping differences among breast cancer patients from an online support group: a cross-sectional study. *J Med Internet Res*. 2014;16(2):e28. doi:10.2196/jmir.2831.
21. Spiegel D. *Living Beyond Limits: New Hope and Help for Facing Life-Threatening Illness*. New York: Random House; 1993.
22. Classen C, Koopman C, Angell K, Spiegel D. Coping styles associated with psychological adjustment to advanced breast cancer. *Health Psychol*. 1996;15(6):434–7. doi:10.1037/0278-6133.15.6.434.
23. Stanton AL, Danoff-Burg S, Cameron CL, et al. Emotionally expressive coping predicts psychological and physical adjustment to breast cancer. *J Consult Clin Psychol*. 2000;68(5):875–82. doi:10.1037//0022-006X.68.5.875.
24. Stanton AL, Danoff-Burg S, Huggins ME. The first year after breast cancer diagnosis: hope and coping strategies as predictors of adjustment. *Psychooncology*. 2002;11(2):93–102. doi:10.1002/pon.574.
25. Stanton AL, Kirk SB, Cameron CL, Danoff-Burg S. Coping through emotional approach: scale construction and validation. *J Pers Soc Psychol*. 2000;78(6):1150–69. doi:10.1037/0278-6133.12.1.16.
26. Lieberman MA, Goldstein BA. Not all negative emotions are equal: the role of emotional expression in online support groups for women with breast cancer. *Psychooncology*. 2006;15(2):160–8. doi:10.1002/pon.932.
27. Lieberman MA. The role of insightful disclosure in outcomes for women in peer-directed breast cancer groups: a replication study. *Psychooncology*. 2007;16(10):961–4. doi:10.1002/pon.1181.
28. Barak A, Boniel-Nissim M, Suler J. Fostering empowerment in online support groups. *Comput Human Behav*. 2008;24(5):1867–1883. doi:10.1016/j.chb.2008.02.004.
29. Brady MJ, Cella DF, Mo F, et al. Reliability and validity of the Functional Assessment of Cancer Therapy-Breast quality-of-life instrument. *J Clin Oncol*. 1997;15(3):974–86.
30. Irwin M, Artin KH, Oxman MN. Screening for depression in the older adult: criterion validity of the 10-item Center for Epidemiological Studies Depression Scale (CES-D). *Arch Intern Med*. 1999;159(15):1701–4.

31. Spencer SM, Lehman JM, Wynings C, et al. Concerns about breast cancer and relations to psychosocial well-being in a multiethnic sample of early-stage patients. *Health Psychol.* 1999;18(2):159–68.
32. Dukes Holland K, Holahan CK. The relation of social support and coping to positive adaptation to breast cancer. *Psychol Health.* 2003;18(1):15–29. doi:10.1080/0887044031000080656.
33. Talley A, Molix L, Schlegel RJ, Bettencourt A. The influence of breast cancer survivors' perceived partner social support and need satisfaction on depressive symptoms: a longitudinal analysis. *Psychol Health.* 2010;25(4):433–49. doi:10.1080/08870440802582682.
34. 2010 Amendments to the 2002 "Ethical principles of psychologists and code of conduct". *Am Psychol.* 65(5):493. doi:10.1037/a0020168.
35. EU legislation on data protection. Available: <http://ec.europa.eu/justice/data-protection/>. Accessed 09 October 2013. Archived by WebCite® at <http://www.webcitation.org/6KEjYUQEE>.
36. Dutch legislation on data protection. Available: http://wetten.overheid.nl/BWBR0011468/geldigheidsdatum_11-07-2013. Archived by WebCite® at <http://www.webcitation.org/6KEjppGbK>.
37. Van Uden-Kraan CF, Drossaert CHC, Taal E, Seydel ER, van de Laar MAFJ. Participation in online patient support groups endorses patients' empowerment. *Patient Educ Couns.* 2009;74(1):61–9. doi:10.1016/j.pec.2008.07.044.
38. Field A. *Discovering Statistics Using SPSS*. London: SAGE Publications Ltd; 2009.
39. Aiken L, West S. *Multiple regression: Testing and interpreting interactions*. London: Sage; 1991.
40. Royston P, Altman DG, Sauerbrei W. Dichotomizing continuous predictors in multiple regression: a bad idea. *Stat Med.* 2006;25:127–141. doi:10.1002/sim.2331.
41. Beutler LE, Moos RH, Lane G. Coping, treatment planning, and treatment outcome: discussion. *J Clin Psychol.* 2003;59(10):1151–67. doi:10.1002/jclp.10216.
42. Bandura A. *Social learning theory*. Englewood Cliffs, NJ: Prentice-Hall; 1977. doi:10.1037/016333.

Appendix 1. Scale Construction - Intensity of Online Support Group Participation

The next questions refer to how frequently and intense you participate on the online forum.

If you think about the last four weeks...

Frequency of visits

How often did you visit the online forum?

- Less than once a month
- Approximately once a month
- Multiple times per month
- Approximately once a week
- Multiple times per week
- Approximately once a day
- Multiple times per day

Average length of visits

What was the average length per visit?

- Less than 10 minutes
- 10 to 30 minutes
- 30 minutes to one hour
- More than one hour

Forum contribution

Did you contribute to the online forum?

- I only read posts from others
- I reacted on (a) post(s) of someone else
- I started a new topic or asked a question
- I both started a new topic/asked a question AND I reacted on (a) post(s) of another

Frequency of posts

How many messages* did you post?

(*initial posts and reactions to questions or stories from others)

- None
- One per week or less
- Multiple posts per week, but not every day
- Every day one post or more

