

VU Research Portal

Remember fast, act skillfully

de Vries, W.

2010

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

de Vries, W. (2010). *Remember fast, act skillfully: Training methods for Basic Life Support; analysis from an educational perspective.*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

7 A web-based micro-simulation program for self-learning BLS skills and the use of an AED. Can laypeople train themselves without a manikin?

Wiebe de Vries^a
Anthony J. Handley^b

^a *Department of Education and Development, Doczero, Rondweg 29, NL-5406 NK Uden, The Netherlands*

^b *40 Queens Road, Colchester, Essex, CO3 3PB, United Kingdom*

Published: Resuscitation 75 (2007) 491–498

Summary

Aim: Various methods, including self-instruction, have been used to try to improve the acquisition of basic life support skills. This is a preliminary report of the effectiveness of a web-based self-training program for BLS and the use of an AED.

Methods: Sixteen volunteers completed on-line training in their own time over a period of 8 weeks. The program included theory, scenario training and self-testing, but without practice on a manikin, or any instructor input. The volunteers were assessed, without prior warning, in a scenario setting. A recording manikin, expert assessors and video recording were used with a modified version of the Cardiff Test.

Results: All 16 volunteers performed the assessed skills in the use of an AED correctly.

Most of the skills of BLS assessed were performed well. Chest compression depth and rate were performed less well (59% and 67% of participants, respectively, performed correctly). Opening the airway and lung inflation were performed poorly (38% and 13% of participants performed correctly), as was checking for safety (19% participants performed correctly). There was no significant correlation between the time a participant spent on-line and the quality of performance. Only 5 of the volunteers had ever attended a BLS course or used a resuscitation manikin before the assessment; their performance scores were not significantly better than those of the other 11 volunteers.

Conclusion: These results suggest that it may be possible to train people in BLS and AED skills using a micro-simulation web-based interactive program but without any practice on a manikin. This seems to be particularly the case for the use of an AED, where performance achieved a uniformly high standard.

Introduction

The more people who are trained in basic life support (BLS) and the use of an automated external defibrillator (AED), potentially the more victims of cardiac arrest can be resuscitated.¹

Unfortunately, BLS skills are poorly acquired,²⁻⁴ so different methods of training have been attempted, including self-instruction using video with a manikin⁵⁻⁹ or without a manikin,^{10,11} using voice assistance,¹²⁻¹⁶ computers¹⁷ and micro simulation.¹⁸ Recent reports have suggested that self-instruction by semi-interactive DVD and the use of a personal manikin can achieve skill acquisition at least as effectively as an instructor-based course.^{19,20} These studies are consistent with the recommendations of ILCOR that instructional

methods should not be limited to traditional techniques.^{21,22}

We hypothesised that with increased public access to a computer and the internet, a web-based training program might be effective, efficient and practical as part of a distance-learning package for BLS and use of an AED. This paper reports the preliminary results of the use of such an internet program.

Materials and methods

The web-based application could be reached from wherever an internet connection was available.

The application consisted of 11 sections covering the theory of BLS and AED use comparable with a manual (Figure 1), a question section for self-assessment of knowledge, a micro-simulation section where the trainee had to react to each scenario by making the correct decisions (Figure 2), and a section for checking if these decisions were correct.




Figure 1.
Example of theory section of on-line program.




Figure 2.
Example of simulation practice section of on-line program.

Volunteer employees were sought from a group of IT implementation and management consultants.

Although the web-based program was developed within the group, the volunteers were from a different department and not involved in any way with this development. They were all computer literate, but were not involved in programming, nor employed for higher computer skills. The volunteers were issued with a licence (in the form of a password) to access the web site and told to visit it as and when they wanted to. They were also told that

they would receive additional training at their place of work 8 weeks, after being given the website licence, that would build on the knowledge they had gained from the web-based program. In the meantime, no checks were made on whether or not they accessed the web site, for how long they did so, or on what progress they were making with self-instruction.

When they attended the expected training session, the volunteers were informed that, instead of further training, they would be assessed on their BLS and AED skill acquisition to date. They were told that management approved of this assessment, and they were asked individually for their permission to be included.

Each participant was taken individually into a room where there was a manikin (AMBU Man C-AMBU AS, Denmark, with CPR Software Kit 2.3), an AED (Lifepak CR Plus training device, Medtronic PhysioControl Corp., Redmond, USA), and a 'bystander' who, they were informed, could not perform CPR or use the AED. They were told to ignore the two assessors standing at the side of the room. Assessment started as each participant entered the room. The AED was programmed for a single-shock scenario, and the assessment was stopped after the 'stand clear' prompt, 2 min after the first shock was delivered.

The Cardiff Test,²³ revised to achieve consistency with Guidelines 2005,²⁴ was used as the assessment tool. The two authors (W.d.V. and A.J.H.) used a score sheet to record steps 1, 2, 3, 4, 6, 12, 13, 15, 20, 24 and 25 of the Cardiff Test (Table 3), the manikin recorded the quality of performance of steps 7, 8, 9, 10 and 11, and a video (Panasonic NV-DS60EG) was taken of each participant's performance from the time of entering the room. This video was used to calculate steps 14, 18 and 21, and for verification of any missing or disputed data.

The overall time each participant had spent online, the percentage of this time on each of the different parts of the program, and the duration of each on-line session were derived from the management system of the application.

All participants were asked if they had followed a BLS or AED course previously, and if they had been familiar with a CPR manikin before the assessment.

Statistics

Significance was accepted when a two-sided *P*-value was <0.05, or confidence intervals did not include unity. All statistics were performed in SPSS® 12.0.1 for Windows (SPSS Inc., Chicago, USA).

Results

Twenty-three participants received a licence and were free to use it as often as they wished, or not at all. Five were unable to attend the assessment meeting; they were not assessed at a later date because they would have been pre-warned of the test.

This left 18 participants to be tested, 11 male and 7 female, of an average age of 44 years (18–54 years). The web management system showed that two of these participants did not access the web based training at any time. They were, therefore, excluded from the subsequent analysis. For the remaining 16 participants the average training time on-line was 189 min (38–408 min). Details of the participants' on-line attendances are given in Table 1. If a participant logged-on, but did not carry out any activities in the program, this was still registered as 'minutes on-line' and influenced the average minutes per session. Some participants visited the web site on consecutive days, whereas others left several days between visits.

	Average ^a (min; max)	Median
Age in years	44,1 (min. 18 max. 55)	47,5
Total minutes on-line	212 (min. 38 max. 408)	202
Minutes per session	103 (min. 9 max. 404)	72
Number of sessions	2.63 (1;5)	2.00
Time span of training in days	8.13 (1;39)	2.00

*: n=16 instead of 18, 2 trainees who did not activate their license were removed.

Table 1.

Results of the trainees on-line.

For this reason, probability distribution was positively skewed. The median was, therefore, also calculated.

Correlations and the results of the tests are shown in Table 2.

There was no significant correlation between the time a participant spent on-line and the quality of performance, nor between correct performance in any one Cardiff step and correct performance in any other step. In other words, performing well in one part of the BLS/AED sequence did not predict good performance overall.

All the steps assessed in the use of an AED were performed correctly by all the participants.

Most of the assessed steps of BLS were performed well (84—100% of the participants performed correctly).

Chest compression depth and rate were performed less well (59% and 67%, respectively, of the participants performed correctly). Opening the airway and lung inflation were performed poorly (38% and 13%, respectively, of the participants performed correctly), as was checking for safety (19% participants performed correctly).

Five of the 16 participants had previously followed a BLS or AED course— one a year beforehand, one 3 years previously and the other 3 more than 12 years previously. These 5 were the only participants who were familiar with a CPR manikin; the other 11 declared that they had never seen one before.

We compared the differences in performance of each assessed step between the group that had previously followed a CPR course and were familiar with a CPR manikin, and the rest of the participants, using a one-way ANOVA, and found no significant differences between them.

Table 3 compares the results of the 16 trainees who accessed the on-line program with those of the two trainees who did no on-line training.

STEP 1; Safe approach	
STEP 2; Check Responsiveness: Talk	
STEP 2; Check Responsiveness: Shake	
STEP 2; Check Responsiveness: Shout for help	
STEP 3; Airway & Breathing: Open Airway	
STEP 3; Airway & Breathing: Check Breathing	
STEP 4; Phone 112	Phone ambulance
STEP 5; Perform CPR as advised by AED	
STEP 6; Hand positioning for chest compressions: Hand position area	
STEP 6; Hand positioning for chest compressions: Total number of correct compressions delivered/minute	
STEP 7; Compression: Average compression rate	90-110
STEP 8; Compression: Average compression depth	38-51 mm
STEP 8; Compression: Return to base line	
STEP 9; Ventilation: Average inflation volume (in litres)	0.4 – 0.6 l
STEP 10; Ventilation: Average number of correct inflations delivered/minute	2-5
STEP 11; Ratio: Compression to Rescue breath ratio 30:2	
STEP 12; Switch on AED	
STEP 13; Remove clothing	
STEP 14; Time interval receipt AED to switch on (in seconds)	
STEP 15; Defibrillator electrode pads attached correctly	
STEP 18; Time interval receipt AED to attach electrodes (in seconds)	
STEP 20; Shock button pushed as directed and shock safety	
STEP 21; Time interval receipt AED to first shock (in seconds)	
STEP 24; Breathing	
STEP 25; Chest compression	

^a Cannot be computed because at least one of the variables is constant, which means that all trainees had the same score.

Table 2.

Correlation between the theoretical and knowledge assessment, and the steps of BLS/AED technique.

Without students without on-line training

Percentage theoretical		Percentage knowledge assessment		Performed, Correct or effective	Mean (min;max)
Pearson Correlation	p	Pearson Correlation	p		
0.124	0.647	0.070	0.796	3/16 (19)	
-0.067	0.806	-0.116	0.669	15/16 (94)	
-0.098	0.719	-0.071	0.793	14/16 (84)	
a		a		16/16 (100)	
0.043	0.875	0.041	0.880	6/16 (38)	
0.188	0.485	0.185	0.492	6/16 (38)	
a		a		16/16 (100)	
a		a		16/16 (100)	
-0.131	0.629	-0.102	0.707	14/16 (88)	
0.324	0.239	-0.146	0.603		28.8 (0;60)
0.293	0.289	-0.082	0.770	10/17 (59)	83.1 (56;115)
0.155	0.582	-0.238	0.394	10/15 (67)	41.6 (12;55)
-0.071	0.800	0.081	0.775	14/15 (93)	
0.268	0.335	-0.159	0.570	2/15 (13)	0.2 (0;6)
0.155	0.582	-0.053	0.850		0.5 (0;3)
-0.067	0.806	-0.138	0.610	15/16 (94)	29:2 (15:0 ; 32;2)
a		a		16/16 (100)	
a		a		16/16 (100)	
a	0.000	0.062	0.849		11.6 (3;48)
-0.098	0.719	-0.071	0.793	16/16 (100)	
.(a)	0.000	0.212	0.508		48.3 (21;126)
0.149	0.582	-0.421	0.104	16/16 (100)	
0.277	0.299	0.308	0.246		70.0 (42;152)
a		a		16/16 (100)	
a		a		16/16 (100)	

Steps of Cardiff Test	Range of acceptable performance
STEP 1; Safe approach	
STEP 2; Check Responsiveness: Talk	
STEP 2; Check Responsiveness: Shake	
STEP 2; Check Responsiveness: Shout for help	
STEP 3; Airway & Breathing: Open Airway	
STEP 3; Airway & Breathing: Check Breathing	
STEP 4; Phone 112	Phone ambulance
STEP 5; Perform CPR as advised by AED	
STEP 6; Hand positioning for chest compressions: Hand position area	
STEP 6; Hand positioning for chest compressions: Total number of correct compressions delivered/minute	
STEP 7; Compression: Average compression rate	90-110
STEP 8; Compression: Average compression depth	38-51 mm
STEP 8; Compression: Return to base line	
STEP 9; Ventilation: Average inflation volume (in litres)	0.4 – 0.6 l
STEP 10; Ventilation: Average number of correct inflations delivered/minute	2-5
STEP 11; Ratio: Compression to Rescue breath ratio 30:2	
STEP 12; Switch on AED	
STEP 13; Remove clothing	
STEP 14; Time interval receipt AED to switch on (in seconds)	
STEP 15; Defibrillator electrode pads attached correctly	
STEP 18; Time interval receipt AED to attach electrodes (in seconds)	
STEP 20; Shock button pushed as directed and shock safety	
STEP 21; Time interval receipt AED to first shock (in seconds)	
STEP 24; Breathing	
STEP 25; Chest compression	

^a One student performed chest-compression-only CPR

^b One student did not ask nor go for an AED

Table 3.

Comparison between the trainees with and without on-line training.

Trainees with on-line training		Trainees without on-line training	
Performance correct or effective	Mean (min;max)	Performance correct or effective	Mean (min;max)
3/16		0/2	
15/16		1/2	
14/16		0/2	
16/16		2/2	
6/16		0/2	
6/16		0/2	
16/16		1/2	
16/16		1/2	
14/16		0/2	
	28.8 (0;60)		58.0 (23;93) ^a
10/17	83.1 (56;115)	0/2	41.0 (3;79)
10/15	41.6 (12;55)	0/2	15.5 (0;31)
14/15		2/2	
2/15	0.2 (0;6)	0/2	0
	0.5 (0;3)	0/2	0
15/16	29:2 (15:0 ; 32;2)	0/2	
16/16		1/2	
16/16		1/2	
	11.6 (3;48)		9.5 (0;19) ^b
16/16 (100)		0/2	
	48.3 (21;126)		33.0 (0;66) ^b
16/16 (100)		1/2	
	70.0 (42;152)		46.0 (0;92) ^b
16/16 (100)		1/2	
16/16 (100)		2/2	

Discussion

Ideally, studies on BLS training should compare the experimental method with an instructor-based method.²⁵ As this was a pilot study to assess the feasibility of on-line training, however, no such control group was used. Manikins are considered necessary when teaching chest compression and ventilation,²⁶ and expert opinion supports the idea that immediate hands-on practice is essential for learning BLS-skills.²⁷ In this study, however, the participants did not use a manikin or AED at any stage of their training, yet, when tested, they achieved 100% score in the use of an AED and performed most BLS skills well.

Each step of the BLS and AED sequence of actions was performed in the correct order by all the 16 participants who had followed the online training. They all checked correctly for consciousness and called for an ambulance. Although most participants were unfamiliar with a BLS manikin, they still provided correct chin lift, placed their hands correctly on the chest to deliver compressions, used the correct compression rate and depth, and returned the chest to a neutral position after each compression.

On the other hand, poor results were recorded for skills related to ventilation, in most cases because participants did not open the airway effectively. This finding is not, however, unique. Previous studies on instructor-led learning and retention have reported poor results for ventilation-related skills;^{28–30} this is a training problem that has yet to be solved. It is of note that the ERC 2005 resuscitation guidelines reduced the emphasis on ventilation during CPR and supported compression-only CPR in certain circumstances.²⁴ The number of correct compressions per minute was low. Of the five participants whose mean depth of compression was outside the accepted range, three were too shallow (<38mm) and two were too deep (>51 mm). Again, this is not unique as even well-trained hospital staff show an inconsistent quality of CPR.^{31,32}

Low scores were achieved for 'safe approach'. This may be partly because it was very clear to the participants on entering the room that there was no physical danger present.

This study has two main limitations. The first is that the only control group consisted of the two participants who did not use the program. It can be seen from Table 3 that their performances were very poor indeed, but the small numbers involved made meaningful statistical analysis impossible. The second limitation is that only tentative conclusions can be drawn from a pilot study with a small number of volunteers.

Conclusions

Overall, the results suggest that it may be possible to achieve reasonable competence in BLS and AED skills using a micro-simulation web-based interactive program, even before practice on a manikin can be provided. This seems to be particularly the case for the use of an AED, where skills were performed to a uniformly high standard. This is consistent with the results of previous studies which show that it is relative easy for lay people to use an AED,^{33–35} and opens the possibility of mass public training via the internet. If this method were to be used, however, it would be important to overcome the problem of ensuring that participants receive sufficient incentive to follow the on-line program.¹⁰ It should not be seen as replacing other methods of training, and practice on a manikin should be encouraged when circumstances permit. It must also be remembered that not everyone is as computer literate as the volunteers in this study.

Conflict of interest

WdV is an employee of Doczero, part of the H2W-group. He is free to perform research, and his management was not involved in designing the study, analysing data or writing the manuscript.

AJH received travel and other expenses for this study without restrictions for research.

Acknowledgements

We wish to thank the employees of H2W for their willingness to be assessed, and Professor Douglas Chamberlain for his kindness in reviewing the manuscript.

References

1. Hazinski MF, Idris AH, Kerber RE, et al. Lay rescuer automated external defibrillator (“Public Access Defibrillation”) programs: lessons learned from an international multicenter trial: advisory Statement from the American Heart Association Emergency Cardiovascular Committee; the Council on Cardiopulmonary, Perioperative, and Critical Care; and the Council on Clinical Cardiology. *Circulation* 2005;111:3336–40.
2. Jansen JJM, Berden HJJM, van der Vleuten CPM, Grol RPTM, Rethans J, Verhoeff ChPM. Evaluation of cardiopulmonary resuscitation skills of general practitioners using different scoring methods. *Resuscitation* 1997;34:35–41.
3. Morgan CLI, Donnelly PD, Lester CA, Assar DH. Effectiveness of the BBC’s 999 training roadshows on cardiopulmonary resuscitation video performance of cohort of unforwarned participants at home six months later. *BMJ* 1996;313(7063):912–6.
4. Kaye W, Mancini ME. Teaching adult resuscitation in the United States—time for a rethink. *Resuscitation* 1998;37:177–87.

5. Braslow A, Brennan RT, Newman MM, Bircher NG, Batcheller AM, Kaye W. CPR training without an instructor: development and evaluation of a video self-instructional system for effective performance of cardiopulmonary resuscitation. *Resuscitation* 1997;34:207–20.
6. Todd KH, Braslow A, Brennan RT, et al. Randomized, controlled trial of video self-instruction versus traditional CPR training. *Ann Emerg Med* 1998;31:364–9.
7. Done ML, Parr M. Teaching basic life support skills using self-directed learning, a self-instructional video, access to practice manikins and learning in pairs. *Resuscitation* 2002;52:287–91.
8. Marco CA, Larkin GL. Public education regarding resuscitation: effects of a multimedia intervention. *Ann Emerg Med* 2003;42:256–60.
9. Batcheller AM, Brennan RT, Braslow A, Urrutia A, Kaye W. Cardiopulmonary resuscitation performance of subjects over forty is better following half-hour video self-instruction compared to traditional four-hour classroom training. *Resuscitation* 2000;43:101–10.
10. Eisenberg M, Damon S, Mandel L, et al. CPR instruction by videotape: results of a community project. *Ann Emerg Med* 1995;25:198–202.
11. Brennan RT, Braslow A. Video self-instruction for cardiopulmonary resuscitation. *Ann Emerg Med* 2000;36: 79–80.
12. Wik L, Thowsen J, Steen PA. An automated voice advisory manikin system for training in basic life support without an instructor. A novel approach to CPR training. *Resuscitation* 2001;50:167–72.
13. Wik L, Myklebust H, Auestad BH, Steen PA. Retention of basic life support skills 6 months after training with an automated voice advisory manikin system without instructor involvement. *Resuscitation* 2002;52:273–9.
14. Handley AJ, Handley SA. Improving CPR performance using an audible feedback system suitable for incorporation into an automated external defibrillator. *Resuscitation* 2003;57:57–62.
15. Wik L, Myklebust H, Auestad BH, Steen PA. Twelve-month retention of CPR skills with automatic correcting verbal feedback. *Resuscitation* 2005;66:27–30.
16. Hostler D, Wang H, Parrish K, Platt TE, Guimond G. The effect of a voice assist manikin (VAM) system on CPR quality among prehospital providers. *Prehosp Emerg Care* 2005;9:53–60.
17. Reder S, Cummings P, Quan L. Comparison of three instructional methods for teaching cardiopulmonary resuscitation and use of an automatic external defibrillator to high school students. *Resuscitation* 2006;69:443–53.
18. Monsieurs KG, Vogels C, Bossaert LL, et al. Learning effect of a novel interactive basic life support CD: the JUST system. *Resuscitation* 2004;62:159–65.
19. Lynch B, Einspruch EL, Nichol G, Becker LB, Aufderheide TP, Idris A. Effectiveness of a 30-min CPR self-instruction program for lay responders: a controlled randomized study. *Resuscitation* 2005;67:31–43.
20. Jones I, Handley AJ, Whitfield R, Newcombe R, Chamberlain D. A preliminary feasibility study of a short DVD-based distance learning package for basic life support. *Resuscitation* 2007;75:350–6.

21. Chamberlain DA, Hazinski MF. ILCOR Advisory Statement; education in resuscitation. *Resuscitation* 2003;59:11—43.
22. International Liaison Committee on Resuscitation. 2005 International consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. Part 8: interdisciplinary topics. *Resuscitation* 2005;67:305—14.
23. Whitfield RH, Newcombe RG, Woollard M. Reliability of the Cardiff Test of basic life support and automated external defibrillation version 3.1. *Resuscitation* 2003;59:291—314.
24. Handley AJ, Koster RW, Monsieurs K, Perkins GD, Davies S, Bossaert LL. European Resuscitation Council Guidelines for Resuscitation 2005. Section 2. Adult basic life support and use of automated external defibrillators. *Resuscitation* 2005;67S1:S7-S23.
25. Hoke RS, Chamberlain DA, Handley AJ. A reference automated external defibrillator provider course for Europe. *Resuscitation* 2006;69:421—33.
26. Baskett PJF, Nolan JP, Handley AJ, Soar J, Biarent D, Richmond S. European Resuscitation Council Guidelines for Resuscitation 2005: Section 9. Principles of training in resuscitation. *Resuscitation* 2005;67S1:S181—9.
27. Bossaert L, Callanan V, Cummins R. Course content. In: Bossaert LL, editor. *European resuscitation council guidelines for resuscitation*. Amsterdam: Elsevier; 1998.
28. Berden HJ, Bierens JJ, Willems FF, Hendrick JM, Pijls NH, Knape JT. Resuscitation skills of lay public after recent training. *Ann Emerg Med* 1994;23(5):1003—8.
29. Wenzel V, Lehmkuhl P, Kubilis PS, Idris AH, Pichlmayr I. Poor correlation or mouth-to-mouth ventilation skills after basic life support training and 6 months later on. *Resuscitation* 1997;35:129—34.
30. Chamberlain D, Smith A, Woollard M, et al. Trials of teaching methods in basic life support (3): comparison of simulated CPR performance after first training and at 6 months, with a note on the value of re-training. *Resuscitation* 2002;53:179—87.
31. Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during in-hospital cardiac arrest. *JAMA* 2005;293:305—10.
32. Einav S, Shleifer A, Kark JD, Landesberg G, Matot I. Performance of department staff in the window between discovery of collapse to cardiac arrest team arrival. *Resuscitation* 2006;69:209—16.
33. Monsieurs KG, Vogels C, Bossaert LL, Meert P, Calle PA. A study comparing the usability of fully automatic versus semi-automatic defibrillation by untrained nursing students. *Resuscitation* 2005;64:41—7.
34. Usatch BR, Cone DC. Automated external defibrillator training and skill retention at a ski patrol. *Prehosp Emerg Care* 2002;6:325—9.
35. Kooij FO, van Alem AP, Koster RW, de Vos R. Training of police officers as first responders with an automated external defibrillator. *Resuscitation* 2004;63:33—41.