

VU Research Portal

Cutaneous Melanoma: Predictors of Patient Survival and the Potential of Priming of the Sentinel Lymph Node

Sluijter, B.J.R.

2011

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Sluijter, B. J. R. (2011). *Cutaneous Melanoma: Predictors of Patient Survival and the Potential of Priming of the Sentinel Lymph Node*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Table of Contents

Introduction

- Chapter 1 Introduction, adapted from “The Dermis as a Portal for Dendritic Cell targeted Immunotherapy of Cutaneous Melanoma” and outline of this thesis **11**
Current Topics in Microbiology and Immunology, in press

Clinical observations in melanoma patients

- Chapter 2 Non-radical diagnostic biopsies do not negatively influence melanoma patient survival **37**
Ann Surg Oncol. 2007 Apr;14(4):1424-30
- Chapter 3 10-year follow-up results; clinical outcome of stage I/II melanoma patients **51**

Immunohistochemistry of primary tumors and SLN from stage I/II melanoma patients

- Chapter 4 Favorable outcome in clinically stage II melanoma patients is associated with the presence of activated tumor infiltrating T-lymphocytes and preserved MHC class I antigen expression **69**
Int J Cancer 2008 Aug 1;123(3):609-15
- Chapter 5 Absence of Granzyme B positive tumor-infiltrating lymphocytes in primary melanoma excisional biopsies is strongly associated with the presence of sentinel lymph node metastasis **83**
Cell Oncol. 2009;31(5):407-13

Priming of the SLN of melanoma patients with GM-CSF and/or PF-3512676 CpG-B

- Chapter 6 Local administration of PF-3512676 CpG-B instigates tumor-specific CD8⁺ T-cell reactivity in melanoma patients **97**
Clin Cancer Res. 2008 Jul 15;14(14): 4532-42
- Chapter 7 Activation and recruitment of BDCA3/CD141⁺ Dendritic Cell subsets to the Sentinel Lymph Node by delivery of CpG-B and GM-CSF at the melanoma excision site **117**
- Chapter 8 Effects of the method of T cell expansion for primed SLN cells from melanoma patients on the quantity and quality of the CD8⁺ cytotoxic T cell fraction **137**
Clin Immunol. 2010 Nov;137(2):221-33.

- Chapter 9 Summarizing Discussion and Future Prospectives **157**
Dutch summary **171**

- Chapter 10 Curriculum vitae **183**
List of publications **184**
Dankwoord **185**

- Chapter 11 Color figures **187**