

# VU Research Portal

## The role of hydrophobicity in protein folding and aggregation

Dijk, E.

2017

### **document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

### **citation for published version (APA)**

Dijk, E. (2017). *The role of hydrophobicity in protein folding and aggregation*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# Contents

|          |   |            |
|----------|---|------------|
| <b>1</b> | <b>Introduction</b>   | <b>1</b>   |
| <b>2</b> | <b>The Hydrophobic Temperature Dependence of Amino Acids Directly Calculated from Protein Structures</b>                              | <b>9</b>   |
| 2.1      | Supplemental information -The Hydrophobic Temperature Dependence of Amino Acids Directly Calculated from Protein Structures . . . . . | 25         |
| <b>3</b> | <b>Consistent treatment of hydrophobicity in protein lattice models accounts for cold denaturation</b>                                | <b>27</b>  |
| 3.1      | Supplementary Material – Consistent treatment of hydrophobicity in protein lattice models accounts for cold denaturation . . .        | 38         |
| <b>4</b> | <b>BICEP: Heat Capacity Baseline Prediction</b>   | <b>53</b>  |
| 4.1      | Supporting Information - BICEP: Heat Capacity Baseline Prediction . . . . .   | 65         |
| <b>5</b> | <b>Predicting the hydrophobic surface area of native protein structures from sequence</b>   | <b>71</b>  |
| 5.1      | Supplementary Information - Predicting the hydrophobic surface area of native protein structures from sequence . . . . .              | 86         |
| <b>6</b> | <b>Cold denaturation of amyloid fibrils explained through the hydrophobic temperature dependence</b>                                  | <b>91</b>  |
| 6.1      | Supplemental information - Cold denaturation of amyloid fibrils explained through the hydrophobic temperature dependence . .          | 104        |
| <b>7</b> | <b>Discussion</b>   | <b>107</b> |
|          | <b>Bibliography</b>   | <b>113</b> |