

VU Research Portal

The role of presynaptic proteins in maintaining neuronal viability

Cerveira Tavares dos Santos, T.

2017

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Cerveira Tavares dos Santos, T. (2017). *The role of presynaptic proteins in maintaining neuronal viability*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

- The Journal of biological chemistry* 2010; **285**(37): 28708-28714.
76. Koch M, Holt M. Coupling exo- and endocytosis: an essential role for PIP(2) at the synapse. *Biochimica et biophysica acta* 2012; **1821**(8): 1114-1132.
 77. Auger KR, Serunian LA, Soltoff SP, Libby P, Cantley LC. PDGF-dependent tyrosine phosphorylation stimulates production of novel polyphosphoinositides in intact cells. *Cell* 1989; **57**(1): 167-175.
 78. Maehama T, Dixon JE. The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate. *The Journal of biological chemistry* 1998; **273**(22): 13375-13378.
 79. Viaud J, Mansour R, Antkowiak A, Mujalli A, Valet C, Chicanne G, Xuereb JM, Terrisse AD, Severin S, Gratacap MP, Gaits-Iacovoni F, Payrastre B. Phosphoinositides: Important lipids in the coordination of cell dynamics. *Biochimie* 2015; **125**: 250-258.
 80. Czech MP. Dynamics of phosphoinositides in membrane retrieval and insertion. *Annu Rev Physiol* 2003; **65**: 791-815.
 81. Katso R, Okkenhaug K, Ahmadi K, White S, Timms J, Waterfield MD. Cellular function of phosphoinositide 3-kinases: implications for development, homeostasis, and cancer. *Annual review of cell and developmental biology* 2001; **17**: 615-675.
 82. Idevall-Hagren O, De Camilli P. Detection and manipulation of phosphoinositides. *Biochimica et biophysica acta* 2015; **1851**(6): 736-745.
 83. Rusten TE, Stenmark H. Analyzing phosphoinositides and their interacting proteins. *Nat Methods* 2006; **3**(4): 251-258.
 84. Bunney TD, Katan M. Phosphoinositide signalling in cancer: beyond PI3K and PTEN. *Nat Rev Cancer* 2010; **10**(5): 342-352.
 85. Vanden Berghe T, Linkermann A, Jouan-Lanhout S, Walczak H, Vandenabeele P. Regulated necrosis: the expanding network of non-apoptotic cell death pathways. *Nature reviews Molecular cell biology* 2014; **15**(2): 135-147.
 86. Schweichel JU, Merker HJ. The morphology of various types of cell death in prenatal tissues. *Teratology* 1973; **7**(3): 253-266.
 87. Nikolettou V, Markaki M, Palikaras K, Tavernarakis N. Crosstalk between apoptosis, necrosis and autophagy. *Biochimica et biophysica acta* 2013; **1833**(12): 3448-3459.
 88. Festjens N, Vanden Berghe T, Cornelis S, Vandenabeele P. RIP1, a kinase on the crossroads of a cell's decision to live or die. *Cell death and differentiation* 2007; **14**(3): 400-410.
 89. Holler N, Zaru R, Micheau O, Thome M, Attinger A, Valitutti S, Bodmer JL, Schneider P, Seed B, Tschopp J. Fas triggers an alternative, caspase-8-independent cell death pathway using the kinase RIP as effector molecule. *Nat Immunol* 2000; **1**(6): 489-495.
 90. Fulda S, Gorman AM, Hori O, Samali A. Cellular stress responses: cell survival and cell death. *Int J Cell Biol* 2010; **2010**: 214074.
 91. Nixon RA, Yang DS. Autophagy and neuronal cell death in neurological disorders. *Cold Spring Harbor perspectives in biology* 2012; **4**(10).
 92. Son JH, Shim JH, Kim KH, Ha JY, Han JY. Neuronal autophagy and neurodegenerative diseases. *Exp Mol Med* 2012; **44**(2): 89-98.
 93. Clarke PG. Developmental cell death: morphological diversity and multiple mechanisms. *Anat Embryol (Berl)* 1990; **181**(3): 195-213.
 94. Zhu XC, Yu JT, Jiang T, Tan L. Autophagy modulation for Alzheimer's disease therapy. *Molecular neurobiology* 2013; **48**(3): 702-714.
 95. Rujano MA, Bosveld F, Salomons FA, Dijk F, van Waarde MA, van der Want JJ, de Vos RA, Brunt ER, Sibon OC, Kampinga HH. Polarised asymmetric inheritance of accumulated protein damage in higher eukaryotes. *PLoS Biol* 2006; **4**(12): e417.
 96. Lindner AB, Madden R, Demarez A, Stewart EJ, Taddei F. Asymmetric segregation of protein aggregates is associated with cellular aging and rejuvenation. *Proceedings of the National Academy of Sciences of the United States of America* 2008; **105**(8): 3076-3081.
 97. Huang Y, Mucke L. Alzheimer mechanisms and therapeutic strategies. *Cell* 2012; **148**(6): 1204-1222.
 98. Beitz JM. Parkinson's disease: a review. *Front Biosci (Schol Ed)* 2014; **6**: 65-74.
 99. Lucassen PJ, Chung WC, Kamphorst W, Swaab DF. DNA damage distribution in the human brain as shown by in situ end labeling: area-specific differences in aging and Alzheimer disease in the absence of apoptotic morphology. *Journal of neuropathology and experimental neurology* 1997; **56**(8): 887-900.
 100. Stadelmann C, Bruck W, Bancher C, Jellinger K, Lassmann H. Alzheimer disease: DNA fragmentation indicates increased neuronal vulnerability, but not apoptosis. *Journal of neuropathology and experimental neurology* 1998;

- 57(5): 456-464.
101. Su JH, Anderson AJ, Cummings BJ, Cotman CW. Immunohistochemical evidence for apoptosis in Alzheimer's disease. *Neuroreport* 1994; **5**(18): 2529-2533.
 102. Smale G, Nichols NR, Brady DR, Finch CE, Horton WE, Jr. Evidence for apoptotic cell death in Alzheimer's disease. *Experimental neurology* 1995; **133**(2): 225-230.
 103. Troncoso JC, Sukhov RR, Kawas CH, Koliatsos VE. In situ labeling of dying cortical neurons in normal aging and in Alzheimer's disease: correlations with senile plaques and disease progression. *Journal of neuropathology and experimental neurology* 1996; **55**(11): 1134-1142.
 104. Rohn TT, Head E, Nesse WH, Cotman CW, Cribbs DH. Activation of caspase-8 in the Alzheimer's disease brain. *Neurobiology of disease* 2001; **8**(6): 1006-1016.
 105. Pompl PN, Yemul S, Xiang Z, Ho L, Haroutunian V, Purohit D, Mohs R, Pasinetti GM. Caspase gene expression in the brain as a function of the clinical progression of Alzheimer disease. *Arch Neurol* 2003; **60**(3): 369-376.
 106. Gervais FG, Xu D, Robertson GS, Vaillancourt JP, Zhu Y, Huang J, LeBlanc A, Smith D, Rigby M, Shearman MS, Clarke EE, Zheng H, Van Der Ploeg LH, Ruffolo SC, Thornberry NA, Xanthoudakis S, Zamboni RJ, Roy S, Nicholson DW. Involvement of caspases in proteolytic cleavage of Alzheimer's amyloid-beta precursor protein and amyloidogenic A beta peptide formation. *Cell* 1999; **97**(3): 395-406.
 107. Gamblin TC, Chen F, Zambrano A, Abraha A, Lagalwar S, Guillozet AL, Lu M, Fu Y, Garcia-Sierra F, LaPointe N, Miller R, Berry RW, Binder LI, Cryns VL. Caspase cleavage of tau: linking amyloid and neurofibrillary tangles in Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United States of America* 2003; **100**(17): 10032-10037.
 108. O'Brien RJ, Wong PC. Amyloid precursor protein processing and Alzheimer's disease. *Annu Rev Neurosci* 2011; **34**: 185-204.
 109. de Calignon A, Fox LM, Pitstick R, Carlson GA, Bacskai BJ, Spire-Jones TL, Hyman BT. Caspase activation precedes and leads to tangles. *Nature* 2010; **464**(7292): 1201-1204.
 110. Rissman RA, Poon WW, Blurton-Jones M, Oddo S, Torp R, Vitek MP, LaFerla FM, Rohn TT, Cotman CW. Caspase-cleavage of tau is an early event in Alzheimer disease tangle pathology. *J Clin Invest* 2004; **114**(1): 121-130.
 111. Nixon RA, Wegiel J, Kumar A, Yu WH, Peterhoff C, Cataldo A, Cuervo AM. Extensive involvement of autophagy in Alzheimer disease: an immuno-electron microscopy study. *Journal of neuropathology and experimental neurology* 2005; **64**(2): 113-122.
 112. Yu WH, Cuervo AM, Kumar A, Peterhoff CM, Schmidt SD, Lee JH, Mohan PS, Mercken M, Farmery MR, Tjernberg LO, Jiang Y, Duff K, Uchiyama Y, Naslund J, Mathews PM, Cataldo AM, Nixon RA. Macroautophagy—a novel Beta-amyloid peptide-generating pathway activated in Alzheimer's disease. *The Journal of cell biology* 2005; **171**(1): 87-98.
 113. Spilman P, Podlutska N, Hart MJ, Debnath J, Gorostiza O, Bredesen D, Richardson A, Strong R, Galvan V. Inhibition of mTOR by rapamycin abolishes cognitive deficits and reduces amyloid-beta levels in a mouse model of Alzheimer's disease. *PLoS one* 2010; **5**(4): e9979.
 114. Caccamo A, Majumder S, Richardson A, Strong R, Oddo S. Molecular interplay between mammalian target of rapamycin (mTOR), amyloid-beta, and Tau: effects on cognitive impairments. *The Journal of biological chemistry* 2010; **285**(17): 13107-13120.
 115. Venderova K, Park DS. Programmed cell death in Parkinson's disease. *Cold Spring Harb Perspect Med* 2012; **2**(8).
 116. Yonekawa Y, Harada A, Okada Y, Funakoshi T, Kanai Y, Takei Y, Terada S, Noda T, Hirokawa N. Defect in synaptic vesicle precursor transport and neuronal cell death in KIF1A motor protein-deficient mice. *The Journal of cell biology* 1998; **141**(2): 431-441.
 117. Golgi C. Sur la structure des cellules nerveuse ganglions spinaux. *Arch Ital Biol* 1898; **30**: 278-286.
 118. Klumperman J. Architecture of the mammalian Golgi. *Cold Spring Harbor perspectives in biology* 2011; **3**(7).
 119. Cooper GM. *The Cell a molecular approach*. 2000.
 120. Sundaramoorthy V, Sultana JM, Atkin JD. Golgi fragmentation in amyotrophic lateral sclerosis, an overview of possible triggers and consequences. *Front Neurosci* 2015; **9**: 400.
 121. Glick BS, Luini A. Models for Golgi traffic: a critical assessment. *Cold Spring Harbor perspectives in biology* 2011; **3**(11): a005215.
 122. Hong W, Lev S. Tethering the assembly of SNARE complexes. *Trends in cell biology* 2014; **24**(1): 35-43.
 123. Malsam J, Sollner TH. Organization of SNAREs within the Golgi stack. *Cold Spring Harbor perspectives in biology* 2011; **3**(10): a005249.
 124. Jahn R, Scheller RH. SNAREs—engines for membrane fusion. *Nature reviews Molecular cell biology* 2006; **7**(9):

- 631-643.
125. Lazaro-Dieguez F, Jimenez N, Barth H, Koster AJ, Renau-Piqueras J, Llopis JL, Burger KN, Egea G. Actin filaments are involved in the maintenance of Golgi cisternae morphology and intra-Golgi pH. *Cell motility and the cytoskeleton* 2006; **63**(12): 778-791.
 126. Valderrama F, Duran JM, Babia T, Barth H, Renau-Piqueras J, Egea G. Actin microfilaments facilitate the retrograde transport from the Golgi complex to the endoplasmic reticulum in mammalian cells. *Traffic* 2001; **2**(10): 717-726.
 127. Dippold HC, Ng MM, Farber-Katz SE, Lee SK, Kerr ML, Peterman MC, Sim R, Wiharto PA, Galbraith KA, Madhavarapu S, Fuchs GJ, Meerloo T, Farquhar MG, Zhou H, Field SJ. GOLPH3 bridges phosphatidylinositol-4-phosphate and actomyosin to stretch and shape the Golgi to promote budding. *Cell* 2009; **139**(2): 337-351.
 128. Ng MM, Dippold HC, Buschman MD, Noakes CJ, Field SJ. GOLPH3L antagonizes GOLPH3 to determine Golgi morphology. *Molecular biology of the cell* 2013; **24**(6): 796-808.
 129. Jaarsma D, Hoogenraad CC. Cytoplasmic dynein and its regulatory proteins in Golgi pathology in nervous system disorders. *Front Neurosci* 2015; **9**: 397.
 130. Palmer KJ, Hughes H, Stephens DJ. Specificity of cytoplasmic dynein subunits in discrete membrane-trafficking steps. *Molecular biology of the cell* 2009; **20**(12): 2885-2899.
 131. Yadav S, Puthenveedu MA, Linstedt AD. Golgin160 recruits the dynein motor to position the Golgi apparatus. *Dev Cell* 2012; **23**(1): 153-165.
 132. Glick BS, Nakano A. Membrane traffic within the Golgi apparatus. *Annual review of cell and developmental biology* 2009; **25**: 113-132.
 133. Fujiwara T, Oda K, Yokota S, Takatsuki A, Ikehara Y, Brefeldin A causes disassembly of the Golgi complex and accumulation of secretory proteins in the endoplasmic reticulum. *The Journal of biological chemistry* 1988; **263**(34): 18545-18552.
 134. Robbins E, Gonatas NK. The Ultrastructure of a Mammalian Cell during the Mitotic Cycle. *The Journal of cell biology* 1964; **21**: 429-463.
 135. Lowe M, Lane JD, Woodman PG, Allan VJ. Caspase-mediated cleavage of syntaxin 5 and giantin accompanies inhibition of secretory traffic during apoptosis. *Journal of cell science* 2004; **117**(Pt 7): 1139-1150.
 136. Sesso A, Fujiwara DT, Jaeger M, Jaeger R, Li TC, Monteiro MM, Correa H, Ferreira MA, Schumacher RI, Belisario J, Kachar B, Chen EJ. Structural elements common to mitosis and apoptosis. *Tissue & cell* 1999; **31**(3): 357-371.
 137. Dascher C, Balch WE. Dominant inhibitory mutants of ARF1 block endoplasmic reticulum to Golgi transport and trigger disassembly of the Golgi apparatus. *The Journal of biological chemistry* 1994; **269**(2): 1437-1448.
 138. Wilson BS, Nuoffer C, Meinkoth JL, McCaffery M, Feramisco JR, Balch WE, Farquhar MG. A Rab1 mutant affecting guanine nucleotide exchange promotes disassembly of the Golgi apparatus. *The Journal of cell biology* 1994; **125**(3): 557-571.
 139. Thayer DA, Jan YN, Jan LY. Increased neuronal activity fragments the Golgi complex. *Proceedings of the National Academy of Sciences of the United States of America* 2013; **110**(4): 1482-1487.
 140. Mourelatos Z, Adler H, Hirano A, Donnenfeld H, Gonatas JO, Gonatas NK. Fragmentation of the Golgi apparatus of motor neurons in amyotrophic lateral sclerosis revealed by organelle-specific antibodies. *Proceedings of the National Academy of Sciences of the United States of America* 1990; **87**(11): 4393-4395.
 141. Haase G, Rabouille C. Golgi Fragmentation in ALS Motor Neurons. New Mechanisms Targeting Microtubules, Tethers, and Transport Vesicles. *Front Neurosci* 2015; **9**: 448.
 142. Mourelatos Z, Gonatas NK, Stieber A, Gurney ME, Dal Canto MC. The Golgi apparatus of spinal cord motor neurons in transgenic mice expressing mutant Cu,Zn superoxide dismutase becomes fragmented in early, preclinical stages of the disease. *Proceedings of the National Academy of Sciences of the United States of America* 1996; **93**(11): 5472-5477.
 143. Gonatas NK, Stieber A, Mourelatos Z, Chen Y, Gonatas JO, Appel SH, Hays AP, Hickey WF, Haw JJ. Fragmentation of the Golgi apparatus of motor neurons in amyotrophic lateral sclerosis. *The American journal of pathology* 1992; **140**(3): 731-737.
 144. Nassif M, Matus S, Castillo K, Hetz C. Amyotrophic lateral sclerosis pathogenesis: a journey through the secretory pathway. *Antioxidants & redox signaling* 2010; **13**(12): 1955-1989.
 145. Persson R, Schnell CR, Borg LA, Fries E. Accumulation of Golgi-processed secretory proteins in an organelle of high density upon reduction of ATP concentration in rat hepatocytes. *The Journal of biological chemistry* 1992; **267**(4): 2760-2766.
 146. Zolov SN, Lupashin VV. Cog3p depletion blocks vesicle-mediated Golgi retrograde trafficking in HeLa cells. *The Journal of cell biology* 2005; **168**(5): 747-759.

147. Stieber A, Mourelatos Z, Gonatas NK. In Alzheimer's disease the Golgi apparatus of a population of neurons without neurofibrillary tangles is fragmented and atrophic. *The American journal of pathology* 1996; **148**(2): 415-426.
148. Huse JT, Liu K, Pijak DS, Carlin D, Lee VM, Doms RW. Beta-secretase processing in the trans-Golgi network preferentially generates truncated amyloid species that accumulate in Alzheimer's disease brain. *The Journal of biological chemistry* 2002; **277**(18): 16278-16284.
149. Sun KH, de Pablo Y, Vincent F, Johnson EO, Chavers AK, Shah K. Novel genetic tools reveal Cdk5's major role in Golgi fragmentation in Alzheimer's disease. *Molecular biology of the cell* 2008; **19**(7): 3052-3069.
150. Joshi G, Chi Y, Huang Z, Wang Y. Abeta-induced Golgi fragmentation in Alzheimer's disease enhances Abeta production. *Proceedings of the National Academy of Sciences of the United States of America* 2014; **111**(13): E1230-1239.
151. Fujita Y, Ohama E, Takatama M, Al-Sarraj S, Okamoto K. Fragmentation of Golgi apparatus of nigral neurons with alpha-synuclein-positive inclusions in patients with Parkinson's disease. *Acta neuropathologica* 2006; **112**(3): 261-265.
152. Milosevic I, Giovedi S, Lou X, Raimondi A, Collesi C, Shen H, Paradise S, O'Toole E, Ferguson S, Cremona O, De Camilli P. Recruitment of endophilin to clathrin-coated pit necks is required for efficient vesicle uncoating after fission. *Neuron* 2011; **72**(4): 587-601.
153. Cremona O, Di Paolo G, Wenk MR, Luthi A, Kim WT, Takei K, Daniell L, Nemoto Y, Shears SB, Flavell RA, McCormick DA, De Camilli P. Essential role of phosphoinositide metabolism in synaptic vesicle recycling. *Cell* 1999; **99**(2): 179-188.
154. Raimondi A, Ferguson SM, Lou X, Armbruster M, Paradise S, Giovedi S, Messa M, Kono N, Takasaki J, Cappello V, O'Toole E, Ryan TA, De Camilli P. Overlapping role of dynamin isoforms in synaptic vesicle endocytosis. *Neuron* 2011; **70**(6): 1100-1114.
155. Pevsner J, Hsu SC, Scheller RH. n-Sec1: a neural-specific syntaxin-binding protein. *Proceedings of the National Academy of Sciences of the United States of America* 1994; **91**(4): 1445-1449.
156. Hata Y, Slaughter CA, Sudhof TC. Synaptic vesicle fusion complex contains unc-18 homologue bound to syntaxin. *Nature* 1993; **366**(6453): 347-351.
157. Garcia EP, Gatti E, Butler M, Burton J, De Camilli P. A rat brain Sec1 homologue related to Rop and UNC18 interacts with syntaxin. *Proceedings of the National Academy of Sciences of the United States of America* 1994; **91**(6): 2003-2007.
158. Tellam JT, McIntosh S, James DE. Molecular identification of two novel Munc-18 isoforms expressed in non-neuronal tissues. *The Journal of biological chemistry* 1995; **270**(11): 5857-5863.
159. Khvotchev M, Dulubova I, Sun J, Dai H, Rizo J, Sudhof TC. Dual modes of Munc18-1/SNARE interactions are coupled by functionally critical binding to syntaxin-1 N terminus. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2007; **27**(45): 12147-12155.
160. Zhou P, Pang ZP, Yang X, Zhang Y, Rosenmund C, Bacaj T, Sudhof TC. Syntaxin-1 N-peptide and Habc-domain perform distinct essential functions in synaptic vesicle fusion. *The EMBO journal* 2013; **32**(1): 159-171.
161. Novick P, Schekman R. Secretion and cell-surface growth are blocked in a temperature-sensitive mutant of *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences of the United States of America* 1979; **76**(4): 1858-1862.
162. Brenner S. The genetics of *Caenorhabditis elegans*. *Genetics* 1974; **77**(1): 71-94.
163. Salzberg A, Cohen N, Halachmi N, Kimchie Z, Lev Z. The *Drosophila* Ras2 and Rop gene pair: a dual homology with a yeast Ras-like gene and a suppressor of its loss-of-function phenotype. *Development* 1993; **117**(4): 1309-1319.
164. Harrison SD, Broadie K, van de Goor J, Rubin GM. Mutations in the *Drosophila* Rop gene suggest a function in general secretion and synaptic transmission. *Neuron* 1994; **13**(3): 555-566.
165. Weimer RM, Richmond JE, Davis WS, Hadwiger G, Nonet ML, Jorgensen EM. Defects in synaptic vesicle docking in unc-18 mutants. *Nature neuroscience* 2003; **6**(10): 1023-1030.
166. Voets T, Toonen RF, Brian EC, de Wit H, Moser T, Rettig J, Sudhof TC, Neher E, Verhage M. Munc18-1 promotes large dense-core vesicle docking. *Neuron* 2001; **31**(4): 581-591.
167. Toonen RF, Kochubey O, de Wit H, Gulyas-Kovacs A, Konijnenburg B, Sorensen JB, Klingauf J, Verhage M. Dissecting docking and tethering of secretory vesicles at the target membrane. *The EMBO journal* 2006; **25**(16): 3725-3737.
168. de Wit H, Cornelisse LN, Toonen RF, Verhage M. Docking of secretory vesicles is syntaxin dependent. *PLoS one* 2006; **1**: e126.
169. Deak F, Xu Y, Chang WP, Dulubova I, Khvotchev M, Liu X, Sudhof TC, Rizo J. Munc18-1 binding to the

- neuronal SNARE complex controls synaptic vesicle priming. *The Journal of cell biology* 2009; **184**(5): 751-764.
170. Han GA, Bin NR, Kang SY, Han L, Sugita S. Domain 3a of Munc18-1 plays a crucial role at the priming stage of exocytosis. *Journal of cell science* 2013; **126**(Pt 11): 2361-2371.
 171. Rodkey TL, Liu S, Barry M, McNew JA. Munc18a scaffolds SNARE assembly to promote membrane fusion. *Molecular biology of the cell* 2008; **19**(12): 5422-5434.
 172. Shen J, Tareste DC, Paumet F, Rothman JE, Melia TJ. Selective activation of cognate SNAREpins by Sec1/Munc18 proteins. *Cell* 2007; **128**(1): 183-195.
 173. Tareste D, Shen J, Melia TJ, Rothman JE. SNAREpin/Munc18 promotes adhesion and fusion of large vesicles to giant membranes. *Proceedings of the National Academy of Sciences of the United States of America* 2008; **105**(7): 2380-2385.
 174. Toonen RF, de Vries KJ, Zalm R, Sudhof TC, Verhage M. Munc18-1 stabilizes syntaxin 1, but is not essential for syntaxin 1 targeting and SNARE complex formation. *Journal of neurochemistry* 2005; **93**(6): 1393-1400.
 175. Rowe J, Corradi N, Malosio ML, Taverna E, Halban P, Meldolesi J, Rosa P. Blockade of membrane transport and disassembly of the Golgi complex by expression of syntaxin 1A in neurosecretion-incompetent cells: prevention by rbSEC1. *Journal of cell science* 1999; **112** (Pt 12): 1865-1877.
 176. Rowe J, Calegari F, Taverna E, Longhi R, Rosa P. Syntaxin 1A is delivered to the apical and basolateral domains of epithelial cells: the role of munc-18 proteins. *Journal of cell science* 2001; **114**(Pt 18): 3323-3332.
 177. Han L, Jiang T, Han GA, Malintan NT, Xie L, Wang L, Tse FW, Gaisano HY, Collins BM, Meunier FA, Sugita S. Rescue of Munc18-1 and -2 double knockdown reveals the essential functions of interaction between Munc18 and closed syntaxin in PC12 cells. *Molecular biology of the cell* 2009; **20**(23): 4962-4975.
 178. Arunachalam L, Han L, Tassew NG, He Y, Wang L, Xie L, Fujita Y, Kwan E, Davletov B, Monnier PP, Gaisano HY, Sugita S. Munc18-1 is critical for plasma membrane localization of syntaxin1 but not of SNAP-25 in PC12 cells. *Molecular biology of the cell* 2008; **19**(2): 722-734.
 179. Dulubova I, Sugita S, Hill S, Hosaka M, Fernandez I, Sudhof TC, Rizo J. A conformational switch in syntaxin during exocytosis: role of munc18. *The EMBO journal* 1999; **18**(16): 4372-4382.
 180. Yang B, Steegmaier M, Gonzalez LC, Jr., Scheller RH. nSec1 binds a closed conformation of syntaxin1A. *The Journal of cell biology* 2000; **148**(2): 247-252.
 181. Misura KM, Scheller RH, Weis WI. Three-dimensional structure of the neuronal-Sec1-syntaxin 1a complex. *Nature* 2000; **404**(6776): 355-362.
 182. Dulubova I, Khvotchev M, Liu S, Huryeva I, Sudhof TC, Rizo J. Munc18-1 binds directly to the neuronal SNARE complex. *Proceedings of the National Academy of Sciences of the United States of America* 2007; **104**(8): 2697-2702.
 183. Burkhardt P, Hattendorf DA, Weis WI, Fasshauer D. Munc18a controls SNARE assembly through its interaction with the syntaxin N-peptide. *The EMBO journal* 2008; **27**(7): 923-933.
 184. Heeroma JH, Plomp JJ, Roubos EW, Verhage M. Development of the mouse neuromuscular junction in the absence of regulated secretion. *Neuroscience* 2003; **120**(3): 733-744.
 185. Fujiwara T, Mishima T, Kofuji T, Chiba T, Tanaka K, Yamamoto A, Akagawa K. Analysis of knock-out mice to determine the role of HPC-1/syntaxin 1A in expressing synaptic plasticity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2006; **26**(21): 5767-5776.
 186. Kofuji T, Fujiwara T, Sanada M, Mishima T, Akagawa K. HPC-1/syntaxin 1A and syntaxin 1B play distinct roles in neuronal survival. *Journal of neurochemistry* 2014; **130**(4): 514-525.
 187. Mishima T, Fujiwara T, Sanada M, Kofuji T, Kanai-Azuma M, Akagawa K. Syntaxin 1B, but not syntaxin 1A, is necessary for the regulation of synaptic vesicle exocytosis and of the readily releasable pool at central synapses. *PLoS one* 2014; **9**(2): e90004.
 188. Sato M, Yoshimura S, Hirai R, Goto A, Kunii M, Atik N, Sato T, Sato K, Harada R, Shimada J, Hatabu T, Yorifuji H, Harada A. The role of VAMP7/TI-VAMP in cell polarity and lysosomal exocytosis in vivo. *Traffic* 2011; **12**(10): 1383-1393.
 189. Danglot L, Zylbersztein K, Petkovic M, Gauberti M, Meziane H, Combe R, Champy MF, Birling MC, Pavlovic G, Bizot JC, Trovero F, Della Ragione F, Proux-Gillardeaux V, Sorg T, Vivien D, D'Esposito M, Galli T. Absence of TI-VAMP/Vamp7 leads to increased anxiety in mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2012; **32**(6): 1962-1968.
 190. Geppert M, Goda Y, Hammer RE, Li C, Rosahl TW, Stevens CF, Sudhof TC. Synaptotagmin I: a major Ca²⁺ sensor for transmitter release at a central synapse. *Cell* 1994; **79**(4): 717-727.
 191. Reim K, Mansour M, Varoqueaux F, McMahon HT, Sudhof TC, Brose N, Rosenmund C. Complexins regulate a late step in Ca²⁺-dependent neurotransmitter release. *Cell* 2001; **104**(1): 71-81.
 192. Fernandez-Chacon R, Wolfel M, Nishimune H, Tabares L, Schmitz F, Castellano-Munoz M, Rosenmund C,

- Montesinos ML, Sanes JR, Schneggenburger R, Sudhof TC. The synaptic vesicle protein CSP alpha prevents presynaptic degeneration. *Neuron* 2004; **42**(2): 237-251.
193. Sharma M, Burre J, Bronk P, Zhang Y, Xu W, Sudhof TC. CSPalpha knockout causes neurodegeneration by impairing SNAP-25 function. *The EMBO journal* 2012; **31**(4): 829-841.
 194. Speidel D, Bruederle CE, Enk C, Voets T, Varoqueaux F, Reim K, Becherer U, Fornai F, Ruggieri S, Holighaus Y, Weihe E, Bruns D, Brose N, Rettig J. CAPS1 regulates catecholamine loading of large dense-core vesicles. *Neuron* 2005; **46**(1): 75-88.
 195. Jockusch WJ, Speidel D, Sigler A, Sorensen JB, Varoqueaux F, Rhee JS, Brose N. CAPS-1 and CAPS-2 are essential synaptic vesicle priming proteins. *Cell* 2007; **131**(4): 796-808.
 196. Ferguson SM, Brasnjo G, Hayashi M, Wolfel M, Collesi C, Giovedi S, Raimondi A, Gong LW, Ariel P, Paradise S, O'Toole E, Flavell R, Cremona O, Miesenbock G, Ryan TA, De Camilli P. A selective activity-dependent requirement for dynamin 1 in synaptic vesicle endocytosis. *Science* 2007; **316**(5824): 570-574.
 197. Ferguson SM, Raimondi A, Paradise S, Shen H, Mesaki K, Ferguson A, Destaing O, Ko G, Takasaki J, Cremona O, E OT, De Camilli P. Coordinated actions of actin and BAR proteins upstream of dynamin at endocytic clathrin-coated pits. *Dev Cell* 2009; **17**(6): 811-822.
 198. Park RJ, Shen H, Liu L, Liu X, Ferguson SM, De Camilli P. Dynamin triple knockout cells reveal off target effects of commonly used dynamin inhibitors. *Journal of cell science* 2013; **126**(Pt 22): 5305-5312.
 199. Hosono R, Hekimi S, Kamiya Y, Sassa T, Murakami S, Nishiwaki K, Miwa J, Taketo A, Kodaira KI. The unc-18 gene encodes a novel protein affecting the kinetics of acetylcholine metabolism in the nematode *Caenorhabditis elegans*. *Journal of neurochemistry* 1992; **58**(4): 1517-1525.
 200. Bouwman J, Maia AS, Camoletto PG, Posthuma G, Roubos EW, Oorschot VM, Klumperman J, Verhage M. Quantification of synapse formation and maintenance in vivo in the absence of synaptic release. *Neuroscience* 2004; **126**(1): 115-126.
 201. Patzke C, Han Y, Covy J, Yi F, Maxeiner S, Wernig M, Sudhof TC. Analysis of conditional heterozygous STXBP1 mutations in human neurons. *J Clin Invest* 2015; **125**(9): 3560-3571.
 202. Barnstable CJ, Hofstein R, Akagawa K. A marker of early amacrine cell development in rat retina. *Brain research* 1985; **352**(2): 286-290.
 203. Bennett MK, Calakos N, Scheller RH. Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science* 1992; **257**(5067): 255-259.
 204. Blasi J, Chapman ER, Yamasaki S, Binz T, Niemann H, Jahn R. Botulinum neurotoxin C1 blocks neurotransmitter release by means of cleaving HPC-1/syntaxin. *The EMBO journal* 1993; **12**(12): 4821-4828.
 205. Kushima Y, Fujiwara T, Sanada M, Akagawa K. Characterization of HPC-1 antigen, an isoform of syntaxin-1, with the isoform-specific monoclonal antibody, 14D8. *Journal of molecular neuroscience : MN* 1997; **8**(1): 19-27.
 206. Noakes PG, Chin D, Kim SS, Liang S, Phillips WD. Expression and localisation of dynamin and syntaxin during neural development and neuromuscular synapse formation. *The Journal of comparative neurology* 1999; **410**(4): 531-540.
 207. Miya F, Yamamoto A, Akagawa K, Kawamoto K, Tashiro Y. Localization of HPC-1/syntaxin 1 in developing rat cerebellar cortex. *Cell Struct Funct* 1996; **21**(6): 525-532.
 208. Aalto MK, Ronne H, Keranen S. Yeast syntaxins Sso1p and Sso2p belong to a family of related membrane proteins that function in vesicular transport. *The EMBO journal* 1993; **12**(11): 4095-4104.
 209. Bennett MK, Garcia-Ararras JE, Elferink LA, Peterson K, Fleming AM, Hazuka CD, Scheller RH. The syntaxin family of vesicular transport receptors. *Cell* 1993; **74**(5): 863-873.
 210. Schulze KL, Broadie K, Perin MS, Bellen HJ. Genetic and electrophysiological studies of *Drosophila* syntaxin-1A demonstrate its role in nonneuronal secretion and neurotransmission. *Cell* 1995; **80**(2): 311-320.
 211. Broadie K, Prokop A, Bellen HJ, O'Kane CJ, Schulze KL, Sweeney ST. Syntaxin and synaptobrevin function downstream of vesicle docking in *Drosophila*. *Neuron* 1995; **15**(3): 663-673.
 212. Gerber SH, Rah JC, Min SW, Liu X, de Wit H, Dulubova I, Meyer AC, Rizo J, Arancillo M, Hammer RE, Verhage M, Rosenmund C, Sudhof TC. Conformational switch of syntaxin-1 controls synaptic vesicle fusion. *Science* 2008; **321**(5895): 1507-1510.
 213. Oyler GA, Polli JW, Wilson MC, Billingsley ML. Developmental expression of the 25-kDa synaptosomal-associated protein (SNAP-25) in rat brain. *Proceedings of the National Academy of Sciences of the United States of America* 1991; **88**(12): 5247-5251.
 214. Tafoya LC, Mameli M, Miyashita T, Guzowski JF, Valenzuela CF, Wilson MC. Expression and function of SNAP-25 as a universal SNARE component in GABAergic neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2006; **26**(30): 7826-7838.

215. Bronk P, Deak F, Wilson MC, Liu X, Sudhof TC, Kavalali ET. Differential effects of SNAP-25 deletion on Ca²⁺-dependent and Ca²⁺-independent neurotransmission. *Journal of neurophysiology* 2007; **98**(2): 794-806.
216. Bark IC. Structure of the chicken gene for SNAP-25 reveals duplicated exon encoding distinct isoforms of the protein. *J Mol Biol* 1993; **233**(1): 67-76.
217. Ravichandran V, Chawla A, Roche PA. Identification of a novel syntaxin- and synaptobrevin/VAMP-binding protein, SNAP-23, expressed in non-neuronal tissues. *The Journal of biological chemistry* 1996; **271**(23): 13300-13303.
218. Schiavo G, Matteoli M, Montecucco C. Neurotoxins affecting neuroexocytosis. *Physiological reviews* 2000; **80**(2): 717-766.
219. Williamson LC, Neale EA. Syntaxin and 25-kDa synaptosomal-associated protein: differential effects of botulinum neurotoxins C1 and A on neuronal survival. *Journal of neuroscience research* 1998; **52**(5): 569-583.
220. Berliocchi L, Fava E, Leist M, Horvat V, Dinsdale D, Read D, Nicotera P. Botulinum neurotoxin C initiates two different programs for neurite degeneration and neuronal apoptosis. *The Journal of cell biology* 2005; **168**(4): 607-618.
221. Peng L, Liu H, Ruan H, Tepp WH, Stoothoff WH, Brown RH, Johnson EA, Yao WD, Zhang SC, Dong M. Cytotoxicity of botulinum neurotoxins reveals a direct role of syntaxin 1 and SNAP-25 in neuron survival. *Nature communications* 2013; **4**: 1472.
222. Schiavo G, Poulain B, Rossetto O, Benfenati F, Tauc L, Montecucco C. Tetanus toxin is a zinc protein and its inhibition of neurotransmitter release and protease activity depend on zinc. *The EMBO journal* 1992; **11**(10): 3577-3583.
223. Schiavo G, Benfenati F, Poulain B, Rossetto O, Polverino de Lauro P, DasGupta BR, Montecucco C. Tetanus and botulinum-B neurotoxins block neurotransmitter release by proteolytic cleavage of synaptobrevin. *Nature* 1992; **359**(6398): 832-835.
224. Morrison R, Kinoshita Y, Johnson M, Ghatan S, Joseph. Neuronal Survival and Cell Death Signaling Pathways. In: Databa MCB (ed). Landes Bioscience; Austin (TX), 2000.
225. Davies CA, Mann DM, Sumpter PQ, Yates PO. A quantitative morphometric analysis of the neuronal and synaptic content of the frontal and temporal cortex in patients with Alzheimer's disease. *Journal of the neurological sciences* 1987; **78**(2): 151-164.
226. Guegan C, Przedborski S. Programmed cell death in amyotrophic lateral sclerosis. *J Clin Invest* 2003; **111**(2): 153-161.
227. Kurokawa Y. [Binding and degenerative effect of Clostridium botulinum type A, C and E toxins on cultured neurons]. *Hokkaido Igaku Zasshi* 1987; **62**(3): 381-391.
228. Zhao LC, Yang B, Wang R, Lipton SA, Zhang D. Type C botulinum toxin causes degeneration of motoneurons in vivo. *Neuroreport* 2010; **21**(1): 14-18.
229. McEwen JM, Kaplan JM. UNC-18 promotes both the anterograde trafficking and synaptic function of syntaxin. *Molecular biology of the cell* 2008; **19**(9): 3836-3846.
230. Farina M, van de Bospoort R, He E, Persoon CM, van Weering JR, Broeke JH, Verhage M, Toonen RF. CAPS-1 promotes fusion competence of stationary dense-core vesicles in presynaptic terminals of mammalian neurons. *Elife* 2015; **4**.
231. Villeneuve J, Scarpa M, Ortega-Bellido M, Malhotra V. MEK1 inactivates Myt1 to regulate Golgi membrane fragmentation and mitotic entry in mammalian cells. *The EMBO journal* 2013; **32**(1): 72-85.
232. Burkel BM, von Dassow G, Bement WM. Versatile fluorescent probes for actin filaments based on the actin-binding domain of utrophin. *Cell motility and the cytoskeleton* 2007; **64**(11): 822-832.
233. Link E, Edelmann L, Chou JH, Binz T, Yamasaki S, Eisel U, Baumert M, Sudhof TC, Niemann H, Jahn R. Tetanus toxin action: inhibition of neurotransmitter release linked to synaptobrevin proteolysis. *Biochemical and biophysical research communications* 1992; **189**(2): 1017-1023.
234. Link E, McMahon H, Fischer von Mollard G, Yamasaki S, Niemann H, Sudhof TC, Jahn R. Cleavage of cellubrevin by tetanus toxin does not affect fusion of early endosomes. *The Journal of biological chemistry* 1993; **268**(25): 18423-18426.
235. Schiavo G, Shone CC, Bennett MK, Scheller RH, Montecucco C. Botulinum neurotoxin type C cleaves a single Lys-Ala bond within the carboxyl-terminal region of syntaxins. *The Journal of biological chemistry* 1995; **270**(18): 10566-10570.
236. Vaidyanathan VV, Yoshino K, Jahnz M, Dorries C, Bade S, Nauenburg S, Niemann H, Binz T. Proteolysis of SNAP-25 isoforms by botulinum neurotoxin types A, C, and E: domains and amino acid residues controlling the formation of enzyme-substrate complexes and cleavage. *Journal of neurochemistry* 1999; **72**(1): 327-337.
237. Foran P, Lawrence GW, Shone CC, Foster KA, Dolly JO. Botulinum neurotoxin C1 cleaves both syntaxin and

- SNAP-25 in intact and permeabilized chromaffin cells: correlation with its blockade of catecholamine release. *Biochemistry* 1996; **35**(8): 2630-2636.
238. Osen-Sand A, Staple JK, Naldi E, Schiavo G, Rossetto O, Petitpierre S, Malgaroli A, Montecucco C, Catsicas S. Common and distinct fusion proteins in axonal growth and transmitter release. *The Journal of comparative neurology* 1996; **367**(2): 222-234.
 239. Galli T, Zahraoui A, Vaidyanathan VV, Raposo G, Tian JM, Karin M, Niemann H, Louvard D. A novel tetanus neurotoxin-insensitive vesicle-associated membrane protein in SNARE complexes of the apical plasma membrane of epithelial cells. *Molecular biology of the cell* 1998; **9**(6): 1437-1448.
 240. Koh JY, Wie MB, Gwag BJ, Sensi SL, Canzoniero LM, Demaro J, Csernansky C, Choi DW. Staurosporine-induced neuronal apoptosis. *Experimental neurology* 1995; **135**(2): 153-159.
 241. van de Bospoort R, Farina M, Schmitz SK, de Jong A, de Wit H, Verhage M, Toonen RF. Munc13 controls the location and efficiency of dense-core vesicle release in neurons. *The Journal of cell biology* 2012; **199**(6): 883-891.
 242. Wierda KD, Toonen RF, de Wit H, Brussaard AB, Verhage M. Interdependence of PKC-dependent and PKC-independent pathways for presynaptic plasticity. *Neuron* 2007; **54**(2): 275-290.
 243. Rosenmund C, Stevens CF. Definition of the readily releasable pool of vesicles at hippocampal synapses. *Neuron* 1996; **16**(6): 1197-1207.
 244. Gulyas-Kovacs A, de Wit H, Milosevic I, Kochubey O, Toonen R, Klingauf J, Verhage M, Sorensen JB. Munc18-1: sequential interactions with the fusion machinery stimulate vesicle docking and priming. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2007; **27**(32): 8676-8686.
 245. de Wit H, Walter AM, Milosevic I, Gulyas-Kovacs A, Riedel D, Sorensen JB, Verhage M. Synaptotagmin-1 docks secretory vesicles to syntaxin-1/SNAP-25 acceptor complexes. *Cell* 2009; **138**(5): 935-946.
 246. Blasi J, Chapman ER, Link E, Binz T, Yamasaki S, De Camilli P, Sudhof TC, Niemann H, Jahn R. Botulinum neurotoxin A selectively cleaves the synaptic protein SNAP-25. *Nature* 1993; **365**(6442): 160-163.
 247. Hu SH, Christie MP, Saez NJ, Latham CF, Jarrott R, Lua LH, Collins BM, Martin JL. Possible roles for Munc18-1 domain 3a and Syntaxin1 N-peptide and C-terminal anchor in SNARE complex formation. *Proceedings of the National Academy of Sciences of the United States of America* 2011; **108**(3): 1040-1045.
 248. Hu SH, Latham CF, Gee CL, James DE, Martin JL. Structure of the Munc18c/Syntaxin4 N-peptide complex defines universal features of the N-peptide binding mode of Sec1/Munc18 proteins. *Proceedings of the National Academy of Sciences of the United States of America* 2007; **104**(21): 8773-8778.
 249. Tellam JT, Macaulay SL, McIntosh S, Hewish DR, Ward CW, James DE. Characterization of Munc-18c and syntaxin-4 in 3T3-L1 adipocytes. Putative role in insulin-dependent movement of GLUT-4. *The Journal of biological chemistry* 1997; **272**(10): 6179-6186.
 250. Khvotchev M, Sudhof TC. Proteolytic processing of amyloid-beta precursor protein by secretases does not require cell surface transport. *The Journal of biological chemistry* 2004; **279**(45): 47101-47108.
 251. Warren G. Membrane partitioning during cell division. *Annu Rev Biochem* 1993; **62**: 323-348.
 252. Valderrama F, Babia T, Ayala I, Kok JW, Renau-Piqueras J, Egea G. Actin microfilaments are essential for the cytoskeletal positioning and morphology of the Golgi complex. *European journal of cell biology* 1998; **76**(1): 9-17.
 253. De Matteis MA, Wilson C, D'Angelo G. Phosphatidylinositol-4-phosphate: the golgi and beyond. *BioEssays : news and reviews in molecular, cellular and developmental biology* 2013; **35**(7): 612-622.
 254. Godi A, Pertile P, Meyers R, Marra P, Di Tullio G, Iurisci C, Luini A, Corda D, De Matteis MA. ARF mediates recruitment of PtdIns-4-OH kinase-beta and stimulates synthesis of PtdIns(4,5)P2 on the Golgi complex. *Nature cell biology* 1999; **1**(5): 280-287.
 255. Balla A, Tuymetova G, Barshishat M, Geiszt M, Balla T. Characterization of type II phosphatidylinositol 4-kinase isoforms reveals association of the enzymes with endosomal vesicular compartments. *The Journal of biological chemistry* 2002; **277**(22): 20041-20050.
 256. Barylko B, Gerber SH, Binns DD, Grichine N, Khvotchev M, Sudhof TC, Albanesi JP. A novel family of phosphatidylinositol 4-kinases conserved from yeast to humans. *The Journal of biological chemistry* 2001; **276**(11): 7705-7708.
 257. Audhya A, Foti M, Emr SD. Distinct roles for the yeast phosphatidylinositol 4-kinases, Stt4p and Pik1p, in secretion, cell growth, and organelle membrane dynamics. *Molecular biology of the cell* 2000; **11**(8): 2673-2689.
 258. Tellam JT, James DE, Stevens TH, Piper RC. Identification of a mammalian Golgi Sec1p-like protein, mVps45. *The Journal of biological chemistry* 1997; **272**(10): 6187-6193.
 259. Rahajeng J, Caplan S, Naslavsky N. Common and distinct roles for the binding partners Rabenosyn-5 and Vps45 in the regulation of endocytic trafficking in mammalian cells. *Experimental cell research* 2010; **316**(5):

- 859-874.
260. Hammond GR, Machner MP, Balla T. A novel probe for phosphatidylinositol 4-phosphate reveals multiple pools beyond the Golgi. *The Journal of cell biology* 2014; **205**(1): 113-126.
 261. Nakatsu F, Baskin JM, Chung J, Tanner LB, Shui G, Lee SY, Pirruccello M, Hao M, Ingolia NT, Wenk MR, De Camilli P. PtdIns4P synthesis by PI4KIIIalpha at the plasma membrane and its impact on plasma membrane identity. *The Journal of cell biology* 2012; **199**(6): 1003-1016.
 262. Brombacher E, Urwyler S, Ragaz C, Weber SS, Kami K, Overduin M, Hilbi H. Rab1 guanine nucleotide exchange factor SidM is a major phosphatidylinositol 4-phosphate-binding effector protein of *Legionella pneumophila*. *The Journal of biological chemistry* 2009; **284**(8): 4846-4856.
 263. Nasuhoglu C, Feng S, Mao J, Yamamoto M, Yin HL, Earnest S, Barylko B, Albanesi JP, Hilgemann DW. Nonradioactive analysis of phosphatidylinositides and other anionic phospholipids by anion-exchange high-performance liquid chromatography with suppressed conductivity detection. *Anal Biochem* 2002; **301**(2): 243-254.
 264. Minin AA. Dispersal of Golgi apparatus in nocodazole-treated fibroblasts is a kinesin-driven process. *Journal of cell science* 1997; **110** (Pt 19): 2495-2505.
 265. Ceridono M, Ory S, Momboisse F, Chasserot-Golaz S, Houy S, Calco V, Haeberle AM, Demais V, Bailly Y, Bader MF, Gasman S. Selective recapture of secretory granule components after full collapse exocytosis in neuroendocrine chromaffin cells. *Traffic* 2011; **12**(1): 72-88.
 266. Cramer LP. Myosin VI: roles for a minus end-directed actin motor in cells. *The Journal of cell biology* 2000; **150**(6): F121-126.
 267. Allan VJ, Thompson HM, McNiven MA. Motoring around the Golgi. *Nature cell biology* 2002; **4**(10): E236-242.
 268. Vogel GF, Klee KM, Janecke AR, Muller T, Hess MW, Huber LA. Cargo-selective apical exocytosis in epithelial cells is conducted by Myo5B, Slp4a, Vamp7, and Syntaxin 3. *The Journal of cell biology* 2015; **211**(3): 587-604.
 269. Bryant NJ, James DE. Vps45p stabilizes the syntaxin homologue Tlg2p and positively regulates SNARE complex formation. *The EMBO journal* 2001; **20**(13): 3380-3388.
 270. Kurps J, de Wit H. The role of Munc18-1 and its orthologs in modulation of cortical F-actin in chromaffin cells. *Journal of molecular neuroscience : MN* 2012; **48**(2): 339-346.
 271. Nichols BJ, Holthuis JC, Pelham HR. The Sec1p homologue Vps45p binds to the syntaxin Tlg2p. *European journal of cell biology* 1998; **77**(4): 263-268.
 272. Dulubova I, Yamaguchi T, Gao Y, Min SW, Huryeva I, Sudhof TC, Rizo J. How Tlg2p/syntaxin 16 'snares' Vps45. *The EMBO journal* 2002; **21**(14): 3620-3631.
 273. Webb GC, Hoedt M, Poole LJ, Jones EW. Genetic interactions between a pep7 mutation and the PEP12 and VPS45 genes: evidence for a novel SNARE component in transport between the *Saccharomyces cerevisiae* Golgi complex and endosome. *Genetics* 1997; **147**(2): 467-478.
 274. Kirkbride KC, Hong NH, French CL, Clark ES, Jerome WG, Weaver AM. Regulation of late endosomal/lysosomal maturation and trafficking by cortactin affects Golgi morphology. *Cytoskeleton (Hoboken)* 2012; **69**(9): 625-643.
 275. Wong K, Meyers dd R, Cantley LC. Subcellular locations of phosphatidylinositol 4-kinase isoforms. *The Journal of biological chemistry* 1997; **272**(20): 13236-13241.
 276. Fruman DA, Meyers RE, Cantley LC. Phosphoinositide kinases. *Annu Rev Biochem* 1998; **67**: 481-507.
 277. Frere SG, Chang-Ileto B, Di Paolo G. Role of phosphoinositides at the neuronal synapse. *Subcell Biochem* 2012; **59**: 131-175.
 278. Lemmon MA. Membrane recognition by phospholipid-binding domains. *Nature reviews Molecular cell biology* 2008; **9**(2): 99-111.
 279. Raucher D, Stauffer T, Chen W, Shen K, Guo S, York JD, Sheetz MP, Meyer T. Phosphatidylinositol 4,5-bisphosphate functions as a second messenger that regulates cytoskeleton-plasma membrane adhesion. *Cell* 2000; **100**(2): 221-228.
 280. Eberhard DA, Cooper CL, Low MG, Holz RW. Evidence that the inositol phospholipids are necessary for exocytosis. Loss of inositol phospholipids and inhibition of secretion in permeabilized cells caused by a bacterial phospholipase C and removal of ATP. *The Biochemical journal* 1990; **268**(1): 15-25.
 281. Kutateladze TG. Translation of the phosphoinositide code by PI effectors. *Nat Chem Biol* 2010; **6**(7): 507-513.
 282. Tanaka K, Adachi H, Konishi H, Iwamatsu A, Ohkawa K, Shirai T, Nagata S, Kikkawa U, Fukui Y. Identification of protein kinase B (PKB) as a phosphatidylinositol 3,4,5-trisphosphate binding protein in *Dictyostelium discoideum*. *Biosci Biotechnol Biochem* 1999; **63**(2): 368-372.
 283. Meier R, Hemmings BA. Regulation of protein kinase B. *J Recept Signal Transduct Res* 1999; **19**(1-4): 121-128.
 284. Kulik G, Klippel A, Weber MJ. Antiapoptotic signalling by the insulin-like growth factor I receptor,

- phosphatidylinositol 3-kinase, and Akt. *Molecular and cellular biology* 1997; **17**(3): 1595-1606.
285. Dudek H, Datta SR, Franke TF, Birnbaum MJ, Yao R, Cooper GM, Segal RA, Kaplan DR, Greenberg ME. Regulation of neuronal survival by the serine-threonine protein kinase Akt. *Science* 1997; **275**(5300): 661-665.
 286. Manning BD, Cantley LC. AKT/PKB signaling: navigating downstream. *Cell* 2007; **129**(7): 1261-1274.
 287. Franke TF, Hornik CP, Segev L, Shostak GA, Sugimoto C. PI3K/Akt and apoptosis: size matters. *Oncogene* 2003; **22**(56): 8983-8998.
 288. Plas DR, Thompson CB. Cell metabolism in the regulation of programmed cell death. *Trends Endocrinol Metab* 2002; **13**(2): 75-78.
 289. van den Bogaart G, Meyenberg K, Diederichsen U, Jahn R. Phosphatidylinositol 4,5-bisphosphate increases Ca²⁺ affinity of synaptotagmin-1 by 40-fold. *The Journal of biological chemistry* 2012; **287**(20): 16447-16453.
 290. Aoyagi K, Sugaya T, Umeda M, Yamamoto S, Terakawa S, Takahashi M. The activation of exocytotic sites by the formation of phosphatidylinositol 4,5-bisphosphate microdomains at syntaxin clusters. *The Journal of biological chemistry* 2005; **280**(17): 17346-17352.
 291. Khuong TM, Habets RL, Kuenen S, Witkowska A, Kaspruwicz J, Swerts J, Jahn R, van den Bogaart G, Verstreken P. Synaptic PI(3,4,5)P₃ is required for Syntaxin1A clustering and neurotransmitter release. *Neuron* 2013; **77**(6): 1097-1108.
 292. Dudok JJ, Groffen AJ, Toonen RF, Verhage M. Deletion of Munc18-1 in 5-HT neurons results in rapid degeneration of the 5-HT system and early postnatal lethality. *PLoS one* 2011; **6**(11): e28137.
 293. Milosevic I, Sorensen JB, Lang T, Krauss M, Nagy G, Haucke V, Jahn R, Neher E. Plasmalemmal phosphatidylinositol-4,5-bisphosphate level regulates the releasable vesicle pool size in chromaffin cells. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2005; **25**(10): 2557-2565.
 294. Lemmon MA, Ferguson KM, O'Brien R, Sigler PB, Schlessinger J. Specific and high-affinity binding of inositol phosphates to an isolated pleckstrin homology domain. *Proceedings of the National Academy of Sciences of the United States of America* 1995; **92**(23): 10472-10476.
 295. Varnai P, Balla T. Visualization of phosphoinositides that bind pleckstrin homology domains: calcium- and agonist-induced dynamic changes and relationship to myo-[³H]inositol-labeled phosphoinositide pools. *The Journal of cell biology* 1998; **143**(2): 501-510.
 296. Stauffer TP, Ahn S, Meyer T. Receptor-induced transient reduction in plasma membrane PtdIns(4,5)P₂ concentration monitored in living cells. *Current biology : CB* 1998; **8**(6): 343-346.
 297. Oatey PB, Venkateswarlu K, Williams AG, Fletcher LM, Foulstone EJ, Cullen PJ, Tavaré JM. Confocal imaging of the subcellular distribution of phosphatidylinositol 3,4,5-trisphosphate in insulin- and PDGF-stimulated 3T3-L1 adipocytes. *The Biochemical journal* 1999; **344 Pt 2**: 511-518.
 298. Klarlund JK, Guilherme A, Holik JJ, Virbasius JV, Chawla A, Czech MP. Signaling by phosphoinositide-3,4,5-trisphosphate through proteins containing pleckstrin and Sec7 homology domains. *Science* 1997; **275**(5308): 1927-1930.
 299. Gray A, Van Der Kaay J, Downes CP. The pleckstrin homology domains of protein kinase B and GRP1 (general receptor for phosphoinositides-1) are sensitive and selective probes for the cellular detection of phosphatidylinositol 3,4-bisphosphate and/or phosphatidylinositol 3,4,5-trisphosphate in vivo. *The Biochemical journal* 1999; **344 Pt 3**: 929-936.
 300. Bravo-Cordero JJ, Magalhaes MA, Eddy RJ, Hodgson L, Condeelis J. Functions of cofilin in cell locomotion and invasion. *Nature reviews Molecular cell biology* 2013; **14**(7): 405-415.
 301. Witke W. The role of profilin complexes in cell motility and other cellular processes. *Trends in cell biology* 2004; **14**(8): 461-469.
 302. Daly RJ. Cortactin signalling and dynamic actin networks. *The Biochemical journal* 2004; **382**(Pt 1): 13-25.
 303. Ferguson SM, De Camilli P. Dynamin, a membrane-remodelling GTPase. *Nature reviews Molecular cell biology* 2012; **13**(2): 75-88.
 304. Rauch ME, Ferguson CG, Prestwich GD, Cafiso DS. Myristoylated alanine-rich C kinase substrate (MARCKS) sequesters spin-labeled phosphatidylinositol 4,5-bisphosphate in lipid bilayers. *The Journal of biological chemistry* 2002; **277**(16): 14068-14076.
 305. Neisch AL, Fehon RG. Ezrin, Radixin and Moesin: key regulators of membrane-cortex interactions and signaling. *Current opinion in cell biology* 2011; **23**(4): 377-382.
 306. Hemmings BA, Restuccia DF. PI3K-PKB/Akt pathway. *Cold Spring Harbor perspectives in biology* 2012; **4**(9): a011189.
 307. James DJ, Khodthong C, Kowalchuk JA, Martin TF. Phosphatidylinositol 4,5-bisphosphate regulates SNARE-dependent membrane fusion. *The Journal of cell biology* 2008; **182**(2): 355-366.
 308. Horiguchi K, Hanada T, Fukui Y, Chishti AH. Transport of PIP₃ by GAKIN, a kinesin-3 family protein,

- regulates neuronal cell polarity. *The Journal of cell biology* 2006; **174**(3): 425-436.
309. Menager C, Arimura N, Fukata Y, Kaibuchi K. PIP3 is involved in neuronal polarization and axon formation. *Journal of neurochemistry* 2004; **89**(1): 109-118.
 310. Ohtsu M, Sakai N, Fujita H, Kashiwagi M, Gasa S, Shimizu S, Eguchi Y, Tsujimoto Y, Sakiyama Y, Kobayashi K, Kuzumaki N. Inhibition of apoptosis by the actin-regulatory protein gelsolin. *The EMBO journal* 1997; **16**(15): 4650-4656.
 311. Geng YJ, Azuma T, Tang JX, Hartwig JH, Muszynski M, Wu Q, Libby P, Kwiatkowski DJ. Caspase-3-induced gelsolin fragmentation contributes to actin cytoskeletal collapse, nucleolysis, and apoptosis of vascular smooth muscle cells exposed to proinflammatory cytokines. *European journal of cell biology* 1998; **77**(4): 294-302.
 312. Di Paolo G, Moskowitz HS, Gipson K, Wenk MR, Voronov S, Obayashi M, Flavell R, Fitzsimonds RM, Ryan TA, De Camilli P. Impaired PtdIns(4,5)P₂ synthesis in nerve terminals produces defects in synaptic vesicle trafficking. *Nature* 2004; **431**(7007): 415-422.
 313. Achiriloaie M, Barylko B, Albanesi JP. Essential role of the dynamin pleckstrin homology domain in receptor-mediated endocytosis. *Molecular and cellular biology* 1999; **19**(2): 1410-1415.
 314. Zoncu R, Perera RM, Sebastian R, Nakatsu F, Chen H, Balla T, Ayala G, Toomre D, De Camilli PV. Loss of endocytic clathrin-coated pits upon acute depletion of phosphatidylinositol 4,5-bisphosphate. *Proceedings of the National Academy of Sciences of the United States of America* 2007; **104**(10): 3793-3798.
 315. Vallis Y, Wigge P, Marks B, Evans PR, McMahon HT. Importance of the pleckstrin homology domain of dynamin in clathrin-mediated endocytosis. *Current biology : CB* 1999; **9**(5): 257-260.
 316. Stocker H, Andjelkovic M, Oldham S, Laffargue M, Wymann MP, Hemmings BA, Hafen E. Living with lethal PIP3 levels: viability of flies lacking PTEN restored by a PH domain mutation in Akt/PKB. *Science* 2002; **295**(5562): 2088-2091.
 317. Watton SJ, Downward J. Akt/PKB localisation and 3' phosphoinositide generation at sites of epithelial cell-matrix and cell-cell interaction. *Current biology : CB* 1999; **9**(8): 433-436.
 318. Stokoe D, Stephens LR, Copeland T, Gaffney PR, Reese CB, Painter GF, Holmes AB, McCormick F, Hawkins PT. Dual role of phosphatidylinositol-3,4,5-trisphosphate in the activation of protein kinase B. *Science* 1997; **277**(5325): 567-570.
 319. Seshacharyulu P, Pandey P, Datta K, Batra SK. Phosphatase: PP2A structural importance, regulation and its aberrant expression in cancer. *Cancer letters* 2013; **335**(1): 9-18.
 320. Andjelkovic M, Jakubowicz T, Cron P, Ming XF, Han JW, Hemmings BA. Activation and phosphorylation of a pleckstrin homology domain containing protein kinase (RAC-PK/PKB) promoted by serum and protein phosphatase inhibitors. *Proceedings of the National Academy of Sciences of the United States of America* 1996; **93**(12): 5699-5704.
 321. Dan HC, Sun M, Kaneko S, Feldman RI, Nicosia SV, Wang HG, Tsang BK, Cheng JQ. Akt phosphorylation and stabilization of X-linked inhibitor of apoptosis protein (XIAP). *The Journal of biological chemistry* 2004; **279**(7): 5405-5412.
 322. Hussain I, Powell D, Howlett DR, Tew DG, Meek TD, Chapman C, Gloger IS, Murphy KE, Southan CD, Ryan DM, Smith TS, Simmons DL, Walsh FS, Dingwall C, Christie G. Identification of a novel aspartic protease (Asp 2) as beta-secretase. *Mol Cell Neurosci* 1999; **14**(6): 419-427.
 323. Yan R, Bienkowski MJ, Shuck ME, Miao H, Tory MC, Pauley AM, Brashier JR, Stratman NC, Mathews WR, Buhl AE, Carter DB, Tomasselli AG, Parodi LA, Heinrikson RL, Gurney ME. Membrane-anchored aspartyl protease with Alzheimer's disease beta-secretase activity. *Nature* 1999; **402**(6761): 533-537.
 324. Anderson JP, Esch FS, Keim PS, Sambamurti K, Lieberburg I, Robakis NK. Exact cleavage site of Alzheimer amyloid precursor in neuronal PC-12 cells. *Neurosci Lett* 1991; **128**(1): 126-128.
 325. Wang R, Meschia JF, Cotter RJ, Sisodia SS. Secretion of the beta/A4 amyloid precursor protein. Identification of a cleavage site in cultured mammalian cells. *The Journal of biological chemistry* 1991; **266**(25): 16960-16964.
 326. Ho CS, Marinescu V, Steinhilb ML, Gaut JR, Turner RS, Stuenkel EL. Synergistic effects of Munc18a and X11 proteins on amyloid precursor protein metabolism. *The Journal of biological chemistry* 2002; **277**(30): 27021-27028.
 327. Borg JP, Straight SW, Kaech SM, de Taddeo-Borg M, Kroon DE, Karnak D, Turner RS, Kim SK, Margolis B. Identification of an evolutionarily conserved heterotrimeric protein complex involved in protein targeting. *The Journal of biological chemistry* 1998; **273**(48): 31633-31636.
 328. Ishiguro K, Kobayashi S, Omori A, Takamatsu M, Yonekura S, Anzai K, Imahori K, Uchida T. Identification of the 23 kDa subunit of tau protein kinase II as a putative activator of cdk5 in bovine brain. *FEBS letters* 1994; **342**(2): 203-208.
 329. Lew J, Huang QQ, Qi Z, Winkfein RJ, Aebersold R, Hunt T, Wang JH. A brain-specific activator of cyclin-

- dependent kinase 5. *Nature* 1994; **371**(6496): 423-426.
330. Patrick GN, Zukerberg L, Nikolic M, de la Monte S, Dikkes P, Tsai LH. Conversion of p35 to p25 deregulates Cdk5 activity and promotes neurodegeneration. *Nature* 1999; **402**(6762): 615-622.
 331. Tsai LH, Delalle I, Caviness VS, Jr., Chae T, Harlow E. p35 is a neural-specific regulatory subunit of cyclin-dependent kinase 5. *Nature* 1994; **371**(6496): 419-423.
 332. Ramos-Miguel A, Hercher C, Beasley CL, Barr AM, Bayer TA, Falkai P, Leurgans SE, Schneider JA, Bennett DA, Honer WG. Loss of Munc18-1 long splice variant in GABAergic terminals is associated with cognitive decline and increased risk of dementia in a community sample. *Mol Neurodegener* 2015; **10**: 65.
 333. Milh M, Villeneuve N, Chouchane M, Kaminska A, Laroche C, Barthez MA, Gitiaux C, Bartoli C, Borges-Correia A, Cacciagli P, Mignon-Ravix C, Cuberos H, Chabrol B, Villard L. Epileptic and nonepileptic features in patients with early onset epileptic encephalopathy and STXBP1 mutations. *Epilepsia* 2011; **52**(10): 1828-1834.
 334. Okamoto M, Sudhof TC. Mints, Munc18-interacting proteins in synaptic vesicle exocytosis. *The Journal of biological chemistry* 1997; **272**(50): 31459-31464.
 335. Tomita S, Ozaki T, Taru H, Oguchi S, Takeda S, Yagi Y, Sakiyama S, Kirino Y, Suzuki T. Interaction of a neuron-specific protein containing PDZ domains with Alzheimer's amyloid precursor protein. *The Journal of biological chemistry* 1999; **274**(4): 2243-2254.
 336. Okamoto M, Sudhof TC. Mint 3: a ubiquitous mint isoform that does not bind to munc18-1 or -2. *European journal of cell biology* 1998; **77**(3): 161-165.
 337. Borg JP, Ooi J, Levy E, Margolis B. The phosphotyrosine interaction domains of X11 and FE65 bind to distinct sites on the YENPTY motif of amyloid precursor protein. *Molecular and cellular biology* 1996; **16**(11): 6229-6241.
 338. Mueller HT, Borg JP, Margolis B, Turner RS. Modulation of amyloid precursor protein metabolism by X11alpha/Mint-1. A deletion analysis of protein-protein interaction domains. *The Journal of biological chemistry* 2000; **275**(50): 39302-39306.
 339. Shuang R, Zhang L, Fletcher A, Groblewski GE, Pevsner J, Stuenkel EL. Regulation of Munc-18/syntaxin 1A interaction by cyclin-dependent kinase 5 in nerve endings. *The Journal of biological chemistry* 1998; **273**(9): 4957-4966.
 340. Ho A, Liu X, Sudhof TC. Deletion of Mint proteins decreases amyloid production in transgenic mouse models of Alzheimer's disease. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2008; **28**(53): 14392-14400.
 341. Haass C, Lemere CA, Capell A, Citron M, Seubert P, Schenk D, Lannfelt L, Selkoe DJ. The Swedish mutation causes early-onset Alzheimer's disease by beta-secretase cleavage within the secretory pathway. *Nat Med* 1995; **1**(12): 1291-1296.
 342. Bauereiss A, Welzel O, Jung J, Grosse-Holz S, Lelental N, Lewczuk P, Wenzel EM, Kornhuber J, Groemer TW. Surface Trafficking of APP and BACE in Live Cells. *Traffic* 2015; **16**(6): 655-675.
 343. Buggia-Prevot V, Fernandez CG, Riordan S, Vetrivel KS, Roseman J, Waters J, Bindokas VP, Vassar R, Thinakaran G. Axonal BACE1 dynamics and targeting in hippocampal neurons: a role for Rab11 GTPase. *Mol Neurodegener* 2014; **9**: 1.
 344. Das U, Scott DA, Ganguly A, Koo EH, Tang Y, Roy S. Activity-induced convergence of APP and BACE-1 in acidic microdomains via an endocytosis-dependent pathway. *Neuron* 2013; **79**(3): 447-460.
 345. Groemer TW, Thiel CS, Holt M, Riedel D, Hua Y, Huve J, Wilhelm BG, Klingauf J. Amyloid precursor protein is trafficked and secreted via synaptic vesicles. *PLoS one* 2011; **6**(4): e18754.
 346. Cirrito JR, Kang JE, Lee J, Stewart FR, Verges DK, Silverio LM, Bu G, Mennerick S, Holtzman DM. Endocytosis is required for synaptic activity-dependent release of amyloid-beta in vivo. *Neuron* 2008; **58**(1): 42-51.
 347. Eehalt R, Keller P, Haass C, Thiele C, Simons K. Amyloidogenic processing of the Alzheimer beta-amyloid precursor protein depends on lipid rafts. *The Journal of cell biology* 2003; **160**(1): 113-123.
 348. Kinoshita A, Fukumoto H, Shah T, Whelan CM, Irizarry MC, Hyman BT. Demonstration by FRET of BACE interaction with the amyloid precursor protein at the cell surface and in early endosomes. *Journal of cell science* 2003; **116**(Pt 16): 3339-3346.
 349. van Bospoort R. Molecular mechanisms of neuronal dense core vesicle release. *PhD Thesis* 2013.
 350. Heeroma JH. The role of regulated secretion in neurite outgrowth, synapse formation and neuronal survival (Doctoral Dissertation). 2005.
 351. Binz T, Blasi J, Yamasaki S, Baumeister A, Link E, Sudhof TC, Jahn R, Niemann H. Proteolysis of SNAP-25 by types E and A botulinum neurotoxins. *The Journal of biological chemistry* 1994; **269**(3): 1617-1620.
 352. Bankaitis VA, Garcia-Mata R, Mousley CJ. Golgi membrane dynamics and lipid metabolism. *Current biology : CB* 2012; **22**(10): R414-424.

References

353. Schweizer A, Fransén JA, Matter K, Kreis TE, Ginsel L, Hauri HP. Identification of an intermediate compartment involved in protein transport from endoplasmic reticulum to Golgi apparatus. *European journal of cell biology* 1990; **53**(2): 185-196.
354. Spang A. On vesicle formation and tethering in the ER-Golgi shuttle. *Current opinion in cell biology* 2009; **21**(4): 531-536.
355. Spang A. Retrograde traffic from the Golgi to the endoplasmic reticulum. *Cold Spring Harbor perspectives in biology* 2013; **5**(6).
356. Gaynor EC, Graham TR, Emr SD. COPI in ER/Golgi and intra-Golgi transport: do yeast COPI mutants point the way? *Biochimica et biophysica acta* 1998; **1404**(1-2): 33-51.
357. Euro E-RESC, Epilepsy Phenome/Genome P, Epi KC. De novo mutations in synaptic transmission genes including DNMI1 cause epileptic encephalopathies. *Am J Hum Genet* 2014; **95**(4): 360-370.
358. Saitu H, Kato M, Mizuguchi T, Hamada K, Osaka H, Tohyama J, Uruno K, Kumada S, Nishiyama K, Nishimura A, Okada I, Yoshimura Y, Hirai S, Kumada T, Hayasaka K, Fukuda A, Ogata K, Matsumoto N. De novo mutations in the gene encoding STXBPI (MUNC18-1) cause early infantile epileptic encephalopathy. *Nat Genet* 2008; **40**(6): 782-788.
359. Stamberger H, Nikanorova M, Willemsen MH, Accorsi P, Angriman M, Baier H, Benkel-Herrenbrueck I, Benoit V, Budetta M, Caliebe A, Cantalupo G, Capovilla G, Casara G, Courage C, Deprez M, Destree A, Dilena R, Erasmus CE, Fannemel M, Fjaer R *et al.* STXBPI encephalopathy: A neurodevelopmental disorder including epilepsy. *Neurology* 2016; **86**(10): 954-962.
360. Law C, Schaan Profes M, Levesque M, Kaltschmidt JA, Verhage M, Kania A. Normal Molecular Specification and Neurodegenerative Disease-Like Death of Spinal Neurons Lacking the SNARE-Associated Synaptic Protein Munc18-1. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2016; **36**(2): 561-576.
361. Trovo L, Ahmed T, Callaerts-Vegh Z, Buzzi A, Bagni C, Chuah M, Vandendriessche T, D'Hooge R, Balschun D, Dotti CG. Low hippocampal PI(4,5)P(2) contributes to reduced cognition in old mice as a result of loss of MARCKS. *Nature neuroscience* 2013; **16**(4): 449-455.
362. Saitu H, Kato M, Matsumoto N. Haploinsufficiency of STXBPI and Ohtahara syndrome. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV (eds). *Jasper's Basic Mechanisms of the Epilepsies*, 4th edn: Bethesda (MD), 2012.
363. Ohtahara S, Yamatogi Y. Ohtahara syndrome: with special reference to its developmental aspects for differentiating from early myoclonic encephalopathy. *Epilepsy research* 2006; **70 Suppl 1**: S58-67.
364. Djukic A, Lado FA, Shinnar S, Moshe SL. Are early myoclonic encephalopathy (EME) and the Ohtahara syndrome (EIEE) independent of each other? *Epilepsy research* 2006; **70 Suppl 1**: S68-76.
365. Otsuka M, Oguni H, Liang JS, Ikeda H, Imai K, Hirasawa K, Imai K, Tachikawa E, Shimojima K, Osawa M, Yamamoto T. STXBPI mutations cause not only Ohtahara syndrome but also West syndrome--result of Japanese cohort study. *Epilepsia* 2010; **51**(12): 2449-2452.
366. Yamashita S, Chiyonobu T, Yoshida M, Maeda H, Zuike M, Kidowaki S, Isoda K, Morimoto M, Kato M, Saitu H, Matsumoto N, Nakahata T, Saito MK, Hosoi H. Mislocalization of syntaxin-1 and impaired neurite growth observed in a human iPSC model for STXBPI-related epileptic encephalopathy. *Epilepsia* 2016; **57**(4): e81-86.
367. Toonen RF, Wierda K, Sons MS, de Wit H, Cornelisse LN, Brussaard A, Plomp JJ, Verhage M. Munc18-1 expression levels control synapse recovery by regulating readily releasable pool size. *Proceedings of the National Academy of Sciences of the United States of America* 2006; **103**(48): 18332-18337.
368. Chai YJ, Sierrecki E, Tomatis VM, Gormal RS, Giles N, Morrow IC, Xia D, Gotz J, Parton RG, Collins BM, Gambin Y, Meunier FA. Munc18-1 is a molecular chaperone for alpha-synuclein, controlling its self-replicating aggregation. *The Journal of cell biology* 2016.
369. Wilkaniec A, Czapski GA, Adamczyk A. Cdk5 at crossroads of protein oligomerization in neurodegenerative diseases: facts and hypotheses. *Journal of neurochemistry* 2016; **136**(2): 222-233.
370. Cruz JC, Tseng HC, Goldman JA, Shih H, Tsai LH. Aberrant Cdk5 activation by p25 triggers pathological events leading to neurodegeneration and neurofibrillary tangles. *Neuron* 2003; **40**(3): 471-483.
371. Wen Y, Yu WH, Maloney B, Bailey J, Ma J, Marie I, Maurin T, Wang L, Figueroa H, Herman M, Krishnamurthy P, Liu L, Planel E, Lau LF, Lahiri DK, Duff K. Transcriptional regulation of beta-secretase by p25/cdk5 leads to enhanced amyloidogenic processing. *Neuron* 2008; **57**(5): 680-690.
372. Stokes CE, Hawthorne JN. Reduced phosphoinositide concentrations in anterior temporal cortex of Alzheimer-diseased brains. *Journal of neurochemistry* 1987; **48**(4): 1018-1021.
373. Berman DE, Dall'Armi C, Voronov SV, McIntire LB, Zhang H, Moore AZ, Staniszewski A, Arancio O, Kim TW, Di Paolo G. Oligomeric amyloid-beta peptide disrupts phosphatidylinositol-4,5-bisphosphate metabolism.

- Nature neuroscience* 2008; **11**(5): 547-554.
374. Landman N, Jeong SY, Shin SY, Voronov SV, Serban G, Kang MS, Park MK, Di Paolo G, Chung S, Kim TW. Presenilin mutations linked to familial Alzheimer's disease cause an imbalance in phosphatidylinositol 4,5-bisphosphate metabolism. *Proceedings of the National Academy of Sciences of the United States of America* 2006; **103**(51): 19524-19529.
 375. van den Bogaart G, Meyenberg K, Risselada HJ, Amin H, Willig KI, Hubrich BE, Dier M, Hell SW, Grubmuller H, Diederichsen U, Jahn R. Membrane protein sequestering by ionic protein-lipid interactions. *Nature* 2011; **479**(7374): 552-555.