

VU Research Portal

Sports Injuries

Delfino Barboza, S.

2018

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Delfino Barboza, S. (2018). *Sports Injuries: Monitoring & Prevention*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

CHAPTER 5

INJURIES IN FIELD HOCKEY PLAYERS: A SYSTEMATIC REVIEW

Saulo Delfino Barboza

Corey Joseph

Joske Nauta

Willem van Mechelen

Evert Verhagen

Sports Medicine 2018; 48(4):849–866

Digital object identifier (DOI): 10.1007/s40279-017-0839-3

ABSTRACT

Background

To commence injury prevention efforts, it is necessary to understand the magnitude of the injury problem. No systematic reviews have yet investigated the extent of injuries in field hockey, despite the popularity of the sport worldwide.

Objective

Our objective was to describe the rate and severity of injuries in field hockey and investigate their characteristics.

Methods

We conducted electronic searches in PubMed, EMBASE, SPORTDiscus, and CINAHL. Prospective cohort studies were included if they were published in English in a peer-reviewed journal and observed all possible injuries sustained by field hockey players during the period of the study.

Results

The risk-of-bias score of the 22 studies included ranged from three to nine of a possible ten. In total, 12 studies (55%) reported injuries normalized by field hockey exposure. Injury rates ranged from 0.1 injuries (in school-aged players) to 90.9 injuries (in Africa Cup of Nations) per 1,000 player-hours and from one injury (in high-school women) to 70 injuries (in under-21 age women) per 1,000 player-sessions. Studies used different classifications for injury severity, but – within studies – injuries were included mostly in the less severe category. The lower limbs were most affected, and contusions/hematomas and abrasions were common types of injury. Contact injuries are common, but non-contact injuries are also a cause for concern.

Conclusions

Considerable heterogeneity meant it was not possible to draw conclusive findings on the extent of the rate and severity of injuries. Establishing the extent of sports injury is considered the first step towards prevention, so there is a need for a consensus on injury surveillance in field hockey.

INTRODUCTION

Field hockey is an Olympic sport played by men and women at both recreational and professional levels. The five continental and 132 national associations that are members of the International Hockey Federation¹ demonstrate the high level of popularity of field hockey worldwide. Field hockey participation may contribute to players' health through the well-known benefits of regular exercise. However, participation in field hockey also entails a risk of injury.²

In general, sports injuries result in individual and societal costs,³ hamper performance, and compromise teams' success over the sporting season.^{4,5} Therefore, injury prevention strategies are of great importance for teams at both recreational and professional levels. Establishing the extent of the injury problem is considered the first step towards effective prevention.⁶ In field hockey, as well as in other sports, this information can aid researchers and health professionals in developing appropriate strategies to reduce and control injuries.⁶

To the best of our knowledge, no systematic reviews have provided a synthesis of information on injuries sustained by field hockey players. Systematic reviews involve gathering evidence from different sources to enable a synthesis of what is currently known about a specific topic (e.g. injuries) and may facilitate the link between research evidence and optimal strategies for healthcare.⁷ Therefore, the aim of this study was to systematically review the literature on injuries sustained by field hockey players in order to describe the extent of such injuries in terms of rate and severity, as well as to identify injury characteristics according to body location, type, and mechanism of injury.

METHODS

Information sources and search strategy

Electronic searches were conducted in PubMed, Excerpta Medical Database (EMBASE), SPORTDiscus, and Cumulative Index to Nursing and Allied Health Literature (CINAHL) databases with no limits on the publication date. The search strategy

combined keywords for injury, field hockey, and study design: (((((((((((injur*) OR traum*) OR risk*) OR overuse) OR overload) OR acute) OR odds) OR incidence) OR prevalence) OR hazard)) AND (((field AND hockey)) OR (hockey NOT ice)) AND (((prosp*) OR retrosp*) OR case*). The detailed search strategy for each database can be found in the supplementary material (page 106). The last search was conducted on 31 May 2017.

Eligibility criteria

Studies were eligible for inclusion if they were published in the English language in a peer-reviewed academic journal, were prospective cohort studies, and observed all possible injuries sustained by field hockey players during the period of the study (i.e. studies that looked only at specific injuries were not included). To minimize the possibility of recall bias, only prospective cohort studies were included.^{8,9} Studies were not included if they described field hockey injuries together with those from other sports whether specific data on field hockey could not be distinguished. Conference abstracts were not included.

Study selection and data collection process

Two reviewers (SDB and CJ) independently screened all records identified in the search strategy in two steps: title and abstract screening, and full-text screening. References of full texts were also screened for possible additional studies not identified in the four databases. Conflicts between reviewers' decisions were resolved through discussion. A third reviewer (EV) was consulted for consensus rating when needed.

One reviewer (SDB) extracted the following information from the included studies: first author, publication year, country in which the study was conducted, primary objective, setting, follow-up period, number and description of field hockey players, injury definition, injury data collection procedure, number of injured players, number of injuries sustained by players during the study, and severity of injuries (Table 1). The number of injuries normalized by exposure to field hockey (i.e. injury rate) was also

extracted. In addition, information on injury according to body location, type of injury, mechanism, and player position was gathered whenever possible. When different studies used the same dataset (Table 1), the results of such studies were combined in one row in all other tables for simplicity.

Risk-of-bias assessment

Two independent reviewers (SDB and CJ) assessed the risk of bias in the included studies using ten criteria previously used in systematic reviews on sports injury.^{9,10} All criteria were rated as one (i.e. low risk of bias) or zero (i.e. high risk of bias). When insufficient information was presented in a study to rate a specific criterion as one or zero, the rating was categorized as 'unable to determine' (UD) and counted as zero. The assessment of each reviewer was compared and conflicts were resolved through discussion. The 10 criteria are described in Table 2.

RESULTS

Search results

We retrieved 810 records from the four databases. Of those, 193 were duplicates. After screening 617 titles and abstracts and 21 full texts, 10 studies matched the inclusion criteria. Screening the references of the full texts resulted in 12 additional records. In the end, 22 studies were included in the review. The flowchart of the inclusion process is presented in Figure 1.

Description of the included studies

The characteristics of the 22 studies included in this review are presented in Table 1. Studies included in this review were published between 1975 and 2016, with 12 (55%) published before 2000^{11–22} and 10 (45%) from 2000 onwards.^{23–32} Two studies used the same dataset from the National Athletic Trainers' Association (NATA) High School Injury database,^{21,27} and two used the same dataset from the National Collegiate

Athletic Association (NCAA) Injury Surveillance System.^{26,28} One study²⁴ was the follow-up of a previous study²³.

Six studies (27%) focused on describing field hockey injuries only.^{14,18,19,28,29,32} The other 16 studies (73%) described the epidemiology of injuries in field hockey together with those in other sports.^{11–13,15–17,20–27,30,31} The period of follow-up varied between studies from a six-day championship tournament²⁰ to 15 consecutive seasons of field hockey.²⁸ The sample size varied between 26²² and 5,385 participants.²⁸ However, seven studies (32%) did not report the number of field hockey players studied.^{11,13,14,19,21,25,27}

The definition of injury varied across the studies. Common criteria to define an injury as recordable were a musculoskeletal condition requiring medical attention and/ or leading to field hockey time-loss (Table 1). The proportion (%) of injured players varied from 6% (in seven months of high school) to 33% (in six days of university games). Twelve studies (55%) did not report the number or proportion of players who had sustained an injury over the study period.^{11–15,18,19,25,26,28,29,32}

Risk-of-bias assessment

Table 2 shows the risk-of-bias assessment for the 22 included studies. The total score ranged from three to nine of a possible 10 points. The studies published during and since 2000 scored higher (range 7–9).^{23–31} Three studies (14%) did not provide a clear definition of injury,^{11,15,18} and three did not describe any characteristics of the players studied.^{11,15,19} These studies were published before the year 2000.

Nine studies (41%) included a random sample of players or studied the entire target population.^{12,16,20,23–25,30–32} Eighteen studies (82%) collected injury data directly from players or medical professionals, 17 studies (77%) used only one method (i.e., not multiple methods) to collect injury data during the study,^{11,13,15–20,23–26,28–32} and one study (5%) did not describe the data collection procedure at all.¹⁴

Twelve studies (55%) employed a medical professional to diagnose injuries.^{13,16,17,21,25-32} The follow-up period of 13 studies (59%) was over six months,^{11-14,17-19,21,24,26-29} and 12 studies (55%) expressed ratios that represented both the number of injuries and the exposure to field hockey.^{11-13,21,23-29,32}




Figure 1. Flowchart of the studies during the inclusion process. Electronic searches were conducted in PubMed, Exerpta Medical Database (EMBASE), SPORTDiscus, and Cumulative Index to Nursing and Allied Health Literature (CINAHL) databases with no limits on the publication date.

Table 1. Characteristics of prospective studies on field hockey injuries arranged by year of publication (least recent to most recent).

Study (country)	Primary objective	Setting and follow-up period	Description of field hockey players	Injury definition (summary)	Injury data collection	Injured players	Number of injuries	Severity of injuries
Weighman and Browne 1975 (UK) ¹¹	Survey injuries in 11 selected sports	Season (8 months)	Men (25 clubs) and women (36 clubs). Number, age, and level NR	NR	Sport clubs' secretaries form	NR	117	Average TL Women: 1.5 days Men: 6.5 days
Clarke and Buckley 1980 (USA) ¹³	Preliminary overview of injury experiences among collegiate women athletes reported to National Athletic Injury/ Illness Reporting System during its first 3 operational years	Season (3 years)	High-school women from annual average of 16 teams. Number and age NR	An injury causing the athlete to miss at least 1 week of participation (≥ 1-week TL)	Athletic trainer injury report form	NR	NR	TL and consequences >3 weeks: 23% Surgery: 5%
Zaricznyj et al. 1980 (USA) ¹²	Analyse causes and severity of sports injuries in a total school-aged population	School season (1 year)	65+ school-aged players (5–17 years) ^b . Number and sex NR	Any traumatic act against the body sufficiently serious to have required first aid, school and insurance accident reports, or medical treatment (MA)	Principals, coaches, sport supervisors, ERs, school insurance company, local physician's injury form	NR	25	Injury type and consequences NR
Mathur et al. 1981 (Nigeria) ¹⁵	Determine sites and types of common injuries associated with competitive sports popular in Nigeria	Season (8 weeks)	212 players. Sex, level, and age NR	NR	Athlete self-report questionnaires	NR	641	NR
Rose 1981 (USA) ¹⁴	Describe women's field hockey injuries at the California State University in Long Beach	Season (4 years)	University women. Number and age NR	Minor injury: required MA of team physician in some cases but handled mainly by the trainer and produced no or limited disability. Major injury: required MA of team physician and produced definite disability needing follow-up care (medical/trainer attention)	NR	NR	81	Injury type and consequences Minor: 82.7% Major: 17.3% ^c
Martin et al. 1987 (USA) ¹⁶	Detail injury experiences of 1985 Junior Olympics	1985 Junior Olympic games (7 days)	53 women. Age NR	Injuries severe enough to withhold athlete from competition, at least temporarily, and to require formal medical evaluation by the trainer (medical/trainer attention and TL)	Medical staff report form	1.5.1% (8)	9	Tissue damage NR

Table 1. Characteristics of prospective studies on field hockey injuries arranged by year of publication (least recent to most recent).

Study (country)	Primary objective	Setting and follow-up period	Description of field hockey players	Injury definition (summary)	Injury data collection	Injured players	Number of injuries	Severity of injuries
Jamison and Lee 1989 (USA) ¹⁸	Compare injuries during Australian Women's Hockey Championships, 1984 (on grass) and 1985 (on Astroturf)	Championship (2 years)	110 women playing at Australians' state teams. Age NR	NR	Athletes self-report questionnaires	NR	178	NR
McLain and Reynolds 1989 (USA) ¹⁷	Investigate sports injuries at a large high-school	School season (7 months)	46 high-school women. Age NR	Any incident resulting from athletic participation that keeps athletes from completing a practice or game or causes athlete to miss a subsequent practice or game (TL)	Athletic: trainer injury evaluation sheet	6% (3)	NR	Average TL 3.3 days
Fuller 1990 (Country NR) ¹⁹	Study whether a characteristic pattern of injuries and their causation existed at county and territorial competition levels in women's field hockey on synthetic turf pitches	Competitive season (2 years)	Women. Number, level, and age NR	Presence of pain, discomfort, or disability arising during or as consequence of playing in a hockey match and for which physiotherapy treatment, advice, or handling was given (MA)	Researcher observation and contact with athletes	NR	135	TL ≤2 days: 90% >2 days: 10%
Cunningham and Barber-Foss 1996 (Australia) ²⁰	Obtain data relating to frequency, type, mechanism, severity of sports injuries incurred during or related to competition	1994 Australian University Games (6 days)	466 ^a university players, aged 17–47 years. Sex NR ^b	Any incident during warm-up or competition that required MA, on-field management to enable continued participation, or removal from the playing field (MA)	Attending officer injury surveillance form	33.5% (156)	181	Required treatment and injury NR
Fawkner et al. 1999 (Australia) ²²	Examine relationship between hassles and athletic injury	Season (13 weeks)	26 professional women aged 26 years on average ^b	Medical problem that resulted from either participation in training or competition, required MA, and restricted further participation in either training or a competition for at least one day post occurrence (MA and ≥ 1-day TL)	Coach recording form	23% (6)	NR	NR
Powell and Barber-Foss 1999 ^f (USA) ²¹	Describe injury patterns in ten high school sports	Season (2 years)	High-school women, number, age NR	(1) injury causing cessation of participation in current game or practice and prevented player's return to that session, (2) injury causing cessation of a player's customary participation on the day following the day of onset, (3) any fracture, even though athlete did not miss any regularly scheduled session, (4) any dental injury, including fillings, luxations, and fractures, and (5) any mild brain injury requiring cessation of player's participation for observation before returning, either in current or next session (MA or ≥ 1-day TL)	Athletic: trainer injury form	(445)	510	TL <8 days: 79.6% 8–21 days: 13.3% >21 days: 7.1%
Stevenson et al. 2000 (Australia) ²³	Describe trends in recreational sports injury in Perth, Western Australia	Winter season (5 months)	393 non-professional men (170) and women (223) aged 25 years on average	Injury occurring while participating in sport and leading to one of the following consequences: reduction in amount or level of sports activity, need for advice or treatment, and/or adverse economic or social effects (TL or MA and/or adverse economic/social effects)	Assisted telephone interview with athletes	28% (198)	279	Injury treatment NR

Table 1. Characteristics of prospective studies on field hockey injuries arranged by year of publication (least recent to most recent).

Study (country)	Primary objective	Setting and follow-up period	Description of field hockey players	Injury definition (summary)	Injury data collection	Injured players	Number of injuries	Severity of injuries
Finch et al. 2002 ^a (Australia) ²⁴	Describe the incidence of injury over two consecutive sporting seasons in a prospective cohort of community-level sporting participants within Australian football, hockey, basketball, netball	Two consecutive winter seasons (10 months)	280 non-professional men (116) ^a and women (164) ^b aged 25 years on average	One that occurred while participating in sport and that led to reduction in the amount or level of sport activity and/or need for advice or treatment and/or adverse economic or social effects (TL or MA and/or adverse economic/social effects)	Assisted telephone interview with athletes	31% (87)	445	Injury treatment NR
Lunge et al. 2006 (Greece) ²⁵	Analyse and compare the incidence, characteristics, and causes of injuries in all team sport tournaments during 2004 Olympic Games	2004 Olympic Games (19 days)	Olympic men and women. Number and age NR	Any physical complaint incurred during the match that received MA from the team physician, regardless of the consequences with respect to absence from the match or training (MA)	Physician injury report form	NR	44	Estimated TL None: 50% ^c 1-3 days: 27.3% ^c 4-7 days: 9.1% ^c >1 month: 2.3% ^c Unspecified: 2.3% ^c Missing: 9.1% ^c
Dick et al. 2007 (USA) ²⁸	Review 15 years of NCAA injury surveillance data for women's field hockey	Season (15 years)	5385 high-school women. Age NR	One that (1) occurred due to participation in an organized intercollegiate practice or competition and (2) required MA by a team-certified athletic trainer or physician and (3) resulted in restriction of the student athlete's participation or performance for one or more calendar days beyond the day of injury (MA and \geq 1-day TL)	Athletic trainer injury report form	NR	3286	>10 TL days Game injuries: 15% Practice injuries: 13%
Hootman et al. 2007 (USA) ²⁶	Summarize 16 years of NCAA injury surveillance data for 15 sports	Season (15 years for field hockey)	5385 high-school women. Age NR	One that (1) occurred due to participation in an organized intercollegiate practice or competition and (2) required MA by a team-certified athletic trainer or physician and (3) resulted in restriction of the student athlete's participation or performance for one or more calendar days beyond the day of injury (MA and \geq 1-day TL)	Athletic trainer injury report form	NR	3286	>10 TL days Game injuries: 15% Practice injuries: 13%
Rauh et al. 2007 ^d (USA) ²⁷	Determine patterns of new and subsequent injuries among female athletes participating in interscholastic sport	Season (2 years)	High-school women. Number and age NR	(1) Any injury causing cessation of participation in current game or practice and prevented player's return to that session; (2) any injury causing cessation of player's customary participation on the day following the day of onset; (3) any fracture, even though the athlete did not miss any regularly scheduled session; (4) any dental injury, including fillings, luxations, and fractures; (5) any mild brain injury requiring cessation of player's participation for observation before returning, either in the current or next session (MA or \geq 1 day TL)	Athletic trainer injury form	(445)	510	TL <8 days: 79.6% 8-21 days: 13.3% >21 days: 7.1%

Table 1. Characteristics of prospective studies on field hockey injuries arranged by year of publication (least recent to most recent).

Study (country)	Primary objective	Setting and follow-up period	Description of field hockey players	Injury definition (summary)	Injury data collection	Injured players	Number of injuries	Severity of injuries
Junge et al. 2009 (China) ³⁰	Analyse the frequency, characteristics, and causes of injuries incurred in competitions and/or training during 2008 Olympic Games	2008 Olympic Games (16 days)	382 Olympic men and women aged 26 years on average ^b	Any musculoskeletal complaint newly incurred due to competition and/or training during the XXIXth Olympiad in Beijing that received MA regardless of consequences with respect to a absence from competition or training (MA)	Physician injury report form	20.4% (78)	78	Estimated TL 3.5% of players
Rishiraj et al. 2009 (Canada) ²⁹	Identify rates, profiles, and severity of injuries associated with participating in under-21 age representative field hockey team	Season (5 years)	75 women aged 18 years on average representing British Columbia Women's Field Hockey Federation	Any event during team or team-related game, practice, or activity (on or off the playing surface) requiring attention by team's therapist or physician and subsequent game/practice TL (MA and \geq 1-day TL)	Athletic therapist injury reporting system	NR	198	TL <7 days: 81% 8–12 days: 17% >21 days: 2%
Engelbrechtsen et al. 2013 (UK) ³¹	Analyse injuries and illnesses during 2012 Olympic Games	2012 Olympic games (19 days)	388 Olympic men (196) and women (192). Age NR	New or recurring musculoskeletal complaints or concussions (injuries) incurred during competition or training during London Olympic Games receiving MA, regardless of consequences regarding absence from competition or training (MA)	Physician injury report form	17% (66)	66	TL \geq 1 days: 37.9% \geq 7 days: 15.2%
Theilen et al. 2016 (multiple countries (Table 3)) ³²	Investigate the incidence and severity of injuries during international field hockey tournaments in 2013	16 International Hockey Federation tournaments ⁸	Professional men and women. Number and age NR	A new musculoskeletal symptom or concussion that led to time stoppage when player was unable to continue playing during competition (TL)	Medical officer injury form	NR	236 ^c	NR

MA: medical attention; ERs: Emergency rooms; NATA: National Athletic Trainers' Association; NCAA: National Collegiate Athletic Association; NR: not reported; TL: time-loss.

^aPlayers participating in school teams. Does not include physical education, non-organized, and community practice (that are reported in the study)

^bData from the whole cohort (not only from field hockey players)

^c Calculated from presented data

^d Studies using the same data from 1995–1997 NATA High School Injury database

^e Finch et al.²⁴ is a follow-up study of Stevenson et al.²³

^f Studies using the same data from 1988–2003 NCAA Injury Surveillance System

⁸ Tournament durations in 2013 ranged from three to 10 days. The specific duration of each tournament can be found at <https://tms.fih.ch/fih/home>

Injury extent in field hockey

Injury rates

In total, 12 studies (55%) reported the number of injuries normalised by player exposure (i.e. injury rate). The injury rates reported in each of these studies are presented in Table 3 and were divided into two categories: (1) number of injuries per 1,000 player-hours of field hockey exposure (i.e. time at risk)^{11,12,23–25,32} and (2) number of injuries per 1,000 player-sessions (i.e. sessions at risk).^{13,21,25–29} One study reported the number of injuries according to both player-hours and player-sessions at risk.²⁵

In the studies describing injuries according to players' time at risk, injury rates ranged from 0.1 injuries (in school-aged players)¹² to 90.9 injuries (in Africa Cup of Nations)³² per 1,000 player-hours of field hockey (Table 3). The injury rate in the studies describing injuries according to players' sessions at risk varied from one injury (in high-school women)¹³ to 70 injuries (in under- 21 age women)²⁹ per 1,000 player-sessions. The injury rates were higher in games than in training sessions in two^{21,28} of the three studies that investigated this outcome.^{21,28,29} In major tournaments, injury rates were higher in men.^{25,32}

Injury severity

Table 1 presents the classification of injuries according to severity. Most of the studies (55%) used field hockey time-loss to report the severity of injuries,^{11,13,17,19,21,25–31} but reported the days of time-loss differently. Some studies reported the average days of time-loss^{11,17} and others used diverse cut-off points to report injury-related days of time-loss, such as two days,¹⁹ eight days,^{21,27} and 10 days.^{26,28} The majority of injuries were in the less severe category in all studies reporting days of time-loss due to injury, regardless of the cut-off points used.^{13,14,19,21,25,28,29,31} Six studies (27%) included severity measures in the methodology but did not specify the number or proportion of injuries according to severity in the results.^{13,16,20,23,24,32} Three studies (14%) did not mention severity of injury at all.^{15,18,22}

Table 2. Risk-of-bias assessment of studies on field hockey injuries according to 10 criteria.

Study	Criteria										Score
	1	2	3	4	5	6	7	8	9	10	
Weightman and Browne 1975 ¹¹	0	1	0	0	1	0	1	0	1	1	5
Clarke and Buckley 1980 ¹³	1	1	1	0	UD	1	1	1	1	1	8
Zaricznyj et al. 1980 ¹²	1	1	1	1	1	0	0	0	1	1	7
Mathur et al. 198 ¹⁵	0	1	0	0	UD	1	1	0	0	0	3
Rose 198 ¹⁴	1	1	1	0	UD	UD	UD	0	1	0	4
Martin et al. 1987 ¹⁶	1	1	1	1	1	1	1	1	0	0	8
Jamison and Lee 1989 ¹⁸	0	1	1	0	1	1	1	0	1	0	6
McLain and Reynolds 1989 ¹⁷	1	1	1	0	1	1	1	1	1	0	8
Fuller 1990 ¹⁹	1	1	0	0	UD	1	1	0	1	0	5
Cunningham and Cunningham 1996 ²⁰	1	1	1	1	UD	1	1	0	0	0	6
Fawkner et al. 1999 ²²	1	1	1	0	1	0	0	0	0	0	4
Powell and Barber-Foss 1999 ²¹	1	1	1	0	UD	1	0	1	1	1	7
Stevenson et al. 2000 ²³	1	1	1	1	0	1	1	0	0	1	7
Finch et al. 2002 ²⁴	1	1	1	1	0	1	1	0	1	1	8
Junge et al. 2006 ²⁵	1	1	1	1	0	1	1	1	0	1	8
Dick et al. 2007 ²⁸	1	1	1	0	UD	1	1	1	1	1	8
Hootman et al. 2007 ²⁶	1	1	1	0	UD	1	1	1	1	1	8
Rauh et al. 2007 ²⁷	1	1	1	0	UD	1	0	1	1	1	7
Junge et al. 2009 ³⁰	1	1	1	1	1	1	1	1	0	0	8
Rishiraj et al. 2009 ²⁹	1	1	1	0	UD	1	1	1	1	1	8
Engelbretsen et al. 2013 ³¹	1	1	1	1	1	1	1	1	0	0	8
Theilen et al. 2016 ³²	1	1	1	1	1	1	1	1	0	1	9
Total, n (%) of studies	19 (86)	22 (100)	19 (86)	9 (41)	9 (41)	18 (82)	17 (77)	12 (55)	13 (59)	12 (55)	

Low risk of bias = 1. High risk of bias = 0. Unable to determine fields (UD) were counted as zero in the score.

- 1 Definition of injury clearly described.
- 2 Prospective design that presents incidence or prevalence data.
- 3 Description of field hockey players (e.g. recreational or professional level).
- 4 The process of inclusion of athletes in the study was at random (i.e. not by convenience) or the data collection was performed with the entire target population.
- 5 Data analysis performed with at least 80% of the athletes included in the study.
- 6 Injury data reported by players or by a healthcare professional.
- 7 Same mode of injury data collection used.
- 8 Injury diagnosis conducted by medical professional.
- 9 Follow-up period of at least six months.
- 10 Incidence or prevalence rates of injury expressed by a ratio that represents both the number of injuries as well as the exposure to field hockey (i.e. number of injuries/hours of field hockey exposure, or number of injuries/sessions of field hockey exposure).

Table 3. Number of field hockey injuries (and 95% confidence intervals) per 1,000 player-hours and player-sessions arranged by players' characteristics.

Study	Players' characteristics	Setting	Injury definition summary	Players' exposure (hours)	Number of injuries per 1,000 player-hours	Players' exposure (sessions)	Number of injuries per 1,000 player-sessions
Weighman and Browne 1975 ¹¹	NR	Season	NR	122,074 ^a	1.0 (0.8–1.1) ^a		
	Men			70,874 ^a	1.0 (0.8–1.3) ^a		
	Women			51,200 ^a	1.3 (0.9–1.6) ^a		
Zaricznyj et al. 1980 ¹²	School players aged 5–17 years. Sex NR	Season	MA	14,286 ^a	0.1 (0.0–1.4) ^a		
Clarke and Buckley 1980 ¹³	High-school women. Age NR	Season	≥ 1-week TL				1.0 ^b
Powell and Barber-Foss ^c 1999 ²¹	High-school women. Age NR	Season	MA or ≥ 1-day TL	138,073		138,073	3.7 (3.4–4.0) ^b
		Game		66,122 ^a		66,122 ^a	4.9 (4.4–5.4) ^a
		Practice		58,125 ^a		58,125 ^a	3.2 (2.7–3.7) ^a
Dick et al. ^d 2007 ²⁸	High-school women. Age NR	Season	MA or ≥ 1-day TL	716,910 ^a		716,910 ^a	4.6 (4.4–4.7) ^a
		Game		155,370 ^a		155,370 ^a	7.9 (7.4–8.3)
		Practice		561,540 ^a		561,540 ^a	3.7 (3.5–3.9)
Rishiraj et al. 2009 ²⁹	Under-21 age women from the BCWFHF	Season	MA or ≥ 1-day TL	2,828		2,828	70.0 (30.2–79.8) ^a
		Game		578		578	67.5 (45.6–89.3) ^a
		Practice		2,250		2,250	68.0 (57.1–78.9) ^a
Finch et al. ^e 2002 ³⁴	Non-professional men and women age 25 years on average	Winter season	TL or MA and/or adverse economic/social effects	29,276 ^a	15.2 (13.8–16.7)		
Junge et al. 2006 ³⁵	Olympic players. Age NR	2004 Olympic Games	MA	1,322 ^a	33 (23–43) ^a	1,133 ^a	39 (27–50)
	Men			770	47 (32–62)	660	55 (37–72)
	Women			552	14 (4–24)	473	17 (5–29)
Theillen et al. 2016 ³²	Professional players. Age NR	2013 FIH tournaments	TL	6,519 ^a	36.2 (31.6–40.8) ^a		
	Men			4,825	48.3 (30.9–68.8)		

Table 3. Number of field hockey injuries (and 95% confidence intervals) per 1,000 player-hours and player-sessions arranged by players' characteristics.

Study	Players' characteristics	Setting	Injury definition summary	Players' exposure (hours)	Number of injuries per 1,000 player-hours	Players' exposure (sessions)	Number of injuries per 1,000 player-sessions
		Africa Cup of Nations (Kenya)		154	90.9 (38.4–143.4)		
		East Asia Games (China)		154	90.9 (38.4–143.4) ^a		
		Junior World Cup (India)		1,129	27.4 (17.4–37.5) ^a		
		Oceania Cup (New Zealand)		154	77.9 (28.4–127.4) ^a		
		Sultan of Johor Cup (Malaysia)		462	28.1 (11.1–45.1) ^a		
		World League Round 2 (India)		385	44.2 (21.5–66.9) ^a		
		World League Round 2 (Russia)		385	44.2 (21.5–66.9) ^a		
		World League Round 2 (France)		385	26 (7.4–44.6) ^a		
		World League Round 2 (Brazil)		385	20.8 (3.4–38.2) ^a		
		World League Semi-final (Malaysia)		616	42.2 (25.2–59.3) ^a		
		World League Semi-final (Netherlands)		616	39.0 (22.5–55.4) ^a		
Women				1,694	29.1 (18.6–39.7)		
		4 Nations Tournament (New Zealand)		154	26.0 (0.0–67.3) ^a		
		East Asia Games (China)		154	26.0 (0.0–67.3) ^a		
		World League Final (Argentina)		616	26.0 (12.1–39.8) ^a		
		World League Round 2 (India)		385	44.2 (21.5–66.9) ^a		
		World League Round 2 (Brazil)		385	23.4 (5.4–41.4) ^a		

BCWFHF: British Columbia Women's Field Hockey Federation; FIH: International Hockey Federation; MA: medical attention; NR: not reported; TL: time-loss.

^a Calculated according to presented data.

^b Impossible to calculate 95% confidence interval.

^c Same data as Kauh et al. 2007.²⁷

^d Same data as Hootman et al. 2007.²⁶

^e A follow-up study of Stevenson et al. 2000.²³

Injury characteristics in field hockey

Body location and types of injury

Fifteen studies (68%) described injuries according to the affected body location.^{12-16,18,19,21,24,25,27-29,31,32} Table 4 presents the proportion (%) of injuries according to body location reported in these studies. The most common site of injury was the lower limbs (ranging from 13%²⁵ to 77%¹⁸ of all injuries), followed by head (2%¹³ to 50%²⁵), upper limbs (0%¹⁶ to 44%¹²) and trunk (0%¹⁸ to 16%²⁸). In the lower limbs, injuries were more frequent in the knee, ankle, lower leg, and thigh (Table 4).

Thirteen studies (59%) described the types of injury sustained by field hockey players.^{13-16,18,20,21,24,25,27-29,31} Table 5 presents the proportion (%) of injuries according to their type. Contusions and hematomas were the most common types of injury (ranging from 14%³¹ to 64%¹⁸ of all injuries), followed by abrasions and lacerations (5%¹⁴ to 51%¹⁵), sprains (2%¹⁸ to 37%¹³), and strains (0%²⁵ to 50%²⁸). Concussions ranged from 0%²⁵ to 25%²⁵.

Injury according to mechanism and player position

Eight studies (36%) described injuries according to their mechanism.^{18-20,25,28,29,31,32} Table 6 presents the proportion (%) of injuries according to their mechanism. Non-contact injuries ranged from 12%¹⁸ to 64%.²⁸ Contact with the ball (ranging from 2%²⁹ to 52%³²) and stick (9%²⁹ to 27%¹⁸) were also common mechanisms, as well as contact with another player (2%¹⁹ to 45%²⁰) or with the ground (9%²⁸ to 15%²⁰).

Three studies (14%) reported injuries according to the injured player's position.^{19,28,29} Goalkeepers sustained fewer injuries in all three studies that reported injuries by playing position (4%¹⁹ to 19%²⁸). Defenders sustained 16%¹⁹ to 36%²⁹ of injuries, while midfielders and forwards sustained 22%²⁸ to 37%¹⁹ (Table 7).

Table 4. Proportion (%) of field hockey injuries by body location. **Bold** formatting indicates the highest values for the main body areas in each study.

Study	Upper limbs						Lower limbs								
	Head, neck, face	Hand, finger, wrist	Upper arm, forearm	Elbow	Shoulder	Total upper limbs	Trunk, upper and lower back	Ankle	Foot, toes	Lower leg	Thigh	Knee	Hip, groin, pelvis	Total lower limbs	Other, unspecified
Clarke and Buckley 1980 ³¹	2											32		72	26
Zaricznyj et al. 1980 ³²	12	40		4		44		16	12	8		8		44	
Mathur et al. 1981 ¹⁵	5	17	4	2	1	24	14	12	11	46	2	6		58	
Rose 1981 ¹⁴	11				1	5	4	27	15	9	11	11	2	75	5
Martin et al. 1987 ¹⁶	22					0								44	33
Jamison and Lee 1989 (astro turf) ¹⁶	15	16	2	1	1	21	5	3	4	15	13	18	7	58	
Jamison and Lee 1989 (grass) ¹⁶	9	9	2	1	1	13	1	9	14	9	13	31	2	77	
Fuller 1990 ¹⁹	10	18	2			20	9	5	9	5	17	24		60	
Powell Barber-Foss' 1999 ³¹	17		13		3	16	5	23				14	22	59	3
Finch et al. ^b 2002 ²⁴		31			4			28	12	19	30	31			
Junge et al. 2006 (men) ²⁵	22	6	9	3	3	16	8	14	3	3	8	22	0	50	
Junge et al. 2006 (women) ²⁵	50	25	0	0	0	25	13	13	0	0	0	0	0	13	
Dick et al. ^c 2007 (game) ²⁶	25	10				21	7	15	3	3	10	18	3	43	4
Dick et al. ^c 2007 (practice) ²⁶	8	2				8	16	15	2	8	27	17	12	60	7
Rishiraj et al. 2009 ²⁹	7	23			6	29	14	14			10	13	15	53	
Engelbreitsen et al. 2013 ³¹	20	16	3		5	23	11	8	8	5	9	11	8	47	
Theilen et al. 2016 (men) ³²	27	19				19	4			13	28			31	9
Theilen et al. 2016 (women) ³²	40	14				14	0			16	12			28	18

^a Same data as Rauh et al. 2007.²⁷

^b Values represent percentages of injured players (i.e. not injuries) and do not add to 100% as some players sustained more than one injury.

^c Same data as Hootman et al. 2007.²⁶

Table 5. Proportion (%) of field hockey injuries by injury type. **Bold** formatting indicates the highest values for each study.

Study	Sprains	Strains	Dislocation	Fracture	Abrasion, laceration	Contusion, hematoma	Swelling, blistering	Concussion	Tendinopathy	Other, unspecified
Clarke and Buckley 1980 ¹³	37	21		7				4		32
Mathur et al. 1981 ¹⁵	20 ^a			6 ^b	51		22	1		
Rose 1981 ¹⁴	32	16	1	1	5	33		4	2	5
Martin et al. 1987 ¹⁶	11				11	33				44
Jamison and Lee 1989 (astroturf) ¹⁸	2	12			26	49	2	3		6
Jamison and Lee 1989 (grass) ¹⁸	2	5			16	64	7	1		5
Cunningham and Cunningham 1996 ²⁰	15	19		2	22	28	3	4		9
Powell Barber-Foss' 1999 ²¹	26	20		6		37		3		8
Finch et al. ^d 2002 ²⁴	28	55	2	14	15	80		1	2	
Junge et al. 2006 (men) ²⁵	11	8	0	8	19	42		0	8	3
Junge et al. 2006 (women) ²⁵	13	0	0	0	25	38		25	0	0
Dick et al. ^e 2007 (game) ²⁶	24	13		15	11	20		9	4	3
Dick et al. ^e 2007 (practice) ²⁶	23	50		5			3 ^f	3	7	8
Rishiraj et al. 2009 ²⁹	10	40	1	1	8	17			24	
Engelbreitson et al. 2013 ³¹	18	14	6	8	21	14			3	18

^a Sprains and strains reported together.

^b Fractures and dislocations reported together.

^c Same data as Rauh et al. 2007.²⁷

^d Values represent percentages of injured players (i.e. not injuries) and do not add to 100% as some players sustained more than one injury.

^e Same data as Hootman et al. 2007.²⁶

^f Reported as inflammation.

Table 6. Proportion (%) of field hockey injuries by injury mechanism. **Bold** formatting indicates the highest values for each study.

Study	Ball-contact	Stick-contact	Player-contact	Ground-contact	Object-contact	Unspecified contact	Non-contact	Unspecified
Jamison and Lee 1989 (astroturf) ¹⁶	32	27	11	12			18	
Jamison and Lee 1989 (grass) ¹⁸	42	23	9	14			12	
Fuller 1990 ¹⁹	30	17	2	10			41	
Cunningham and Cunningham 1996 ⁵⁰			45	15			36	4
Junge et al. 2006 (men) ²⁵						58	36	6
Junge et al. 2006 (women) ²⁵						75	13	13
Dick et al., 2007 (game) ²⁶	29	18	14	9			28	2
Dick et al., 2007 ^a (practice) ²⁶			5			26	64	5
Rishiraj et al. 2009 ²⁹	2	9	12	12	3 ^b		62	
Engelbreitsen et al. 2013 ³¹			8		44^c		41	7
Theilen et al. 2016 (men) ³²	37	25	23	15				
Theilen et al. 2016 (women) ³²	52	14	12	20				2

^a Same data as Hootman et al. 2007.²⁶

^b Contact with the goal.

^c Contact with unspecified moving or stagnant object.

Table 7. Proportion (%) of field hockey injuries by player position. **Bold** formatting indicates the highest values for each study.

Study	Forwards	Midfielders	Defenders	Goalkeepers	Other, unknown
Rishiraj et al. 2009 ²⁹	32	22	36	10	
Fuller 1990 ¹⁹	37	37	16	4	6
Dick et al., 2007 ^a (game) ²⁶	22	28	24	19	7

^a Player position at time of injury. Same data as Hootman et al. 2007.²⁶

DISCUSSION

To the best of our knowledge, the present study is the first systematic review to summarize the descriptive evidence of injuries sustained by field hockey players. We included only prospective studies to ensure we gathered the most reliable information available on the extent of injuries in field hockey in terms of rate and severity, as well as injury characteristics according to body location, type, and mechanism of injury. To reduce and control field hockey injuries, as for all sports, we must first establish the extent of the injury problem.⁶ The substantial heterogeneity between studies included in this review prevented conclusive findings on the extent of the rate and severity of injuries in field hockey (Tables 1 and 2). Such heterogeneity may be caused by the different definitions and methods employed to record and report injuries and the different characteristics and levels of players studied.

This systematic review shows that, despite the long history of field hockey and its popularity worldwide, prospective studies focusing on overall field hockey injuries are still lacking. The majority of the studies investigated field hockey injuries together with injuries in other sports.^{11–13,15–17,20–27,30,31} Within such studies, injury rates in field hockey were comparable to those in other team sports, such as basketball,^{23,24,26} netball,^{23,24} lacrosse,²⁶ and softball.^{21,27} The injury rate in field hockey can be considered low compared with football (soccer).^{21,25–27} However, in major tournaments, the rate of time-loss injuries in field hockey³² can be considered higher than football (soccer).⁴ These findings confirm that the risk of sustaining an injury in field hockey should not be neglected.

Despite the considerable heterogeneity between studies, it is still possible to observe similar characteristics of injuries with regard to body location, type, and mechanism of injury. Most of the injuries described in the studies included in this review were to the lower limbs (Table 4), affecting mainly the knee and the ankle. This is in line with previous studies on team sports involving running and stepping manoeuvres, such as football (soccer)³³ and lacrosse,³⁴ and justifies a focus on preventive efforts in this body area. Interestingly, the majority of injuries sustained by women during major

tournaments were to the head.^{25,32} A specific analysis of head injuries in collegiate women's field hockey showed that 48% of these injuries occurred due to contact with an elevated ball.³⁵ Most (39%) of the concussions were due to direct contact with another player, and 25% were due to contact with an elevated ball.³⁵

Contusions and hematomas were common types of injury, as were abrasions and lacerations, which might be due to players' contact with the ball, stick, and playing surface.^{2,28} A specific analysis of ball-contact injuries in 11 collegiate sports showed that injury rates were the highest in women's softball, followed by women's field hockey and men's baseball.³⁶ In field hockey, the common activities associated with ball-contact injuries were defending, general play, and blocking shots.³⁶ To reduce the injury burden, the International Hockey Federation stated that goalkeepers must wear protective equipment comprising at least headgear, leg guards, and kickers.³⁷ Field players are recommended to use shin, ankle, and mouth protection,³⁷ and other research suggested that the use of such equipment should be mandatory.² Accordingly, some national associations have updated their rules to make shin, ankle, and mouth protection obligatory.^{38,39}

It is important to note that non-contact injuries are also a cause for concern in field hockey (Table 6). Although protective equipment has a fundamental role in injury prevention, it may not prevent most of the non-contact injuries. During the last decades, different studies have shown that it is possible to prevent injuries in team sports with structured exercise.⁴⁰⁻⁴⁴ Yet, to our knowledge, evidence showing the implementation of such programs in field hockey is lacking. Nevertheless, exercise programs that have proven effective in preventing sports injury can be introduced as part of the regular training schedule of the field hockey team, especially programs focusing on the prevention of lower limb injuries.⁴⁰⁻⁴² While there is no structured exercise program for field hockey, stakeholders can also use open source resources for overall and specific injury prevention that are supported by the International Olympic Committee, such as exercise programs and guidelines on load management and youth athletic development.⁴⁵⁻⁴⁷

Future recommendations

The present systematic review shows that studies have used different definitions and methods to record and classify injuries and their severity, and this prevents conclusive findings on the extent of the injury problem in field hockey. As establishing the extent of sports injury is considered the first step toward effective prevention,⁶ one of the main findings of this review is the recognition of the need for a consensus on the methodology of injury surveillance in field hockey. Consensus statements on the methodology of injury surveillance have been made available for a variety of sports.^{8,48-54} A consensus statement represents the result of a comprehensive collective analysis, evaluation, and opinion of a panel of experts regarding a specific subject (e.g. methodology of injury surveillance in field hockey).⁵⁵ Consequently, consensus statements enable investigators from different settings to access and employ the same definitions and methods to collect and report injury data. Comparisons among different studies as well as data pooling for meta-analyses are then facilitated.

The common goal in field hockey is to promote players' safety while maintaining the traditions of the sport.³⁵ Protecting the health of the athletes is also a priority of the International Olympic Committee,⁵⁶ and resources for injury prevention have been made available for the public in general.⁴⁵⁻⁴⁷ The field hockey community would benefit from studies investigating the implementation of such resources and from strategies that have been proven to be effective in other sports.⁴⁰⁻⁴⁴ Until there is consensus on the methodology of injury surveillance in field hockey, investigators may use consensus from other team sports in future studies as an example.^{8,52,53} Based on the gaps identified in the studies included in this review, the authors also suggest that future studies adhere to the reporting guidelines from the Enhancing the Quality and Transparency of Health Research (EQUATOR) Network. The EQUATOR Network provides comprehensive documentation on what information needs to be reported in scientific manuscripts depending on the study design.⁵⁷ By following an appropriate guideline such as that of the EQUATOR Network, future investigators will facilitate assessment of the generalizability, strengths, and limitations of studies on field hockey injuries.

Limitations

Electronic searches were conducted in four databases that were considered relevant for this systematic review. This does not rule out the possibility of eligible articles published in journals that were not indexed in any of these databases. To minimize this limitation, we screened the references of the full texts assessed for eligibility and included additional studies that were not identified in the database search. In addition, this systematic review included only scientific manuscripts published in English, although studies on field hockey injuries have been published in other languages. These were not included because the authors were unable to translate the papers accurately enough to extract their data.

CONCLUSION

The present systematic review shows that, despite the long history and the popularity of field hockey worldwide, few prospective studies have investigated the overall injury problem in field hockey. Most of the information on field hockey injuries registered prospectively comes from studies conducted in multi-sport settings. The range of definitions, methods, and reporting employed by studies prevents conclusive findings on the rate and severity of injuries in field hockey. To facilitate the development of evidence-based strategies for injury prevention, field hockey may benefit from a consensus on the methodology of injury surveillance. While no specific consensus is available for field hockey, future studies may use widely accepted consensus from other sports, such as football (soccer). In addition, future studies on field hockey injuries are encouraged to adhere to the reporting guidelines from the EQUATOR Network.

Despite the considerable heterogeneity, it is clear that most of the injuries sustained by field hockey players affect the lower limbs, justifying efforts to develop preventive strategies for this body area. Contact injuries, such as contusions/hematomas, and abrasions, are frequent, and the use of protective equipment for the ankle, shin, hand, mouth, and eye/face has been recommended. Nevertheless, non-contact injuries are

also common in field hockey, and most of these may not be prevented by protective gear. To reduce the burden of injuries, field hockey stakeholders may implement exercise-based injury prevention programs and guidelines on load management and youth athletic development that have been supported by the International Olympic Committee.

SUPPLEMENTARY MATERIAL

Literature search strategy and searched databases	
Search strategy	Terms
Studies about injury	1. injur* 2. traum* 3. risk* 4. overuse 5. overload 6. acute 7. odds 8. incidence 9. prevalence 10. hazard 11. OR/1–10
Studies about field hockey	12. field 13. hockey 14. ice 15. 12 AND 13 OR 12 NOT 14
Prospective and retrospective studies	16. prosp* 17. retrosp* 18. case* 19. OR/16–18
Databases	
PubMed	11 AND 15 AND 19
Excerpta Medical Database (EMBASE)	11 AND 15 AND 19 AND [embase]/lim NOT [medline]/lim
SPORTDiscus	11 AND 15 AND 19 AND [limit: academic journals]
Cumulative Index to Nursing and Allied Health Literature (CINAHL)	11 AND 15 AND 19 AND [limit: academic journals]

REFERENCES

1. International Hockey Federation. <http://www.fih.ch/hockey-basics/history>. Accessed August 2, 2017.
2. Murtaugh K. Field hockey injuries. *Curr Sports Med Rep*. 2009;8(5):267-272. doi:10.1249/JSR.0b013e3181b7f1f4.
3. van Mechelen W. The severity of sports injuries. *Sport Med*. 1997;24(3):176-180. doi:10.2165/00007256-199724030-00006.
4. Hägglund M, Waldén M, Magnusson H, Kristenson K, Bengtsson H, Ekstrand J. Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med*. 2013;47(12):738-742. doi:10.1136/bjsports-2013-092215.
5. Eirale C, Tol JL, Farooq A, Smiley F, Chalabi H. Low injury rate strongly correlates with team success in Qatari professional football. *Br J Sports Med*. 2013;47(12):807-808. doi:10.1136/bjsports-2012-091040.
6. van Mechelen W, Hlobil H, Kemper HCG. Incidence, severity, aetiology and prevention of sports injuries. *Sport Med*. 1992;14(2):82-99. doi:10.2165/00007256-199214020-00002.
7. Cook DJ. Systematic reviews: synthesis of best evidence for clinical decisions. *Ann Intern Med*. 1997;126(5):376. doi:10.7326/0003-4819-126-5-199703010-00006.
8. Fuller CW, Molloy MG, Bagate C, et al. Consensus statement on injury definitions and data collection procedures for studies of injuries in rugby union. *Br J Sports Med*. 2007;41(5):328-331. doi:10.1136/bjism.2006.033282.
9. Weiler R, Van Mechelen W, Fuller C, Verhagen E. Sport injuries sustained by athletes with disability: a systematic review. *Sports Med*. 2016;46(8):1141-1153. doi:10.1007/s40279-016-0478-0.
10. Lopes AD, Hespagnol Júnior LC, Yeung SS, Costa LOP. What are the main running-related musculoskeletal injuries? A systematic review. *Sports Med*. 2012;42(10):891-905. doi:10.2165/11631170-000000000-00000.
11. Weightman D, Browne RC. Injuries in eleven selected sports. *Br J Sports Med*. 1975;9(3):136-141. doi:10.1136/bjism.9.3.136.
12. Zaricznyj B, Shattuck LJM, Mast TA, Robertson R V, D'Elia G. Sports-related injuries in school-aged children. *Am J Sports Med*. 1980;8(5):318-324.

- doi:10.1177/036354658000800504.
13. Clarke KS, Buckley WE. Women's injuries in collegiate sports. A preliminary comparative overview of three seasons. *Am J Sports Med.* 1980;8(3):187-191. doi:10.1177/036354658000800308.
 14. Rose CP. Injuries in women's field hockey: a four-year study. *Phys Sportsmed.* 1981;9(3):97-100. doi:10.1080/00913847.1981.11711034.
 15. Mathur DN, Salokun SO, Uyanga DP. Common injuries among Nigerian games players. *Br J Sports Med.* 1981;15(2):129-132. <http://www.ncbi.nlm.nih.gov/pubmed/7272655>.
 16. Martin RK, Yesalis CE, Foster D, Albright JP. Sports injuries at the 1985 Junior Olympics. An epidemiologic analysis. *Am J Sports Med.* 1987;15(6):603-608. doi:10.1177/036354658701500614.
 17. McLain LG, Reynolds S. Sports injuries in a high school. *Pediatrics.* 1989;84(3):446-450. <http://www.ncbi.nlm.nih.gov/pubmed/2771547>.
 18. Jamison S, Lee C. The incidence of female hockey injuries on grass and synthetic playing surfaces. *Aust J Sci Med Sport.* 1989;21(2):15-17.
 19. Fuller MI. A study of injuries in women's field hockey as played on synthetic turf pitches. *Physiother Sport.* 1990;12(5):3-6.
 20. Cunningham C, Cunningham S. Injury surveillance at a national multi-sport event. *Aust J Sci Med Sport.* 1996;28(2):50-56. <http://www.ncbi.nlm.nih.gov/pubmed/8836476>.
 21. Powell JW, Barber-Foss KD. Injury patterns in selected high school sports: a review of the 1995-1997 seasons. *J Athl Train.* 1999;34(3):277-284. <http://www.ncbi.nlm.nih.gov/pubmed/16558577>.
 22. Fawkner HJ, McMurray NE, Summers JJ. Athletic injury and minor life events: a prospective study. *J Sci Med Sport.* 1999;2(2):117-124. doi:10.1016/S1440-2440(99)80191-1.
 23. Stevenson MR, Hamer P, Finch CF, Elliot B, Kresnow M. Sport, age, and sex specific incidence of sports injuries in Western Australia. *Br J Sports Med.* 2000;34(3):188-194. <http://www.ncbi.nlm.nih.gov/pubmed/10854018>.
 24. Finch C, Costa A Da, Stevenson M, Hamer P, Elliott B. Sports injury experiences from the Western Australian sports injury cohort study. *Aust N Z J Public Health.*

- 2002;26(5):462-467. doi:10.1111/j.1467-842X.2002.tb00348.x.
25. Junge A, Langevoort G, Pipe A, et al. Injuries in team sport tournaments during the 2004 Olympic Games. *Am J Sports Med.* 2006;34(4):565-576. doi:10.1177/0363546505281807.
 26. Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train.* 2007;42(2):311-319. <http://www.ncbi.nlm.nih.gov/pubmed/17710181>.
 27. Rauh MJ, Macera CA, Ji M, Wiksten DL. Subsequent injury patterns in girls' high school sports. *J Athl Train.* 2007;42(4):486-494. <http://www.ncbi.nlm.nih.gov/pubmed/18176621>.
 28. Dick R, Hootman JM, Agel J, Vela L, Marshall SW, Messina R. Descriptive epidemiology of collegiate women's field hockey injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 through 2002-2003. *J Athl Train.* 2007;42(2):211-220. <http://www.ncbi.nlm.nih.gov/pubmed/17710169>.
 29. Rishiraj N, Taunton JE, Niven B. Injury profile of elite under-21 age female field hockey players. *J Sports Med Phys Fitness.* 2009;49(1):71-77. <http://www.ncbi.nlm.nih.gov/pubmed/19188898>.
 30. Junge A, Engebretsen L, Mountjoy ML, et al. Sports injuries during the Summer Olympic Games 2008. *Am J Sports Med.* 2009;37(11):2165-2172. doi:10.1177/0363546509339357.
 31. Engebretsen L, Soligard T, Steffen K, et al. Sports injuries and illnesses during the London Summer Olympic Games 2012. *Br J Sports Med.* 2013;47(7):407-414. doi:10.1136/bjsports-2013-092380.
 32. Theilen T-M, Mueller-Eising W, Wefers Bettink P, Rolle U. Injury data of major international field hockey tournaments. *Br J Sports Med.* 2016;50(11):657-660. doi:10.1136/bjsports-2015-094847.
 33. Faude O, Rösler R, Junge A. Football injuries in children and adolescent players: are there clues for prevention? *Sports Med.* 2013;43(9):819-837. doi:10.1007/s40279-013-0061-x.
 34. Putukian M, Lincoln AE, Crisco JJ. Sports-specific issues in men's and women's lacrosse. *Curr Sports Med Rep.* 2014;13(5):334-340. doi:10.1249/JSR.0000000000000092.

35. Gardner EC. Head, face, and eye injuries in collegiate women's field hockey. *Am J Sports Med.* 2015;43(8):2027-2034. doi:10.1177/0363546515588175.
36. Fraser MA, Grooms DR, Guskiewicz KM, Kerr ZY. Ball-contact injuries in 11 National Collegiate Athletic Association Sports: the injury surveillance program, 2009—2010 through 2014—2015. *J Athl Train.* 2017;1062-6050-52.3.10. doi:10.4085/1062-6050-52.3.10.
37. International Hockey Federation. Rules of hockey. <http://www.fih.ch/inside-fih/our-official-documents/rules-of-hockey>. Accessed June 27, 2017.
38. National Collegiate Athletic Association. *Field hockey rules modifications*. <http://www.ncaa.org/championships/playing-rules/field-hockey-rules-game>.
39. Koninklijke Nederlandse Hockey Bond. *Spelreglement veldhockey*.; 2017. <https://www.knhb.nl/kenniscentrum/scheidsrechters/alles-over-de-spelregels>.
40. Olsen O-E, Myklebust G, Engebretsen L, Holme I, Bahr R. Exercises to prevent lower limb injuries in youth sports: cluster randomised controlled trial. *BMJ.* 2005;330(7489):449. doi:10.1136/bmj.38330.632801.8F.
41. Verhagen E, van der Beek A, Twisk J, Bouter L, Bahr R, van Mechelen W. The effect of a proprioceptive balance board training program for the prevention of ankle sprains: a prospective controlled trial. *Am J Sports Med.* 2004;32(6):1385-1393. doi:10.1177/0363546503262177.
42. Al Attar WSA, Soomro N, Sinclair PJ, Pappas E, Sanders RH. Effect of injury prevention programs that include the Nordic hamstring exercise on hamstring injury rates in soccer players: a systematic review and meta-analysis. *Sports Med.* 2017;47(5):907-916. doi:10.1007/s40279-016-0638-2.
43. Thorborg K, Krommes KK, Esteve E, Clausen MB, Bartels EM, Rathleff MS. Effect of specific exercise-based football injury prevention programmes on the overall injury rate in football: a systematic review and meta-analysis of the FIFA 11 and 11+ programmes. *Br J Sports Med.* 2017;51(7):562-571. doi:10.1136/bjsports-2016-097066.
44. Longo UG, Loppini M, Berton A, Marinozzi A, Maffulli N, Denaro V. The FIFA 11+ program is effective in preventing injuries in elite male basketball players: a cluster randomized controlled trial. *Am J Sports Med.* 2012;40(5):996-1005. doi:10.1177/0363546512438761.

45. Verhagen E. Get Set: prevent sports injuries with exercise! *Br J Sports Med.* 2015;49(11):762-762. doi:10.1136/bjsports-2015-094644.
46. Soligard T, Schweltnus M, Alonso J-M, et al. How much is too much? (Part 1) International Olympic Committee consensus statement on load in sport and risk of injury. *Br J Sports Med.* 2016;50(17):1030-1041. doi:10.1136/bjsports-2016-096581.
47. Bergeron MF, Mountjoy M, Armstrong N, et al. International Olympic Committee consensus statement on youth athletic development. *Br J Sports Med.* 2015;49(13):843-851. doi:10.1136/bjsports-2015-094962.
48. Yamato TP, Saragiotto BT, Lopes AD. A consensus definition of running-related injury in recreational runners: a modified Delphi approach. *J Orthop Sports Phys Ther.* 2015;45(5):375-380. doi:10.2519/jospt.2015.5741.
49. Mountjoy M, Junge A, Alonso JM, et al. Consensus statement on the methodology of injury and illness surveillance in FINA (aquatic sports). *Br J Sports Med.* 2016;50(10):590-596. doi:10.1136/bjsports-2015-095686.
50. Turner M, Fuller CW, Egan D, et al. European consensus on epidemiological studies of injuries in the thoroughbred horse racing industry. *Br J Sports Med.* 2012;46(10):704-708. doi:10.1136/bjsports-2011-090312.
51. Pluim BM, Fuller CW, Batt ME, et al. Consensus statement on epidemiological studies of medical conditions in tennis, April 2009. *Br J Sports Med.* 2009;43(12):893-897. doi:10.1136/bjism.2009.064915.
52. Fuller CW, Ekstrand J, Junge A, et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Br J Sports Med.* 2006;40(3):193-201. doi:10.1136/bjism.2005.025270.
53. Orchard J, Newman D, Stretch R, Frost W, Mansingh A, Leipus A. Methods for injury surveillance in international cricket. *J Sci Med Sport.* 2005;8(1):1-14. doi:10.1016/S1440-2440(05)80019-2.
54. Timpka T, Alonso J-M, Jacobsson J, et al. Injury and illness definitions and data collection procedures for use in epidemiological studies in Athletics (track and field): consensus statement. *Br J Sports Med.* 2014;48(7):483-490. doi:10.1136/bjsports-2013-093241.
55. Consensus statements. *Diabetes Care.* 2002;25(Supplement 1):S139-S139. doi:10.2337/diacare.25.2007.S139.

CHAPTER 5

56. Engebretsen L, Bahr R, Cook JL, et al. The IOC Centres of Excellence bring prevention to sports medicine. *Br J Sports Med.* 2014;48(17):1270-1275. doi:10.1136/bjsports-2014-093992.
57. The EQUATOR Network. Enhancing the QUALity and Transparency Of Health Research. <http://www.equator-network.org>. Accessed July 2, 2017.