

VU Research Portal

The cholesterol-raising diterpenes from coffee beans increase serum lipid transfer protein activity levels in humans

van Tol, A.; Urgert, R.; de Jong-Caesar, R.; van Gent, T.; Scheek, L. M.; de Roos, B.; Katan, M.B.

published in

Atherosclerosis
1997

DOI (link to publisher)

[10.1016/S0021-9150\(97\)00099-3](https://doi.org/10.1016/S0021-9150(97)00099-3)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

van Tol, A., Urgert, R., de Jong-Caesar, R., van Gent, T., Scheek, L. M., de Roos, B., & Katan, M. B. (1997). The cholesterol-raising diterpenes from coffee beans increase serum lipid transfer protein activity levels in humans. *Atherosclerosis*, 132(2), 251-254. [https://doi.org/10.1016/S0021-9150\(97\)00099-3](https://doi.org/10.1016/S0021-9150(97)00099-3)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Acetylsalicylate and salicylates in foods

P.L.T.M. Karin Janssen^{a,*}, Martijn B. Katan^a, Wija A. van Staveren^a, Peter C.H. Hollman^b,
Dini P. Venema^b

^aDepartment of Human Nutrition, Agricultural University, Bomenweg 2, 6703 HD, Wageningen, The Netherlands

^bDLO State Institute for Quality Control of Agricultural Products, Wageningen, The Netherlands

Abstract

Acetylsalicylic acid is effective in the prevention of cardiovascular disease. It was suggested that fruits and vegetables provide unknown amounts of acetylsalicylic acid. We could not find any acetylsalicylic acid in 30 foods using HPLC with fluorescence detection (detection limits: 0.02 mg/kg for fresh, and 0.2 mg/kg for dried products). We showed that urinary excretion of salicylates is a valid indicator for intake, and found a median salicylate excretion of 10 μmol (1.4 mg) in 24 h urine of 17 volunteers eating a variety of diets. Our data suggest that the content of (acetyl)salicylic acid of diets may be too low to affect disease risk. © 1997 Elsevier Science Ireland Ltd.

Keywords: Acetylsalicylate; Diet; Excretion; Salicylate

Acetylsalicylic acid is effective in the prevention of cardiovascular disease [1,2], and is associated with a decreased risk for colon cancer [3,4]. Levels of salicylates in foods are thus of interest, but data are scarce and controversial. Swain et al. suggested that a normal mixed Western diet provides 72–1448 μmol (10–200 mg) of natural salicylates, and 17 μmol (3 mg) acetylsalicylate daily [5].

We measured levels of total salicylates and acetylsalicylate in 30 foods using a specific and sensitive HPLC method with fluorescence detection. Detection limits were 0.1 $\mu\text{mol}/\text{kg}$ for fresh, and 1.4 $\mu\text{mol}/\text{kg}$ for dried products. Levels of total salicylate were 10–100 times lower than published previously: we found total salicylate levels of 0–0.7 $\mu\text{mol}/100\text{ g}$ in vegetables

and fruits, and 2–20 $\mu\text{mol}/100\text{ g}$ in herbs and spices. Levels of acetylsalicylate were lower than the limit of detection in all foods. Based on these results, we estimate that a Western diet provides about 0–15 $\mu\text{mol}/\text{day}$ of total salicylates.

We subsequently gave 66 μmol of pure salicylic or 58 μmol of acetylsalicylic acid to six healthy volunteers in a placebo controlled cross-over study, and recovered 77–80% in 24-h urine. Thus, urinary excretion is a valid indicator for intake.

We studied 24-h urinary excretion levels of salicylates in 17 volunteers eating a wide variety of diets, in order to estimate the true salicylate content of the diets. Median excretion of salicylates in urine was 10 $\mu\text{mol}/24\text{ h}$ (range 6–12). These results are in accordance with our food analyses data.

We conclude that levels of total salicylates and acetylsalicylate in diets are low, and probably insufficient to affect disease risk.

* Corresponding author. Tel.: +31 317 484421; fax: +31 317 483342; e-mail: karin.janssen@et3.voed.wau.nl

Acknowledgements

Funding was received from the Netherlands Heart Foundation (93.084) and Foundation for Nutrition and Health Sciences, The Netherlands.

References

- [1] Fuster, V., Dyken, M.L., Vokonas, P.S. and Hennekens, C. (1993) Aspirin as a therapeutic agent in cardiovascular disease. *Circulation*, 87, 659–675.
- [2] Badimon, J.J., Fuster, V., Chesebro, J.H. and Badimon, L. (1993) Coronary atherosclerosis; a multifactorial disease. *Circulation*, 87 (suppl. II), 3–16.
- [3] Giovannucci, E., Rimm, E.B. and Stampfer, M.J. et al. (1994) Aspirin use and the risk for colorectal cancer and adenoma in male health professionals. *Ann. Intern. Med.*, 121, 241–246.
- [4] Giovannucci, E., Egan, K.M., Hunter, D.J., Stampfer, M.J., Colditz, G.A., Willet, W.C. and Speizer, F.E. (1995) Aspirin and the risk of colorectal cancer in women. *N. Engl. J. Med.*, 333, 609–614.
- [5] Swain, A.R., Dutton, S.P. and Truswell, A.S. (1985) Salicylates in foods. *J. Am. Diet Assoc.*, 85, 950–960.