

VU Research Portal

Towards Doppler-free two-photon spectroscopy of trapped and cooled HD⁺ ions

Patra, S.

2019

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Patra, S. (2019). *Towards Doppler-free two-photon spectroscopy of trapped and cooled HD⁺ ions*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Contents

1	Introduction	1
1.1	Prologue	1
1.2	Determination of the proton-electron mass ratio μ_{pe}	3
1.3	The hydrogen molecular ion HD^+ : Benchmark system for testing molecular theory	4
1.4	Fundamental constants from HD^+	6
1.5	Test of the Standard Model of particle physics and search for physics beyond it	8
1.6	Vibrational spectroscopy of the HD^+ ion	10
1.7	Outline and summary of this thesis	11
2	Necessary background information	15
2.1	Advances in <i>ab initio</i> calculations of the energy levels of HD^+	15
2.1.1	Summary of recent advances in the calculations of the ro-vibrational transition frequencies of HD^+	15
2.1.2	Hyperfine structure calculation	16
2.2	Summary of spectroscopy of the $(v, L) : (0, 2) \rightarrow (8, 3)$ overtone in HD^+	17
2.3	Doppler-free two-photon spectroscopy of ro-vibrational transitions of HD^+	23
2.3.1	Feasibility study	23
2.3.2	Limitations of the OBE model	30
3	Experimental setup	33
3.1	Laser-cooling of Be^+ ions and the ion trap setup	33
3.1.1	Ion trap setup	33
3.1.2	Laser-cooling setup for Be^+ ions	35
3.1.3	Experimental control and automation	39
3.1.4	Performance of the ion trap and laser-cooling setup	43
3.2	Frequency stabilisation of the spectroscopy lasers	44
		iii

3.2.1	The optical frequency comb (OFC) and the ultra-stable laser	45
3.2.2	The transfer-oscillator locking scheme	46
	Principle and implementation	46
	Performance of the frequency stabilisation scheme	54
	Effect of intra-office optical fibre on the spectra purity of the lasers	58
	Selection of f_{rep} of the OFC for multiple lasers	60
4	<i>A priori</i> determination of required experimental parameters	63
4.1	Introduction	63
4.2	Choice of hyperfine components	64
4.2.1	Zeeman shifts	64
4.2.2	Suppression of Doppler-broadened background	66
4.3	Modelling the Doppler-free two-photon spectrum: <i>a priori</i> determination of required experimental parameters	68
4.3.1	Brief review of the rate equation model	69
4.3.2	Calculation of the $(v, L) : (0, 3) \rightarrow (4, 2) \rightarrow (9, 3)$ Doppler-free two-photon spectrum: results	72
4.3.3	Estimate of frequency shift due to underlying Doppler-broadened background	77
4.4	Zeeman and AC Stark shifts	79
4.4.1	Zeeman shift	79
4.4.2	AC Stark shift due to the cooling laser at 313 nm and the dissociation laser at 532 nm	81
	Appendices	84
4.A	AC Stark shift due to the cooling laser and the dissociation laser	84
5	Ghost features in Doppler-broadened spectra of ro-vibrational transitions in trapped HD^+ ions	87
5.1	Introduction	87
5.2	Brief review of the rate equation model	88
5.3	Calculation of the $(v, L) : (0, 3) \rightarrow (4, 2)$ spectrum	90
5.4	Ghost features in $(v, L) : (0, 3) \rightarrow (4, 2)$ single-photon spectrum	92
5.5	Significance of ghost features for spectroscopy of trapped ions	99
5.6	Summary and conclusion	100
6	Proton-electron mass ratio from HD^+ revisited	103
6.1	Introduction	103
6.2	Improvement in theory	104
6.3	Comparison between experiment and theory	107

6.4	Determination of the proton-electron mass ratio μ_{pe}	110
6.5	Outlook and conclusion	115
6.6	Acknowledgments	116
7	Observation of the $(v, L) : (0, 3) \rightarrow (9, 3)$ two-photon transition in HD^+	117
7.1	Doppler-broadened $(v, L) : (0, 3) \rightarrow (9, 3)$ ro-vibrational transition in HD^+	117
7.2	Doppler-free two-photon $(v, L) : (0, 3) \rightarrow (9, 3)$ transition in HD^+	119
7.2.1	Experimental protocol and measured spectrum	119
7.2.2	Statistical test to distinguish “signal” from “noise”	123
	The statistical test	123
	Welch’s t -test	125
	Statistical analysis of the measured spectrum	126
	Appendices	130
7.A	Choice of optimum REMPD duration for the Doppler-free two-photon spectroscopy experiment	130
8	Conclusion and Outlook	135
	Bibliography	137
	Acknowledgement	153