

VU Research Portal

Contextual correlates of dietary behaviours in adults across Europe

Matias de Pinho, M.G.

2019

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Matias de Pinho, M. G. (2019). *Contextual correlates of dietary behaviours in adults across Europe*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Chapter 2

Exploring the relationship between perceived barriers to healthy eating and dietary behaviours in European adults

Published as:

Pinho MGM, Mackenbach JD, Charreire H, Oppert JM, Bárdos H, Glonti K, Rutter H, Compernelle S, De Bourdeaudhuij I, Beulens JWJ, Brug J, Lakerveld J.

Exploring the relationship between perceived barriers to healthy eating and dietary behaviours in European adults. *European journal of nutrition*. 2018 Aug 1;57(5):1761-70.

Abstract

Purpose: Dietary behaviours may be influenced by perceptions of barriers to healthy eating. Using data from a large cross-European study (n=5,900), we explored associations between various perceived barriers to healthy eating and dietary behaviours among adults from urban regions in five European countries, and examined whether associations differed across regions and socio-demographic characteristics, such as age and sex. **Methods:** Frequency of consumption of fruit, vegetables, fish, fast food, sugar-sweetened beverages, sweets, breakfast and home-cooked meals were split by the median into higher and lower consumption. We tested associations between barriers (irregular working hours; giving up preferred foods; busy lifestyle; lack of willpower; price of healthy food; taste preferences of family and friends; lack of healthy options and unappealing foods) and dietary variables using multilevel logistic regression models. We explored whether associations differed by age, sex, education, urban region, weight status, household composition or employment. **Results:** Respondents who perceived any barrier were less likely to report higher consumption of healthier foods and more likely to report higher consumption of fast food. 'Lack of willpower', 'time constraints' and 'taste preferences' were most consistently associated with consumption. For example, those perceiving lack of willpower ate less fruit (odds ratio (OR) 0.57; 95% confidence interval (CI) 0.50 – 0.64), and those with a busy lifestyle ate less vegetables (OR 0.54; 95% CI 0.47 – 0.62). Many associations differed in size, but not in direction, by region, sex, age, and household composition. **Conclusion:** Perceived 'lack of willpower', 'time constraints' and 'taste preferences' were barriers most strongly related to dietary behaviours, but the association between various barriers and lower intake of fruit and vegetables was somewhat more pronounced among younger participants and women.

Background

Maintaining healthy dietary behaviours (e.g. diet that is rich in fruit and vegetables and low consumption of foods that are high in saturated fat and sugar) is crucial for population health and the prevention of non-communicable disease [1-7]. Both contextual ('midstream' and 'upstream') and individual ('downstream') factors can influence dietary behaviours [8]. We recently studied the interactions between neighbourhood characteristics and the number of individual perceived barriers with obesity-related behaviour [9]. Perceived barriers to healthy eating are an important individual-level factor [10,11] and people who perceive a greater number of barriers are more likely to report to consume less healthy diets [12]. As suggested by health behaviour theories (i.e. *Social Cognitive Theory* and the *Theory of Planned Behaviour*), individuals who perceive more barriers have lower motivation, lower levels of self-efficacy and possibly lower behavioural control required to maintain a healthy diet [13,14]. Across studies, the most frequently reported barriers to healthy eating relate to time constraints, taste preferences and monetary costs [15-18].

The majority of previous studies were conducted within specific populations [12,16,18-22], from which we have learned that these barriers may differ across subgroups. For instance, older women are more likely to report that healthy food is easily available and are less likely to perceive lack of time as a barrier to healthy eating [20]. Although these studies have provided an indication of some of the most common barriers that individuals perceive and their relationship to diet, they often provided only descriptive analysis [15,23], and thus far have stopped short of comparing the relation of perceived barriers to food consumption in a diverse sample across different countries [12,22,24]. We aimed to address this gap by focusing on the association between perceived barriers to healthy eating and different types of dietary behaviours among adults within urban regions in five different European countries. Considering that barriers to healthy eating may differ across subgroups and the lack of evidence available for more general populations, we also wanted to explore whether associations, if present, differed by age groups, sex, educational attainment, weight status, household composition, employment status and across urban regions.

Methods

Study design, sampling and participants

This study was part of the SPOTLIGHT project [25]. A survey was conducted in five urban regions across Europe: Ghent and suburbs (Belgium), Paris and inner suburbs (France), Budapest and suburbs (Hungary), the Randstad (a conurbation including the cities of Amsterdam, Rotterdam, The Hague and Utrecht in the Netherlands) and Greater London (UK). Neighbourhood sampling was based on a combination of residential density and socioeconomic status (SES) data at the neighbourhood level. This resulted in four types of pre-specified neighbourhoods: low SES/low residential density, low SES/high residential density, high SES/low residential density and high SES/high residential density. In each country, three neighbourhoods of each type were randomly sampled (i.e. 12 neighbourhoods per country, 60 neighbourhoods in total). Adults (18 years and older) living in the selected neighbourhoods were invited to participate in a web-based survey which included questions on demographics, neighbourhood perceptions, social environmental factors, health, motivations for and barriers to healthy behaviour, obesity-related behaviours, as well as weight and height. A total of 6,037 - 10.8%, out of 55,893 invited - subjects participated in the survey which took place between February and September 2014. For more details on sampling, design and participant recruitment, please refer to Lakerveld et al. (2015) [26]. Local ethics committees in each of the countries approved the study. All survey participants included in the analysis provided informed consent.

Measures

Individual characteristics

Information on age, sex, height, weight, educational attainment, household composition (total number of adults and children), employment status, urban region of residence, dietary behaviours and perceived barriers to healthy eating was collected through the online survey. Body Mass Index (BMI) was calculated on the basis of self-reported weight and height.

Dietary behaviours

We asked participants how many times a week they consumed fruit, vegetables, fish, fast food, sugar-sweetened beverages and sweets. The respondents also reported how often they ate breakfast and how many times a week they, or someone in their household, prepared home-cooked meals using ingredients, as opposed to eat ready-made or takeaway meals. The response options were: once a week or less; 2 times a week; 3 times a week; 4 times a week; 5 times a week; 6 times a week; 7 times a week (each day); twice a day; more than twice a day. Because the dietary variables were not normally distributed, and our questionnaire focused on frequency of consumption rather than quantity, we were not able to dichotomise the variables with cut-offs based on quantitative dietary guidelines. Therefore, we dichotomised these variables based on observed median values as follows: consumption of fruit, vegetables and breakfast <7 or ≥ 7 times a week; home-cooked meals <6 or ≥ 6 times a week; sweets <3 or ≥ 3 times a week; sugar-sweetened beverages, fish and fast food <2 or ≥ 2 times a week. These dichotomous variables are further referred to as *healthier dietary behaviours* (consumption above the median of fruit, vegetables, fish, home-cooked meals, breakfast) and *less healthy dietary behaviours* (consumption above the median of fast food, sugar-sweetened beverages and sweets).

Perceived barriers to healthy eating

The types of barriers to healthy eating were derived from the pan-European consumer attitudinal study [15], and included eight items: irregular working hours; giving up preferred foods; busy lifestyle; lack of willpower; price of healthy food; taste preferences of family and friends; lack of healthy options and unappealing foods. We asked participants the following question: 'How often do these reasons prevent you from eating a healthy diet?'. The response options were never; seldom; sometimes; often and almost always. Due to the distribution of cases across the five categories of the perceived barrier variables, we created a dichotomous variable for each barrier. The "not perceived as a barrier" category was created by merging the responses options *never and seldom*. The "perceived as a barrier" category was the result of merging the *sometimes, often and almost always* options.

Statistical analysis

A total of 137 individuals were excluded from further analyses as their residential neighbourhood could not be identified, resulting in an analytical sample of 5,900 participants.

Weight status – based on self-reported weight and height – was defined according to the World Health Organization BMI cut-off points: under/normal weight (BMI <25 kg/m²); overweight/obese (BMI ≥25 kg/m²) [27]. Age was classified into three groups: 18-40 years old; 41-64 years old; 65 years old or older. Due to differences in country-specific education systems, educational attainment was classified into two groups: ‘lower’ (higher secondary education or less) and ‘higher’ (college or university level). Household composition was classified into three groups: one person; two people and three people or more. Employment status (which includes people who were employed or in education) was classified into two groups: yes or no.

As the percentage of missing values ranged from 1% (age and sex) to 16.4% (fast food consumption), complete case analysis could potentially lead to biased results. We therefore performed multiple imputation on all variables (including outcomes and determinants) [28]. To that end, data were assumed to be missing at random, i.e. the probability of a value being missing depends on other observed values but not on the missing value itself [29]. Based on the percentage of missing values, we chose to impute 20 datasets, following the recommendation of Rubin [29] and Bodner [30]. This was done via predictive mean matching using STATA® 14.

Descriptive analyses were performed on un-imputed data. To verify the association between the different perceived barriers and food consumption variables, multilevel logistic regression analyses with random intercepts were conducted with individual characteristics as the first level: age (continuous), sex, educational attainment, BMI (continuous), household composition, employment status. Those were nested within residential neighbourhoods, the second level. We tested for effect modification by age, sex, educational attainment, weight status, household composition and employment status. To do so, interaction terms were added to the models. For sensitivity analysis, we checked how many interactions would remain significant by using a stricter value for statistical significance ($p < 0.001$ instead of 0.05). For further sensitivity analysis, we tested which of the barriers remained significantly associated and were most strongly related to dietary behaviour in a multivariable logistic regression model with all

the barriers as predictor variables for each outcome. All analyses were performed using STATA® 14.

Results

The mean age of the participants was 52 years (standard deviation 16.4). Just over half the participants were female (55.9 %) and highly educated (53.5%) (Table 1). The percentage of respondents who were overweight or obese was 45.7%. With regard to dietary behaviours, 80.6% of participants reported having breakfast every day and 37.8% reported to eat fish at least twice a week. The most frequently stated perceived barrier to healthy eating was 'lack of willpower' (44.6%) followed by 'busy lifestyle' (42.9%), 'price of healthy foods' (31.8%) and 'irregular working hours' (31.5%). Descriptive results by urban regions show that a 'lack of willpower' was the most frequently mentioned barrier in France, the Netherlands and the United Kingdom. In Belgium, the most frequently mentioned barrier was having a 'busy lifestyle' and in Hungary, it was 'price of healthy food' (data not shown).

Interaction terms for all potential effect modifiers (age, sex, educational attainment, weight status, household composition, employment status and urban region) were significant for at least one or more associations, and stratified analyses were conducted accordingly. Since there were few differences in the direction of the associations between strata and differences in effect sizes were similar to non-stratified outcomes, we present the overall – non stratified – results in Table 2 and provide all stratified results in Supplementary Tables 1 to 8.

Almost all barriers were inversely and significantly associated with the consumption of healthier foods. The strongest barrier to higher consumption of many food items was self-reported 'lack of willpower'. Respondents reporting this barrier had a lower probability to have higher levels of consumption of home-cooked meals, fruit, vegetables and fish (by 52.0%, 43.0%, 53.0% and 33.0%, respectively). In general, the barriers were most strongly related to vegetable consumption and cooking meals at home. For vegetable consumption, barriers related to willpower, time, price and taste were particularly important, and for home cooked meals the barriers related with willpower and time were most strongly related. Barriers related with time were also important for having breakfast as participants who reported working irregular hours were 38% less likely to report having breakfast seven days a week (Table 2).

Table 1. Descriptive characteristics of the participants to the SPOTLIGHT survey. Analytical sample n = 5,900

| Characteristics | n | Mean (Standard deviation) |
|---|----------|----------------------------------|
| Age | 5,841 | 52 (16.4) |
| BMI | 5,195 | 25.2 (4.51) |
| | | % |
| Sex | 5,841 | |
| Female | | 55.9 |
| Educational attainment | 5,335 | |
| Higher | | 53.5 |
| Weight status (kg/m²) | 5,195 | |
| Overweight/obese | | 45.7 |
| Household composition | 5,330 | |
| 1 person | | 22.6 |
| 2 people | | 39.2 |
| 3 people or more | | 38.2 |
| Employed or in education | 5,878 | |
| Yes | | 58.5 |
| Urban region | 5,900 | |
| Ghent and suburbs (Belgium) | | 30.1 |
| Paris and inner suburbs (France) | | 13.9 |
| Budapest and suburbs (Hungary) | | 14.8 |
| Randstad (conurbation in The Netherlands) | | 27.3 |
| Greater London (United Kingdom) | | 13.9 |
| Perceived barriers – Frequency mentioned | | |
| Irregular working hours | 5,156 | 31.5 |
| Giving up preferred foods | 5,106 | 30.3 |
| Busy lifestyle | 5,149 | 42.9 |
| Lack of willpower | 5,153 | 44.6 |
| Price of healthy foods | 5,148 | 31.8 |
| Taste preferences of family and friends | 5,102 | 28.8 |
| Lack of healthy options | 5,124 | 17.8 |
| Unappealing foods | 5,078 | 20.2 |
| Healthier dietary behaviours | | |
| Higher consumption of fruit | 5,353 | 53.9 |
| Higher consumption of vegetable | 5,412 | 60.1 |
| Higher consumption of fish | 5,337 | 37.8 |
| Higher consumption of breakfast | 5,406 | 80.6 |
| Higher consumption of home-cooked meals | 5,351 | 64.9 |
| Unhealthier dietary behaviours | | |
| Higher consumption of fast food | 4,935 | 6.0 |
| Higher consumption of sugar-sweetened beverages | 5,221 | 53.4 |
| Higher consumption of sweets | 5,297 | 54.0 |

'Higher consumption' means consumption above the correspondent median value for each food item. Median value for fruit and vegetables <7 or ≥7 times a week; For breakfast <7 or equal to 7; For home-cooked meals <6 or ≥6 times a week; Sweets <3 or ≥3 times; For sugar-sweetened beverages, fish and fast food <2 or ≥2 times a week. Educational attainment: Higher (college or university level). Perceived barriers to health eating as dichotomous variable, the percentage indicates participants that answered 'perceived it as a barrier'.

Table 2. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating overall associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900)

| Barriers * | Fruit | Vegetables | Fish | Breakfast |
|---|--|--|--|--|
| | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) |
| Irregular working hours | 0.70 (0.61 – 0.80) ^e | 0.68 (0.59 – 0.79) | 0.91 (0.79 – 1.05) ^c | 0.62 (0.53 – 0.73) |
| Giving up preferred foods | 0.67 (0.59 – 0.76) | 0.55 (0.48 – 0.63) ^{b g} | 0.76 (0.66 – 0.87) ^e | 0.70 (0.60 – 0.81) ^f |
| Busy lifestyle | 0.65 (0.57 – 0.73) ^e | 0.54 (0.47 – 0.62) ^b | 0.85 (0.75 – 0.98) ^{a g} | 0.64 (0.55 – 0.74) ^g |
| Lack of willpower | 0.57 (0.50 – 0.64) ^g | 0.47 (0.42 – 0.54) ^{b c g} | 0.67 (0.59 – 0.76) ^e | 0.63 (0.54 – 0.73) ^{f g} |
| Price of healthy foods | 0.65 (0.58 – 0.75) ^d | 0.53 (0.47 – 0.61) ^{e g} | 0.67 (0.58 – 0.78) ^{e g} | 0.70 (0.59 – 0.82) ^{e f} |
| Taste preferences of family and friends | 0.72 (0.63 – 0.81) ^g | 0.65 (0.57 – 0.74) ^{a e} | 0.81 (0.70 – 0.94) ^{e g} | 0.86 (0.74 – 1.00) |
| Lack of healthy options | 0.82 (0.70 – 0.95) ^{d f g} | 0.74 (0.63 – 0.87) | 0.80 (0.67 – 0.95) ^e | 0.69 (0.58 – 0.82) ^g |
| Unappealing foods | 0.60 (0.52 – 0.69) ^{a d} | 0.54 (0.47 – 0.63) ^{a b d} | 0.72 (0.61 – 0.86) ^g | 0.63 (0.53 – 0.75) |
| Barriers * | Home-cooked meals | Fast food | Sweets | Sugar-sweetened beverages |
| | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) |
| Irregular working hours | 0.53 (0.45 – 0.61) ^g | 1.93 (1.45 – 2.57) | 1.20 (1.05 – 1.38) ^a | 1.40 (1.22 – 1.61) ^g |
| Giving up preferred foods | 0.60 (0.53 – 0.69) ^g | 1.73 (1.34 – 2.24) ^c | 1.28 (1.12 – 1.47) ^{b d} | 1.25 (1.10 – 1.43) |
| Busy lifestyle | 0.52 (0.45 – 0.59) ^{a c f g} | 2.07 (1.57 – 2.73) ^f | 1.13 (0.99 – 1.28) ^g | 1.41 (1.23 – 1.61) |
| Lack of willpower | 0.48 (0.42 – 0.55) ^g | 2.06 (1.60 – 2.66) | 1.45 (1.29 – 1.64) ^{b g} | 1.43 (1.27 – 1.61) ^b |
| Price of healthy foods | 0.66 (0.57 – 0.77) ^{a e g} | 1.59 (1.20 – 2.11) ^g | 0.97 (0.85 – 1.11) ^g | 0.93 (0.81 – 1.07) |
| Taste preferences of family and friends | 0.81 (0.71 – 0.93) ^{e f g} | 1.48 (1.13 – 1.94) | 0.98 (0.86 – 1.13) | 1.21 (1.05 – 1.39) |
| Lack of healthy options | 0.71 (0.60 – 0.83) ^g | 1.67 (1.24 – 2.23) ^{a c} | 0.87 (0.77 – 1.04) | 1.06 (0.91 – 1.24) ^{b g} |
| Unappealing foods | 0.65 (0.56 – 0.76) | 2.07 (1.54 – 2.77) ^e | 1.01 (0.88 – 1.17) ^e | 1.48 (1.28 – 1.72) |

* Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely). All analysis were performed in separated models and adjusted for age, sex educational attainment, BMI, household composition, employment status and urban region (urban region stands for the sampled regions in Belgium, France, Hungary, Netherlands and United Kingdom); Results presented in bold were statistically significant (p<0.05). Effect modification by: ^a age group; ^b sex; ^c education; ^d weight status; ^e household composition; ^f employment; ^g urban regions. Odds Ratios for dietary outcomes refer to 'Higher consumption' of food items which means consumption above the correspondent median value for each food item. Median value for fruit and vegetables <7 or >7 times a week; For breakfast <7 or equal to 7; For home-cooked meals <6 or >6 times a week; Sweets <3 or >3 times; For sugar-sweetened beverages, fish and fast food <2 or >2 times a week.

All the reported barriers to healthy eating were positively and significantly associated with fast food consumption. Respondents who reported having a 'busy lifestyle' and a 'lack of willpower' and who framed 'healthy food as being unappealing' were twice as likely to consume fast food at least twice a week than those who did not report such barriers. Although almost all barriers (with the exception of 'cost of healthy food' and 'lack of healthy options') were significantly associated with higher sugar-sweetened beverage consumption, the effect sizes of the associations with this dietary behaviour were relatively small. Higher consumption of sweets was directly associated with 'irregular working hours', 'giving up preferred foods' and 'lack of willpower'.

The variable that was most often a significant effect modifier was urban region, which appeared to be particularly relevant in relation to the barriers associated with consumption of home-cooked meals and vegetables (Supplementary Table 7). Another often significant effect modifier was household composition: the association between almost all barriers and fish consumption was mainly significant for people living in a two-person household. The exception was the barrier 'taste preference of family and friends' which was significantly associated with fish consumption among respondents in three or more person households (Supplementary Table 5). Age group and sex were also frequently significant effect modifiers, especially in the associations between the barrier 'unappealing food' and consumption of fruit and vegetables. Younger participants who perceived healthy food to be unappealing were 52.0% less likely to report eating fruit every day and 59.0% less likely to report eating vegetables every day (Supplementary Table 1). Most associations between the barriers and vegetables consumption were stronger in women than in men (Supplementary Table 2).

Results from sensitivity analyses showed that even if we were to use a p value below 0.001, about half of the interactions would remain significant. In addition, when we included all the independent variables in the models (Supplementary Table 8), in almost all instances, the odds ratios (OR) became less strong as compared to the models in which each perceived barrier was analysed separately (Table 2). Nonetheless, 'lack of willpower' remained the barrier most strongly related to greater consumption of home-cooked meals, fruit, vegetables and fish. Most of the results remained in the same direction, but interestingly not the association between 'lack of healthy options' and fruit (OR 1.20, 95% confidence interval (CI) 1.01 – 1.42) and vegetables (OR 1.18, 95%CI 0.98 – 1.42) consumption.

Discussion

We explored the association between perceived barriers to healthy eating and dietary behaviours in adults across five urban regions in Europe. Among the participants in this study, the most often-mentioned barriers to healthy eating were 'lack of willpower', 'having a busy lifestyle' and the 'price of healthy foods'. The barriers were associated with frequency of consumption of fruit, vegetables, breakfast, home-cooked meals and fast food. The barriers were in particular strongly associated with vegetable consumption, and the time-related barriers appeared to be especially important for the consumption of home-cooked meals. Many of the associations between barriers and dietary behaviours were modified by socio-demographic factors and urban region.

Our finding that the reported barriers were often associated with self-reported dietary behaviours is in line with the results of Williams et al. (2012). They found that Australian women were more likely to eat healthily if they perceived to have a higher self-efficacy towards adhering to a healthy diet, did not perceive taste as a barrier to eating fruit and vegetables, and did not perceive time or price as barriers to eating healthily [31]. In our study, the barriers studied were less strongly associated with consumption of sweets and sugar-sweetened beverages compared to the other dietary behaviours. This may be due to the fact that consuming sugary foods and drinks requires less effort and planning, as these products are omnipresent in many settings in Western societies. On the other hand, eating foods such as vegetables and fish and the habit of cooking at home require more effort in terms of planning and/or skills, and may therefore be more susceptible to barriers.

We found that self-reported 'lack of willpower' was the most frequently reported barrier and was most strongly associated with a lower likelihood of having home-cooked meals, and eating fruit, vegetables and fish on a regular basis. This may reflect public and media discourse focused on individual level willpower as a key determinant of eating behaviour and obesity. Puhl and Heuer (2009) found in their review that it is common for the media to frame unhealthy eating, and consequently obesity, as a personal responsibility with individual level solutions such as behaviour change from unhealthy dietary habits into healthier ones [32]. Equally, it might be that individuals exposed to obesogenic environments feel that their own level of willpower is not always adequate to overcome the many opportunities and temptations for unhealthy eating their environment offers them. For instance, in another study within the SPOTLIGHT project, we found that the

relations between neighbourhood characteristics and the consumption of fruit and vegetables were often modified by a greater number of perceived barriers to healthy eating [9].

Other important barriers were taste-related (i.e. 'perceiving healthy foods as unappealing', 'taste preferences of family and friends', 'giving up preferred foods'). These appeared to be particularly relevant for vegetable consumption. In a Norwegian study, taste was the barrier most significantly associated with constraints in consumption of fish and vegetables [16]. From our results, taste-related barriers were also associated with less frequent fish consumption, but we found other barriers (price and willpower) to be more strongly related to fish consumption. High cost was also the most cited barrier to fish consumption in an Australian study of older adults [33]. Based on our results, few people reported eating fast food frequently, but those who did were more likely to perceive time and taste as barriers to healthy eating. This is in accordance with previous studies [22,34].

We frequently found time related barriers (i.e. 'irregular working hours' and 'busy lifestyle') being associated with reduced frequency of consumption of vegetables and home-cooked meals. In this sample, people who experienced time constraints were less likely to have home-cooked meals regularly. Similar results were found in a study conducted in the USA in which working adults who cared the most about convenience were those who spent the least time on home cooking [35]. Home cooking, in turn, was associated with having a healthier diet (i.e. greater consumption of fruits and vegetables and less frequent visits to fast-food restaurants) [35]. In previous studies, time related barriers were also associated with reduced frequency of vegetables consumption [22,24].

We also tested whether the associations were modified by urban region and socio-demographic variables. For many of the associations we found effect modification, with stratified analyses indicating that the barrier-consumption relation was stronger in some sub-groups than in others. In general, subgroup analyses showed that the associations between barriers and the consumption of fruit and vegetables were more pronounced in younger participants and women; the relation between barriers and lower consumption of vegetables and higher consumption of sweets were stronger in women; and the relation between barriers and fast food was stronger for the higher educated individuals. The barriers and the barrier-consumption associations were often different between urban regions, most notably in relation to consuming home-cooked meals: in contrast to what was found for other urban regions, in Hungary, no barriers were associated with

the likelihood of consuming home-cooked meals. The associations between barriers and dietary behaviours were also different depending on the type of household. We found that in three-person households, which are likely to contain children, the barrier 'taste preferences of family and friends' was more important for fish consumption than for smaller households. This is concordant with the results from an intervention study with women who were trying to adopt healthier dietary behaviours, which described that preferences of children and family was the most important barrier, especially in trying to increase the consumption of vegetables, lentils and fish [36].

The association between the taste-related barrier (finding healthy foods unappealing) with fruit and vegetable consumption differed across age groups. Our results suggest that taste preferences for fruit and vegetables may be more important for younger adults than for older adults. In line with this, in a ten-year longitudinal study, Larson et al. (2012) found that having a favourable taste preference for fruit and vegetables was an important predictor of increased consumption of both items from adolescence into adulthood [37]. Based on previous studies that have shown that more highly educated individuals exhibit healthier lifestyle behaviour [38,39], and that specific barriers such as taste preferences are more prevalent among lower educated people [40] we expected that education would be an effect modifier in many of the associations studied, but interestingly, this was not the case.

This study needs to be seen in the light of some limitations, for instance, the use of self-reported measures of dietary behaviours to obtain information on the consumption of a limited number of specific foods. Nonetheless, it is known that self-reported measures can provide valuable information on the consumption of foods and beverages in population-based studies [41]. In addition, our study included items that have previously been associated with having a healthy diet and consistent with current dietary recommendations [42,43]. The categorisation of dietary behaviours can also be seen as a limitation, as we were unable to distinguish participants who never consume certain foods from those who consume them at least once a week. This may especially be a limitation for less frequently consumed foods, such as fish. In addition, due to their distribution we decided to dichotomise the perceived barriers variables, potentially losing important nuances such as the difference between those who perceived barriers seldom (categorised as 'no barrier') and sometimes (categorised as barrier). We also may not have been able to account for all important barriers that people might experience. Informed by previous research [15], we chose barriers that appeared to be most important in terms of their relation with healthy eating, but there may be other relevant barriers

that have not been studied before. In this regard, a qualitative study alongside this quantitative analysis to identify additional relevant barriers for further research is warranted. Lastly, a low response rate (around 10%), which is a common problem for population-based studies [44], may have led to a selection bias. Although the number of men and women; lower and higher educated individuals; and age groups are balanced in our sample, generalisation of findings should be done with caution [26].

The study's strengths include our ability to recruit a large sample across different countries in Europe, which contributes to higher external validity and enables comparisons across urban regions. In addition, we were able to link several perceived barriers to healthy eating with the consumption of healthy and unhealthy dietary behaviours in a diverse sample, in which individuals varied in terms of age (younger and older adults), sex and socio-demographics characteristics. Furthermore, most previous studies on perceived barriers to healthy eating have focused on the evaluation of barriers related to the consumption of specific healthy food items, namely, fruit, vegetables and/or fish [16,18,24,45]. A strength of the current study is that we were able to link multiple generic barriers to different dietary habits, thus allowing for a comparison of the importance of these barriers for a number of dietary habits. In doing so, we believe we have contributed to a more comprehensive understanding of the subjective factors that can influence people's dietary behaviours.

In conclusion, we found several associations between perceived barriers to healthy eating and food consumption, of which the most frequent was self-reported lack of willpower. People who perceived any barrier to healthy eating were less likely to report healthier dietary behaviours, especially vegetables consumption, but also consumption of fruit, fish, breakfast and home-cooked meals, and were more likely to report eating fast food. Findings from this study may contribute to the design of interventions that target individual-level barriers to healthy eating since we found that associations between perceived barriers to healthy eating and food consumption were different across urban regions and subgroups. For instance, interventions aiming to increase fruit and vegetables consumption among adults could focus on taste related issues, especially among younger adults and women. However, upstream responses that shift the balance of influences on people's diets through promoting a healthier food environment may well have an important part to play in attenuating some of these negative influences that people perceive.

References - Chapter 2

1. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, Amann M, Anderson HR, Andrews KG, Aryee M, Atkinson C, Bacchus LJ, Bahalim AN, Balakrishnan K, Balmes J, Barker-Collo S, Baxter A, Bell ML, Blore JD, Blyth F, Bonner C, Borges G, Bourne R, Boussinesq M, Brauer M, Brooks P, Bruce NG, Brunekreef B, Bryan-Hancock C, Bucello C, Buchbinder R, Bull F, Burnett RT, Byers TE, Calabria B, Carapetis J, Carnahan E, Chafe Z, Charlson F, Chen H, Chen JS, Cheng AT, Child JC, Cohen A, Colson KE, Cowie BC, Darby S, Darling S, Davis A, Degenhardt L, Dentener F, Des Jarlais DC, Devries K, Dherani M, Ding EL, Dorsey ER, Driscoll T, Edmond K, Ali SE, Engell RE, Erwin PJ, Fahimi S, Falder G, Farzadfar F, Ferrari A, Finucane MM, Flaxman S, Fowkes FG, Freedman G, Freeman MK, Gakidou E, Ghosh S, Giovannucci E, Gmel G, Graham K, Grainger R, Grant B, Gunnell D, Gutierrez HR, Hall W, Hoek HW, Hogan A, Hosgood HD, 3rd, Hoy D, Hu H, Hubbell BJ, Hutchings SJ, Ibeanusi SE, Jacklyn GL, Jasrasaria R, Jonas JB, Kan H, Kanis JA, Kassebaum N, Kawakami N, Khang YH, Khatibzadeh S, Khoo JP, Kok C, Laden F, Lalloo R, Lan Q, Lathlean T, Leasher JL, Leigh J, Li Y, Lin JK, Lipshultz SE, London S, Lozano R, Lu Y, Mak J, Malekzadeh R, Mallinger L, Marcenes W, March L, Marks R, Martin R, McGale P, McGrath J, Mehta S, Mensah GA, Merriman TR, Micha R, Michaud C, Mishra V, Mohd Hanafiah K, Mokdad AA, Morawska L, Mozaffarian D, Murphy T, Naghavi M, Neal B, Nelson PK, Nolla JM, Norman R, Olives C, Omer SB, Orchard J, Osborne R, Ostro B, Page A, Pandey KD, Parry CD, Passmore E, Patra J, Pearce N, Pelizzari PM, Petzold M, Phillips MR, Pope D, Pope CA, 3rd, Powles J, Rao M, Razavi H, Rehfuss EA, Rehm JT, Ritz B, Rivara FP, Roberts T, Robinson C, Rodriguez-Portales JA, Romieu I, Room R, Rosenfeld LC, Roy A, Rushton L, Salomon JA, Sampson U, Sanchez-Riera L, Sanman E, Sapkota A, Seedat S, Shi P, Shield K, Shivakoti R, Singh GM, Sleet DA, Smith E, Smith KR, Stapelberg NJ, Steenland K, Stockl H, Stovner LJ, Straif K, Straney L, Thurston GD, Tran JH, Van Dingenen R, van Donkelaar A, Veerman JL, Vijayakumar L, Weintraub R, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams W, Wilson N, Woolf AD, Yip P, Zielinski JM, Lopez AD, Murray CJ, Ezzati M, AlMazroa MA, Memish ZA (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380 (9859):2224-2260. doi:10.1016/S0140-6736(12)61766-8
2. Bouvard V, Loomis D, Guyton KZ, Grosse Y, Ghissassi FE, Benbrahim-Tallaa L, Guha N, Mattock H, Straif K, International Agency for Research on Cancer

Monograph Working G (2015) Carcinogenicity of consumption of red and processed meat. *Lancet Oncol* 16 (16):1599-1600. doi:10.1016/S1470-2045(15)00444-1

3. Boffetta P, Couto E, Wichmann J, Ferrari P, Trichopoulos D, Bueno-de-Mesquita HB, van Duijnhoven FJ, Buchner FL, Key T, Boeing H, Nothlings U, Linseisen J, Gonzalez CA, Overvad K, Nielsen MR, Tjønneland A, Olsen A, Clavel-Chapelon F, Boutron-Ruault MC, Morois S, Lagiou P, Naska A, Benetou V, Kaaks R, Rohrmann S, Panico S, Sieri S, Vineis P, Palli D, van Gils CH, Peeters PH, Lund E, Brustad M, Engeset D, Huerta JM, Rodriguez L, Sanchez MJ, Dorronsoro M, Barricarte A, Hallmans G, Johansson I, Manjer J, Sonestedt E, Allen NE, Bingham S, Khaw KT, Slimani N, Jenab M, Mouw T, Norat T, Riboli E, Trichopoulou A (2010) Fruit and vegetable intake and overall cancer risk in the European Prospective Investigation into Cancer and Nutrition (EPIC). *J Natl Cancer Inst* 102 (8):529-537. doi:10.1093/jnci/djq072

4. Crowe FL, Roddam AW, Key TJ, Appleby PN, Overvad K, Jakobsen MU, Tjønneland A, Hansen L, Boeing H, Weikert C, Linseisen J, Kaaks R, Trichopoulou A, Misirli G, Lagiou P, Sacerdote C, Pala V, Palli D, Tumino R, Panico S, Bueno-de-Mesquita HB, Boer J, van Gils CH, Beulens JW, Barricarte A, Rodriguez L, Larranaga N, Sanchez MJ, Tormo MJ, Buckland G, Lund E, Hedblad B, Melander O, Jansson JH, Wennberg P, Wareham NJ, Slimani N, Romieu I, Jenab M, Danesh J, Gallo V, Norat T, Riboli E, European Prospective Investigation into C, Nutrition -Heart Study C (2011) Fruit and vegetable intake and mortality from ischaemic heart disease: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heart study. *Eur Heart J* 32 (10):1235-1243. doi:10.1093/eurheartj/ehq465

5. Hooper L, Abdelhamid A, Moore HJ, Douthwaite W, Skeaff CM, Summerbell CD (2012) Effect of reducing total fat intake on body weight: systematic review and meta-analysis of randomised controlled trials and cohort studies. *BMJ* 345:e7666. doi:10.1136/bmj.e7666

6. Imamura F, O'Connor L, Ye Z, Mursu J, Hayashino Y, Bhupathiraju SN, Forouhi NG (2015) Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. *BMJ* 351:h3576. doi:10.1136/bmj.h3576

7. Te Morenga L, Mallard S, Mann J (2013) Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ* 346:e7492. doi:10.1136/bmj.e7492
8. Swinburn B, Egger G, Raza F (1999) Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Preventive medicine* 29 (6 Pt 1):563-570. doi:10.1006/pmed.1999.0585
9. Mackenbach JD, Lakerveld J, Van Lenthe FJ, Teixeira PJ, Compernelle S, De Bourdeaudhuij I, Charreire H, Oppert JM, Bardos H, Glonti K, Rutter H, McKee M, Nijpels G, Brug J (2016) Interactions of individual perceived barriers and neighbourhood destinations with obesity-related behaviours in Europe. *Obes Rev* 17 Suppl 1:68-80. doi:10.1111/obr.12374
10. Sleddens EF, Kroeze W, Kohl LF, Bolten LM, Velema E, Kaspers P, Kremers SP, Brug J (2015) Correlates of dietary behavior in adults: an umbrella review. *Nutr Rev* 73 (8):477-499. doi:10.1093/nutrit/nuv007
11. Kremers SP, De Bruijn G-J, Visscher TL, Van Mechelen W, De Vries NK, Brug J (2006) Environmental influences on energy balance-related behaviors: a dual-process view. *International Journal of Behavioral Nutrition and Physical Activity* 3 (1):9
12. Lara J, McCrum LA, Mathers JC (2014) Association of Mediterranean diet and other health behaviours with barriers to healthy eating and perceived health among British adults of retirement age. *Maturitas* 79 (3):292-298. doi:10.1016/j.maturitas.2014.07.003
13. Ajzen I (1991) The theory of planned behavior. *Organizational Behavior and Human Decision Processes* 50 (179-211)
14. Bandura A (2004) Health promotion by social cognitive means. *Health Educ Behav* 31 (2):143-164. doi:10.1177/1090198104263660
15. Kearney JM, McElhone S (1999) Perceived barriers in trying to eat healthier--results of a pan-EU consumer attitudinal survey. *The British journal of nutrition* 81 Suppl 2:S133-137
16. Skuland SE (2015) Healthy Eating and Barriers Related to Social Class. The case of vegetable and fish consumption in Norway. *Appetite* 92:217-226. doi:10.1016/j.appet.2015.05.008

17. Yeh MC, Ickes SB, Lowenstein LM, Shuval K, Ammerman AS, Farris R, Katz DL (2008) Understanding barriers and facilitators of fruit and vegetable consumption among a diverse multi-ethnic population in the USA. *Health Promot Int* 23 (1):42-51. doi:10.1093/heapro/damo44
18. Pollard C, Miller M, Woodman RJ, Meng R, Binns C (2009) Changes in knowledge, beliefs, and behaviors related to fruit and vegetable consumption among Western Australian adults from 1995 to 2004. *American journal of public health* 99 (2):355-361. doi:10.2105/AJPH.2007.131367
19. Escoto KH, Laska MN, Larson N, Neumark-Sztainer D, Hannan PJ (2012) Work hours and perceived time barriers to healthful eating among young adults. *Am J Health Behav* 36 (6):786-796. doi:10.5993/AJHB.36.6.6
20. Williams L, Ball K, Crawford D (2010) Why do some socioeconomically disadvantaged women eat better than others? An investigation of the personal, social and environmental correlates of fruit and vegetable consumption. *Appetite* 55 (3):441-446. doi:10.1016/j.appet.2010.08.004
21. Trondsen T, Scholderer J, Lund E, Eggen AE (2003) Perceived barriers to consumption of fish among Norwegian women. *Appetite* 41 (3):301-314. doi:10.1016/s0195-6663(03)00108-9
22. Welch N, McNaughton SA, Hunter W, Hume C, Crawford D (2009) Is the perception of time pressure a barrier to healthy eating and physical activity among women? *Public Health Nutr* 12 (7):888-895. doi:10.1017/s1368980008003066
23. Shaikh AR, Yaroch AL, Nebeling L, Yeh MC, Resnicow K (2008) Psychosocial predictors of fruit and vegetable consumption in adults a review of the literature. *Am J Prev Med* 34 (6):535-543. doi:10.1016/j.amepre.2007.12.028
24. Mook K, Laraia BA, Oddo VM, Jones-Smith JC (2016) Food Security Status and Barriers to Fruit and Vegetable Consumption in Two Economically Deprived Communities of Oakland, California, 2013-2014. *Preventing chronic disease* 13:E21. doi:10.5888/pcd13.150402
25. Lakerveld J, Brug J, Bot S, Teixeira PJ, Rutter H, Woodward E, Samdal O, Stockley L, De Bourdeaudhuij I, van Assema P, Robertson A, Lobstein T, Oppert JM, Adany R, Nijpels G (2012) Sustainable prevention of obesity through integrated strategies: The SPOTLIGHT project's conceptual framework and design. *BMC public health* 12:793. doi:10.1186/1471-2458-12-793

26. Lakerveld J, Ben Rebah M, Mackenbach JD, Charreire H, Compernelle S, Glonti K, Bardos H, Rutter H, De Bourdeaudhuij I, Brug J, Oppert JM (2015) Obesity-related behaviours and BMI in five urban regions across Europe: sampling design and results from the SPOTLIGHT cross-sectional survey. *BMJ Open* 5 (10):e008505. doi:10.1136/bmjopen-2015-008505
27. WHO (2000) Obesity: preventing and managing the global epidemic. Part I – The problem of overweight and obesity. WHO Technical Report Series, 894. Geneva
28. Moons KGM, Donders RART, Stijnen T, Harrell FE (2006) Using the outcome for imputation of missing predictor values was preferred. *J Clin Epidemiol* 59 (10):1092-1101. doi:10.1016/j.jclinepi.2006.01.009
29. Rubin DB (1987) Multiple imputation for non-response in surveys. Wiley J & Sons, New York
30. Bodner TE (2008) What improves with increased missing data imputations? *Struct Equ Modeling* 15:651-675
31. Williams LK, Thornton L, Crawford D (2012) Optimising women's diets. An examination of factors that promote healthy eating and reduce the likelihood of unhealthy eating. *Appetite* 59 (1):41-46. doi:10.1016/j.appet.2012.03.014
32. Puhl RM, Heuer CA (2009) The stigma of obesity: a review and update. *Obesity (Silver Spring)* 17 (5):941-964. doi:10.1038/oby.2008.636
33. Grieger JA, Miller M, Cobiac L (2012) Knowledge and barriers relating to fish consumption in older Australians. *Appetite* 59 (2):456-463. doi:10.1016/j.appet.2012.06.009
34. Lucan SC, Barg FK, Long JA (2010) Promoters and barriers to fruit, vegetable, and fast-food consumption among urban, low-income African Americans--a qualitative approach. *American journal of public health* 100 (4):631-635. doi:10.2105/AJPH.2009.172692
35. Monsivais P, Aggarwal A, Drewnowski A (2014) Time spent on home food preparation and indicators of healthy eating. *Am J Prev Med* 47 (6):796-802. doi:10.1016/j.amepre.2014.07.033
36. Raberg Kjollesdal MK, Telle Hjellset V, Bjorge B, Holmboe-Ottesen G, Wandel M (2010) Barriers to healthy eating among Norwegian-Pakistani women

participating in a culturally adapted intervention. *Scand J Public Health* 38 (5 Suppl):52-59. doi:10.1177/1403494810378923

37. Larson N, Laska MN, Story M, Neumark-Sztainer D (2012) Predictors of fruit and vegetable intake in young adulthood. *Journal of the Academy of Nutrition and Dietetics* 112 (8):1216-1222. doi:10.1016/j.jand.2012.03.035

38. Chandola T, Clarke P, Morris JN, Blane D (2006) Pathways between education and health: a causal modelling approach. *J R Statist Soc A* 169 (2):337-359

39. Cutler DM, Lleras-Muney A (2006) Education and Health: Evaluating Theories and Evidence Paper presented at the The Health Effects of Non-Health Policies,

40. de Mestral C, Khalatbari-Soltani S, Stringhini S, Marques-Vidal P (2017) Fifteen-year trends in the prevalence of barriers to healthy eating in a high-income country. *The American journal of clinical nutrition*. doi:10.3945/ajcn.116.143719

41. Subar AF, Freedman LS, Tooze JA, Kirkpatrick SI, Boushey C, Neuhauser ML, Thompson FE, Potischman N, Guenther PM, Tarasuk V, Reedy J, Krebs-Smith SM (2015) Addressing Current Criticism Regarding the Value of Self-Report Dietary Data. *The Journal of nutrition* 145 (12):2639-2645. doi:10.3945/jn.115.219634

42. Pot GK, Richards M, Prynne CJ, Stephen AM (2014) Development of the Eating Choices Index (ECI): a four-item index to measure healthiness of diet. *Public Health Nutr* 17 (12):2660-2666. doi:10.1017/S1368980013003352

43. WHO (2015) Healthy diet. Fact sheet n. 394.

44. O'Neill TW, Marsden D, Matthis C, Raspe H, Silman AJ (1995) Survey response rates: national and regional differences in a European multicentre study of vertebral osteoporosis. *Journal of epidemiology and community health* 49 (1):87-93

45. Dijkstra SC, Neter JE, van Stralen MM, Knol DL, Brouwer IA, Huisman M, Visser M (2015) The role of perceived barriers in explaining socio-economic status differences in adherence to the fruit, vegetable and fish guidelines in older adults: a mediation study. *Public Health Nutr* 18 (5):797-808. doi:10.1017/S1368980014001487

Supplementary Tables of Chapter 2

Supplementary Table 1. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by age groups as a significant effect modification was found for the depicted associations (the table continues on next page).

| Barriers ^{ab} | | Fruit OR (95% CI) | Vegetables OR (95% CI) | Fish OR (95% CI) |
|----------------------------------|---|---------------------------|---------------------------|---------------------------|
| 18-40 years (n=1,681) | Busy lifestyle | - | - | 1.00 (0.79 – 1.29) |
| | Taste preferences of family and friends | - | 0.61 (0.48 – 0.76) | - |
| | Unappealing foods | 0.48 (0.36 – 0.63) | 0.41 (0.32 – 0.54) | - |
| 41-64 years (n=2,801) | Busy lifestyle | - | - | 0.77 (0.64 – 0.92) |
| | Taste preferences of family and friends | - | 0.63 (0.52 – 0.76) | - |
| | Unappealing foods | 0.66 (0.53 – 0.82) | 0.56 (0.44 – 0.70) | - |
| 65 years or more (n=1,402) | Busy lifestyle | - | - | 0.81 (0.58 – 1.13) |
| | Taste preferences of family and friends | - | 0.69 (0.49 – 0.98) | - |
| | Unappealing foods | 0.69 (0.50 – 0.95) | 0.77 (0.55 – 1.08) | - |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by sex, educational attainment, BMI, household composition, employment status and urban regions. Results presented in bold were statistically significant (p<0.05).

Supplementary Table 1. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by age groups as a significant effect modification was found for the depicted associations (table conclusion).

| Barriers ^{ab} | | Home-cooked meals OR (95% CI) | Fast food OR (95% CI) | Sweets OR (95% CI) |
|----------------------------------|-------------------------|----------------------------------|---------------------------|---------------------------|
| 18-40 years (n=1,681) | Irregular working hours | - | - | 1.08 (0.87 – 1.34) |
| | Busy lifestyle | 0.43 (0.34 – 0.56) | - | - |
| | Price of healthy foods | 0.77 (0.61 – 0.98) | - | - |
| | Lack of healthy options | - | 1.37 (0.94 – 1.97) | - |
| 41-64 years (n=2,801) | Irregular working hours | - | - | 1.17 (0.97 – 1.42) |
| | Busy lifestyle | 0.48 (0.39 – 0.58) | - | - |
| | Price of healthy foods | 0.57 (0.46 – 0.71) | - | - |
| | Lack of healthy options | - | 2.57 (1.60 – 4.13) | - |
| 65 years or more (n=1,402) | Irregular working hours | - | - | 1.74 (1.08 – 2.81) |
| | Busy lifestyle | 0.71 (0.49 – 1.04) | - | - |
| | Price of healthy foods | 0.48 (0.33 – 0.68) | - | - |
| | Lack of healthy options | - | 1.14 (0.39 – 3.33) | - |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by sex, educational attainment, BMI, household composition, employment status and urban regions. Results presented in bold were statistically significant (p<0.05).

Supplementary Table 2. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by sex as a significant effect modification was found for the depicted associations.

| | | Vegetables | Sweets | Sugar-sweetened beverages |
|------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| Barriers ^{ab} | | OR (95% CI) | OR (95% CI) | OR (95% CI) |
| Males (n=2,597) | Giving up preferred foods | 0.63 (0.52 – 0.77) | 1.06 (0.86 – 1.30) | - |
| | Busy lifestyle | 0.63 (0.51 – 0.76) | - | - |
| | Lack of willpower | 0.62 (0.51 – 0.75) | 1.29 (1.07 – 1.56) | 1.73 (1.43 – 2.09) |
| | Lack of healthy options | - | - | 0.77 (0.61 – 0.99) |
| | Unappealing foods | 0.71 (0.57 – 0.87) | - | - |
| Females (n=3,293) | Giving up preferred foods | 0.49 (0.41 – 0.58) | 1.50 (1.25 – 1.80) | - |
| | Busy lifestyle | 0.48 (0.40 – 0.58) | - | - |
| | Lack of willpower | 0.38 (0.32 – 0.45) | 1.59 (1.36 – 1.86) | 1.24 (1.06 – 1.45) |
| | Lack of healthy options | - | - | 1.28 (1.04 – 1.57) |
| | Unappealing foods | 0.44 (0.36 – 0.54) | - | - |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by age, educational attainment, BMI, household composition, employment status and urban regions. Results presented in bold were statistically significant (p<0.05).

Supplementary Table 3. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by educational attainment as a significant effect modification was found for the depicted associations.

| | | Vegetables | Fish | Home-cooked meals | Fast food |
|------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| Barriers ^{ab} | | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) |
| Lower (n=2738) | Irregular working hours | - | 0.77 (0.61 – 0.98) | - | - |
| | Giving up preferred foods | - | - | - | 1.29 (0.92 – 1.80) |
| | Busy lifestyle | - | - | 0.59 (0.48 – 0.73) | - |
| | Lack of willpower | 0.42 (0.35 – 0.51) | - | - | - |
| | Lack of healthy options | - | - | - | 1.51 (1.03 – 2.21) |
| Higher (n=3125) | Irregular working hours | - | 1.02 (0.84 – 1.23) | - | - |
| | Giving up preferred foods | - | - | - | 2.46 (1.66 – 3.64) |
| | Busy lifestyle | - | - | 0.44 (0.36 – 0.53) | - |
| | Lack of willpower | 0.53 (0.44 – 0.63) | - | - | - |
| | Lack of healthy options | - | - | - | 1.94 (1.26 – 2.98) |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by age, sex, BMI, household composition, employment status and urban regions. Results presented in bold were statistically significant ($p < 0.05$).

Supplementary Table 4. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by weight status as a significant effect modification was found for the depicted associations.

| Barriers ^{ab} | | Fruit OR (95% CI) | Vegetables OR (95% CI) | Sweets OR (95% CI) |
|----------------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| Under/normal weight (n=2819) | Giving up preferred foods | - | - | 1.49 (1.23 – 1.81) |
| | Price of healthy foods | 0.60 (0.50 – 0.72) | - | - |
| | Lack of healthy options | 0.69 (0.55 – 0.85) | - | - |
| | Unappealing foods | 0.48 (0.39 – 0.60) | 0.43 (0.34 – 0.54) | - |
| Overweight/ obese (n=2376) | Giving up preferred foods | - | - | 1.13 (0.94 – 1.36) |
| | Price of healthy foods | 0.72 (0.59 – 0.87) | - | - |
| | Lack of healthy options | 0.98 (0.78 – 1.24) | - | - |
| | Unappealing foods | 0.73 (0.59 – 0.91) | 0.65 (0.52 – 0.80) | - |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by age, sex, educational attainment, household composition, employment status and urban regions. Results presented in bold were statistically significant (p<0.05).

Supplementary Table 5. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by types of household composition as a significant effect modification was found for the depicted associations (the table continues on next page).

| | Barriers^{ab} | Fruit OR (95% CI) | Vegetables OR (95% CI) | Fish OR (95% CI) | Breakfast OR (95% CI) |
|----------------------------|---|-----------------------------|----------------------------------|----------------------------|---------------------------------|
| 1 person (n=1,328) | Irregular working hours | 0.58 (0.42 – 0.79) | - | - | - |
| | Giving up preferred foods | - | - | 0.80 (0.61 – 1.06) | - |
| | Busy lifestyle | 0.55 (0.41 – 0.73) | - | - | - |
| | Lack of willpower | - | - | 0.75 (0.58 – 0.96) | - |
| | Price of healthy foods | - | 0.54 (0.41 – 0.72) | 0.74 (0.55 – 0.99) | 0.80 (0.57 – 1.11) |
| | Taste preferences of family and friends | - | 0.91 (0.67 – 1.25) | 1.15 (0.83 – 1.59) | - |
| | Lack of healthy options | - | - | 0.92 (0.67 – 1.27) | - |
| 2 people (n=2,292) | Irregular working hours | 0.75 (0.60 – 0.94) | - | - | - |
| | Giving up preferred foods | - | - | 0.59 (0.47 – 0.74) | - |
| | Busy lifestyle | 0.59 (0.48 – 0.73) | - | - | - |
| | Lack of willpower | - | - | 0.53 (0.43 – 0.66) | - |
| | Price of healthy foods | - | 0.49 (0.39 – 0.62) | 0.61 (0.48 – 0.78) | 0.60 (0.45 – 0.80) |
| | Taste preferences of family and friends | - | 0.66 (0.53 – 0.82) | 0.80 (0.64 – 1.00) | - |
| | Lack of healthy options | - | - | 0.66 (0.49 – 0.88) | - |
| 3 or more people (n=2,224) | Irregular working hours | 0.71 (0.58 – 0.87) | - | - | - |
| | Giving up preferred foods | - | - | 0.93 (0.74 – 1.17) | - |
| | Busy lifestyle | 0.75 (0.62 – 0.90) | - | - | - |
| | Lack of willpower | - | - | 0.81 (0.65 – 1.00) | - |
| | Price of healthy foods | - | 0.53 (0.43 – 0.65) | 0.66 (0.52 – 0.83) | 0.66 (0.52 – 0.85) |
| | Taste preferences of family and friends | - | 0.51 (0.42 – 0.63) | 0.68 (0.54 – 0.86) | - |
| | Lack of healthy options | - | - | 0.93 (0.71 – 1.21) | - |

Supplementary Table 5. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by types of household composition as a significant effect modification was found for the depicted associations (table conclusion).

| Barriers ^{ab} | | Home-cooked meals | Fast food | Sweets |
|----------------------------------|---|---------------------------|---------------------------|--------------------|
| | | OR (95% CI) | OR (95% CI) | OR (95% CI) |
| 1 person (n=1,328) | Price of healthy foods | 0.72 (0.53 – 0.99) | - | - |
| | Taste preferences of family and friends | 0.87 (0.64 – 1.19) | - | - |
| | Unappealing foods | - | 1.87 (1.09 – 3.21) | 1.05 (0.77 – 1.43) |
| 2 people (n=2,292) | Price of healthy foods | 0.54 (0.42 – 0.70) | - | - |
| | Taste preferences of family and friends | 0.76 (0.60 – 0.96) | - | - |
| | Unappealing foods | - | 2.78 (1.73 – 4.48) | 1.15 (0.91 – 1.45) |
| 3 or more people (n=2,224) | Price of healthy foods | 0.64 (0.51 – 0.82) | - | - |
| | Taste preferences of family and friends | 0.82 (0.66 – 1.02) | - | - |
| | Unappealing foods | - | 1.87 (1.21 – 2.90) | 0.87 (0.69 – 1.09) |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by age, sex, educational attainment, BMI, employment status and urban regions. Results presented in bold were statistically significant ($p < 0.05$).

Supplementary Table 6. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by employment status as a significant effect modification was found for the depicted associations.

| | | Fruit | Breakfast | Home-cooked meals | Fast food |
|---------------------------------|---|---------------------------|---------------------------|---------------------------|---------------------------|
| Barriers ^{ab} | | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) |
| Not Working nor in education | Giving up preferred foods | - | 0.55 (0.42 – 0.73) | - | - |
| | Busy lifestyle | - | - | 0.70 (0.55 -0.88) | 1.32 (0.78 – 2.23) |
| | Lack of willpower | - | 0.58 (0.44 – 0.76) | - | - |
| | Price of healthy foods | - | 0.59 (0.43 – 0.81) | - | - |
| | Taste preferences of family and friends | - | - | 0.88 (0.69 – 1.13) | - |
| | Lack of healthy options | 0.68 (0.52 – 0.89) | - | - | - |
| Working or in education | Giving up preferred foods | - | 0.74 (0.62 – 0.89) | - | - |
| | Busy lifestyle | - | - | 0.42 (0.36 – 0.50) | 2.68 (1.86 – 3.87) |
| | Lack of willpower | - | 0.65 (0.55 – 0.79) | - | - |
| | Price of healthy foods | - | 0.74 (0.61 – 0.89) | - | - |
| | Taste preferences of family and friends | - | - | 0.77 (0.65 – 0.93) | - |
| | Lack of healthy options | 0.90 (0.75 – 1.09) | - | - | - |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by age, sex, educational attainment, BMI, household composition and urban regions. Results presented in bold were statistically significant ($p < 0.05$)

Supplementary Table 7. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5,900). Results were split by urban regions as a significant effect modification was found for the depicted associations (the table continues on next page).

| Barriers^{ab} | | Fruit OR (95% CI) | Vegetables OR (95% CI) | Fish OR (95% CI) | Breakfast OR (95% CI) |
|------------------------------|---|------------------------------------|---|-----------------------------------|--|
| Belgium (n=17,74) | Giving up preferred foods | - | 0.42 (0.33 – 0.55) | - | - |
| | Busy lifestyle | - | - | 0.76 (0.61 – 0.95) | 0.54 (0.41 – 0.73) |
| | Lack of willpower | - | 0.44 (0.35 – 0.56) | - | 0.62 (0.47 – 0.82) |
| | Price of healthy foods | - | 0.40 (0.31 – 0.52) | 0.89 (0.67 – 1.18) | - |
| | Taste preferences of family and friends | 0.84 (0.66 – 1.06) | - | 0.92 (0.72 – 1.18) | - |
| | Lack of healthy options | 0.76 (0.57 – 1.00) | - | - | 0.58 (0.41 – 0.81) |
| | Unappealing foods | - | - | 0.79 (0.59 – 1.05) | - |
| France (n=8,20) | Giving up preferred foods | - | 0.67 (0.48 – 0.93) | - | - |
| | Busy lifestyle | - | - | 0.74 (0.53 – 1.05) | 1.12 (0.73 – 1.70) |
| | Lack of willpower | - | 0.37 (0.26 – 0.52) | - | 0.86 (0.57 – 1.30) |
| | Price of healthy foods | - | 0.60 (0.44 – 0.81) | 0.71 (0.51 – 0.98) | - |
| | Taste preferences of family and friends | 0.59 (0.43 – 0.83) | - | 0.71 (0.52 – 0.98) | - |
| | Lack of healthy options | 0.80 (0.57 – 1.23) | - | - | 0.89 (0.58 – 1.37) |
| | Unappealing foods | - | - | 0.72 (0.51 – 1.00) | - |
| Hungary (n=875) | Giving up preferred foods | - | 0.75 (0.55 – 1.02) | - | - |
| | Busy lifestyle | - | - | 0.89 (0.53 – 1.50) | 0.75 (0.54 – 1.04) |
| | Lack of willpower | - | 0.59 (0.43 – 0.82) | - | 0.68 (0.49 – 0.95) |
| | Price of healthy foods | - | 0.53 (0.38 – 0.73) | 0.34 (0.21 – 0.56) | - |
| | Taste preferences of family and friends | 0.61 (0.44 – 0.84) | - | 0.56 (0.33 – 0.97) | - |
| | Lack of healthy options | 1.17 (0.84 – 1.63) | - | - | 0.99 (0.70 – 1.42) |
| | Unappealing foods | - | - | 0.49 (0.25 – 0.98) | - |

Supplementary Table 7. Odds ratio's (OR) and 95% confidence intervals (95%CI) as derived from multilevel multivariable logistics analyses indicating associations between perceived barriers to healthy eating and dietary behaviours among adults in five urban regions in Europe. The SPOTLIGHT Project (n=5900). Results were split by urban regions as a significant effect modification was found for the depicted associations (table conclusion).

| Barriers ^{ab} | | Home-cooked meals | Fast food | Sweets | Sugar-sweetened beverages |
|---------------------------|---|---------------------------|---------------------------|---------------------------|---------------------------|
| | | OR (95% CI) | OR (95% CI) | OR (95% CI) | OR (95% CI) |
| Netherlands (n=1,609) | Irregular working hours | 0.33 (0.24 – 0.46) | - | - | 1.47 (1.09 – 1.98) |
| | Giving up preferred foods | 0.49 (0.36 – 0.66) | - | - | - |
| | Busy lifestyle | 0.35 (0.26 – 0.48) | - | 1.24 (0.95 – 1.61) | - |
| | Lack of willpower | 0.36 (0.28 – 0.47) | - | 1.75 (1.39 – 2.20) | - |
| | Price of healthy foods | 0.61 (0.43 – 0.86) | 2.04 (1.04 – 4.02) | 1.03 (0.75 – 1.40) | - |
| | Taste preferences of family and friends | 0.67 (0.49 – 0.92) | - | - | - |
| | Lack of healthy options | 0.67 (0.37 – 1.21) | - | - | 1.20 (0.74 – 1.94) |
| United Kingdom (n=822) | Irregular working hours | 0.63 (0.43 – 0.92) | - | - | 1.18 (0.83 – 1.68) |
| | Giving up preferred foods | 0.59 (0.42 – 0.83) | - | - | - |
| | Busy lifestyle | 0.39 (0.27 – 0.58) | - | 1.18 (0.84 – 1.66) | - |
| | Lack of willpower | 0.57 (0.40 – 0.79) | - | 1.77 (1.28 – 2.45) | - |
| | Price of healthy foods | 0.75 (0.54 – 1.06) | 1.55 (0.96 – 2.52) | 0.92 (0.64 – 1.33) | - |
| | Taste preferences of family and friends | 0.75 (0.54 – 1.06) | - | - | - |
| | Lack of healthy options | 0.62 (0.44 – 0.89) | - | - | 0.88 (0.62 – 1.25) |

^a Reference category in each barrier: Not perceived as a barrier (merged responses options: never and rarely); ^b Empty rows or columns (no significant effect modification across barrier or food items) were omitted. All analysis were performed in separated models and adjusted by age, sex, educational attainment, BMI, household composition and employment status. Results presented in bold were statistically significant (p<0.05).

